



| | |
|--------------------|--|
| Title | Molecular analysis of the microbial community structures in water-flooding petroleum reservoirs with different temperatures |
| Author(s) | Wang, LY; Duan, RY; Liu, JF; Yang, SZ; Gu, J; Mu, BZ |
| Citation | Biogeosciences Discussions, 2012, v. 9 n. 11, p. 5177-5203 |
| Issued Date | 2012 |
| URL | http://hdl.handle.net/10722/196825 |
| Rights | Creative Commons: Attribution 3.0 Hong Kong License |

This discussion paper is/has been under review for the journal Biogeosciences (BG).
Please refer to the corresponding final paper in BG if available.

Molecular analysis of the microbial community structures in water-flooding petroleum reservoirs with different temperatures

L.-Y. Wang¹, R.-Y. Duan¹, J.-F. Liu¹, S.-Z. Yang¹, J.-D. Gu², and B.-Z. Mu¹

¹State Key Laboratory of Bioreactor Engineering and Institute of Applied Chemistry, East China University of Science and Technology, Shanghai 200237, China

²School of Biological Sciences, The University of Hong Kong, Pokfulam Road, Hong Kong SAR, China

Received: 6 April 2012 – Accepted: 16 April 2012 – Published: 27 April 2012

Correspondence to: B.-Z. Mu (bzmu@ecust.edu.cn)

Published by Copernicus Publications on behalf of the European Geosciences Union.

BGD

9, 5177–5203, 2012

The microbial communities of petroleum reservoirs fluids

L.-Y. Wang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

Temperature is one of the most important environmental factors regulating the activity and determining the composition of the microbial community. Analysis of microbial communities from six water-flooding petroleum reservoirs at temperatures from 20 to 63 °C by 16S rRNA gene clone libraries indicates the presence of physiologically diverse and temperature-dependent microorganisms in these subterrestrial ecosystems. In high-temperature petroleum reservoirs, most of the archaeal sequences belong to the thermophilic archaea including the genera *Thermococcus*, *Methanothermobacter* and *Thermoplasma*, most of the bacterial sequences belong to the phyla *Firmicutes*, *Thermotogae* and *Thermodesulfobacteria*; in low-temperature petroleum reservoirs, most of the archaeal sequences are affiliated with the genera *Methanobacterium*, *Methanoculleus* and *Methanocalculus*, most of the bacterial sequences to the phyla *Proteobacteria*, *Bacteroidetes* and *Actinobacteria*. Canonical correspondence analysis (CCA) revealed that temperature, mineralization, ionic type as well as volatile fatty acids showed correlation with the microbial community structures. These organisms may be adapted to the environmental conditions of these petroleum reservoirs over geologic time by metabolizing buried organic matter from the original deep subsurface environment and became the common inhabitants in subsurface environments.

1 Introduction

Petroleum reservoirs represent extreme anaerobic environments because of the temperature, pressure and salinity with multiphase fluids of oil, gas and water. Microorganisms in such subterranean play an important role in energy flow and nutrients cycling. The microbial activity, in particular sulfate-reducing bacteria (SRBs) not only causes issues like reservoir souring and corrosion of drilling equipment with H₂S, but potentially can also be used to our advantage, e.g., microbial-enhanced oil recovery (MEOR) applications. Extraction of currently usable energy from marginal petroleum reservoirs by

BGD

9, 5177–5203, 2012

The microbial communities of petroleum reservoirs fluids

L.-Y. Wang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

The microbial communities of petroleum reservoirs fluids

L.-Y. Wang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

microbial conversion of residual oil to methane (natural gas) has received renewed attention in the past decade (Parkes, 1999; Sufliya et al., 2004; Gieg et al., 2008; Jones et al., 2008; Wang et al., 2010, 2011; Mbadanga et al., 2011, 2012). Microorganisms with diverse physiological and metabolic abilities and phylogenetic affiliations have been recovered from oil reservoirs by culture-dependent and culture-independent approaches since the first sulfate-reducing bacteria (SRB) was isolated from production water (Bastin et al., 1926). Though isolation efforts have identified numerous bacterial and archaeal species that are capable of mediating various metabolic processes occurring in oil fields, culture-independent 16S rRNA gene- and functional genes-based investigations have provided new information on the microbial community composition in such deep-subsurface ecosystem (Li et al., 2010, 2011).

Culture-independent surveys of high-temperature oil reservoirs have been conducted in oil fields in continental and offshore California (Orphan et al., 2000, 2003), an offshore oil field in Qinghuang, China (Li et al., 2007b), Huabei oil field in continental China (Li et al., 2006, 2007a), Dagang oil field in China (Nazina et al., 2006), and in Troll oil formation in the North Sea (Dahle et al., 2008). Bacterial sequences affiliated with *Firmiucutes* are the most frequency detected in these high-temperature oil reservoirs. In addition, the discovery of moderately thermophilic members of the *Bacteroidetes* (genus *Anaerophaga*) in Troll oil formation in the North Sea and sulfate reducing *Nitrospira* (genus *Thermodesulfovibrio*) in the Huabei and Qinghuang oil fields in China have not been previously isolated from oil reservoirs. Most of Archaeal belong to methanogens including methyltrophic, acetoclastic as well as CO₂-reducing methanogens and CO₂-reducing methanogens are the most common members in high-temperature oil reservoirs. Moreover, methanogenesis from acetate driven by syntrophic acetate oxidation has been documented in high-temperature reservoir (Nazina et al., 2006).

Compared with high-temperature oil reservoirs, few 16S-based analysis of the microbial community in low-temperature oil reservoirs have been reported. The bacterial diversity in a low-temperature, low-salinity, non-water flooded oil reservoir (Pelikan

lake oil field) in Western Canada was extremely low with only one phylotype related to the genus *Arcobacter* (*ε-Proteobacteria*) (Grabowski et al., 2005). Several potentially metabolic active fermentative and/or acetogenic microorganisms, sulfide-oxidizers and sulfate-reducers were identified from a low-temperature oil reservoir in Western Canada by 16S rRNA gene clones library analysis (Voordouw et al., 1996).

The ecological processes of these microorganisms play considerable important roles in energy flow and nutrients cycling in subsurface ecosystem. The distribution of different microbial community structures in petroleum reservoirs depends entirely on their adaption to the in situ physical and chemical variables, including temperature, pH, and salinity. Despite of the available studies on microbial population both in high- or low-petroleum reservoirs, microbial populations in reservoirs from low to high temperatures has not been studied in a single investigation. In this report, the distribution of microbial community in petroleum reservoirs from production water of several petroleum reservoirs at temperature of 20, 32, 37, 45, 58 and 63 °C was investigated by 16S rRNA gene library analysis. Microbial community data were also correlated with environmental factors using canonical correspondence analysis (CCA).

2 Materials and methods

2.1 Collection of sample and nucleic acid extraction

The production water samples of six production oil wells (S1–S6) from six Block affiliated with 3 oil fields of China: Well Z3-26 belongs to Zhan 3 of Shengli oil field; Well B18-43 to Ba 19, Well B51-45 to Ba 51 and Well M17-10 to Menggulin of Huabei oil field; Well 7222 to No. 7, Well 6190 to No. 6 of Kelamayi oil field. The temperature and the mineralization of these six petroleum reservoirs sampled range from 20 to 63 °C and 1301 to 11196 mg l⁻¹, respectively and the pH of these production waters were neutral or slightly alkaline. The characterization of the petroleum reservoirs water sampled are listed in Table 1. In order to characterize the microbial community from

BGD

9, 5177–5203, 2012

The microbial communities of petroleum reservoirs fluids

L.-Y. Wang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

the different temperature petroleum reservoir, these samples were grouped into two classes: high-temperature (45 ~ 63 °C) and low-temperature (20 ~ 37 °C).

Ten liters of production water at wellhead were taken from each of the six production oil wells from six petroleum reservoirs. Samples were collected into sterile bottles to full capacity after discarding the initial oil/water mixture. The bottles were tightly sealed to avoid oxygen intrusion and immediately transported to the laboratory and filtered directly to minimize the chance of community changes. During filtration procedures, the residual oil was removed by heating the sample to 50 °C for 15 min and by phase separation in 2 l sterilized separatory funnel. The water samples were filtered through 0.22 µm polycarbonate membranes (25 mm diameter; Millipore, Bedford, USA). The polycarbonate membranes containing the cells was placed in a sterile centrifuge tube containing sterile silica beads for beating to break the cells. Genomic DNA was extracted by a method developed previously in this laboratory (Li et al., 2007b).

2.2 16S rRNA gene amplification and cloning

16S rRNA genes were amplified by PCR using the primers B27F [5'-AGAGTTTGATCCTGGCTCAG-3'] and B1492R [5'-TACGGYTACCTTGTTACGACTT-3'] (Nazina et al., 2006) for bacteria, and the primers A21F [5'-TTCCGGTTGATCCYGCCGGA-3'] (DeLong, 1992) and A1041R [5'-GGCCATGCACCWCCTCTC-3'] (Kolganova et al., 2002) for archaea. The final 50 µl reaction mixture volume contained 2 µl of template DNA, 0.5 µM of each primer, 25 µl of 2 × master mix (Promega, USA), 21 µl of nuclease-free water. Polymerase chain reaction cycles were performed on a Peltier thermal cycler (Bio-Rad, USA) as follows: after 5 min of initial denaturation at 95 °C, nucleic acids were amplified for 30 cycles (45 s of denaturation at 95 °C, 45 s of annealing at 50 °C and 1 min of elongation at 72 °C) followed by a final extension step at 72 °C for 20 min. PCR products were separated on 0.8% (w/v) agarose gel and stained with ethidium bromide. The amplicons were cloned into a pMD[®]19-T Simple vector (Takara, Japan) according to the manufacturer's instructions.

BGD

9, 5177–5203, 2012

The microbial communities of petroleum reservoirs fluids

L.-Y. Wang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

2.3 Sequencing and phylogenetic analysis

Inserts of selected clones were amplified by PCR with forward M13F (5'-GTTTTCC CAGTCACGA-3') and the reverse M13R (5'-CAGGAAACAGCTATGAC-3') plasmid specific primer set. The sequencing was performed on an ABI 377 sequencer (Dye Terminator Cycle Sequencing Ready Reaction FS Kit; PE Applied Biosystems) using M13 universal sequencing primers. Obtained DNA sequences were checked for vector by VecScreen Widget 1.0 software before further analysis. Sequence data were aligned using the the NAST alignment algorithm (DeSantis et al., 2006a) on the GreenGenes website (<http://greengenes.lbl.gov>), with clones having similarities of 98% or above grouped into operational taxonomic units (OTUs). The clones were homology-searched using Ribosomal Database Project II (Wang et al., 2007). The nearest relatives of each OTU were identified using the BLASTN network service (Altschul et al., 1997). Chimeras were detected using Bellerophon, version 3 (Huber et al., 2004; DeSantis et al., 2006) and removed from further examination. Phylogenetic trees were constructed based on the neighbor-joining algorithm (Saitou and Nei, 1987) using the MEGA5 software (Tamura et al., 2011). Bootstrap analysis with 1000 replicates was applied to assign confidence levels to the nodes in the trees.

2.4 Statistical analysis

The coverage of each clone library was calculated by the equation $C = [1 - n_1/N] \times 100$, where n_1 is the number of OTUs represented by only one clone and N is the total number of clones examined (Good, 1953). To examine the temperature distribution of microbial community in production water of petroleum reservoir, 16S rRNA gene sequences were analyzed with the online software UniFrac (<http://bmf2.colorado.edu/unifrac/>) using the principal coordinates analysis (PCoA) as suggested previously (Lozupone and Knight, 2005). Correlations between the microbial communities and environmental factors were determined by the canonical correspondence analysis (CCA) using the

BGD

9, 5177–5203, 2012

The microbial communities of petroleum reservoirs fluids

L.-Y. Wang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

software CANOCO (ver. 4.5, Microcomputer Power, Ithaca, NY, USA) (ter Braak and Šmilauer, 2002).

2.5 Nucleotide sequence accession numbers

Partial 16S rRNA gene sequences for Bacteria and Archaea obtained in this study were deposited in GenBank databases under accession numbers JQ433723-JQ433816 and JF754550-JF754565.

3 Results

3.1 Diversity of microbial community in water-flooding petroleum reservoirs

Six production water samples from six water-flooding petroleum reservoirs with different temperatures were analyzed by PCR amplification with bacterial and archaeal specific primer sets. For the members in the domain bacteria, 93, 226, 254, 80, 185 and 142 clones were randomly selected from the libraries of production water samples S1, S2, S3, S4, S5 and S6, respectively. Of the total sequences screened by MOTHUR software version 1.6, 5, 16, 10, 13, 30 and 31 operational taxonomy units (OTUs) were obtained from S1, S2, S3, S4, S5 and S6, respectively and classified into thirteen different phylogenetic groups (phylum level) (Figs. 1–3). The coverage of the clone library was 100 % for S1, S2, S3 and S5, and 96 % for S4, as well as 99 % for S6 from rarefaction analysis.

For the members in the domain archaea, 125, 56, 79, 60, 24 and 33 clones were randomly selected from the libraries of production water samples S1, S2, S3, S4, S5 and S6, respectively. Of the total sequences screened by MOTHUR software version 1.6, 5, 1, 4, 2, 2 and 2 OTUs were obtained from S1, S2, S3, S4, S5 and S6, respectively and classified into eleven different phylogenetic groups (genus level) (Fig. 4). The coverage of the clone library was all 100 % except 98 % for S1 from rarefaction analysis.

BGD

9, 5177–5203, 2012

The microbial communities of petroleum reservoirs fluids

L.-Y. Wang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

The microbial communities of petroleum reservoirs fluids

L.-Y. Wang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Bacterial and archaeal DNA sequences based on the percentage representation of major phylum or genus in clone libraries from the six different temperature reservoirs are shown in Fig. 5. The bacterial sequences were clustered within thirteen phyla: *Proteobacteria* (α -, β -, γ -, δ -, ϵ -), *Firmicutes*, *Bacteroidetes*, *Actinobacteria*, *Chloroflexi*, *Thermotogae*, *Thermodesulfobacteria*, *Deinococcus-Thermus* and TM7. Compared with low temperature reservoir, bacterial sequences affiliated with the phylum *Firmicutes* account for the highest percentage in S2 (65.9%) and S3 (29.9%) from high temperature petroleum reservoir. In contrast, *Proteobacteria* (α -, β -, γ -, δ -, ϵ -) account for the highest percentage in the S4, S5 and S6 from low temperature petroleum reservoirs. α -*Proteobacteria* increased with the decrease of petroleum reservoir temperature, but the percentage of β -*Proteobacteria* decreased with the reduction of petroleum reservoir temperature. γ -*Proteobacteria* shared similar high percentage (30~40%) in the S4, S5 and S6 as well as in S2 (31.5%). However, it is interesting that all the bacterial sequences affiliated with γ -*Proteobacteria* are found in S1. ϵ -*Proteobacteria* was encountered in S2 and S5 accounting for 2.7% and 5.1%, respectively. γ -*Proteobacteria* was detected in S3 and S6 with 7.1% and 0.7%, respectively. *Bacteroidetes* was another frequently encountered phylum in low temperature petroleum reservoir of S5 and S6 for 9.0% and 17.6%, respectively. *Thermotogae* with higher percentage accounted for 19.5% in a high temperature petroleum reservoir. The remaining bacterial phyla account for relative low abundance in the six petroleum reservoirs.

The archaeal sequences were clustered within eleven phylogenetic groups (genus level): *Methanocella*, *Methanosaeta*, *Methanomethylivorans*, *Methanolinea*, *Methanocalculus*, *Methanoculleus*, *Methanothermobacter*, *Methanobacterium*, *Thermococcus*, *Halogeometricum* and *Thermogymnomonas*. In the high-temperature oil reservoirs, most of the genera belong to the thermophilic archaea. The sequences affiliated with the genus *Thermococcus* and *Methanothermobacter* were account for high percentage, 88.8% of total archaeal clones with the genus *Thermococcus* in S1 and all the archaeal clones affiliated with the genus *Methanothermobacter* in S2. The

sequences affiliated with the genus *Methanomethylovorans* was only found in S3 accounting for 54.4%. The sequences affiliated with the genus *Methanolinea* were detected in S4 with a high percentage (70%). The sequences affiliated with the genus *Methanobacterium* were the most abundance in low temperature petroleum reservoir, accounting for 79.2% in S5 and 66.7% in S6.

3.2 Microbial community classification of water-flooding petroleum reservoirs

PCoA of bacterial and archaeal community structures were carried out by Unifrac based on the phylogenetic tree of 16S rRNA gene sequences in the six investigated petroleum reservoirs indicates that bacteria and archaea display high niche specificity (Fig. 6). S5 and S6 were collected from low temperature petroleum reservoirs grouped together, sharing similar bacterial and archaeal community structures. Although S1 and S2 represented high temperature petroleum reservoirs, both bacteria and archaea community structure did not grouped together because of a great difference in mineralization and the concentration of Cl^- . In addition, S2 and S3 grouped together, sharing similar bacterial community structures in PCoA of bacterial classification while S3 and S4 grouped together, sharing similar archaeal community structures in PCoA of archaeal classification.

3.3 Correlations of microbial communities with environmental factors

To find out the relationships between the distribution of microbial community and the environmental variables of the petroleum reservoirs, canonical correspondence analysis was conducted based on the bacterial and archaeal 16S rRNA gene sequences and the major physiochemical parameters of petroleum reservoirs (Table 1). The first two axes of the CCA analysis explained 69.9% and 55.3% of the total variance for the bacterial and archaeal communities, respectively (Fig. 7). The physiochemical parameters of petroleum reservoirs were divided to three groups to better analyze the relationships. In the first group, the differences in the bacterial and archaeal community structures

BGD

9, 5177–5203, 2012

The microbial communities of petroleum reservoirs fluids

L.-Y. Wang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The microbial communities of petroleum reservoirs fluids

L.-Y. Wang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

(Fig. 7A, a) were related to temperature, mineralization, average permeability, oil viscosity, effective porosity, effective thickness and water flooding operation years. The bacterial phylogenetic group of *Firmicutes* and *Thermotogae* had positive correlation with the temperature, γ -*Proteobacteria* and *Chloroflexi* with the effective porosity and oil viscosity, others with the effective thickness and water flooding operation years; most of the archaeal phylogenetic group (*Methanocella*, *Methanosaeta*, *Methanomethylovorans*, *Methanolinea*, *Methanothermobacter*, *Thermococcus*, *Halo geometricum* and *Thermogymnomonas*) correlated positively with the temperature, others with the water flooding operation years, mineralization and effective thickness. In the second group, the differences in the bacterial and archaeal community structure (Fig. 7B, b) were related to differences in the concentration of Cl^- , SO_4^{2-} , PO_4^{3-} , NO_3^- , Na^+ , K^+ , Ca^{2+} , Mg^{2+} , and Mn^{2+} . The bacterial phylogenetic group of *Bacteroidetes*, α -*Proteobacteria*, *Actinobacteria*, and *Deinococcus-Thermus* showed positive correlation with the concentration of NO_3^- , β -*Proteobacteria* with the concentration of Cl^- , γ -*Proteobacteria* and *Chloroflexi* with the concentration of SO_4^{2-} and PO_4^{3-} ; the archaeal phylogenetic group of *Methanosaeta*, *Thermococcus*, *Halo geometricum* and *Thermogymnomonas* correlated positively with the concentration of SO_4^{2-} , *Methanocalculus*, *Methanobacterium* and *Methanoculleus* with the concentration of NO_3^- . In the third group, the differences in the bacterial and archaeal community structure (Fig. 7C, c) were related to differences in the concentration of volatile fatty acid including acetate, propionate, isobutyrate and butyrate. The bacterial phylogenetic group of *Thermotogae* showed positive correlation with the concentration of propionate, *Firmicutes* with the concentration of acetate, except for γ -*Proteobacteria* and *Chloroflexi* others were with the concentration of isobutyrate; the archaeal phylogenetic group of *Methanomethylovorans*, *Methanothermobacter*, *Methanolinea* and *Methanocella* had positive correlation with the concentration of butyrate, *Methanoculleus* with the concentration of isobutyrate.

4 Discussion

It has been widely accepted that the combination of temperature, salinity and pressure in subsurface petroleum reservoirs drastically reduces microbial populations and metabolic activity. The microbial population differs from one oil reservoir to another because of the variations in their physical, chemical, and geochemical entities. The physiologically diverse microbial assemblage of microorganisms are distributed in such ecosystems. From the about 1000 bacterial and 400 archaeal 16S rRNA gene sequences retrieved from production water samples of six different oil reservoirs with temperature from 20 to 63°C and salinity from 1.12 to 15.73 gL⁻¹, the bacterial sequences affiliated with the phylum *Firmicutes* had the greatest proportion in high temperature reservoirs. Dominant groups within the *Firmicutes* were the family *Pep-
toco*ccaceae, *Thermoanaerobacteraceae*, *Syntrophomonadaceae*, *Lachnospiraceae*, *Erysipelotrichaceae* and Incertae Sedis XII. Members of these groups are obligate anaerobic and thermophilic microorganisms. The sequence type (S2 B clone: 94) retrieved from oil reservoirs with temperature of 58°C in this study is most closely related to *Pelotomaculum thermopropionicum*, a member of *Clostridiales*, and a thermophilic propionate-oxidizing anaerobic bacterium, isolated from anaerobic sludge blanket reactor in Niigata, Japan (Imachi et al., 2002). Such microorganism was once postulated to be associated with hydrocarbon-degradation (Abu Laban et al., 2009) (Gieg et al., 2008). The sequence type (S2 B clone: 127) is most closely related to *Thermacetogenium phaeum*, is a strictly anaerobic, thermophilic, syntrophic acetate-oxidizing bacterium, which can oxidize acetate in co-culture with a thermophilic hydrogenotrophic methanogen (Hattori et al., 2000). The sequence types (S2 B clone: 210 and S3 B clone: 14) are most closely related to *Syntrophothermus lipocalidus*, a novel thermophilic, syntrophic isobutyrate-oxidizing bacterium (Sekiguchi et al., 2000). According to the physicochemical data from production water samples and CCA analysis, the bacterial group *Firmicutes* showed higher values for acetate, propionate, isobutyrate and butyrate (Fig. 7C), suggesting that VFA components within the production water serve

The microbial communities of petroleum reservoirs fluids

L.-Y. Wang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The microbial communities of petroleum reservoirs fluids

L.-Y. Wang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

as their major carbon sources, and demonstrated a higher support between the phylogenetic data and the physicochemical measurements. Other sequence type within the phylum *Firmicutes* also had a greater proportion in high temperature reservoirs, such as sequence type (S3B clone: 98), most closely related to *Fusibacter* spp., an anaerobic, thiosulfate-reducing bacterium isolated from an oil-producing well (Ravot et al., 1999). It is not a typical thermophilic microorganism with optimal growth temperature at 37 °C and such type sequences were also found in low temperature petroleum reservoir represented as S6 B clone 70 with petroleum temperature at 20 °C. CCA analysis showed that the phylum *Thermotogae* is another bacterial group correlated positively with the temperature. The sequence types (S2 B clone: 210 and S3 B clone: 14) are most closely related to *Thermotoga hypogea*, a xylanolytic, thermophilic, strictly anaerobic bacterium isolated from an oil-producing well (Fardeau et al., 1997). Many thermophilic bacteria with optimum growth temperatures from 45 to 80 °C and have been isolated from oil fields (Beeder et al., 1995; Cayll et al., 1995; Jeanthon et al., 1995; Ravot et al., 1995; Rees et al., 1995; Fardeau et al., 1996, 1997; Nilsen et al., 1996). In this study, although S1, S2, S3 represent thermophilic temperature, not all of the bacterial sequences belong to thermophilic microorganisms, especially the bacterial sequences affiliated with *Pseudomonas* spp. within the phylum γ -*Proteobacteria* in S1 with temperature at 63 °C. It may be that the sample was contaminated during handling judging by the single type of bacterial diversity and non-temperature dependent of physiological characteristics. However, it is a general trend that microbial populations are greatly reduced in petroleum reservoirs with the combination of high-temperature, high-mineralization and high concentration of SO_4^{2-} . High-temperature oil reservoirs contained bacterial sequences affiliated with the α -, β -, γ -, δ -, and ϵ -*Proteobacteria*, but low-temperature oil reservoirs by the γ -*Proteobacteria*. This is best illustrated by the no correlation between physicochemistry and phylogeny, where γ -*Proteobacterial* sequences related to strict aerobes and facultative anaerobes.

In addition, CCA analysis indicated that temperature exhibited the greatest influence on the archaeal community. The archaea identified from the petroleum reservoirs

The microbial communities of petroleum reservoirs fluids

L.-Y. Wang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

are overwhelmingly methanogens including methyltrophic (*Methanomethylovorans*), acetoclastic (*Methanosaeta*) and CO₂-reducing methanogens (*Methanothermobacter*, *Methanoculleus*, *Methanobacteria*, *Methanocalculus*, *Methanocella* and *Methanolinea*), possibly being mesophilic or thermophilic. In contrast, in sample S1 with the highest temperature (63°C), besides *Methanosaeta* (0.8% of total clones), majority of the clones were phylogenetically related to the genus *Thermococcus* (88.8% of total clones) and to the species *T. sibiricus* (99% sequence similarity) in these reservoirs (Fig. 4). Clones pertaining to *Thermogymnomonas* (9.6% of total clones) and *Halogeotricum* (0.8% of total clones) were also detected, but at a less extent in sample S1. In this respect, the ecological significance of methanogenic archaea in petroleum reservoirs is to serve as terminal electron acceptor, e.g., hydrogenotrophic ones in the complete oxidation of hydrocarbons. Most of the sequences assigned to CO₂-reducing methanogens in present research is in line with the view that CO₂-reducing methanogens being the most commonly encountered in both cultivation and culture-independent studies of oilfield archaea (Head et al., 2010). Moreover, methanogenesis from acetate driven via syntrophic acetate oxidation has been documented in high-temperature oil reservoirs and is responsible by *Methanothermobacter-Thermoanaerobacter* co-culture (Shestakova et al., 2011) as well as in methanogenic alkanes degradation enrichment derived from production water of high-temperature oil reservoir (Mbadinga et al., 2012). Furthermore, the high apparent abundance of thermophilic, syntrophic acetate, propionate, isobutyrate and butyrate-oxidizing bacterium *Firmicutes* (*Thermoanaerobacter*) as well as thermophilic CO₂-reducing methanogens *Methanothermobacter*, coupled with the transiently high levels of corresponding substrate metabolites detected in high-temperature oil reservoir (Table 1), further demonstrating that syntrophic acetate oxidation is the main methanogenic pathway in a high-temperature petroleum reservoirs in situ. In addition to the well-known CO₂-reduction pathways, the type of methyl/methanol-utilizing methanogens as the second most common group in the present research contribute significantly in subsurface biogas formation (Strapoc et al., 2011). Although obligate acetate utilizers are represented by small

populations of *Methanosaeta*, the potential contribution of acetate pathways cannot be ignored. In addition, metabolic types of the active methanogens vary substantially between reservoirs and appear to be controlled by local geochemical conditions within the reservoirs.

5 Conclusions

Taken together, the presence of physiologically diverse and temperature-dependent microbial community inhabiting in petroleum reservoirs. The high abundance thermophilic archaea including the genus *Thermococcus*, *Methanothermobacter* and *Thermoplasma*, and the bacterial sequences belong to the phylum *Firmicutes*, *Thermotogae* and *Thermodesulfobacteria* in high-temperature petroleum reservoirs relative to low-temperature petroleum reservoirs. The high abundance of the archaeal sequences belong to the genus *Methanobacterium*, *Methanoculleus* and *Methanocalculus*, and bacterial sequences belong to the phylum *Proteobacteria*, *Bacteroidetes* and *Actinobacteria* in low-temperature petroleum reservoirs relative to high-temperature petroleum reservoirs. The results of canonical correspondence analysis (CCA) and principal coordinates analysis (PCoA) showed a consistency between the phylogenetic data and the physicochemical measurements for the sampled environment and further demonstrated that syntrophic acetate oxidation is the main methanogenic pathway in a high-temperature petroleum reservoirs.

Acknowledgements. We thank Hui Li for discussion on correlation analyses. This work was supported by the National Natural Science Foundation of China (Grant No. 41073055) and the NSFC/RGC Joint Research Fund (No. 41161160560).

References

Abu Laban, N., Selesi, D., Jobelius, C., and Meckenstock, R. U.: Anaerobic benzene degradation by Gram-positive sulfate-reducing bacteria, *FEMS Microbiol. Ecol.*, 68, 300–311, 2009.

BGD

9, 5177–5203, 2012

The microbial communities of petroleum reservoirs fluids

L.-Y. Wang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The microbial communities of petroleum reservoirs fluids

L.-Y. Wang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Altschul, S., Madden, T., Schaffer, A., Zhang, J., Zhang, Z., Miller, W., and Lipman, D.: Gapped BLAST and PSI-BLAST: a new generation of protein database search programs, *Nucl. Acids. Res.*, 25, 3389–3402, 1997.

Bastin, E. S., Greer, F. E., Merritt, C. A., and Moulton, G.: The presence of sulphate reducing bacteria in oil field waters, *Science*, 63, 21–24, 1926.

Beeder, J., Torsvik, T., and Lien, T.: *Thermodesulforhabdus norvegicus* gen. nov., sp. nov., a novel thermophilic sulfate-reducing bacterium from oil field water, *Arch. Microbiol.*, 164, 331–336, 1995.

ter Braak, C. J. F. and Šmilauer, P.: CANOCO Reference manual and CanoDraw for Windows User's guide: Software for Canonical Community Ordination (version 4.5), Microcomputer Power, Ithaca NY, USA, 2002.

Cayol, J.-L., Ollivier, B., Patel, B. K. C., Ravot, G., Magot, M., Ageron, E., Grimont, P. A. D., and Garcia, J.-L.: Description of *Thermoanaerobacter brockii* subsp. *lactiethylicus* subsp. nov., isolated from a deep subsurface French oil well, a proposal to reclassify *Thermoanaerobacter finnii* as *Thermoanaerobacter brockii* subsp. *finnii* comb. nov., and an emended description of *Thermoanaerobacter brockii*, *Int. J. Syst. Bacteriol.*, 45, 783–789, 1995.

Dahle, H., Garshol, F., Madsen, M., and Birkeland, N.-K.: Microbial community structure analysis of produced water from a high-temperature North Sea oil-field, *Anton. Leeuw. Int. J. G.*, 93, 37–49, 2008.

DeLong, E. F.: Archaea in coastal marine environments, *P. Natl. Acad. Sci. USA*, 89, 5685–5689, 1992.

DeSantis, T. Z., Hugenholtz, P., Keller, K., Brodie, E. L., Larsen, N., Piceno, Y. M., Phan, R., and Andersen, G. L.: NAST: a multiple sequence alignment server for comparative analysis of 16S rRNA genes, *Nucl. Acids. Res.*, 34, 394–399, 2006a.

DeSantis, T. Z., Hugenholtz, P., Larsen, N., Rojas, M., Brodie, E. L., Keller, K., Huber, T., Dalevi, D., Hu, P., and Andersen, G. L.: Greengenes, a chimera-checked 16S rRNA gene database and workbench compatible with ARB, *Appl. Environ. Microbiol.*, 72, 5069–5072, 2006.

Fardeau, M. L., Faudon, C., Cayol, J. L., Magot, M., Patel, B. K. C., and Ollivier, B.: Effect of thiosulphate as electron acceptor on glucose and xylose oxidation by *Thermoanaerobacter finnii* and a *Thermoanaerobacter* sp. isolated from oil field water, *Res. Microbiol.*, 147, 159–165, 1996.

The microbial communities of petroleum reservoirs fluids

L.-Y. Wang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Fardeau, M.-L., Ollivier, B., Patel, B. K. C., Magot, M., Thomas, P., Rimbault, A., Rocchiccioli, F., and Garcia, J.-L.: *Thermotoga hypogea* sp. nov., a xylanolytic, thermophilic bacterium from an oil-producing well, *Int. J. Syst. Bacteriol.*, 47, 1013–1019, 1997.
- Gieg, L. M., Duncan, K. E., and Suflita, J. M.: Bioenergy production via microbial conversion of residual oil to natural gas, *Appl. Environ. Microbiol.*, 74, 3022–3029, 2008.
- Good, I. J.: The population frequencies of species and the estimation of population parameters, *Biometrika*, 40, 237–264, 1953.
- Grabowski, A., Nercessian, O., Fayolle, F., Blanchet, D., and Jeanthon, C.: Microbial diversity in production waters of a low-temperature biodegraded oil reservoir, *FEMS Microbiol. Ecol.*, 54, 427–443, 2005.
- Hattori, S., Kamagata, Y., Hanada, S., and Shoun, H.: *Thermacetogenium phaeum* gen. nov., sp. nov., a strictly anaerobic, thermophilic, syntrophic acetate-oxidizing bacterium, *Int. J. Syst. Evol. Microbiol.*, 50, 1601–1609, 2000.
- Head, I. M., Larter, S. R., Gray, N. D., Sherry, A., Adams, J. J., Aitken, C. M., Jones, D. M., Rowan, A. K., Huang, H., and Röling, W. F. M.: Hydrocarbon degradation in petroleum reservoirs, in: *Handbook of Hydrocarbon and Lipid Microbiology*, edited by: Timmis, K. N., Springer, Berlin Heidelberg, 3097–3109, 2010.
- Huber, T., Faulkner, G., and Hugenholz, P.: Bellerophon: a program to detect chimeric sequences in multiple sequence alignments, *Bioinformatics*, 20, 2317–2319, 2004.
- Imachi, H., Sekiguchi, Y., Kamagata, Y., Hanada, S., Ohashi, A., and Harada, H.: *Pelotomaculum thermopropionicum* gen. nov., sp. nov., an anaerobic, thermophilic, syntrophic propionate-oxidizing bacterium, *Int. J. Syst. Evol. Microbiol.*, 52, 1729–1735, 2002.
- Jeanthon, C., Reysenbach, A.-L., L'Haridon, S., Gambacorta, A., Pace, N., Glénat, P., and Prieur, D.: *Thermotoga subterranea* sp. nov., a new thermophilic bacterium isolated from a continental oil reservoir, *Arch. Microbiol.*, 164, 91–97, 1995.
- Jones, D. M., Head, I. M., Gray, N. D., Adams, J. J., Rowan, A. K., Aitken, C. M., Bennett, B., Huang, H., Brown, A., Bowler, B. F. J., Oldenburg, T., Erdmann, M., and Larter, S. R.: Crude-oil biodegradation via methanogenesis in subsurface petroleum reservoirs, *Nature*, 451, 176–180, 2008.
- Kolganova, T. V., Kuznetsov, B. B., and Turova, T. P.: Selection and testing of oligonucleotide primers for amplification sequencing of archaeal 16S rRNA genes, *Mikrobiologiya*, 71, 283–285, 2002.

The microbial communities of petroleum reservoirs fluids

L.-Y. Wang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Li, H., Yang, S.-Z., Mu, B.-Z., Rong, Z.-F., and Zhang, J.: Molecular analysis of the bacterial community in a continental high-temperature and water-flooded petroleum reservoir, *FEMS Microbiol. Lett.*, 257, 92–98, 2006.
- Li, H., Yang, S.-Z., and Mu, B.-Z.: Phylogenetic diversity of the archaeal community in a continental high-temperature, water-flooded petroleum reservoir, *Curr. Microbiol.*, 55, 382–388, 2007a.
- Li, H., Yang, S.-Z., Mu, B.-Z., Rong, Z.-F., and Zhang, J.: Molecular phylogenetic diversity of the microbial community associated with a high-temperature petroleum reservoir at an offshore oilfield, *FEMS Microbiol. Ecol.*, 60, 74–84, 2007b.
- Li, H., Chen, S., Mu, B.-Z., and Go, J.-D.: Molecular detection of anaerobic ammonium-oxidizing (anammox) bacteria in high temperature petroleum reservoirs, *Microb. Ecol.*, 60, 771–783, 2010.
- Li, H., Mu, B.-Z., Jiang, Y., and Gu, J.-D.: Production processes affected prokaryotic *amoA* gene abundance and distribution in high-temperature petroleum reservoirs, *Geomicrobiol. J.*, 28, 692–704, 2011.
- Lozupone, C. and Knight, R.: UniFrac: a new phylogenetic method for comparing microbial communities, *Appl. Environ. Microbiol.*, 71, 8228–8235, 2005.
- Mbadinga, S. M., Wang, L.-Y., Zhou, L., Liu, J.-F., Gu, J.-D., and Mu, B.-Z.: Microbial communities involved in anaerobic degradation of alkanes, *Int. Biodeterior. Biodegrad.*, 65, 1–13, 2011.
- Mbadinga, S. M., Li, K. P., Zhou, L., Wang, L. Y., Yang, S. Z., Liu, J.-F., Gu, J.-D., and Mu, B.-Z.: Analysis of alkane-dependent methanogenic community derived from production water of high temperature petroleum reservoir, *Applied. Microbiol. Biotechnol.*, doi:10.1007/s00253-011-3828-8, 2012.
- Nazina, T. N., Shestakova, N. M., Grigor'yan, A. A., Mikhailova, E. M., Tourova, T. P., Poltraus, A. B., Feng, C.-X., Ni, F.-T., and Belyaev, S. S.: Phylogenetic diversity and activity of anaerobic microorganisms of high-temperature horizons of the Dagang oil field (P.R. China), *Microbiology*, 75, 55–65, 2006.
- Nilsen, R. K., Torsvik, T., and Lien, T.: *Desulfotomaculum thermocisternum* sp. nov., a sulfate reducer isolated from a hot North Sea oil reservoir, *Int. J. Syst. Evol. Microbiol.*, 46, 397–402, 1996.

The microbial communities of petroleum reservoirs fluids

L.-Y. Wang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Orphan, V. J., Taylor, L. T., Hafenbradl, D., and Delong, E. F.: Culture-dependent and culture-independent characterization of microbial assemblages associated with high-temperature petroleum reservoirs, *Appl. Environ. Microbiol.*, 66, 700–711, 2000.

Orphan, V. J., Goffredi, S. K., Delong, E. F., and Boles, J. R.: Geochemical influence on diversity and microbial processes in high temperature oil reservoirs, *Geomicrobiol. J.*, 20, 295–311, 2003.

Parkes, J.: Cracking anaerobic bacteria, *Nature*, 401, 217–218, 1999.

Ravot, G., Magot, M., Fardeau, M.-L., Patel, B. K. C., Prensier, G., Egan, A., Garcia, J.-L., and Ollivier, B.: *Thermotoga elfii* sp. nov., a novel thermophilic bacterium from an African oil-producing well, *Int. J. Syst. Evol. Microbiol.*, 45, 308–314, 1995.

Ravot, G., Magot, M., Fardeau, M.-L., Patel, B. K. C., Thomas, P., Garcia, J.-L., and Ollivier, B.: *Fusibacter paucivorans* gen. nov., sp. nov., an anaerobic, thiosulfate-reducing bacterium from an oil-producing well, *Int. J. Syst. Evol. Microbiol.*, 49, 1141–1147, 1999.

Rees, G. N., Grassia, G. S., Sheehy, A. J., Dwivedi, P. P., and Patel, B. K. C.: *Desulfacinum infernum* gen. nov., sp. nov., a thermophilic sulfate-reducing bacterium from a petroleum reservoir, *Int. J. Syst. Evol. Microbiol.*, 45, 85–89, 1995.

Saitou, N. and Nei, M.: The neighbor-joining method: a new method for reconstructing phylogenetic trees, *Mol. Biol. Evol.*, 4, 406–425, 1987.

Sekiguchi, Y., Kamagata, Y., Nakamura, K., Ohashi, A., and Harada, H.: *Syntrophothermus lipocalidus* gen. nov., sp. nov., a novel thermophilic, syntrophic, fatty-acid-oxidizing anaerobe which utilizes isobutyrate, *Int. J. Syst. Evol. Microbiol.*, 50, 771–779, 2000.

Shestakova, N. M., Ivoilov, V. S., Tourova, T. P., Belyaev, S. S., Poltarau, A. B., and Nazina, T. N.: Which microbial communities are present? Application of clone libraries: syntrophic acetate degradation to methane in a high-temperature petroleum reservoir-culture-based and 16S rRNA genes characterisation, in: *Applied Microbiology and Molecular Biology in Oilfield Systems*, edited by: Whitby, C. and Skovhus, T. L., Springer, Netherlands, 45–53, 2011.

Strapoć, D., Ashby, M., Wood, L., Levinson, R., and Huizinga, B.: How specific microbial communities benefit the oil industry: significant contribution of methyl/methanol-utilising methanogenic pathway in a subsurface biogas environment, in: *Applied Microbiology and Molecular Biology in Oilfield Systems*, edited by: Whitby, C. and Skovhus, T. L., Springer, Netherlands, 211–216, 2011.

The microbial communities of petroleum reservoirs fluids

L.-Y. Wang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Suflita, J. M., Davidova, I. A., Gieg, L. M., Nanny, M., and Prince, R. C.: Anaerobic hydrocarbon biodegradation and the prospects for microbial enhanced energy production, in: *Studies in Surface Science and Catalysis*, edited by: Rafael, V. D. and Rodolfo, Q.-R., Elsevier, Amsterdam, The Netherlands, 283–305, 2004.

5 Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M., and Kumar, S.: MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods, *Mol. Biol. Evol.*, 28, 2731–2739, 2011.

Voordouw, G., Armstrong, S., Reimer, M., Fouts, B., Telang, A., Shen, Y., and Gevertz, D.: Characterization of 16S rRNA genes from oil field microbial communities indicates the presence of a variety of sulfate-reducing, fermentative, and sulfide-oxidizing bacteria, *Appl. Environ. Microbiol.*, 62, 1623–1629, 1996.

10 Wang, L. Y., Mbadinga, S. M., Li, H., Liu, J. F., Yang, S. Z., and Mu, B. Z.: Anaerobic degradation of petroleum hydrocarbons and enlightenment of the prospects for microbial bio-gasification of residual oil, *Microbiology*, 37, 96–102, 2010 (in Chinese with English abstract).

15 Wang, L. Y., Gao, C. X., Mbadinga, S. M., Zhou, L., Liu, J. F., Gu, J. D., and Mu, B. Z.: Characterization of an alkane-degrading methanogenic enrichment culture from production water an oil reservoir after 274 days of incubation, *Int. Biodeterior. Biodegrada.*, 65, 444–450, 2011.

20 Wang, Q., Garrity, G. M., Tiedje, J. M., and Cole, J. R.: Naive bayesian classifier for rapid assignment of rRNA sequences into the new bacterial taxonomy, *Appl. Environ. Microbiol.*, 73, 5261–5267, 2007.

The microbial communities of petroleum reservoirs fluids

L.-Y. Wang et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)
Table 1. Characterization of the water samples collected from different petroleum reservoirs.

| | S1 Z3-26 | S2 B18-43 | S3 B51-45 | S4 M17-10 | S5 7222 | S6 6190 |
|---|-------------|--------------|--------------|--------------|------------|------------|
| Depth (m) | ~ 1300 | ~ 1490 | ~ 1101 | ~ 802 | ~ 1088 | ~ 480 |
| Temp (°C) | 63 | 58 | 45 | 37 | 32 | 21 |
| pH | 7.1 | 7.2 | 7.2 | 7.2 | 7.1 | 7.0 |
| Effective thickness (m) | 4.2 | 5.0 | 5.2 | 14.4 | 15.7 | 18.4 |
| Effective porosity (%) | 30 | 17.3 | 22.2 | 24.7 | 17.4 | 20.5 |
| Average permeability ($\times 10^{-3} \mu\text{m}^2$) | 800 | 691 | 12.6 | 675.3 | 274 | 466 |
| Oil viscosity (mPas) | 1720 | 13.7 | 402 | 179.1 | 44.8 | 417 |
| Water flooding operation (yr) | 22 | 10 | 4 | 22 | 46 | 38 |
| Mineralization (mg l^{-1}) | 8425 | 2891 | 4091 | 1121 | 15728 | 4212 |
| Cl^{-} (mg l^{-1}) | 3850 | 361 | 819 | 447 | 2000 | 3864 |
| SO_4^{2-} (mg l^{-1}) | 2244 | 12.1 | 32.4 | 6.8 | 7.7 | 124.8 |
| PO_4^{3-} (mg l^{-1}) | 0.1 | Nd | Nd | 0.08 | Nd | Nd |
| NO_3^{-} (mg l^{-1}) | Nd | Nd | Nd | Nd | 1.4 | 34.1 |
| Na^{+} (mg l^{-1}) | 3313 | 1629 | 1064 | 618.3 | 5399 | 4196 |
| K^{+} (mg l^{-1}) | 94.2 | 28.1 | 22.3 | 4.2 | 45.6 | 35.1 |
| Ca^{2+} (mg l^{-1}) | 195.6 | 3.6 | 53.0 | 19.2 | 128.2 | 103.3 |
| Mg^{2+} (mg l^{-1}) | 46.1 | 1.4 | 17.6 | 0.15 | 64.0 | 44.7 |
| Mn^{2+} (mg l^{-1}) | 0.3 | Nd | 0.1 | Nd | 0.4 | 0.3 |
| Acetate (mg l^{-1}) | 32 | 856 | 57.9 | 5.3 | 6.97 | 344 |
| Propionate (mg l^{-1}) | 1.2 | 8.0 | Nd | Nd | Nd | Nd |
| Isobutyrate (mg l^{-1}) | Nd | 13.8 | Nd | 9.8 | Nd | 32.7 |
| Butyrate (mg l^{-1}) | 0.2 | 2.3 | 0.5 | 2.3 | Nd | Nd |

(Nd: not detected)

The microbial communities of petroleum reservoirs fluids

L.-Y. Wang et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 2. Phylogenetic tree of the *Firmicutes* 16S rRNA gene phylotypes retrieved from six different petroleum reservoirs (shown in bold) and closely related sequences from GenBank database. Alignments to related sequences (shown with accession number) were performed with MEGA 5 software. The topology of the tree was obtained with the neighbor-joining method. Bootstrap values ($n = 1000$ replicates) of $\geq 75\%$ are reported. Scale bar represents nucleotide changes per site. Sampling locations are as named in Table 1.




Fig. 3. Phylogenetic tree of the *Actinobacteria*, *Thermodesulfobacteria*, *Thermotogae*, *Deinococcus-Thermus*, *Chloroflexi*, *Bacteroidetes* and TM7 16S rRNA gene phylotypes retrieved from six different petroleum reservoirs (shown in bold) and closely related sequences from GenBank database. Alignments to related sequences (shown with accession number) were performed with MEGA 5 software. The topology of the tree was obtained with the neighbor-joining method. Bootstrap values ($n = 1000$ replicates) of $\geq 75\%$ are reported. Scale bar represents nucleotide changes per site. Sampling locations are as named in Table 1.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 4. Phylogenetic tree of the archaeal 16S rRNA gene phylotypes retrieved from six different petroleum reservoirs (shown in bold) and closely related sequences from GenBank database. Alignments to related sequences (shown with accession number) were performed with MEGA 4 software. The topology of the tree was obtained with the neighbor-joining method. Bootstrap values ($n = 1000$ replicates) of $\geq 75\%$ are reported. Scale bar represents nucleotide changes per site. Sampling locations are as named in Table 1.

The microbial communities of petroleum reservoirs fluids

L.-Y. Wang et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The microbial communities of petroleum reservoirs fluids

L.-Y. Wang et al.




Fig. 5. Relative proportion of bacterial (**A**) and archaeal (**B**) taxa from 16S rRNA gene sequence clone libraries constructed from DNA extracted from production waters collected from oil reservoir with temperature 63, 58, 45, 37, 32 and 20 °C. Sampling locations are as named in Table 1.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

The microbial communities of petroleum reservoirs fluids

L.-Y. Wang et al.




Fig. 6. PCoA ordination diagrams of the bacterial (**A**) and archaeal (**B**) assemblages calculated with 16S rRNA gene sequences from production water of petroleum reservoirs. Shown are the plots of the first two principal coordinate axes (P1 and P2) for PCoA and the distributions of the bacterial and archaeal assemblages (designated with the sampling wells) in response to these axes.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 7. CCA ordination plots for the first two dimensions to show the relationship between the bacterial and archaeal diversity with environmental factors analyzed using a 16S rRNA gene sequence in the production water of petroleum reservoirs. Correlations between environmental variables and CCA axes are represented by the length and angle of arrows (environmental factor vectors), designated with the bacterial group was showed in (A, B and C) and designated with the archaeal group was showed in (a, b and c).