



<b>Title</b>	<b>Influenza-like illness among Hong Kong Chinese pregnant women</b>
<b>Author(s)</b>	<b>Wong, VWY; Tarrant, AM</b>
<b>Citation</b>	<b>The Journal of Maternal - Fetal &amp; Neonatal Medicine, 2015, v. 28 n. 1, p. 116-119</b>
<b>Issued Date</b>	<b>2015</b>
<b>URL</b>	<b><a href="http://hdl.handle.net/10722/197867">http://hdl.handle.net/10722/197867</a></b>
<b>Rights</b>	<b>The Journal of Maternal - Fetal &amp; Neonatal Medicine. Copyright © Informa Healthcare.</b>

## **SHORT REPORT**

### **Influenza-like illness among Hong Kong Chinese pregnant women**

Valerie W.Y. Wong<sup>1</sup>, Marie Tarrant<sup>1</sup>

#### **Affiliations:**

<sup>1</sup> School of Nursing, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Hong Kong Special Administrative Region, China.

#### **Author for correspondence and reprint requests:**

Valerie W.Y. Wong, School of Nursing, Li Ka Shing Faculty of Medicine, The University of Hong Kong, 21 Sassoon Road, Pokfulam, Hong Kong.

Tel: +852 2816 0137; Fax: +852 2872 6079; email: valw@hku.hk

Word count (abstract): 100

Word count (main text): 1,635

Running head: Influenza and Pregnancy

## **ABSTRACT**

We assessed the self-reported prevalence of influenza-like illness (ILI) during pregnancy in two samples of 546 and 2,764 new mothers who were pregnant during the 2009 – 10 and 2010 – 11 peak influenza seasons. During pregnancy, 11% of participants experienced an ILI. Cough, sore throat and nasal congestion were the most common reported symptoms. Only 4.6% and 9% of participants in sample 1 and 2 had an underlying chronic illness respectively and 3.3% of mothers in both groups were smokers. Conducting regular surveillance on influenza prevalence during pregnancy is essential to evaluate the costs and benefits of influenza vaccination programmes.

## **INTRODUCTION**

The latest position paper from the World Health Organization (WHO) on seasonal influenza vaccination states that the highest priority for vaccination should be pregnant women (1). The recommendation is based on the excess risk of influenza-related morbidity and mortality in this group, the safety of the influenza vaccine throughout the pregnancy, and the vaccine's effectiveness in reducing the risk of influenza and its complications among pregnant women and infants up to six months of age. However, existing evidence on the burden of seasonal influenza infection during pregnancy is largely based on studies from Western countries, and there is little evidence from other regions such as Asia. Pregnant women in various countries may have different susceptibility to influenza infection due to variation in the prevalence of risk factors in the population, such as underlying chronic illnesses, obesity, and smoking patterns. Therefore, assessing the prevalence of influenza and influenza-like-illness among pregnant women in different populations is important to evaluate the population level burden of influenza infection among pregnant women.

Moreover, community-based influenza surveillance focusing on pregnant women is essential to provide baseline information regarding influenza activity and in evaluating the potential benefits of influenza vaccination programmes. To our knowledge, only one previous study conducted in 2002 has actually reported the prevalence of influenza-like-illness among pregnant women (2). Therefore, the aims of this analysis were to assess the prevalence of influenza-like illness (ILI) among Chinese pregnant women and to examine its association with potential risk factors and adverse pregnancy outcomes.

## **METHODS**

This paper included two large samples of postpartum women from the postnatal wards of public hospitals providing obstetric services in Hong Kong. Briefly, eligible participants were Hong Kong Chinese mothers, 18 years old or above, and had just given birth to a live newborn. The first sample of participants were recruited from February to June 2010 during the second wave of A/H1N1 influenza pandemic and included only those mothers with no serious medical and obstetric complications (i.e., infant admitted to the neonatal intensive care unit, congenital anomalies or birth defects, born at <37 weeks, or birth weight <2,500g). In this study, we recruited 549 participants from four study sites. Between April and June 2011, we further recruited 2,846 postpartum participants from the postnatal wards of all eight public hospitals. A detailed description of the study methods of both studies is available elsewhere (3-4). Participants in both samples were pregnant throughout the winter influenza season, which theoretically increases the sensitivity and specificity of the case definition. Data were collected using a self-completed questionnaire assessing socio-demographic characteristics, maternal influenza vaccination status, maternal and paternal smoking patterns, pre-existing chronic illnesses, medical conditions arising during pregnancy, and the occurrence of respiratory and febrile illnesses during pregnancy.

The primary outcomes were the prevalence of ILI during the 2009 – 2010 and 2010 – 2011 influenza seasons. ILI was defined based on the febrile acute respiratory illness definition for influenza surveillance from the WHO, which defines ILI as the presence of fever plus cough or sore throat (5). We also examined at potential risk factors for ILI (i.e., maternal and paternal smoking status, seasonal and H1N1 influenza vaccination status, and underlying chronic illnesses and adverse pregnancy outcomes, defined as emergency caesarean section

(yes/no), preterm birth (<37 weeks gestation) and low birth weight (<2500 grams). Due to the historical low prevalence rate of obesity in Hong Kong, we did not assess BMI among pregnant women in either sample (6,7). Since we only included mothers without pregnancy complications in the first sample, no data on preterm birth or low birth weight is available for this group.

Descriptive statistics were used to assess the prevalence of self-reported ILI and logistic regression was used to assess the unadjusted and fully adjusted associations between ILI and both background risk factors and adverse pregnancy outcomes. All data analysis was conducted using Stata 13 (Stata Corp, College Station, TX). Ethical approval was obtained from the Institutional Review Boards of all participating institutions and informed written consent was obtained from all participants.

## **RESULTS**

Complete data were available for 546 (99.5%) of the 549 recruited participants in the 2009 – 2010 sample and 2,764 (97.1%) of the 2,846 participants in the 2010 – 2011 sample. The demographic characteristics of the study participants are presented in Table 1. Overall, sample 1 and 63% in sample 2 were 25 to 34 years of age, 67% in sample 1 and 63% in sample 2 were born in Hong Kong, only 3% were current smokers in both groups. Five percent (n = 25) and 9% (n = 254) had underlying chronic illnesses respectively, of which 36% (n = 9) and 28% (n = 72) were chronic respiratory illnesses. Of all participants, 11% of participants in both groups reported having an ILI in pregnancy. Since the participants in sample 1 were pregnant during the second wave of 2009 A/H1N1 influenza pandemic, their

seasonal influenza vaccination rate was marginally higher than the second sample (5.0% vs 1.7%). Cough, sore throat and nasal congestion were the three most common respiratory symptoms reported. In the fully adjusted analysis, only paternal smoking status during pregnancy was significantly associated with the risk of ILI (Table 2) while there were no significant associations between ILI and pregnancy outcomes (Table 3).

## **DISCUSSION**

To our knowledge, this is the first report on the population-based prevalence of ILI among Chinese pregnant women and only the second report on this topic in any population. Our results consistently show a low rate of ILI (11%) among Hong Kong women during pregnancy across two influenza seasons in 2009 – 10 and 2010 – 11. In comparison, Tuyishime reported that 64% of Canadian women experienced a febrile ILI during pregnancy (2), an almost six-fold difference. However, Tuyishime’s study adopted a somewhat broader ILI definition (any combination of respiratory symptoms with fever). Upon reclassification of our data using Tuyishime’s broader ILI definition, the prevalence remained unchanged at 11%. The substantial differences in the ILI rates between the two populations can possibly be explained by the presence of other risk factors such as pre-existing health conditions, obesity, and smoking rates among pregnant women. One US study found that 27% of pregnant women reported at least one chronic illness (7) compared with 5 to 9% in Hong Kong in our studies. In addition, the prevalence of obesity and maternal smoking in Asia are much lower than most Western countries. WHO data show that age-standardized obesity (BMI >30.0) estimates among adult women are 33% in the US, 25% in Australia and the United Kingdom, 24% in Canada, and 15% in France and Italy compared with 7% in China and Indonesia, 6% in Singapore, and only 4% in Japan. In addition, the age-standardized prevalence of smoking

among adult females was 32% in France, 22% in the UK, and 17% in the US, while Asia Pacific rates were substantially lower at 11% in Japan, 4% in Singapore, 2% in China and Thailand, and 1% in Malaysia and Vietnam (8).

Furthermore, while current evidence suggests that pregnant women are at higher risk of seasonal influenza-related morbidity, mortality, and adverse obstetric outcomes, it is not clear how much of this evidence is from studies conducted in Asia. During the 2009 A/H1N1 influenza pandemic, outcomes for infected pregnant women in Asia were much more favourable than in Western countries. In Hong Kong, up until May 2010, 87 pregnant women infected with 2009 A/H1N1 influenza were hospitalized, and only one required mechanical ventilation and none died (9). In Singapore, 211 A/H1N1 laboratory-confirmed pregnant women presented to a hospital between May and September, 2009; only two cases were complicated by pneumonia and both recovered uneventfully (10). In Japan, 181 pregnant women required hospitalization and none died (11) and only five Korean pregnant women were admitted to hospital and one died (12).

This study has several limitations. The data reported here was a secondary analysis of two studies designed to examine uptake and predictors of maternal influenza vaccination and was not designed specifically to measure the burden of influenza infection of ILI during pregnancy. Recall bias may also be an issue since the study outcome data was collected retrospectively through self-report in the first 24 to 48 hours after birth when participants may be less able to recall events that happened earlier in the pregnancy. Research has shown, however, that maternal recall of events during pregnancy is reliable and accurate (13) and this is especially likely to be true with regard to febrile illness during pregnancy. In addition, this



study did not assess laboratory-confirmed influenza and therefore, the true prevalence of influenza infection may actually be lower. Although our results show no association between ILI during pregnancy and adverse pregnancy outcomes, it is possible that mothers with preterm or low birth weight infants may have been less willing to participate in the study and thus were under sampled.

Determining the actual burden of influenza infection in high-risk populations is important to facilitate the planning of effective influenza vaccination programmes and other primary preventive interventions. WHO recommends that regional governments establish sustainable and effective surveillance systems to assess the epidemiology of influenza in high-risk groups, such as pregnant women and chronically ill individuals (1). In regions where the burden of influenza among pregnant women may be low, this type of surveillance is necessary so that resources are allocated to the groups with the highest needs. In the case of maternal vaccination programmes, it is also important to assess the burden of influenza infection in newborns under 6 months of age as the benefits of maternal vaccination for this group may provide a more substantial rationale for vaccination programmes.

## **FUNDING**

Funding was provided by the Health and Medical Research Fund for the Control of Infectious Diseases, Government of the Hong Kong Special Administrative Region, Hong Kong SAR (Grant: #10090982).

## **DECLARATION OF INTERESTS**

The authors report no declarations of interest.

## REFERENCES

- [1] Vaccines against influenza WHO position paper - November 2012. Weekly Epidemiological Record 2012;87(47):461-476.
- [2] Tuyishime JD, De Wals P, Moutquin JM, Frost E. Influenza-like illness during pregnancy: results from a study in the eastern townships, Province of Quebec. Journal of Obstetrics and Gynaecology Canada 2003;25(12):1020-1025.
- [3] Tarrant M, Wu KM, Yuen CY, Cheung KL, Chan VH. Determinants of 2009 A/H1N1 influenza vaccination among pregnant women in Hong Kong. Maternal and Child Health Journal 2013;17(1):23-32.
- [4] Yuen YSC, Fong YTD, Lee LYI, Sing C, Siu SME, Tarrant M. Prevalence and predictors of maternal seasonal influenza vaccination in Hong Kong. Vaccine 2013;31(2013):5281-5288.
- [5] World Health Organization (WHO). WHO Interim Global Epidemiological Surveillance Standards for Influenza. Geneva: World Health Organization, 2012.
- [6] Ko GT, Tang JS. Prevalence of obesity, overweight and underweight in a Hong Kong community: the United Christian Nethersole Community Health Service (UCNCHS) primary health care program 1996-1997. Asia Pacific Journal of Clinical Nutrition. 2006;15(2):236-241.
- [7] Chatterjee S, Kotelchuck M, Sambamoorthi U. Prevalence of chronic illness in pregnancy, access to care, and health care costs: implications for interconception care. Womens Health Issues 2008;18(6 Suppl):S107-116.
- [8] World Health Organization (WHO). WHO report on the global tobacco epidemic, 2013: enforcing bans on tobacco advertising, promotion and sponsorship. Geneva: 2013.
- [9] Hospital Authority. Pandemic H1N1 influenza admissions to public hospitals in Hong Kong 2010 (unpublished data). Hong Kong 2010.

- [10] Lim ML, Chong CY, Tee WSN, Lim WY, Chee JJ. Influenza A/H1N1 (2009) infection in pregnancy -- an Asian perspective. *British Journal of Obstetrics and Gynaecology* 2010;117(5):551-556.
- [11] Nakai A, Saito S, Unno N, Kubo T, Minakami H. Review of the pandemic (H1N1) 2009 among pregnant Japanese women. *Journal of Obstetric and Gynaecology Research* 2012;38(5):757-762.
- [12] An JH, Kim HN, Choi OJ, Kim GS, Kim UJ, Jang MO, Kang SJ, Park KH, Jung SI, Kwon YS, Jang HC. Was 2009 Pandemic Influenza A (H1N1) Mild Among Pregnant Korean Women? *Chonnam Medical Journal* 2013;49(2):96-99.
- [13] Bat-Erdene U, Metcalfe A, McDonald SW, Tough S. Validation of Canadian mothers' recall of events in labour and delivery with electronic health records. *BioMed Central Pregnancy and Childbirth* 2013;13(Suppl 1):S3.

**Table 1 Characteristics of participants**

Variable	Sample 1		Sample 2	
	N	wt %	N	wt %
<b>Total</b>	<b>546</b>	<b>100.0</b>	<b>2,764</b>	<b>100.0</b>
<b>Age</b>				
18 - 24 years	41	7.5	263	9.5
25 - 29 years	131	24.0	716	25.9
30 - 34 years	247	45.2	1,027	37.2
≥ 35 years	127	23.3	758	27.4
<b>Education</b>				
Compulsory secondary	276	50.6	1,649	59.7
Upper secondary	107	19.6	488	17.7
University degree or above	163	29.9	627	22.7
<b>Place of birth</b>				
Hong Kong	367	67.2	1,735	62.8
Mainland China	167	30.6	968	35.0
Other	12	2.2	61	2.2
<b>Smoking status during pregnancy</b>				
No	528	96.7	2,674	96.7
Yes	18	3.3	90	3.3
<b>Partner's smoking status during pregnancy</b>				
No	374	68.5	1,855	67.1
Yes	172	31.5	909	32.9
<b>Seasonal influenza vaccination status during pregnancy</b>				
No	519	95.1	2,716	98.3
Yes	27	5.0	48	1.7
<b>H1N1 influenza vaccination status during pregnancy</b>				
No	512	93.8	NA	NA
Yes	34	6.2	NA	NA
<b>Pre-existing chronic illness</b>				
No	521	95.4	2,510	90.8
Yes	25	4.6	254	9.1
<i>Types: (some participants have more than one illness)</i>				
<i>Respiratory/ pulmonary disease</i>	9	36.0	72	28.4
<i>Thyroid disease</i>	4	16.0	42	16.5
<i>Hepatitis B</i>	7	28.0	38	15.0
<i>Others</i>	6	24.0	120	47.2
<b>Emergency caesarian section</b>				
No	498	91.2	2,412	87.3
Yes	48	8.8	352	12.7
<b>Preterm birth (&lt;37 weeks gestation)</b>				
No	NA	NA	2,550	92.3
Yes	NA	NA	214	7.7
<b>Low birth weight infant (&lt;2500 gms)</b>				
No	NA	NA	2,548	92.2
Yes	NA	NA	216	7.8

NA=not available

<b>Table 2 Unadjusted and adjusted odds ratios for risk of ILI with regards to potential risk factors</b>												
<b>Variable</b>	<b>Sample 1</b>						<b>Sample 2</b>					
	<b>Unadjusted ILI</b>			<b>Adjusted ILI<sup>1</sup></b>			<b>Unadjusted ILI</b>			<b>Adjusted ILI<sup>1</sup></b>		
	<b>OR</b>	<b>95% CI</b>	<b>Sig.</b>	<b>AOR</b>	<b>95% CI</b>	<b>Sig.</b>	<b>OR</b>	<b>95% CI</b>	<b>Sig.</b>	<b>AOR</b>	<b>95%CI</b>	<b>Sig.</b>
<b>Smoking status during pregnancy</b>												
No	1			1			1			1		
Yes	0.99	0.22, 4.43	0.99	0.95	0.21, 4.39	0.95	1.33	0.73, 2.42	0.36	1.15	0.62, 2.16	0.66
<b>Partner's smoking status during pregnancy</b>												
No	1			1			1			1		
Yes	0.82	0.46, 1.49	0.52	0.85	0.45, 1.60	0.61	1.11	0.87, 1.42	0.42	<b>1.36</b>	<b>1.04, 1.79</b>	<b>0.03</b>
<b>Seasonal influenza vaccine status in pregnancy</b>												
No	1			1			1			1		
Yes	0.99	0.29, 3.40	0.99	0.65	0.16, 2.56	0.54	0.70	0.25, 1.97	0.50	0.62	0.22, 1.75	0.36
<b>H1N1 influenza vaccine status in pregnancy</b>												
No	1			1								
Yes	2.20	0.91, 5.29	0.08	2.20	0.82, 5.92	0.12	NA	NA	NA	NA	NA	NA
<b>Pre-existing chronic illness</b>												
No	1			1			1			1		
Yes	1.09	0.32, 3.75	0.89	1.00	0.28, 3.57	1.00	1.05	0.70, 1.56	0.83	1.06	0.71, 1.59	0.78

<sup>1</sup>Adjusted for age group, educational level, place of birth, and the remaining variables in the table  
NA=not available

**Table 3 Unadjusted and adjusted ORs for the risk of selected maternal and neonatal outcomes with regard to ILI in sample 2**

Variable	Emergency caesarian section			Preterm			Low birth weight		
	OR	95% CI	Sig.	OR	95% CI	Sig.	OR	95% CI	Sig.
<b>Unadjusted ILI</b>									
No	1			1			1		
Yes	1.03	0.73, 1.46	0.87	1.24	0.82, 1.87	0.30	1.23	0.81, 1.85	0.33
<b>Adjusted ILI<sup>1</sup></b>									
No	1			1			1		
Yes	0.99	0.70, 1.42	0.93	1.23	0.81, 1.87	0.33	1.14	0.75, 1.73	0.55

<sup>1</sup>Adjusted for age group, educational level, place of birth, family income, maternal smoking status, seasonal influenza vaccination status, pre-existing chronic illness, and medical conditions arise during pregnancy  
 NA = not available