



Title	Unraveling the molecular basis of temperature-dependent genetic regulation in <i>Penicillium marneffei</i>
Author(s)	Yang, E; Wang, G; Woo, PCY; Lau, SKP; Chow, WN; Chong, KTK; Tse, H; Kao, RYT; Chan, CM; Che, X; Yuen, KY; Cai, JJ
Citation	Eukaryotic Cell , 2013, v. 12 n. 9, p. 1214-1224
Issued Date	2013
URL	http://hdl.handle.net/10722/200746
Rights	Eukaryotic Cell . Copyright © American Society for Microbiology.; This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License.

Unraveling the Molecular Basis of Temperature-Dependent Genetic Regulation in *Penicillium marneffeii*

Ence Yang,^a Gang Wang,^a Patrick C. Y. Woo,^b Susanna K. P. Lau,^b Wang-Ngai Chow,^b Ken T. K. Chong,^b Herman Tse,^b Richard Y. T. Kao,^b Che-Man Chan,^b Xiaoyan Che,^c Kwok-Yung Yuen,^b James J. Cai^a

Department of Veterinary Integrative Biosciences, College of Veterinary Medicine and Biomedical Sciences, Texas A&M University, College Station, Texas, USA^a; Department of Microbiology, University of Hong Kong, Hong Kong^b; Center for Clinical Laboratory, Zhujiang Hospital, Southern Medical University, Guangzhou, People's Republic of China^c

Penicillium marneffeii is an opportunistic fungal pathogen endemic in Southeast Asia, causing lethal systemic infections in immunocompromised patients. *P. marneffeii* grows in a mycelial form at the ambient temperature of 25°C and transitions to a yeast form at 37°C. The ability to alternate between the mycelial and yeast forms at different temperatures, namely, thermal dimorphism, has long been considered critical for the pathogenicity of *P. marneffeii*, yet the underlying genetic mechanisms remain elusive. Here we employed high-throughput sequencing to unravel global transcriptional profiles of *P. marneffeii* PM1 grown at 25 and 37°C. Among ~11,000 protein-coding genes, 1,447 were overexpressed and 1,414 were underexpressed at 37°C. Counter-intuitively, heat-responsive genes, predicted in *P. marneffeii* through sequence comparison, did not tend to be overexpressed at 37°C. These results suggest that *P. marneffeii* may take a distinct strategy of genetic regulation at the elevated temperature; the current knowledge concerning fungal heat response, based on studies of model fungal organisms, may not be applicable to *P. marneffeii*. Our results further showed that the tandem repeat sequences (TRSs) are overrepresented in coding regions of *P. marneffeii* genes, and TRS-containing genes tend to be overexpressed at 37°C. Furthermore, genomic sequences and expression data were integrated to characterize gene clusters, multigene families, and species-specific genes of *P. marneffeii*. In sum, we present an integrated analysis and a comprehensive resource toward a better understanding of temperature-dependent genetic regulation in *P. marneffeii*.

With over one million new infections every year and latent infections worldwide growing to the tens of millions, dimorphic fungal infection presents many significant public health challenges to date (1–3). As the only known dimorphic species in the *Penicillium* genus, *Penicillium marneffeii* (recently renamed *Talaromyces marneffeii*) is of particular concern. *P. marneffeii* primarily infects immunocompromised individuals, causing lethal systemic infection, or penicilliosis (4–6). Over the last 20 years, there has been a marked increase in the number of penicilliosis cases, with a concurrent rise in immunosuppression due to the global spread of HIV infection (7–15). Penicilliosis is the third most prevalent opportunistic infection among HIV patients in Southeast Asia (16). Thus, infection by *P. marneffeii* has become an AIDS-defining illness (6, 17).

P. marneffeii, *Blastomyces dermatitidis*, *Histoplasma capsulatum*, *Coccidioides immitis*, *Paracoccidioides brasiliensis*, and *Sporothrix schenckii*, are collectively regarded as thermally dimorphic fungi, which undergo temperature-dependent transition between two distinct growth forms: mycelia and yeast. For instance, *P. marneffeii* grows vegetatively as mycelia at 25°C, showing the typical multinuclear mold morphology; at 37°C, it undergoes the phase transition with concomitant coupling of nuclear and cellular division to form uninucleate, single-celled yeasts. The mycelium-to-yeast transition is considered to be a requisite for pathogenesis of *P. marneffeii*, as the yeast cell is the *in vivo* form that capably evades the host immune system (18, 19). These features well place *P. marneffeii* as a model experimental system for investigation of fungal growth processes and their contribution to pathogenicity (19).

Early dimorphic fungal studies were limited to morphological examination or clinical isolates and the cellular events accompa-

nying the phase transition processes. Only in the past decade have studies focused on the molecular mechanisms of phase transition with the application of novel genetic approaches in *P. marneffeii* (20–22). To date, more than 40 *P. marneffeii* genes have been cloned or experimentally characterized (23–27). However, except for several transcriptional regulators and signal transduction factors, most of these genes are involved in vegetative growth and asexual differentiation (28). Thus, very little is known about genes responsible for phase-specific growth.

In a previous publication (29), we briefly announced the sequencing of the genome of *P. marneffeii* strain PM1, without providing in-depth genomic and transcriptomic analyses. In the present study, we use comparative genomic and transcriptomic approaches to systematically characterize *P. marneffeii* genomic sequences and global gene expression. We refine the annotation of protein-coding genes and then use high-throughput mRNA sequencing (RNA-seq) to measure gene expression in mycelia at 25°C and yeast cells at 37°C. Through comparative genomics between *P. marneffeii* and several model fungal species, we gain new insights into the evolutionary history of the *P. marneffeii* genome. Our transcriptomic analysis suggests the existence of uncharacter-

Received 2 July 2013 Accepted 5 July 2013

Published ahead of print 12 July 2013

Address correspondence to James J. Cai, jcai@tamu.edu.

Supplemental material for this article may be found at <http://dx.doi.org/10.1128/EC.00159-13>.

Copyright © 2013, American Society for Microbiology. All Rights Reserved.

doi:10.1128/EC.00159-13

ized regulatory pathways that might be essential for thermal adaptation in *P. marneffeii*.

MATERIALS AND METHODS

DNA preparation, sequencing, and assembly. Genomic DNA was prepared from the arthroconidia of *P. marneffeii* PM1 grown at 37°C. A single colony of the fungus grown on Sabouraud's dextrose agar (SDA) at 37°C was inoculated into yeast peptone broth and incubated in a shaker for 3 days. Cells were cooled in ice for 10 min, harvested by centrifugation at 2,000 × g for 10 min, washed twice, and resuspended in ice-cold 50 mM EDTA buffer (pH 7.5). Novozyme 234 (20 mg/ml) was added, and the mixture was incubated at 37°C for 1 h followed by digestion in a mixture of 1 mg/ml proteinase K, 1% *N*-lauroylsarcosine, and 0.5 M EDTA (pH 9.5) at 50°C for 2 h. Genomic DNA was then extracted by phenol and phenol-chloroform and finally precipitated and washed in ethanol. After digestion with RNase A, a second ethanol precipitation was followed by washing with 70% ethanol, and then the DNA was air dried and dissolved in 500 μl of Tris-EDTA (TE [pH 8.0]) (30). Two genomic DNA libraries were made in pUC18 carrying insert sizes from 2.0 to 3.0 kb and from 7.5 to 8.0 kb, respectively. DNA inserts were prepared by physical shearing using the sonication method (31). The genome sequence was assembled from deep whole-genome shotgun (WGS) coverage obtained by paired-end sequencing from a variety of clone types (i.e., all inserts were sequenced from both ends to generate paired reads). A total of 190.3 Mb of sequence data, which is equivalent to ~6.6× coverage of the genome, was generated by random shotgun sequencing. The Phred/Phrap/Consed (<http://www.phrap.org/>) package was used for base calling, contig assembly, and quality assessment (32, 33). Contigs were ordered into supercontigs (also known as scaffolds) by using the scaffold building program Bambus (<http://www.cbcb.umd.edu/software/bambus/>) (34).

RNA-seq analysis. At the beginning of the experiment, conidia of strain PM1 were inoculated in SDA plates and cultured at 25 and 37°C for a week. The germinated cells were transferred into new SDA plates every week for 2 weeks to establish stable colonies in either the mycelial or yeast growth form. One week before the extraction of total RNA, in order to obtain fresh cells, the homogenous cells were cultured in new SDA plates. The total RNAs were extracted from each condition for two independent biological replicates using the E.Z.N.A. fungal RNA kit (Omega Bio-Tek). The effect of global transcriptional amplification or suppression might lead to erroneous interpretation of gene expression experiments (35). To control for this effect, we adjusted the total RNA concentration according to the DNA content before the standard poly(A)⁺ RNA-seq was performed. DNA was quantified using Qubit 2.0 fluorometer (Life Technologies, Grand Island, NY). We obtained ~13 million 90-bp paired-end reads for each sample. The reads were trimmed basing on their qualities by FastQC and fastx-trimmer. Filtered reads were mapped to the annotated genomes with Tophat v2.0.2 (36). For each sample, ~2.0 Gb of reads was mapped, representing ~100× coverage of *P. marneffeii* transcriptome. Cufflinks v2.0.2 (36) was used to calculate the fragments per kilobase of exon per million fragments mapped (FPKM) (37) and the confidence internals of the estimation for each gene. For differential expression analysis, mapped reads were counted using SAMMATE (<http://sammate.sourceforge.net/>) (38). Differentially expressed genes were detected by jointly using three R packages: edgeR (39), DESeq (40), and baySeq (41), which all are available at the Bioconductor open source bioinformatics software repository (<http://bioconductor.org/>). Genes with a false discovery rate (FDR) of <0.05 as reported by all three packages were considered to be differentially expressed.

Gene prediction and identification of orthologs. *Ab initio* gene predictions were performed using FGENESH (SoftBerry, Mount Kisco, NY). The original prediction was manually refined with assistance from GenomeScan (<http://genes.mit.edu/genomescan.html>), another gene prediction program that combines sequence similarity and exon-intron composition. The putative ortholog pairs were predicted by using InParanoid (<http://inparanoid.sbc.su.se/>) (42). Synteny was identified us-

ing ADHoRe (<http://bioinformatics.psb.ugent.be/software/details/ADHoRe>) (43), with an *r*² cutoff of 0.8, maximum gap size of 35 genes, and minimum number of pairs of 3. The ADHoRe results were filtered such that the maximum probability for a segment to be generated by chance was <0.01. Gene families in each genome were identified by using NCBI BLASTClust (with 30% of identical residues and aligned over at least 80% of their lengths).

IFA staining with Mp1p-specific antibodies. *P. marneffeii* was first cultured on SDA plates at 37 and 25°C for 7 days, respectively. Culture supernatants were obtained by inoculating fungal cells from plates into RPMI 1640 and shaken at 37 and 25°C for 5 days. Tissues were collected by centrifugation at 5,000 rpm for 20 min. The fungal mycelium and yeast cells were washed twice, resuspended in phosphate-buffered saline (PBS), and placed on wells of Teflon-coated slides. The immobilized cells were fixed for 10 min with prechilled fixed buffer containing acetone (70% [wt/vol]) and methanol (30% [vol/vol]) and stored at 70°C until they were used. The slides were incubated with the monoclonal antibodies (MAbs) against Mp1p at a concentration of 10 μg/ml in a humid chamber for 60 min at 37°C. The MAb was prepared by a procedure described previously (44, 45). Briefly, BALB/c mice were immunized intradermally with 50 mg of purified Mp1p emulsified with Freund's adjuvant at 10-day intervals for 8 weeks followed by intravenously administered boosters for 3 days before fusion. Antibodies specific for *P. marneffeii* in the hybridoma cell culture supernatant were screened by indirect enzyme-linked immunosorbent assay (ELISA) using Mp1p and indirect fluorescent antibody (IFA) staining using slides with immobilized fungal cells killed in 3.7% formalin solution overnight at 4°C. The isotypes of the MAbs were determined using a commercially available mouse MAb isotyping kit (Zymed Laboratories, Carlsbad, CA). MAbs were purified from ascitic fluid by ammonium sulfate precipitation and were conjugated with biotin using the EZLink Sulfo-NHS-biotinylation kit following the manufacturer's instructions (Pierce Biotechnology, Rockford, IL). After the slides were washed with PBS containing 0.1% (vol/vol) Tween 20, fluorescein isothiocyanate-conjugated goat anti-mouse IgG (Zymed Laboratories) was added and the slides were incubated for 40 min at 37°C. After several washes, a 0.25% (wt/vol) solution of Evans blue in PBS was added for counterstaining; fluorescence was determined by fluorescent laser confocal microscopy with a Leica TCS SP2 AOBs microscope.

GO analysis. The orthologs of *P. marneffeii* genes in *Aspergillus fumigatus*, *Aspergillus nidulans*, *Candida albicans*, *Neurospora crassa*, or *Saccharomyces cerevisiae* were identified, and then the Gene Ontology (GO) terms of orthologs of *P. marneffeii* in all five model fungal species were taken together. Nonredundant terms were assigned to corresponding *P. marneffeii* genes. In GO term enrichment tests, we grouped GO terms in the resulting lists into major classes based on the hierarchical structure between GO terms using REVIGO (46). Significance of functional enrichment of GO terms in differentially expressed gene sets was assessed using the chi-square test. All resulting *P* values were adjusted for multiple hypothesis testing using the method of Storey and Tibshirani (47), limiting the FDR to 0.05.

Identification of coding tandem repeats. The intragenic tandem repeats in the *P. marneffeii* genome and other fungal genomes were identified by a previously described method (48) based on EMBOSS ETANDEM (49). The threshold score was set to 20. All genes were scanned for long repeats (>40 nucleotides [nt]) or short repeats (3 to 39 nt). A sequence was considered to be an intragenic repeat if it met two conditions: (i) repeat conservation was at least 85%, and (ii) the numbers of repeats were at least 20 for trinucleotide repeats, 16 for repeats between 4 and 10 nt, 10 for repeats between 11 and 39 nt, and 3 for repeats of at least 40 nt.

Data sources for model fungal organisms. The genome and protein sequences of model fungal organisms were obtained from the following sources: AspGD (50) for *A. fumigatus* and *A. nidulans*, the *Candida* Genome Database (CGD) (51) for *C. albicans*, the *Neurospora crassa* Database (<http://www.broadinstitute.org/annotation/genome/neurospora>) for *N. crassa*, and the *Saccharomyces* Genome Database (SGD) (52) for *S.*




FIG 1 Temperature-dependent dimorphism in *P. marneffeii*. (Top panels) One-week *P. marneffeii* grown on SDA plates. Soluble red pigments were produced and diffused into the agar at 25°C; a yeast-type colony formed with decreased production of pigments at 37°C. (Bottom panels) $\times 400$ microscopic views of long mycelia, conidia, and conidiophore (white arrow) at 25°C and fission yeast-like cells (arthroconidia) at 37°C.

cerevisiae. The phenotypic information of single mutants, used to identify heat-responsive genes, was collected from AspGD (50) for *A. fumigatus* and *A. nidulans*, and PhenomeBLAST for *S. cerevisiae*.

Data accession number. Expression data have been submitted to the GEO database under accession number [GSE48898](https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE48898).

RESULTS

Under controlled laboratory conditions, *P. marneffeii* PM1 displayed two temperature-dependent morphologies: the mycelial form at 25°C and the yeast form at 37°C (Fig. 1). We took a systematic approach, coupling genome sequence and transcriptome analyses, to investigate the genetic regulation of thermal dimorphism in *P. marneffeii*. The rest of the article is organized as follows. First we outline the basic features of the genome sequence of *P. marneffeii* PM1 and present the global results of RNA-seq analysis defining its transcriptome. Subsequently, we describe the annotation of gene functions and detect *P. marneffeii* genes expressed differentially between mycelial cells at 25°C and yeast cells at 37°C, emphasizing the expression of heat-responsive genes. Finally, we integrate the genome sequencing and expression data to describe gene clusters, genes with tandem repeat sequences in the coding region, and multigene families and species-specific genes in *P. marneffeii*.

Genome sequences and transcriptome profile of *P. marneffeii*. The draft genome of *P. marneffeii* PM1 was obtained using the whole-genome shotgun (WGS) approach (29). A total of 311,029 WGS reads of an average length of 690 bp were assembled into 2,780 contigs, which were subsequently ordered into 273 supercontigs (Table 1). The assembly produces a $6.6\times$ coverage of the *P. marneffeii* genome of an estimated size of ~ 31 million bp. The genome size of *P. marneffeii* is comparable to those of *Aspergillus fumigatus*, *A. nidulans*, and *Penicillium chrysogenum* (29.4, 30.5, and 32.2 Mb, respectively) (53–55). A total of 10,060 protein-coding genes were initially predicted *in silico*. The genic regions

occupy 62.1% of the total genome sequences, while the protein-coding sequences occupy 51.2%. The average gene density is one gene per 2.8 kb, and the average gene length is 1.75 kb. A total of 28,180 introns were found in 91% of *P. marneffeii* genes. The size of introns ranges from 15 to 1,617 bp, with a mean of 111 bp. The telomere tandem repeat TTAGGG, along with telomere-associated helicases, was identified at the end of 16 contigs, suggesting 8 chromosomes in *P. marneffeii*, as in *A. fumigatus* and *A. nidulans*.

We precultured isogenic *P. marneffeii* PM1 on solid media at 25 and 37°C. Mycelial and yeast cells were collected for RNA-seq analysis. Over 13 million paired-end reads 90 bp long for each sample were generated with two biological replicates. After short-read mapping and transcript calling, the expressed transcripts were obtained and compared against the predicted genes. Through the comparison, 982 novel transcripts, overlapping with at least a 50-bp open reading frame (ORF), were identified. Accordingly, the total number of predicted protein-coding genes was updated to 11,042. Among these genes, 26.34% (2,908 genes) are expressed into more than one isoform through alternative splicing, making a total of 15,567 unique transcripts. The expression level was estimated at the whole-gene level in fragments per kilobase of exon per million fragments mapped (FPKM) (37). In mycelial cells at 25°C, 95.2% of genes (10,508) and 89.8% of transcripts (13,993) were expressed (FPKM > 0.05); in yeast cells at 37°C, 97.2% of genes (10,730) and 91.6% of transcripts (14,256) were expressed. Jointly, 98.5% of genes (10,882) and 95.6% of transcripts (14,886) were expressed at either temperature.

To identify *P. marneffeii* genes expressed at a significantly different level between mycelial cells at 25°C and yeast cells at 37°C, we conducted differential expression analysis with RNA-seq data. Here we followed the general practice of assuming that all samples placed into the analysis platform contain similar amounts of total RNA (cf. 35). We required that the differentially expressed (DE) genes be those detected as significant by all three commonly used algorithms—namely, edgeR (39), DESeq (40), and baySeq (41)—at the level of a false discovery rate (FDR) of <0.05 simultaneously. Using such conservative criteria, a total of 2,861 DE genes were identified, including 1,447 overexpressed and 1,414 underexpressed in yeast cells at 37°C, compared with mycelial cells at 25°C (see Table S1 in the supplemental material).

There are 32 *P. marneffeii* genes whose expressions were previ-

TABLE 1 General assembly features of the *P. marneffeii* genome

Feature	Value
Coverage	6.6
Estimated genome size, Mb	~ 31
Assembly length (excluding gaps), Mb	28.98
Total contig length, Mb	28.89
Scaffolds, no.	273
Contigs, no.	2,780
GC genome, %	47
Predicted protein-coding genes, no.	10,060
Portion of coding, %	62
GC coding, %	50
Mean gene length, bp	1,753
Mean intergenic distance, bp	1,051
Mean exon length, bp	380
Mean intron length, bp	111
tRNAs, no.	110

TABLE 2 Comparison of gene expression results from previously published studies with those from our RNA-seq study

Gene	Function	Previously reported result		RNA-seq result in this study				Expression change at 37°C	Replication
		Expression change at 37°C	Reference	25°C		37°C			
				Replicate 1	Replicate 2	Replicate 1	Replicate 2		
<i>abaA</i>	Transcriptional regulator	↓	27	163.3	170.3	20.1	18.4	↓	Yes
<i>actA</i>	Actin	—	79	1313.2	1,350.2	688.8	890.4	—	Yes
<i>acuD</i>	Isocitrate lyase	↑	80	198.5	486.8	217.3	225.0	—	No
<i>areA</i>	Transcriptional activator	—	81	30.1	44.3	37.1	50.6	—	Yes
<i>brlA</i>	Transcription factor	↓	82	1,407.6	1,340.8	5.1	4.2	↓	Yes
<i>cflA</i>	Rho family GTPase	—	83	153.3	149.1	140.0	122.6	—	Yes
<i>cflB</i>	Rho family GTPase	—	84	110.4	116.0	45.2	73.0	—	Yes
<i>cpeA</i>	Catalase-peroxidase	—	85	595.3	934.8	329.8	105.9	—	Yes
<i>drkA</i>	Two-component histidine kinase	—	86	72.5	73.5	40.1	46.8	—	Yes
<i>gasA</i>	G-protein complex a subunit	—	87	71.5	64.3	100.1	69.5	—	Yes
<i>gasC</i>	G-protein complex a subunit	↓	23	102.3	108.9	39.2	41.8	↓	Yes
<i>gpdA</i>	Glyceraldehyde-3-phosphate dehydrogenase	↓	88	3,280.2	3,146.5	2,701.4	2,209.3	—	No
<i>hsp30</i>	Heat shock protein	↓	89	1,059.1	764.4	2,645.7	3,081.7	—	No
<i>hsp70</i>	Heat shock protein	↑	90	2,401.0	2,503.0	1,259.7	1,757.3	—	No
<i>mp1</i>	Antigenic mannoprotein	↑	91	1.6	3.1	1,551.1	493.6	↑	Yes
<i>mplp6</i>	Antigenic mannoprotein	↑	92	0.0	0.0	1,081.4	177.5	↑	Yes
<i>myoB</i>	Type II myosin	—	93	32.8	18.0	17.8	18.6	—	Yes
<i>pakA</i>	p21-activated kinase	—	94	46.2	47.5	42.4	46.5	—	Yes
<i>pakB</i>	p21-activated kinase	↓	26	68.1	82.3	44.9	38.1	—	No
<i>hbk1</i>	Histidine kinase	—	95	11.8	10.3	9.0	15.4	—	Yes
<i>rsr1</i>	RAS small monomeric GTPase	↑	96	22.1	18.3	39.8	17.4	—	No
<i>rasA</i>	Ras GTPase	—	97	45.9	38.7	24.8	20.7	—	Yes
<i>rfxA</i>	Transcriptional regulator	—	21	82.3	68.1	52.1	65.7	—	Yes
<i>sakA</i>	Stress-activated protein kinase	—	98	201.3	215.8	165.6	103.0	—	Yes
<i>skn7</i>	Stress response transcription factor	—	99	72.0	80.6	47.5	35.8	—	Yes
<i>slnA</i>	Two-component histidine kinase	—	86	11.1	10.9	23.6	19.8	—	Yes
<i>sodA</i>	Cu, Zn superoxide dismutase	—	100	409.1	674.6	500.5	785.9	—	Yes
<i>stlA</i>	Control of mating and yeast-hyphal transitions	—	101	43.3	55.9	30.6	49.9	—	Yes
<i>stuA</i>	APSES transcription factor	↓	24	256.8	403.5	25.7	49.9	↓	Yes
<i>thpA</i>	RNA polymerase I and III transcription factor	—	25	379.0	674.6	507.5	532.7	—	Yes
<i>tupA</i>	Transcriptional repressor	—	102	76.2	49.9	41.8	38.9	—	Yes

^a ↑, expression is higher at 37°C than at 25°C; ↓, expression is lower at 37°C than at 25°C; —, no difference in expression.

ously measured mostly by using microarrays at 25 and 37°C or in mycelial and yeast cells (Table 2). We made a comparison of gene expression measures between our RNA-seq results and these previously published results. We found that 26 of the 32 genes showed a consistent pattern in the direction of expression change, suggesting a high reproducibility of DE analysis across different measurement methods and *P. marneffei* strains.

Biological functions of 14 *P. marneffei* genes in the morphogenetic control for conidial germination, hyphal growth, asexual development, and yeast morphogenesis so far have been elucidated (as reviewed in reference 28). Among six genes known to play a role at 25°C exclusively, three (*brlA*, *gasC*, and *stuA*) were found to be expressed more abundantly at 25°C than at 37°C (Fig. 2). Among eight genes functioning at both 25°C and 37°C, *abaA* is required in asexual development and yeast growth and is known to be overexpressed at 25°C (27). Our RNA-seq data showed that *abaA* is indeed expressed significantly higher at 25°C, while the other seven genes are expressed at similar levels at the two temperatures (Fig. 2).

Functional annotation of *P. marneffei* genes. To predict the function of *P. marneffei* genes, we employed *in silico* annotation through comparative genomics across fungal species. For each *P. marneffei* gene, functional annotation information (e.g., GO terms) of orthologous genes in other model fungal species were transferred. The model fungal species included *Aspergillus fumigatus*, *Aspergillus nidulans*, *Candida albicans*, *Neurospora crassa*, and *Saccharomyces cerevisiae*. A total of 8,161 protein-coding genes of *P. marneffei* were annotated with at least one GO term. We found

that 261 GO terms are overrepresented in *P. marneffei* genes overexpressed at 37°C (FDR < 0.05) (see Table S2 in the supplemental material). These terms include “RNA-directed DNA polymerase activity,” “misfolded RNA binding,” “group I intron splicing,” and “snRNA localization” (Table 3). On the other hand, we found that 453 GO terms are overrepresented in *P. marneffei* genes underexpressed at 37°C (i.e., overexpressed at 25°C) (FDR < 0.05) (see Table S3 in the supplemental material). These terms include “α-1,4-glucan synthase activity,” “response to light stimulus,” and “secondary metabolic processes.”

Expression analysis of heat-responsive genes in *P. marneffei*. To study the expression regulation of the heat response, we compiled a list of 52 heat-responsive genes whose function has been characterized using gene knockout experiments in *A. fumigatus*, *A. nidulans*, or *S. cerevisiae*. Using this collection of genes, we identified 168 potential heat-responsive genes in *P. marneffei*. Our RNA-seq results showed that only 6 (3.6%) *P. marneffei* heat-responsive genes are overexpressed at 37°C, which is significantly fewer than expected (FDR < 0.05). We also collected another list of genes related to a broad sense of heat shock response identified by using two-dimensional (2D) fluorescence difference gel electrophoresis in *A. fumigatus* (56). Our RNA-seq results showed that *P. marneffei* orthologs of most of these genes are not expressed differentially at 25 and 37°C (see Table S4 in the supplemental material). Furthermore, none of the heat shock proteins and heat shock factors in *P. marneffei* were found to be overexpressed at 37°C (Table 4).

Next we started from a list of 12 GO terms concerning heat




FIG 2 Expression levels of morphogenesis genes of *P. marneffei*. The left panel depicts the life cycle and dimorphic switch of *P. marneffei*. The blue region (bottom left) represents processes in the mycelium form, the red region (top right) represents processes in the yeast form, and the green region (middle) represents processes involved in the dimorphic switching. The right panel shows FPKM values of gene expression measured at 25 and 37°C. Each bar represents an independent replicate. ***, significant difference in levels of expression of the gene at 25 versus 37°C (see Materials and Methods for details).

response and found none of these GO terms are enriched in genes overexpressed at 37°C (Table 5). Notably, GO:0036168 (with the term “filamentous growth of a population of unicellular organisms in response to heat”) is underrepresented in genes overexpressed at 37°C and overrepresented in genes underexpressed at 37°C. Among 370 *P. marneffei* genes whose annotation is linked with any of the 12 heat response GO terms, 16 are overexpressed at 37°C, which is significantly fewer than 52 genes underexpressed at 37°C ($P < 0.001$, Fisher’s exact test).

Taken together, these results suggest the limited value of using

the existing gene annotations available in other fungal species in annotating *P. marneffei* genes, especially for genes that might play a role in heat response. The reason for this might be that *P. marneffei* has an uncharacterized regulatory network and signaling pathways in response to heat, which differ considerably from those currently known in other fungal species.

Gene clusters in *P. marneffei* and other filamentous fungi. To identify gene clusters conserved across species, we compared the genome sequences of *P. marneffei* with those of *A. nidulans*, *A. fumigatus*, and *N. crassa*. We revealed extensive regions of con-

TABLE 3 Top representatives of GO terms enriched in *P. marneffei* genes upregulated or downregulated at 37°C

Function	GO term in genes with responses at 37°C	
	Upregulated	Downregulated
Cell components	GO:0030686—90S preribosome GO:0005730—nucleolus	GO:0000329—fungal-type vacuole membrane GO:0005887—integral to plasma membrane GO:0051286—cell tip GO:0016020—membrane GO:0032153—cell division site
Biological processes	GO:0003964—RNA-directed DNA polymerase activity GO:0008026—ATP-dependent helicase activity GO:0000293—ferric-chelate reductase activity GO:0034336—misfolded RNA binding	GO:0004497—monooxygenase activity GO:0000036—acyl carrier activity GO:0033201— α -1,4-glucan synthase activity GO:0004553—hydrolase activity, hydrolyzing O-glycosyl compounds
Molecular	GO:0015116—sulfate transmembrane transporter activity GO:0042254—ribosome biogenesis GO:0000372—group I intron splicing GO:0048254—snoRNA localization GO:0040010—positive regulation of growth rate GO:0006675—mannosyl-inositol phosphorylceramide metabolic process	GO:0016208—AMP binding GO:0045461—sterigmatocystin biosynthetic process GO:0009416—response to light stimulus GO:0019748—secondary metabolic process GO:0055114—oxidation-reduction process GO:0015707—nitrite transport

TABLE 4 Gene expression of putative heat shock proteins and heat shock factors in *P. marneffeii*

Gene ID	Homolog(s)	FPKM at:				Expression change at 37°C
		25°C		37°C		
		Replicate 1	Replicate 2	Replicate 1	Replicate 2	
pm1g_01498	<i>hsp12</i> of <i>S. cerevisiae</i>	1,594.7	1,015.9	17.5	12.1	↓
pm1g_02064	<i>hsp60</i> of <i>A. fumigatus</i>	262.2	215.8	199.9	476.7	—
pm1g_03635	<i>ssc70</i> (mitochondrial <i>hsp70</i>) of <i>A. fumigatus</i>	392.4	389.7	591.2	1,225.2	—
pm1g_05682	<i>hsp30</i> of <i>A. fumigatus</i>	988.1	825.0	2,520.4	3,467.3	—
pm1g_06129	<i>hsp104</i> of <i>S. cerevisiae</i>	159.9	120.9	150.2	152.3	—
pm1g_06571	<i>hsp78</i> of <i>A. fumigatus</i>	58.7	38.7	64.3	87.0	—
pm1g_08006	<i>hsp70</i> of <i>A. fumigatus</i>	2,224.6	2,401.0	1,175.3	1,977.2	—
pm1g_10441	<i>hsp90</i> of <i>A. fumigatus</i>	591.2	591.2	629.3	941.3	—
pm1g_10905	<i>hsp88</i> of <i>A. fumigatus</i>	285.0	339.1	301.3	406.3	—
pm1g_01128	<i>aha1</i> (<i>hsp82</i> cochaperone) of <i>S. cerevisiae</i>	138.1	156.6	221.9	283.0	—
pm1g_05408	<i>hsf1</i> of <i>A. fumigatus</i>	51.0	55.5	52.1	37.1	—
pm1g_05918	<i>sti1</i> (<i>hsp90</i> cochaperone) of <i>S. cerevisiae</i>	138.1	150.2	167.9	316.4	—
pm1g_08054	<i>sis1</i> (<i>hsp40</i> cochaperone) of <i>S. cerevisiae</i>	232.9	232.9	120.1	100.1	↓
pm1g_08850	<i>hjl1</i> (<i>hsp40</i> cochaperone) of <i>S. cerevisiae</i>	42.4	43.3	37.9	22.8	—

served synteny, as well as a considerable extent of genome reorganization between these fungal species. For example, there are 1,340 regions containing four or more genes colinear between *P. marneffeii* and *A. nidulans* and 1,273 between *P. marneffeii* and *A. fumigatus*. Totals of 3,188 and 3,716 *P. marneffeii* genes were found to be involved in the two cases, respectively. The largest syntenic cluster contains 27 gene pairs, appearing in *P. marneffeii* and *A. nidulans*.

Melanin is made in various pathogenic fungi and has been implicated in the pathogenesis of a number of fungal infections (57). Melanin may play a role in inhibiting cytokine production in the host cells (58) and apoptosis in macrophages (59). In *A. fumigatus*, the two kinds of melanin, dihydroxynaphthalene-melanin (DHN-melanin) and pyomelanin (60–63), are generated. Six genes related to the biosynthesis of DHN-melanin were clustered together (60, 61) and genetically regulated by the expression of

brlA (64). Our previous study revealed the existence of this gene cluster in *P. marneffeii* (22). RNA-seq data from this study showed that all six genes in the cluster are significantly downregulated at 37°C (Fig. 3).

Furthermore, we identified the pyomelanin biosynthesis gene cluster, which is also conserved between *P. marneffeii* and *A. fumigatus*. Pyomelanin biosynthesis pathway is tightly linked with the pathway of tyrosine degradation: two genes, *hpdD* and *hmgX*, known to be specifically involved in the pyomelanin biosynthesis pathway, are collocated in the same cluster with the other four genes in the tyrosine degradation pathway (62, 63) (Fig. 3). The identified cluster in *P. marneffeii* spans a 17-kb genomic region in a single contig. Half of the genes in the cluster were found to be expressed significantly higher at 37°C. The *P. marneffeii* cluster, 6 kb longer than that of *A. fumigatus*, contains a gene between *hmgR* and *maiA*, whose ortholog in *A. fumigatus* is relocated. The ex-

TABLE 5 Enrichment analysis of GO terms related to heat response for *P. marneffeii* genes

GO no.	GO term	No. of genes with GO term (FDR) ^a		
		All genes (<i>n</i> = 8,161)	Genes overexpressed at 37°C (<i>n</i> = 1,447)	Genes underexpressed at 37°C (<i>n</i> = 1,414)
GO:0009408	Response to heat	148	1 (1.96 × 10⁻⁴)	22 (0.292)
GO:0031072	Heat shock protein binding	44	3 (0.173)	4 (0.294)
GO:0031990	mRNA export from nucleus in response to heat stress	68	0 (6.52 × 10⁻³)	7 (0.279)
GO:0034605	Cellular response to heat	259	11 (5.33 × 10⁻⁴)	45 (8.52 × 10⁻²)
GO:0036165	Invasive growth in response to heat	3	0 (0.194)	2 (3.13 × 10⁻²)
GO:0036168	Filamentous growth of a population of unicellular organisms in response to heat	175	7 (3.21 × 10⁻³)	34 (3.57 × 10⁻²)
GO:0061408	Positive regulation of transcription from RNA polymerase II promoter in response to heat stress	13	1 (0.206)	1 (0.293)
GO:0070370	Cellular heat acclimation	83	0 (3.32 × 10⁻³)	13 (0.313)
GO:0070414	Trehalose metabolism in response to heat stress	2	0 (0.187)	0 (0.260)
GO:1900432	Negative regulation of filamentous growth of a population of unicellular organisms in response to heat	62	0 (8.51 × 10⁻³)	5 (0.301)
GO:1900433	Positive regulation of filamentous growth of a population of unicellular organisms in response to heat	60	1 (2.08 × 10⁻²)	11 (0.268)
GO:2000728	Regulation of mRNA export from nucleus in response to heat stress	5	0 (0.192)	0 (0.296)

^a Significant values where FDR is <0.05 are in boldface.




FIG 3 Gene cluster organization and expression of two melanin biosynthesis clusters in *P. marneffei*. The gene cluster of DHN-melanin biosynthesis (A) and the gene cluster of tyrosine degradation (B) in *A. fumigatus* and *P. marneffei* are shown. The expression level (FPKM) is shown in the log scale for DHN-melanin genes and the linear scale for tyrosine degradation genes. ***, significant difference in levels of expression of the gene at 25 versus 37°C (see Materials and Methods for details).

pression of the gene, with potential catalytic activity, is downregulated at 37°C.

A. fumigatus DHN-melanin biosynthesis genes are highly expressed at 37°C, suggesting that DHN-melanin is implicated in *A. fumigatus* infection (65–67). Our results showed that *P. marneffei* produces less DHN-melanin and more pyomelanin at 37°C, suggesting that pyomelanin, instead of DHN-melanin, plays a role in the pathogenicity of *P. marneffei*.

Abundant TRSs in coding regions of *P. marneffei* genes. Tandem repeat sequences (TRSs) are two or more adjacent copies of the same sequence of nucleotides resulting from tandem duplication events. TRSs located in the coding region of genes produce repeated units of amino acids in the genes’ protein products. These protein polymorphisms may create quantitative alterations in fungal phenotypes, such as adhesion, flocculation, and biofilm formation (48). We compared the relative abundance of coding TRSs between *P. marneffei* and three other filamentous fungi (Table 6). *P. marneffei* shows the highest B/G ratio, where B represents the bases in coding TRSs and G represents the genome size. That is

TABLE 6 TRSs in coding regions of the fungal genomes

Species	TRSs in coding region (kb) (B)	Genome size (Mb) (G)	B/G ratio
<i>P. marneffei</i>	23.8	30	0.79
<i>A. fumigatus</i>	12.7	28	0.45
<i>A. nidulans</i>	16.8	30	0.56
<i>N. crassa</i>	22.1	40	0.55

to say, *P. marneffei* accommodates more coding TRSs per unit of genome than other fungal species under consideration. We hypothesized that the presence of abundant coding TRSs in *P. marneffei* genes might allow the fungal cells to “disguise” themselves in order to evade the host immune system’s defenses. If true, the expression of TRS-containing genes should be upregulated at 37°C. Indeed, among 66 genes that contain coding TRSs, 15 were found to be overexpressed at 37°C (see Table S5 in the supplemental material). The number is significantly higher than expected ($P < 0.01$, Fisher’s exact test). No specific DNA sequence motif was identified in 5’ regulatory regions of these genes, suggesting that the upregulated expression of these genes may not be controlled by a master transcription factor.

Multigene families and species-specific genes in *P. marneffei*. Among protein-coding genes of *P. marneffei*, 2,283 of them belong to 665 multigene families (see Materials and Methods for details about gene family classification). The most expanded gene families include the following: the MFS multidrug transporter family (34 genes), the short-chain dehydrogenase or reductase family (31 genes), the hexose transporter family (27 genes), the pepsin-type protease family (24 genes), and the major facilitator superfamily (23 genes). Notably, the expansion of the MP1 domain protein (Mp1p, a cell wall antigen) has been proposed to be implicated in *P. marneffei* pathogenesis (68). Twelve genes containing the MP1 domain have been identified in *P. marneffei*; in contrast to that, only two MP1 domain genes have been identified in *A. fumigatus* (53, 69–71). Among these 12 genes, 7 were found to be overexpressed and one underexpressed at 37°C (Fig. 4A).




FIG 4 Expression levels of *P. marneffii* MP1 domain genes and their regulatory regions. (A) Expression levels of 12 MP1 domain genes at 25 versus 37°C (see Materials and Methods for details). (B) Putative transcription factor binding sites in 5' regulatory regions. The positions of binding motifs are indicated with the blue and green bars and labeled with the subtype name of transcription factors: (i) MADS, (ii) NFY CCAAT binding, and (iii) fungal Zn cluster. The position of the TATA box is indicated by the red bar. (C) Indirect immunofluorescent staining with antibodies to Mp1p (pm1g_08423) at 37°C.

Using a motif-discovering tool, MEME (72), we identified conserved DNA sequence motifs in the 5' regulatory regions of these MP1 domain genes (Fig. 4B). These motifs are the putative binding sites of subtypes of transcription factors: (i) MADS, (ii) NFY CCAAT binding, and (iii) fungal Zn cluster (73). These results indicate that these MP1 domain genes are likely to be regulated by shared transcription factors. Next, to confirm the cellular localization of Mp1p protein products, we performed the indirect fluorescent antibody (IFA) staining assay with antibodies for Mp1p (pm1g_08423) (see Materials and Methods). The results confirm that the Mp1p gene is expressed at 37°C and that the gene product is located on the cell wall of yeast cells (Fig. 4C). No expression and protein product were detected at 25°C. Furthermore, we used between-supercontig sequence comparison to search for conserved gene order. No evidence for large-genomic-segment duplication in *P. marneffii* was identified. Compared with *S. cerevisiae*, the *P. marneffii* genome contains relatively fewer recently duplicated gene pairs, as indicated by the small nonsynonymous-to-synonymous substitution ratio (K_a/K_s) between two copies of duplicate genes ($P < 0.01$, Fisher's exact test) (74). These differences can be explained by the whole-genome duplication in *S. cerevisiae* (75).

Finally, we identified 1,009 *P. marneffii* genes whose protein sequences do not share any sequence similarity with other protein sequences in the NCBI nonredundant database (BLASTP E value cutoff of $1e-10$). Among these orphan genes, 107 are not expressed at either 25 or 37°C. In the remaining 902 expressed ones, 117 (13.0%) are overexpressed at 37°C ($FDR > 0.05$) and only 63

(7.0%) are underexpressed at 37°C ($FDR < 0.05$). Thus, *P. marneffii*-specific genes are not likely to be overexpressed but are less likely to be underexpressed in yeast cells at 37°C.

DISCUSSION

P. marneffii is an emerging model system for investigating fungal growth processes and their contribution to pathogenicity (28). To understand the genetics of *P. marneffii*, especially in response to elevated temperature, we jointly used genome sequences and transcriptomic profiles to unravel the molecular basis of genetic regulation in *P. marneffii*. Previous studies of *P. marneffii* transcriptome were based on microarrays (76, 77). The adaptation of the RNA-seq approach, which measures gene expression in a much higher dynamic range, allowed us to produce an unbiased, high-resolution-expression measurement of *P. marneffii* genes for the first time.

Globally, we identified 1,414 and 1,447 *P. marneffii* protein-coding genes as being up- or downregulated at 37°C, respectively, compared with at 25°C. That is, more than a quarter of genes in the *P. marneffii* genome are expressed at different levels at two temperatures. This figure is likely to be a lower bound of the estimation as we adapted highly stringent statistical criteria for asserting DE genes. Next the myriad of genes presented a great challenge to gene function annotation. We leveraged comparative genomics across fungal species and detected a variety of annotation features over- or underrepresented in heat-responsive genes in *P. marneffii*. Some of these discoveries seemed to fit our expectations well.

For example, the family of Mp1 domain protein (Mp1p) is expanded significantly in *P. marneffei*, and the majority of members of this family are overexpressed in yeast cells at 37°C. A second example is the expansion of the Mp1p protein family in *P. marneffei*, suggesting that duplicate genes confer novel phenotypic changes and facilitate organismal adaptation. Moreover, we hypothesized that *P. marneffei* genes whose coding region contains TRSs are likely to be overexpressed at 37°C, because products of these genes may play a role in helping the fungus escape the attack of immune system of the host cell. Indeed, *P. marneffei* has more abundant coding TRSs in its genes, and these genes are more likely to be overexpressed at 37°C. In addition, genes involved in biosynthesis of pyomelanin were found to be upregulated at 37°C, while genes involved in biosynthesis of DHN-melanin were found to be downregulated at 37°C, which suggests conditional regulation for specific melanin biosynthesis pathways.

Many genes and their expression patterns negated our expectations. For example, we expected that genes whose annotation contains GO terms related to heat should tend to be expressed higher than other genes at 37°C. However, we found that significantly fewer “heat-related” genes in *P. marneffei* were upregulated at 37°C. We also expected that *P. marneffei*-specific genes are likely to be overexpressed at 37°C. This is because species-specific genes are likely to define the unique biology feature of the species and play a role in response to environmental changes (47). However, we found that *P. marneffei*-specific genes were not enriched in genes overexpressed at 37°C.

These unexpected results suggest that functional annotation of *P. marneffei* genes through mapping of annotation from orthologous genes remains unreliable. Given that *P. marneffei* is a unique species in its genus, it is possible that our current knowledge about genetic regulation of heat response is inapplicable to *P. marneffei*. Furthermore, we might have to update our view with regard to the link between thermal dimorphism and pathogenesis in *P. marneffei*. The new idea is that steadily low expression of heat-responsive genes may confer virulence to *P. marneffei* grown at the temperature of 37°C in mammalian host cells. In *Phytophthora sojae*, for example, the virulence of the plant-pathogenic fungus is attained by the silencing instead of activation of a gene (78). We hypothesize that the pathogenicity of *P. marneffei* is associated with the suppressed expression of heat-responsive genes at 37°C, and the phase transition of *P. marneffei* may result from gene silencing. In other words, the yeast form is an impaired form of the filamentous form.

ACKNOWLEDGMENTS

We thank Smriti Shankar for assisting with data processing. We acknowledge the TAMU Supercomputing Facility and the WSGI Compute Cluster for providing computing resources and systems administration support.

This study is supported by 2012 Whole Systems Genomics Initiative (WSGI) Catalyst Research Grant 248307-12001 from Texas A&M University (TAMU) to J.J.C. E.Y. is supported by TAMU CVM Postdoctoral Trainee Research Grant 02-144002-03504, and G.W. is supported by TAMU CVM Graduate Trainee Research Grant 02-291039-00002.

REFERENCES

- Klein BS, Tebbets B. 2007. Dimorphism and virulence in fungi. *Curr. Opin. Microbiol.* 10:314–319.
- Rappleye CA, Goldman WE. 2006. Defining virulence genes in the dimorphic fungi. *Annu. Rev. Microbiol.* 60:281–303.
- Nemecek JC, Wuthrich M, Klein BS. 2006. Global control of dimorphism and virulence in fungi. *Science* 312:583–588.
- Supparatpinyo K, Khamwan C, Baosoung V, Nelson KE, Sirisanthana T. 1994. Disseminated *Penicillium marneffei* infection in Southeast Asia. *Lancet* 344:110–113.
- Hsu LY, Wijaya L, Ng EST, Gotuzzo E. 2012. Tropical fungal infections. *Infect. Dis. Clin. North Am.* 26:497–512.
- Duong TA. 1996. Infection due to *Penicillium marneffei*, an emerging pathogen: review of 155 reported cases. *Clin. Infect. Dis.* 23:125–130.
- Yap FB, Thevarajah S, Asmah J. 2010. *Penicillium marneffei* infection in an African man. *Dermatol. Online J.* 16:2. <http://escholarship.org/uc/item/6j96f72f>.
- Le T, Wolbers M, Chi NH, Quang VM, Chinh NT, Lan NP, Lam PS, Kozal MJ, Shikuma CM, Day JN, Farrar J. 2011. Epidemiology, seasonality, and predictors of outcome of AIDS-associated *Penicillium marneffei* infection in Ho Chi Minh City, Viet Nam. *Clin. Infect. Dis.* 52:945–952.
- Devi KR, Singh LR, Rajkumari R, Usharani M, Devi Kh S, Singh TB. 2007. *Penicillium marneffei*—an indicator disease of AIDS: a case report. *Indian J. Pathol. Microbiol.* 50:674–676.
- Nguyen K, Taylor S, Wanger A, Ali A, Rapini RP. 2006. A case of *Penicillium marneffei* in a US hospital. *J. Am. Acad. Dermatol.* 54:730–732.
- Deesomchok A, Tanprawate S. 2006. A 12-case series of *Penicillium marneffei* pneumonia. *J. Med. Assoc. Thai.* 89:441–447.
- Mens H, Hojlyng N, Arendrup MC. 2004. Disseminated *Penicillium marneffei* sepsis in a HIV-positive Thai woman in Denmark. *Scand. J. Infect. Dis.* 36:507–509.
- Hien TV, Loc PP, Hoa NT, Duong NM, Quang VM, McNeil MM, Dung NT, Ashford DA. 2001. First cases of disseminated penicilliosis marneffei infection among patients with acquired immunodeficiency syndrome in Vietnam. *Clin. Infect. Dis.* 32:e78–e80.
- Depraetere K, Colebunders R, Ieven M, De Droogh E, Pelgrom Y, Hauben E, Van Marck E, Devroey C. 1998. Two imported cases of *Penicillium marneffei* infection in Belgium. *Acta Clin. Belg.* 53:255–258.
- Chang CC, Liao ST, Huang WS, Liu JD, Shih LS. 1995. Disseminated *Penicillium marneffei* infection in a patient with acquired immunodeficiency syndrome. *J. Formos. Med. Assoc.* 94:572–575.
- Ustianowski AP, Sieu TPM, Day JN. 2008. *Penicillium marneffei* infection in HIV. *Curr. Opin. Infect. Dis.* 21:31–36.
- Angus BJ, Schmid ML, Dockrell DH, Grant AD. 2011. Travel-related opportunistic infections. *HIV Med.* 12:88–101.
- McGinnis MR. 1994. *Penicillium marneffei*, dimorphic fungus of increasing importance. *Clin. Microbiol. Newsl.* 16:29–31.
- Cánovas D, Andrianopoulos A. 2007. The biology of the thermally dimorphic fungal pathogen *Penicillium marneffei*, p 213–226. In Kavanagh K (ed), *New insights in medical mycology*. Springer, Dordrecht, Netherlands.
- Kummasook A, Cooper CR, Vanittanakom N. 2010. An improved *Agrobacterium*-mediated transformation system for the functional genetic analysis of *Penicillium marneffei*. *Med. Mycol.* 48:1066–1074.
- Bugeja HE, Hynes MJ, Andrianopoulos A. 2010. The RFX protein RfxA is an essential regulator of growth and morphogenesis in *Penicillium marneffei*. *Eukaryot. Cell* 9:578–591.
- Woo PCY, Tam EWT, Chong KTK, Cai JJ, Tung ETK, Ngan AHY, Lau SKP, Yuen KY. 2010. High diversity of polyketide synthase genes and the melanin biosynthesis gene cluster in *Penicillium marneffei*. *FEBS J.* 277:3750–3758.
- Zuber S, Hynes MJ, Andrianopoulos A. 2003. The G-protein alpha-subunit GasC plays a major role in germination in the dimorphic fungus *Penicillium marneffei*. *Genetics* 164:487–499.
- Borneman AR, Hynes MJ, Andrianopoulos A. 2002. A basic helix-loop-helix protein with similarity to the fungal morphological regulators, Phd1p, Efg1p and StuA, controls conidiation but not dimorphic growth in *Penicillium marneffei*. *Mol. Microbiol.* 44:621–631.
- Pongsunk S, Andrianopoulos A, Chaiyaraj SC. 2005. Conditional lethal disruption of TATA-binding protein gene in *Penicillium marneffei*. *Fungal Genet. Biol.* 42:893–903.
- Boyce KJ, Schreider L, Andrianopoulos A. 2009. In vivo yeast cell morphogenesis is regulated by a p21-activated kinase in the human pathogen *Penicillium marneffei*. *PLoS Pathog.* 5:e1000678. doi:10.1371/journal.ppat.1000678.
- Borneman AR, Hynes MJ, Andrianopoulos A. 2000. The *abaA* homologue of *Penicillium marneffei* participates in two developmental programmes: conidiation and dimorphic growth. *Mol. Microbiol.* 38:1034–1047.

28. Boyce KJ, Andrianopoulos A. 2013. Morphogenetic circuitry regulating growth and development in the dimorphic pathogen *Penicillium marneffe*. *Eukaryot. Cell* 12:154–160.
29. Woo PCY, Lau SKP, Liu B, Cai JJ, Chong KTK, Tse H, Kao RYT, Chan CM, Chow WN, Yuen KY. 2011. Draft genome sequence of *Penicillium marneffe* strain PM1. *Eukaryot. Cell* 10:1740–1741.
30. Yuen K, Pascal G, Wong SSS, Glaser P, Woo PCY, Kunst F, Cai JJ, Cheung EYL, Medigue C, Danchin A. 2003. Exploring the *Penicillium marneffe* genome. *Arch. Microbiol.* 179:339–353.
31. Frangeul L, Nelson KE, Buchrieser C, Danchin A, Glaser P, Kunst F. 1999. Cloning and assembly strategies in microbial genome projects. *Microbiology* 145:2625–2634.
32. Ewing B, Hillier L, Wendl MC, Green P. 1998. Base-calling of automated sequencer traces using phred. I. Accuracy assessment. *Genome Res.* 8:175–185.
33. Ewing B, Green P. 1998. Base-calling of automated sequencer traces using phred. II. Error probabilities. *Genome Res.* 8:186–194.
34. Pop M, Kosack DS, Salzberg SL. 2004. Hierarchical scaffolding with Bambus. *Genome Res.* 14:149–159.
35. Marcais G, Kingsford C. 2011. A fast, lock-free approach for efficient parallel counting of occurrences of k-mers. *Bioinformatics* 27:764–770.
36. Trapnell C, Roberts A, Goff L, Pertea G, Kim D, Kelley DR, Pimentel H, Salzberg SL, Rinn JL, Pachter L. 2012. Differential gene and transcript expression analysis of RNA-seq experiments with TopHat and Cufflinks. *Nat. Protoc.* 7:562–578.
37. Trapnell C, Williams BA, Pertea G, Mortazavi A, Kwan G, van Baren MJ, Salzberg SL, Wold BJ, Pachter L. 2010. Transcript assembly and quantification by RNA-Seq reveals unannotated transcripts and isoform switching during cell differentiation. *Nat. Biotechnol.* 28:511–515.
38. Xu G, Deng N, Zhao Z, Judeh T, Flemington E, Zhu D. 2011. SAMMate: a GUI tool for processing short read alignments in SAM/BAM format. *Source Code Biol. Med.* 6:2. doi:10.1186/1751-0473-6-2.
39. Robinson MD, McCarthy DJ, Smyth GK. 2010. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* 26:139–140.
40. Anders S, Huber W. 2010. Differential expression analysis for sequence count data. *Genome Biol.* 11:R106. doi:10.1186/gb-2010-11-10-r106.
41. Hardcastle TJ, Kelly KA. 2010. baySeq: empirical Bayesian methods for identifying differential expression in sequence count data. *BMC Bioinformatics* 11:422. doi:10.1186/1471-2105-11-422.
42. O'Brien KP, Remm M, Sonnhammer ELL. 2005. Inparanoid: a comprehensive database of eukaryotic orthologs. *Nucleic Acids Res.* 33:D476–D480.
43. Vandepoele K, Saeys Y, Simillion C, Raes J, Van de Peer Y. 2002. The automatic detection of homologous regions (ADHoRe) and its application to microcolinearity between *Arabidopsis* and rice. *Genome Res.* 12:1792–1801.
44. Hao W, Pan YX, Ding YQ, Xiao S, Yin K, Wang YD, Qiu LW, Zhang QL, Woo PCY, Lau SKP, Yuen KY, Che XY. 2008. Well-characterized monoclonal antibodies against cell wall antigen of *Aspergillus* species improve immunoassay specificity and sensitivity. *Clin. Vaccine Immunol.* 15:194–202.
45. Qiu LW, Di BA, Wen K, Wang XSA, Liang WH, Wang YD, Pan YX, Wang M, Ding YQ, Che XY. 2009. Development of an antigen capture immunoassay based on monoclonal antibodies specific for dengue virus serotype 2 nonstructural protein 1 for early and rapid identification of dengue virus serotype 2 infections. *Clin. Vaccine Immunol.* 16:88–95.
46. Supek F, Bosnjak M, Skunca N, Smuc T. 2011. REVIGO summarizes and visualizes long lists of Gene Ontology terms. *PLoS One* 6:e21800. doi:10.1371/journal.pone.0021800.
47. Storey JD, Tibshirani R. 2003. Statistical significance for genomewide studies. *Proc. Natl. Acad. Sci. U. S. A.* 100:9440–9445.
48. Verstrepen KJ, Jansen A, Lewitter F, Fink GR. 2005. Intragenic tandem repeats generate functional variability. *Nat. Genet.* 37:986–990.
49. Rice P, Longden I, Bleasby A. 2000. EMBOSS: The European Molecular Biology Open Software Suite. *Trends Genet.* 16:276–277.
50. Arnaud MB, Cerqueira GC, Inglis DO, Skrzypek MS, Binkley J, Chibucos MC, Crabtree J, Howarth C, Orvis J, Shah P, Wymore F, Binkley G, Miyasato SR, Simison M, Sherlock G, Wortman JR. 2012. The *Aspergillus* Genome Database (AspGD): recent developments in comprehensive multispecies curation, comparative genomics and community resources. *Nucleic Acids Res.* 40:D653–D659.
51. Inglis DO, Arnaud MB, Binkley J, Shah P, Skrzypek MS, Wymore F, Binkley G, Miyasato SR, Simison M, Sherlock G. 2012. The *Candida* Genome Database incorporates multiple *Candida* species: multispecies search and analysis tools with curated gene and protein information for *Candida albicans* and *Candida glabrata*. *Nucleic Acids Res.* 40:D667–D674.
52. Cherry JM, Hong EL, Amundsen C, Balakrishnan R, Binkley G, Chan ET, Christie KR, Costanzo MC, Dwight SS, Engel SR, Fisk DG, Hirschman JE, Hitz BC, Karra K, Krieger CJ, Miyasato SR, Nash RS, Park J, Skrzypek MS, Simison M, Weng S, Wong ED. 2012. *Saccharomyces* Genome Database: the genomics resource of budding yeast. *Nucleic Acids Res.* 40:D700–D705.
53. Nierman WC, Pain A, Anderson MJ, Wortman JR, Kim HS, Arroyo J, Berriman M, Abe K, Archer DB, Bermejo C, Bennett J, Bowyer P, Chen D, Collins M, Coulsen R, Dvorsky R, Dyer PS, Farman M, Fedorova N, Fedorova N, Feldblyum TV, Fischer R, Fosker N, Fraser A, Garcia JL, Garcia MJ, Goble A, Goldman GH, Gomi K, Griffith-Jones S, Gwilliam R, Haas B, Haas H, Harris D, Horiuchi H, Huang J, Humphray S, Jimenez J, Keller N, Khouri H, Kitamoto K, Kobayashi T, Konzack S, Kulkarni R, Kumagai T, Lafton A, Latge JP, Li WX, Lord A, Majoros WH, May GS, Miller BL, Mohamoud Y, Molina M, Monod M, Mouyna I, Mulligan S, Murphy L, O'Neil S, Paulsen I, Penalva MA, Pertea M, Price C, Pritchard BL, Quail MA, Rabinowitz E, Rawlins N, Rajandream MA, Reichard U, Renauld H, Robson GD, de Cordoba SR, Rodriguez-Pena JM, Ronning CM, Rutter S, Salzberg SL, Sanchez M, Sanchez-Ferrero JC, Saunders D, Seeger K, Squares R, Squares S, Takeuchi M, Tekaia F, Turner G, de Aldana CRV, Weidman J, White O, Woodward J, Yu JH, Fraser C, Galagan JE, Asai K, Machida M, Hall N, Barrell B, Denning DW. 2005. Genomic sequence of the pathogenic and allergenic filamentous fungus *Aspergillus fumigatus*. *Nature* 438:1151–1156.
54. Galagan JE, Calvo SE, Cuomo C, Ma LJ, Wortman JR, Batzoglu S, Lee SI, Basturkmen M, Spevak CC, Clutterbuck J, Kapitonov V, Jurka J, Scacciochio C, Farman M, Butler J, Purcell S, Harris S, Braus GH, Draht O, Busch S, D'Enfert C, Bouchier C, Goldman GH, Bell-Pedersen D, Griffiths-Jones S, Doonan JH, Yu J, Vienken K, Pain A, Freitag M, Selker EU, Archer DB, Penalva MA, Oakley BR, Momany M, Tanaka T, Kumagai T, Asai K, Machida M, Nierman WC, Denning DW, Caddick M, Hynes M, Paoletti M, Fischer R, Miller B, Dyer P, Sachs MS, Osmani SA, Birren BW. 2005. Sequencing of *Aspergillus nidulans* and comparative analysis with *A. fumigatus* and *A. oryzae*. *Nature* 438:1105–1115.
55. van den Berg MA, Albang R, Albermann K, Badger JH, Daran JM, Driessen AJM, Garcia-Estrada C, Fedorova ND, Harris DM, Heijnen WHM, Joardar V, Kiel JAKW, Kovalchuk A, Martin JF, Nierman WC, Nijland JG, Pronk JT, Roubos JA, van der Klei IJ, van Peij NNME, Veenhuis M, von Dohren H, Wagner C, Wortman J, Bovenberg RAL. 2008. Genome sequencing and analysis of the filamentous fungus *Penicillium chrysogenum*. *Nat. Biotechnol.* 26:1161–1168.
56. Albrecht D, Guthke R, Brakhage AA, Kniemeyer O. 2010. Integrative analysis of the heat shock response in *Aspergillus fumigatus*. *BMC Genomics* 11:32. doi:10.1186/1471-2164-11-32.
57. Jacobson ES. 2000. Pathogenic roles for fungal melanins. *Clin. Microbiol. Rev.* 13:708–717.
58. Chai LYA, Netea MG, Sugui J, Vonk AG, van de Sande WWJ, Warris A, Kwon-Chung KJ, Kullberg BJ. 2010. *Aspergillus fumigatus* conidial melanin modulates host cytokine response. *Immunobiology* 215:915–920.
59. Volling K, Thywissen A, Brakhage AA, Saluz HP. 2011. Phagocytosis of melanized *Aspergillus* conidia by macrophages exerts cytoprotective effects by sustained PI3K/Akt signalling. *Cell. Microbiol.* 13:1130–1148.
60. Tsai HF, Wheeler MH, Chang YC, Kwon-Chung KJ. 1999. A developmentally regulated gene cluster involved in conidial pigment biosynthesis in *Aspergillus fumigatus*. *J. Bacteriol.* 181:6469–6477.
61. Fujii I, Yasuoka Y, Tsai HF, Chang YC, Kwon-Chung KJ, Ebizuka Y. 2004. Hydrolytic polyketide shortening by Ayglp, a novel enzyme involved in fungal melanin biosynthesis. *J. Biol. Chem.* 279:44613–44620.
62. Schmalder-Ripcke J, Sugareva V, Gebhardt P, Winkler R, Kniemeyer O, Heinekamp T, Brakhage AA. 2009. Production of pyomelanin, a second type of melanin, via the tyrosine degradation pathway in *Aspergillus fumigatus*. *Appl. Environ. Microbiol.* 75:493–503.
63. Keller S, Macheleidt J, Scherlach K, Schmalder-Ripcke J, Jacobsen ID, Heinekamp T, Brakhage AA. 2011. Pyomelanin formation in *Aspergillus fumigatus* requires HmgX and the transcriptional activator HmgR but is

- dispensable for virulence. *PLoS One* 6:e26604. doi:10.1371/journal.pone.0026604.
64. Twumasi-Boateng K, Yu Y, Chen D, Gravelat FN, Nierman WC, Sheppard DC. 2009. Transcriptional profiling identifies a role for BrlA in the response to nitrogen depletion and for StuA in the regulation of secondary metabolite clusters in *Aspergillus fumigatus*. *Eukaryot. Cell* 8:104–115.
 65. Langfelder K, Streibel M, Jahn B, Haase G, Brakhage AA. 2003. Biosynthesis of fungal melanins and their importance for human pathogenic fungi. *Fungal Genet. Biol.* 38:143–158.
 66. Tsai HF, Chang YC, Washburn RG, Wheeler MH, Kwon-Chung KJ. 1998. The developmentally regulated *alb1* gene of *Aspergillus fumigatus*: its role in modulation of conidial morphology and virulence. *J. Bacteriol.* 180:3031–3038.
 67. Jahn B, Koch A, Schmidt A, Wanner G, Gehringer H, Bhakdi S, Brakhage AA. 1997. Isolation and characterization of a pigmentless-conidium mutant of *Aspergillus fumigatus* with altered conidial surface and reduced virulence. *Infect. Immun.* 65:5110–5117.
 68. Wong LP, Woo PCY, Wu AYY, Yuen KY. 2002. DNA immunization using a secreted cell wall antigen Mplp is protective against *Penicillium marneffeii* infection. *Vaccine* 20:2878–2886.
 69. Yuen KY, Chan CM, Chan KM, Woo PCY, Che XY, Leung ASP, Cao L. 2001. Characterization of AFMP1: a novel target for serodiagnosis of aspergillosis. *J. Clin. Microbiol.* 39:3830–3837.
 70. Chong KTK, Woo PCY, Lau SKP, Huang Y, Yuen KY. 2004. AFMP2 encodes a novel immunogenic protein of the antigenic mannoprotein superfamily in *Aspergillus fumigatus*. *J. Clin. Microbiol.* 42:2287–2291.
 71. Fedorova ND, Khaldi N, Joardar VS, Maiti R, Amedeo P, Anderson MJ, Crabtree J, Silva JC, Badger JH, Albarraq A, Angiuoli S, Bussey H, Bowyer P, Cotty PJ, Dyer PS, Egan A, Galens K, Fraser-Liggett CM, Haas BJ, Inman JM, Kent R, Lemieux S, Malavazi I, Orvis J, Roemer T, Ronning CM, Sundaram JP, Sutton G, Turner G, Venter JC, White OR, Whitty BR, Youngman P, Wolfe KH, Goldman GH, Wortman JR, Jiang B, Denning DW, Nierman WC. 2008. Genomic islands in the pathogenic filamentous fungus *Aspergillus fumigatus*. *PLoS Genet.* 4:e1000046. doi:10.1371/journal.pgen.1000046.
 72. Bailey TL, Boden M, Buske FA, Frith M, Grant CE, Clementi L, Ren JY, Li WW, Noble WS. 2009. MEME SUITE: tools for motif discovery and searching. *Nucleic Acids Res.* 37:W202–W208.
 73. Samson RA, Yilmaz N, Houbraken J, Spierenburg H, Seifert KA, Peterson SW, Varga J, Frisvad JC. 2011. Phylogeny and nomenclature of the genus *Talaromyces* and taxa accommodated in *Penicillium* subgenus *Biverticillium*. *Stud. Mycol.* 70:159–183.
 74. Yang E, Hulse AM, Cai JJ. 2012. Evolutionary analysis of sequence divergence and diversity of duplicate genes in *Aspergillus fumigatus*. *Evol. Bioinformatics Online* 8:623–644.
 75. Seoighe C, Wolfe KH. 1998. Extent of genomic rearrangement after genome duplication in yeast. *Proc. Natl. Acad. Sci. U. S. A.* 95:4447–4452.
 76. Liu HF, Xi LY, Zhang JM, Li XQ, Liu XY, Lu CM, Sun JF. 2007. Identifying differentially expressed genes in the dimorphic fungus *Penicillium marneffeii* by suppression subtractive hybridization. *FEMS Microbiol. Lett.* 270:97–103.
 77. Lin X, Ran Y, Gou L, He F, Zhang R, Wang P, Dai Y. 2012. Comprehensive transcription analysis of human pathogenic fungus *Penicillium marneffeii* in mycelial and yeast cells. *Med. Mycol.* 50:835–842.
 78. Qutob D, Patrick Chapman B, Gijzen M. 2013. Transgenerational gene silencing causes gain of virulence in a plant pathogen. *Nat. Commun.* 4:1349. doi:10.1038/ncomms2354.
 79. Kummasook A, Tzarphmaag A, Thirach S, Pongpom M, Cooper CR, Vanittanakom N. 2011. *Penicillium marneffeii* actin expression during phase transition, oxidative stress, and macrophage infection. *Mol. Biol. Rep.* 38:2813–2819.
 80. Canovas D, Andrianopoulos A. 2006. Developmental regulation of the glyoxylate cycle in the human pathogen *Penicillium marneffeii*. *Mol. Microbiol.* 62:1725–1738.
 81. Bugeja HE, Hynes MJ, Andrianopoulos A. 2012. AreA controls nitrogen source utilisation during both growth programs of the dimorphic fungus *Penicillium marneffeii*. *Fungal Biol.* 116:145–154.
 82. Andrianopoulos A. 2002. Control of morphogenesis in the human fungal pathogen *Penicillium marneffeii*. *Int. J. Med. Microbiol.* 292:331–347.
 83. Boyce KJ, Hynes MJ, Andrianopoulos A. 2001. The CDC42 homolog of the dimorphic fungus *Penicillium marneffeii* is required for correct cell polarization during growth but not development. *J. Bacteriol.* 183:3447–3457.
 84. Boyce KJ, Hynes MJ, Andrianopoulos A. 2003. Control of morphogenesis and actin localization by the *Penicillium marneffeii* RAC homolog. *J. Cell Sci.* 116:1249–1260.
 85. Pongpom P, Cooper CR, Vanittanakom N. 2005. Isolation and characterization of a catalase-peroxidase gene from the pathogenic fungus, *Penicillium marneffeii*. *Med. Mycol.* 43:403–411.
 86. Boyce KJ, Schreider L, Kirszenblat L, Andrianopoulos A. 2011. The two-component histidine kinases DrkA and SlnA are required for in vivo growth in the human pathogen *Penicillium marneffeii*. *Mol. Microbiol.* 82:1164–1184.
 87. Zuber S, Hynes MJ, Andrianopoulos A. 2002. G-protein signaling mediates asexual development at 25°C but has no effect on yeast-like growth at 37°C in the dimorphic fungus *Penicillium marneffeii*. *Eukaryot. Cell* 1:440–447.
 88. Thirach S, Cooper CR, Vanittanakom N. 2008. Molecular analysis of the *Penicillium marneffeii* glyceraldehyde-3-phosphate dehydrogenase-encoding gene (*gpdA*) and differential expression of *gpdA* and the isocitrate lyase-encoding gene (*acuD*) upon internalization by murine macrophages. *J. Med. Microbiol.* 57:1322–1328.
 89. Vanittanakom N, Pongpom M, Praparattanapan J, Cooper CR, Siritanathana T. 2009. Isolation and expression of heat shock protein 30 gene from *Penicillium marneffeii*. *Med. Mycol.* 47:521–526.
 90. Kummasook A, Pongpom P, Vanittanakom N. 2007. Cloning, characterization and differential expression of an *hsp70* gene from the pathogenic dimorphic fungus, *Penicillium marneffeii*. *DNA Seq.* 18:385–394.
 91. Cao L, Chan CM, Lee C, Wong SSY, Yuen KY. 1998. *MP1* encodes an abundant and highly antigenic cell wall mannoprotein in the pathogenic fungus *Penicillium marneffeii*. *Infect. Immun.* 66:966–973.
 92. Pongpom M, Vanittanakom N. 2011. Characterization of an MPLP6, a gene coding for a yeast phase specific, antigenic mannoprotein in *Penicillium marneffeii*. *Med. Mycol.* 49:32–39.
 93. Canovas D, Boyce KJ, Andrianopoulos A. 2011. The fungal type II myosin in *Penicillium marneffeii*, MyoB, is essential for chitin deposition at nascent septation sites but not actin localization. *Eukaryot. Cell* 10:302–312.
 94. Boyce KJ, Andrianopoulos A. 2007. A p21-activated kinase is required for conidial germination in *Penicillium marneffeii*. *PLoS Pathog.* 3:e162. doi:10.1371/journal.ppat.0030162.
 95. Wang F, Tao JH, Qian Z, You S, Dong H, Shen H, Chen XX, Tang SR, Ren SX. 2009. A histidine kinase PmHHK1 regulates polar growth, sporulation and cell wall composition in the dimorphic fungus *Penicillium marneffeii*. *Mycol. Res.* 113:915–923.
 96. Feng PY, Xie Z, Sun JF, Zhang JM, Li XQ, Lu CM, Xi LY. 2010. Molecular cloning, characterization and expression of PmRsr1, a Ras-related gene from yeast form of *Penicillium marneffeii*. *Mol. Biol. Rep.* 37:3533–3540.
 97. Boyce KJ, Hynes MJ, Andrianopoulos A. 2005. The Ras and Rho GTPases genetically interact to co-ordinately regulate cell polarity during development in *Penicillium marneffeii*. *Mol. Microbiol.* 55:1487–1501.
 98. Nimmanee P, Vanittanakom N. 2012. Characterization of *Penicillium marneffeii* sakA gene and the mutant generating by split-marker recombination. Abstracts of the 18th Congress of the International Society for Human and Animal Mycology/Poster presentation, 11–15 June 2012. *Mycoses* 55(Suppl 4):132.
 99. Cao CW, Liu W, Li RY. 2009. *Penicillium marneffeii* SKN7, a novel gene, could complement the hypersensitivity of *S. cerevisiae* *skn7* disruptant strain to oxidative stress. *Mycopathologia* 168:23–30.
 100. Thirach S, Cooper CR, Vanittanakom P, Vanittanakom N. 2007. The copper, zinc superoxide dismutase gene of *Penicillium marneffeii*: cloning, characterization, and differential expression during phase transition and macrophage infection. *Med. Mycol.* 45:409–417.
 101. Borneman AR, Hynes MJ, Andrianopoulos A. 2001. An STE12 homolog from the asexual, dimorphic fungus *Penicillium marneffeii* complements the defect in sexual development of an *Aspergillus nidulans* *steA* mutant. *Genetics* 157:1003–1014.
 102. Todd RB, Greenhalgh JR, Hynes MJ, Andrianopoulos A. 2003. TupA, the *Penicillium marneffeii* Tup1p homologue, represses both yeast and spore development. *Mol. Microbiol.* 48:85–94.