



<b>Title</b>	<b>Aberrant large tumor suppressor 2 (LATS2) gene expression correlates with EGFR mutation and survival in lung adenocarcinomas</b>
<b>Author(s)</b>	<b>LUO, SY; Sit, AKY; Sihoe, DLA; Suen, WS; Au, TWK; Tang, X; Ma, ESK; Chan, WK; Wistuba, II; Minna, JD; Tsao, GSW; Lam, CLD</b>
<b>Citation</b>	<b>Lung Cancer, 2014, v. 85 n. 2, p. 282-292</b>
<b>Issued Date</b>	<b>2014</b>
<b>URL</b>	<b><a href="http://hdl.handle.net/10722/202501">http://hdl.handle.net/10722/202501</a></b>
<b>Rights</b>	<b>© &lt;year&gt;. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <a href="http://creativecommons.org/licenses/by-nc-nd/4.0/">http://creativecommons.org/licenses/by-nc-nd/4.0/</a></b>

## Accepted Manuscript

Title: Aberrant Large Tumor Suppressor 2 (*LATS2*) gene expression correlates with *EGFR* mutation and survival in lung adenocarcinomas

Author: Susan Y. Luo A.K. Sit Alan D.L. Sihoe Wai-Sing Suen Wing-Kuk Au Ximing Tang Edmond S.K. Ma Wai-Kong Chan Ignacio I. Wistuba John D. Minna George S.W. Tsao David C.L. Lam



PII: S0169-5002(14)00257-8  
DOI: <http://dx.doi.org/doi:10.1016/j.lungcan.2014.05.025>  
Reference: LUNG 4623

To appear in: *Lung Cancer*

Received date: 24-1-2014  
Revised date: 14-5-2014  
Accepted date: 30-5-2014

Please cite this article as: Luo SY, Sit AK, Sihoe ADL, Suen W-S, Au W-K, Tang X, Ma ESK, Chan W-K, Wistuba II, Minna JD, Tsao GSW, Lam DCL, Aberrant Large Tumor Suppressor 2 (*LATS2*) gene expression correlates with *EGFR* mutation and survival in lung adenocarcinomas, *Lung Cancer* (2014), <http://dx.doi.org/10.1016/j.lungcan.2014.05.025>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Aberrant Large Tumor Suppressor 2 (*LATS2*) gene expression correlates with *EGFR* mutation and survival in lung adenocarcinomas**

*Susan Y Luo*<sup>1</sup>, *AK Sit*<sup>2</sup>, *Alan DL Sihoe*<sup>2</sup>, *Wai-Sing Suen*<sup>2</sup>, *Wing-Kuk Au*<sup>2</sup>, *Ximing Tang*<sup>3</sup>, *Edmond SK Ma*<sup>4</sup>, *Wai-Kong Chan*<sup>4</sup>, *Ignacio I Wistuba*<sup>3</sup>, *John D Minna*<sup>5</sup>, *George SW Tsao*<sup>6</sup>, *David CL Lam*<sup>1</sup>

*Departments of Medicine*<sup>1</sup>, *Cardiothoracic Surgery*<sup>2</sup> and *Anatomy*<sup>6</sup>, *Queen Mary Hospital, University of Hong Kong*; *Department of Translational Molecular Pathology, University of Texas MD Anderson Cancer Center*<sup>3</sup>; *Hamon Center for Therapeutic Oncology Research, University of Texas Southwestern Medical Center at Dallas*<sup>5</sup>, *Texas, USA*; *Department of Pathology, Hong Kong Sanatorium and Hospital, Hong Kong*<sup>4</sup>.

Correspondence to:

Dr David CL Lam

Department of Medicine, Queen Mary Hospital, University of Hong Kong

Tel: (852) 2255 5814

Fax: (852) 2816 2863

Email: [dcllam@hku.hk](mailto:dcllam@hku.hk)

Keywords: Lung cancer, large tumor suppressor gene expression, gene silencing

Text Word count: 3504

Abstract Word Count: 250

Number of Figures and Tables: 6

Disclosure: This study was supported in part by the Hong Kong SK Yee Medical Grant and the University of Hong Kong Seed Funding.

## Abstract

**Background:** Large Tumor Suppressor 2(*LATS2*) gene is a putative tumor suppressor gene with potential roles in regulation of cell proliferation and apoptosis in lung cancer. The aim of this study is to explore the association of aberrant *LATS2* expression with *EGFR* mutation and survival in lung adenocarcinoma(AD), and the effects of *LATS2* silencing in both lung AD cell lines.

**Methods:** *LATS2* mRNA and protein expression in resected lung AD were correlated with demographic characteristics, *EGFR* mutation and survival. *LATS2*-specific siRNA was transfected into four *EGFR* wild-type(WT) and three *EGFR* mutant AD cell lines and the changes in *LATS2* expression and relevant signaling molecules before and after *LATS2* knockdown were assayed.

**Results:** Fifty resected lung AD were included(M:F=23:27, Smokers:non-smokers=19:31, *EGFR* mutant:wild-type=21:29) with *LATS2* mRNA levels showed no significant difference between gender, age, smoking and pathological stages while *LATS2* immunohistochemical staining on an independent set of 79 lung AD showed similar trend. *LATS2* mRNA level was found to be a significant independent predictor for survival status(Disease-free survival RR=0.217; p=0.003; Overall survival RR=0.238; p=0.036). siRNA-mediated suppression of *LATS2* expression resulted in augmentation of *ERK* phosphorylation in *EGFR* wild-type AD cell lines with high basal *LATS2* expression, discriminatory modulation of *Akt* signaling between *EGFR*

wild-type and mutant cells, and induction of *p53* accumulation in AD cell lines with low baseline *p53* levels.

**Conclusions:** *LATS2* expression level is predictive of survival in patients with resected lung AD. *LATS2* may modulate and contribute to tumor growth via different signaling pathways in *EGFR* mutant and wild-type tumors. (Abstract word count: 250)

Accepted Manuscript

## Introduction

*LATS2* (Large Tumor Suppressor 2), one of the two human homologues of *Drosophila* warts, is a putative tumor suppressor gene which encodes for a serine/threonine kinase<sup>1</sup>. As a component in the Hippo signaling pathway, *LATS2* kinase plays a critical role in controlling organ size development and in coordinating cell proliferation and apoptosis<sup>2,3</sup>. As a putative tumor suppressor gene, *LATS2* displays multiple mechanisms of actions in different cancer cell types, including cell cycle regulation by controlling G1/S and G2/M transition<sup>4,5</sup>, induction of apoptosis by down-regulating anti-apoptotic proteins Bcl-2 and Bcl-X<sub>L</sub><sup>6</sup>, as well as maintenance of mitotic fidelity and genetic stability by interacting with other regulators of cell division, such as *p53*<sup>7</sup> and Aurora kinases<sup>8-10</sup>.

Dysregulation of *LATS2* functions has been found in different types of tumors. The chromosomal location of *LATS2* is mapped to 13q11-q12 where there is frequent loss of heterozygosity<sup>11</sup>. In breast cancer, *LATS2* mRNA expression was down-regulated by promoter hypermethylation and this alteration was associated with large tumor size, high rate of metastasis and estrogen receptor and progesterone receptor negativity<sup>12</sup>. *LATS2* may also play a role in the development of prostate cancer based on findings that reduced *LATS2* expression occurred in prostate tumors and *LATS2* negatively modulated androgen receptor-regulated gene transcription<sup>13</sup>. In malignant mesothelioma (MM), *LATS2* was found to be inactivated in MM cells and this inactivation will lead to deregulated cell growth by allowing constitutive activation of

the downstream transcription factor of *LATS2* in the Hippo pathway, YAP<sup>14,15</sup>. The expression of *LATS2* exhibits suppressive effects on mesothelioma cells.

In non-small cell lung cancer (NSCLC), occasional information has been reported on the effects of aberrant expression of *LATS2* except its being one potential targets for microRNA-135b action to promote lung cancer metastasis<sup>16</sup>. *LATS2* mutations are rare<sup>17,18</sup> but other mechanisms have been reported to cause down-regulation of its expression such as promoter hypermethylation<sup>19</sup> and micro-RNA regulation<sup>20,21</sup>. The relation between reduced *LATS2* expression and lung cancer progression, and the underlying mechanisms remain unknown. In addition, we have previously found from expression profiling experiments that *LATS2* showed differential expression between pulmonary adenocarcinomas with wild-type *EGFR* and ones bearing *EGFR* mutations at exons 18 to 21<sup>22</sup>. In this study, we further validated the differential expression of *LATS2* in lung adenocarcinoma (AD) tissues in relation to *EGFR* mutation status, as well as other clinicopathological factors including as smoking history and survival status. Furthermore, we silenced *LATS2* expression by siRNA inference in several lung adenocarcinoma cell lines and examined *LATS2* knockdown effects on *EGFR* downstream signaling pathways, *Ras/Erk*, *PI3K/Akt*, as well as *p53* network.

## Materials and Methods

### Human lung adenocarcinoma tissue

Newly diagnosed of lung adenocarcinoma patients were recruited prospectively before planned surgical resection. The protocol for lung cancer surgical specimen collection was approved by the Institutional Review Board Research Ethics Committee of the University of Hong Kong and Hong Kong Hospital Authority Hong Kong West Cluster. Resected lung adenocarcinoma (AD) tissues from Chinese patients were included in this study. Upon resection, these tissue samples were immediately submerged in *RNAlater* RNA Stabilization Reagent (Qiagen), frozen in liquid nitrogen, and stored at  $-80^{\circ}\text{C}$  until RNA extraction. An independent set of 79 archival paraffin blocks of lung adenocarcinomas with known patient demographics including age, gender and *EGFR* mutation status (but not smoking habits) collected successively at the Histopathology Laboratory of the Hong Kong and Sanatorium Hospital was used for immunohistochemical studies. All the tumor tissues used were collected from patients who underwent resection of lung tumors without prior treatment with *EGFR* tyrosine kinase inhibitors or any other form of anti-cancer treatment.

### Human adenocarcinoma cell lines

Seven AD cell lines were cultured in RPMI 1640(Gibco, USA) supplemented with 1% Penicillin-Streptomycin(Gibco, USA) and 2.5% or 10% fetal bovine



serum(Gibco, USA). AD cell lines used in this study were HKULC-2, HKULC-4<sup>23</sup>, and H1648, H1650, H1975 and H2023 were from John D Minna M.D., University of Texas Southwestern Medical Center at Dallas; while PC9 was from PC Yang M.D., National Taiwan University. Three AD cell lines with *EGFR* mutations, which were H1975 with double mutations L858R and T790M, PC9 and H1650 bearing *EGFR* deletions at exon 19.

### **Direct sequencing for *EGFR* mutations**

We utilized the historical standard for *EGFR* mutation testing, directing sequencing<sup>24,25</sup>. In order to enrich tumor cell content, we performed micro-dissection on formalin-fixed paraffin-embedded sections before DNA extraction. Then, exons 18-21 of *EGFR* were PCR-amplified by applying respective primers. Purified PCR product was analyzed by ABI 3730xl DNA analyzer and sequence data was reviewed by Sequence Scanner Software.

### **Real time PCR**

Total RNA was extracted from frozen resected lung adenocarcinoma tissues by using the RNeasy kit (Qiagen, UK). After reverse transcription of total RNA via QuantiTect<sup>®</sup> kit (Qiagen, UK), 150ng cDNA templates were used to detect *LATS2* mRNA expression by using quantitative real time PCR (RT-qPCR) with the SYBR Green I method (Qiagen, UK). The primer sequences for *LATS2* mRNA were:

Forward, 5'-TGGCACCTACTCCCACAG-3', and Reverse, 5'-

CCAAGGGCTTTCTTCATCT-3'<sup>26</sup>. The ribosomal 18S gene was chosen as an internal control, and the primer sequences were: Forward, 5'-AGGAATTGACGGAAGGGCAC-3', and Reverse, 5'-GGACATCTAAGGGCATCACA-3'. Thermal cycle conditions were: 95°C for 5 minutes followed by 40 cycles of amplification at 95°C for 15 second per cycle, 58°C (*LATS2*) or 60°C (18S) for 45 seconds and 72°C for 45 seconds. The dissociation curve analysis was carried out to exclude non-specific primer dimers. The relative expression level was determined as  $2^{-\Delta\Delta Ct}$  with relative to a reference sample (BS65.2N-KT, an immortalized normal bronchial epithelial cell line). The logarithmic values to base of two of these relative expression levels were used in the following statistical analysis.

### **Immunohistochemistry**

Immunohistochemistry(IHC) of the *LATS2* protein was performed on formalin-fixed, paraffin-embedded sections of an independent set of 79 paraffin blocks of lung AD different from those used for real-time PCR assay above.

Antigen retrieval was conducted in Tris-EDTA buffer(10mM Tris, 1mM EDTA, pH=9) at 95°C for 30 minutes. The goat polyclonal antibody against human *LATS2*(dilution 1:200; Santa Cruz, USA) or negative control mouse IgG1(dilution: 1:200; Dako, Denmark) was incubated with the sections overnight at 4°C. After treating with the rabbit anti-goat/mouse secondary antibodies(dilution 1:400; Dako,

Denmark) respectively at 37°C for 30 minutes, the specimens were stained with DAB substrate chromogen system(Dako, Denmark) for 5 minutes. Scoring of immunohistochemical staining was performed by independent pathologists(Ximing Tang and Ignacio I Wistuba) without knowing the clinical annotations of the specimens. *LATS2* expression H-scores based on the extension (0-100) and intensity (0, none; 1+, weak; 2+, moderate; and 3+, strong) of IHC staining was performed by two pathologists (X.T. and I.W.). If there is discrepancy between the two pathologists concerned, an independent third pathologist in the same laboratory who were not aware of the clinical details and previous H-scores by the first two pathologists. A final H score (0–300) was obtained by multiplying the intensity (1-3) and reactivity extension (0-100) for each case. Mean score was taken for the whole group of samples and samples with H scores above mean would be classified as of higher expression levels whereas those with H scores below mean would be classified as of lower expression levels.

### **Western Blotting**

Whole-cell lysates were prepared in 1X RIPA lysis buffer (Rockland) with addition of 1% protease inhibitor cocktail (Sigma). Total protein (80 ug for *Akt* and 50 ug for other proteins) was fractionated using SDS-PAGE and then transferred to nitrocellulose membranes (ExPro). After blocking with 5% non-fat milk in Tris buffered saline, membranes were incubated overnight at 4°C with various primary antibodies and then probed with HRP-conjugated secondary antibodies (anti-rabbit or anti-mouse, Abcam) for 2 hours at 4°C. Primary antibodies to *LATS2*, *ERK 1/2*, *p-*

*ERK1/2*, *Akt*, *p-Akt* (T308), *p-Akt* (S473) and *p53* were from Cell Signaling.  $\beta$ -actin (Sigma) was used as a loading control. Immune complexes were visualized by using ECL detection kit (GE Healthcare, Japan), and the band intensity was quantified by Image J software.

### **LATS2 siRNA Transfection**

Cells were transfected using the HiPerfect reagent according to the manufacturer's instructions (Qiagen, UK). The target sequence of *LATS2* specific siRNA (siLATS2) is 5'-CTCCGCAAAGGGTACTCAA-3', and 5nM siRNA was added to silence *LATS2*. One negative control siRNA (Qiagen, UK) was also included. Cultured cells were harvested for RNA or protein extraction at baseline and at 48 hours.

### **Statistical Analysis**

Data were analyzed with SPSS 18.0 software. Differences between groups were estimated using the Chi-squared test, the Student's t test, or the log-rank test. Disease-free survival (DFS) and overall survival (OS) curves were calculated by the Kaplan-Meier method. Stepwise multiple regression models were built to determine the clinical parameters that independently predict either PFS or OS. Log rank tests were used to compare cumulative survival between different groups. All p values were two-sided and  $p < 0.05$  was considered statistically significant. Cox proportional

regression model was applied for multivariate analysis. A probability level of 0.05 was used to determine statistical significance.

## Results

### Demographics of lung cancer patients

Fifty patients were recruited prospectively before they underwent surgical resection, with 23 (46.9%) male and 26 (53.1%) female patients and with age range of 38 to 88 years (mean  $\pm$  S.D.; 64.1 $\pm$ 9.2 years). The ratio of non-smokers (61%) to ex- or current smokers (39%) was around 2 to 1. 28 tumors were *EGFR* wild-type (57%) with no mutation at exons 18 - 21 while 21 were *EGFR* mutants that showed at least one *EGFR* mutation in exons 18 to 21 (43%). The details can be found in Table 1.

### LATS2 mRNA Expression in lung AD patients

AD tissue samples expressing *LATS2* mRNA at levels above the mean expression level (6.3; range 0-13.1) were assigned to the high expression group (mean expression value 9.4, n=23), and samples with expression less than the mean value were considered as the low expression group (mean expression value 3.6, n=26).

*LATS2* mRNA levels showed no significant differences between different clinical parameters, including gender, age, smoking history, pathological stage and *EGFR* mutation status (Table 1). Survival analysis indicated that high *LATS2*

expression group had significantly longer disease-free survival (DFS) ( $p=0.002$ , Figure 1A) and overall survival (OS) ( $p=0.041$ , Figure 1B) than the low *LATS2* expression group. Furthermore, *LATS2* mRNA levels correlated with DFS (partial correlation ratio, 0.37;  $p=0.012$ ). Multivariate analysis further confirmed that *LATS2* mRNA level was a significant prognostic factor for survival status (DFS: hazard ratio, 0.221;  $p=0.003$ ; OS: hazard ratio, 0.238;  $p=0.036$ ) independent of age, gender, smoking history and staging (Table 2). Meanwhile, pathological stage (hazard ratio, 5.102;  $p=0.009$ ), as well as the presence of *EGFR* mutations (hazard ratio, 0.207;  $p=0.006$ ) were also significant prognostic factor of DFS (Table 2). When dividing samples based on both *EGFR* mutation status and *LATS2* level, patients with wild-type *EGFR* as well as expressing low *LATS2* expression displayed poor DFS (Figure 1C). Inferior DFS was also observed in non-smokers exhibiting low *LATS2* levels (Figure 1D).

### **LATS2 protein expression in lung AD tumors**

Immunohistochemistry (IHC) staining on the independent set of 79 paraffin blocks of AD indicated that *LATS2* protein was expressed at relatively low levels in both the cytoplasm and the nucleus (Figure 2C). After excluding 16 cases with zero H-score in both compartments, paired test revealed that *LATS2* protein expression was significantly higher in the cytoplasm than in the nucleus ( $p<0.001$ ). There was an association between *LATS2* IHC scores in the cytoplasm and gender with *EGFR* mutation status, in which female patients with *EGFR* mutations displayed modestly

low levels of *LATS2* cytoplasm expression ( $p=0.031$ ). In males, *EGFR* wild-type cohort expressed slightly higher cytoplasmic staining of *LATS2*. These observations with *LATS2* IHC were consistent with the findings of *LATS2* expression at mRNA levels among gender with *EGFR* mutation status groups in a different set of tumor specimens with real-time PCR assay done as described in the previous section. Nuclear *LATS2* expression levels were generally low and no remarkable difference was found between all the clinical parameters examined.

### ***LATS2* mRNA expression in AD cell lines**

Before transfection, *LATS2* protein levels were quantified in each cell line (Figure 2A). Basal *LATS2* expression in H2023, HKULC-2 and H1650 were much higher, while H1648, HKULC-4, H1975 and PC9 exhibited relatively lower expression levels of *LATS2*. After being transfected with *LATS2*-specific siRNA (si*LATS2*) or Negative Control siRNA (siCont), *LATS2* expression was successfully silenced in all cell lines confirmed with reduced levels of *LATS2* protein (Figure 2B).

### ***LATS2* modulates *ERK* pathway in the *EGFR* wild-type AD cell lines with high *LATS2* expression**

After successfully silencing *LATS2* expression in AD cell lines, we first examined for changes in phosphorylation of mitogen-activated protein kinase, *ERK1/2* (Figure 3A). In most of the AD cell lines studied except for H1975 in which we failed to detect the activation of *ERK1/2*, the levels of phosphorylated *ERK1/2* did not alter significantly

with *LATS2* silencing. Nevertheless, a modest increase in the phosphorylation status of *ERK1/2* was observed in H2023.

### ***LATS2* knockdown differentially influences *Akt* activation in different lung AD cell lines**

As another *EGFR*-activated signaling cascade, *Akt* pathway could also be modulated by *LATS2* kinase. siRNA silencing of *LATS2* increased levels of Thr308-phosphorylated *Akt* in H2023 (Figure 3B). Since phosphorylation of *Akt* at Thr308 could enhance *Akt* activity, relatively high basal *LATS2* expression in this cell lines would inhibit *Akt* activation. Similar to the situation of *ERK* activity, *LATS2* knockdown did not affect *Akt* phosphorylation in neither HKULC-4 which expressed very low level of *LATS2* nor cell lines, PC9 and H1975, with *EGFR* mutations.

Phosphorylation at Ser473, which contributes to maximal *Akt* Activity, was also improved by *LATS2* knockdown in HKULC-2 cells (Figure 3C). Together with above results, *LATS2* might diminish *Akt* activation in H2023 and HKULC-2 cells. Both cell lines showed high *LATS2* expression at baseline. In H1650, the addition of *LATS2* siRNA slightly reduced the level of phosphorylated *Akt* at Ser473. Although the difference was not significant, it was still plausible that *LATS2* may differentially influence *Akt* activation and the regulation might be independent of *EGFR* mutation status.



### **Silencing of *LATS2* induces *p53* accumulation in lung AD cell lines with low basal *p53* levels**

Previous reports have indicated that *LATS2* is able to stabilize p53 thus facilitating p53-dependent checkpoint response to mitotic stress in breast cancer and osteosarcoma cells. However, in HKULC-2 cells, *LATS2* knockdown markedly increased total *p53* levels (Figure 3D), suggesting that, instead of inducing *p53* accumulation, *LATS2* may actually downregulate *p53* protein expression in these cells. Noticeably, HKULC-2 cells expressed very low amount of p53 at baseline (Figure 3D). Another two cell lines, HKULC-4 and H1650, which also exhibited low basal *p53* expression, *LATS2* silencing likewise enhanced *p53* levels although the differences were not statistically significant.

### **Discussion**

In this study, we demonstrated that low *LATS2* expression was an independent and significant predictor of poor overall survival. Furthermore, expression of *LATS2* correlated with DFS of these lung adenocarcinoma patients (partial correlation ratio, 0.37;  $p=0.012$ ). These findings suggested that *LATS2* may express tumor suppressive effects in lung adenocarcinoma.

In this study, patients with tumors harboring EGFR mutations exhibited longer median DFS than those with wild-type EGFR (median DFS: 33.4 vs. 27.2 months,

respectively;  $p=0.006$ ). None of the recruited patients received TKI treatment. Other than complete resection for most of these early stage tumors, no treatment intervention was apparent. The influence of EGFR gene mutation on prognosis as shown in this study deserves further investigation. This unusual observation may stem from differences in ethnicity and pathological stage of recruited patients, as well as different detection methods of EGFR mutations, compared to reported data<sup>27-29 30-32</sup>. Since wild type *EGFR* also significantly predicted inferior DFS, it is not surprising that tumors with wild type *EGFR* and low *LATS2* expression exhibited the shortest DFS (overall  $p=0.004$ , Figure 1C). In non-smokers, in which *EGFR* mutations are more common, significant difference in DFS was still observed between two *LATS2* expression groups ( $p = 0.010$ , Figure 1D). Similar observation has been made before, that knockdown of YAP (Yes kinase-associated protein), whose functions can be inhibited by *LATS2*, sensitizes cancer cells to EGFR-TKI erlotinib<sup>33</sup>. Further studies are warranted to investigate the relationships between *LATS2* expression levels and therapeutic responses of these patients.

Immunohistochemistry suggested that AD tumors expressed *LATS2* in both cytoplasm and nucleus. During interphase, *LATS2* mainly remained in the cytoplasm, especially localized to the centrosome<sup>8</sup>. After phosphorylated by *Aurora A* kinase during mitosis, *LATS2* could translocate to the chromosome and the central spindle, thus modulating chromosome segregation and cytokinesis<sup>9,10,34</sup>. To date, the best characterized function of *LATS2* is the regulation of *YAP/TAZ* proto-proteins via

canonical Hippo signaling mainly in breast, colon and hepatic cancers as well as in malignant pleural mesothelioma<sup>15,35,36</sup> but not lung adenocarcinomas. *LATS2* might still play critical roles in non-canonical Hippo signaling and even in Hippo independent pathways, such as the G-protein-coupled receptor (GPCR) signaling<sup>37,38</sup> and the *K-ras/Raf-1* axis<sup>39,40</sup>. The exact roles, as well as the relationship between the subcellular localization and the biological functions of *LATS2*, in lung adenocarcinomas need further exploration.

Our data suggested that *LATS2* can suppress *ERK* signaling in *EGFR* wild-type lung adenocarcinoma cell lines with high basal *LATS2* expression. Another study conducted in HeLa cells identified three down-regulated genes in siRNA-*LATS1/2*-treated cells, namely *SPRED1*, *SPRY2* and *SPRY4*<sup>41</sup>. These three proteins act as negative regulators of the pathway<sup>42-45</sup>, indicating that *LATS2* might inhibit *ERK* activation through regulating some members of the Sprouty proteins. Additionally, as an upstream activator of *ERK1/2*, any change in *Ras* protein levels after silencing of *LATS2* may contribute to observed alterations in *ERK1/2* activities. However, in si*LATS2*-treated H2023, no detectable change in *Ras* protein expression was observed (Figure 4). This implies that *LATS2* regulate *ERK* phosphorylation via other mechanisms in H2023. It is mentionable that, in most of lung AD cell lines studied *LATS2* knockdown induced *Ras* protein expression (Figure 4). The consequence of this modulated *Ras* expression by *LATS2* needs further investigation.

The regulation of *LATS2* on *Akt* signaling was more complicated and independent of *EGFR* mutation status. In two *EGFR* wild-type cell lines (H2023 and HKULC-2), *LATS2* could reduce *Akt* activation by inhibiting *Akt* phosphorylation at either Thr308 or Ser473. This alternation may probably result from the ability of *LATS2* to maintain the protein phosphatase-2A catalytic subunit (PP2A-C) level<sup>39</sup> which in turn leads to dephosphorylation of *Akt* Thr308<sup>46</sup>. On the other hand, the interaction between *LATS2* and *Akt* may be reciprocal and linked by mammalian sterile 20-like kinase-1 (*Mst1*). Apart from an upstream activator of *LATS2*, *Mst1* has been reported to function as an inhibitor of *Akt*<sup>47</sup>, and, meanwhile, *Akt* signaling leads to phosphorylation of *Mst1* at Thr120 and limits *Mst1*-mediated tumor suppressive functions<sup>48,49</sup>. The role of *LATS2* in this network deserves further study. However, *LATS2* may also positively modulate *Akt* activity since silencing of *LATS2* in H1650 triggered a mild decrease in the levels of phosphorylated *Akt*. As H1650 exhibits loss of *PTEN* gene<sup>50</sup> and constitutive activation of *Akt*, downregulation of *LATS2* in this cell line, which may in turn inhibit *Akt* activity, is possible to compensate this abnormal signaling transduction.

In this study, *LATS2* may suppress *p53* expression, which is contradictory to previous studies that *LATS2* can cause *p53* induction in response to mitotic apparatus damage<sup>7</sup> as well as oncogenic activation<sup>51</sup>. That may be partially explained by the absence of mitotic or oncogenic stress introduced to cells in this study. On the other hand, in NSCLC tumors, downregulated *LATS2* mRNA levels have been found to be

highly correlated with lower expression of *Mdm2* which leads to *p53* ubiquitination and degradation<sup>17,18</sup>, implying that diminished *LATS2* expression might positively regulate *p53* protein level, as seen in HKULC-2, HKULC-4 and H1650. All of three cell lines express very low amount of *p53* at baseline, while two of them, HKULC-2 and H1650, show relatively high basal *LATS2* levels. So it is reasonable to deduce that the low basal *p53* expression may partially result from the corresponding high basal expression of *LATS2*. Additionally, based on two studies that *Snail* has been found to suppress *p53* in A549 lung carcinoma cell line<sup>52</sup> and *LATS2* can act as a positive regulator of Snail1 protein<sup>53</sup>, we could speculate that *LATS2* may negatively mediate *p53* activity through *Snail*. Further research can include these two candidates, *MDM2* and *Snail*, to disclose the comprehensive interaction between *LATS2* and *p53* in lung adenocarcinoma cells. Moreover, since silencing of *LATS2* would induce *p53* accumulation, we might postulate that *LATS2* is linked to drug sensitivity through regulation of *p53*-mediated processes<sup>54</sup>.

In conclusion, relatively higher level of *LATS2* gene correlates with the better survival in patients with early stage resected lung AD. *In vitro* *LATS2* knockdown suggested that this differential expression could impact *EGFR*-activated signaling network, particularly in high *LATS2* expression group, and *p53* pathway. *LATS2* also appeared to modulate different signaling pathways via phosphorylated ERK or Akt in *EGFR* wild-type and *EGFR* mutant lung AD cell lines respectively. Thus, *LATS2* gene expression in lung AD patients warrant further investigations for being a

potential biomarker for survival status in lung cancer patients of both early and advanced stages as well as the role of LATS2 expression in modulating different signaling pathways in either EGFR mutant or EGFR wild-type lung cancer.

### **Acknowledgement**

We would like to thank Professor PC Yang, National Taiwan University, for providing the lung cancer cell line (PC9) used in this study.

This study was supported in part by the Hong Kong SK Yee Medical Grant and the University of Hong Kong Seed Funding.

### **Conflict of Interest Statement**

All the authors in this manuscript declare no conflict of interest that could inappropriately influence or bias this piece of work.

## Reference

1. Hori T, Takaori-Kondo A, Kamikubo Y, et al: Molecular cloning of a novel human protein kinase, kpm, that is homologous to warts/lats, a Drosophila tumor suppressor. *Oncogene* 19:3101-9, 2000
2. Matsui Y, Nakano N, Shao D, et al: Lats2 is a negative regulator of myocyte size in the heart. *Circ Res* 103:1309-18, 2008
3. Pan D: Hippo signaling in organ size control. *Genes Dev* 21:886-97, 2007
4. Kamikubo Y, Takaori-Kondo A, Uchiyama T, et al: Inhibition of cell growth by conditional expression of kpm, a human homologue of Drosophila warts/lats tumor suppressor. *J Biol Chem* 278:17609-14, 2003
5. Li Y, Pei J, Xia H, et al: Lats2, a putative tumor suppressor, inhibits G1/S transition. *Oncogene* 22:4398-405, 2003
6. Ke H, Pei J, Ni Z, et al: Putative tumor suppressor Lats2 induces apoptosis through downregulation of Bcl-2 and Bcl-x(L). *Exp Cell Res* 298:329-38, 2004
7. Aylon Y, Michael D, Shmueli A, et al: A positive feedback loop between the p53 and Lats2 tumor suppressors prevents tetraploidization. *Genes Dev* 20:2687-700, 2006
8. Toji S, Yabuta N, Hosomi T, et al: The centrosomal protein Lats2 is a phosphorylation target of Aurora-A kinase. *Genes Cells* 9:383-97, 2004
9. Yabuta N, Mukai S, Okada N, et al: The tumor suppressor Lats2 is pivotal in Aurora A and Aurora B signaling during mitosis. *Cell Cycle* 10:2724-36, 2011
10. Yabuta N, Okada N, Ito A, et al: Lats2 is an essential mitotic regulator required for the coordination of cell division. *J Biol Chem* 282:19259-71, 2007
11. Yabuta N, Fujii T, Copeland NG, et al: Structure, expression, and chromosome mapping of LATS2, a mammalian homologue of the Drosophila tumor suppressor gene lats/warts. *Genomics* 63:263-70, 2000
12. Takahashi Y, Miyoshi Y, Takahata C, et al: Down-regulation of LATS1 and LATS2 mRNA expression by promoter hypermethylation and its association with biologically aggressive phenotype in human breast cancers. *Clin Cancer Res* 11:1380-5, 2005
13. Powzaniuk M, McElwee-Witmer S, Vogel RL, et al: The LATS2/KPM tumor suppressor is a negative regulator of the androgen receptor. *Mol Endocrinol* 18:2011-23, 2004
14. Mizuno T, Murakami H, Fujii M, et al: YAP induces malignant mesothelioma cell proliferation by upregulating transcription of cell cycle-promoting genes. *Oncogene* 31:5117-22, 2012
15. Murakami H, Mizuno T, Taniguchi T, et al: LATS2 is a tumor suppressor gene of malignant mesothelioma. *Cancer Res* 71:873-83, 2011
16. Lin CW, Chang YL, Chang YC, et al: MicroRNA-135b promotes lung cancer metastasis by regulating multiple targets in the Hippo pathway and LZTS1. *Nat Commun* 4:1877, 2013

17. Strazisar M, Mlakar V, Glavac D: LATS2 tumour specific mutations and down-regulation of the gene in non-small cell carcinoma. *Lung Cancer* 64:257-62, 2009
18. Strazisar M, Mlakar V, Glavac D: The expression of COX-2, hTERT, MDM2, LATS2 and S100A2 in different types of non-small cell lung cancer (NSCLC). *Cell Mol Biol Lett* 14:442-56, 2009
19. Sasaki H, Hikosaka Y, Kawano O, et al: Hypermethylation of the large tumor suppressor genes in Japanese lung cancer. *Oncol Lett* 1:303-307, 2010
20. Lai JH, She TF, Juang YM, et al: Comparative proteomic profiling of human lung adenocarcinoma cells (CL 1-0) expressing miR-372. *Electrophoresis* 33:675-88, 2012
21. Liu X, Sempere LF, Ouyang H, et al: MicroRNA-31 functions as an oncogenic microRNA in mouse and human lung cancer cells by repressing specific tumor suppressors. *J Clin Invest* 120:1298-309, 2010
22. Lam DC, Girard L, Beer DG, et al: Expression of LATS2 and STK17B distinguishes epidermal growth factor receptor (EGFR) gene mutations in exons 19 and 21 from those in 18 and 20 in lung adenocarcinomas. *Journal of Thoracic Oncology* 6:S734, 2011
23. Lam DC, Girard L, Suen WS, et al: Establishment and expression profiling of new lung cancer cell lines from Chinese smokers and lifetime never-smokers. *J Thorac Oncol* 1:932-42, 2006
24. Ellison G, Zhu G, Moulis A, et al: EGFR mutation testing in lung cancer: a review of available methods and their use for analysis of tumour tissue and cytology samples. *J Clin Pathol* 66:79-89, 2013
25. Ma ES, Ng WK, Wong CL: EGFR gene mutation study in cytology specimens. *Acta Cytol* 56:661-8, 2012
26. Zhang Y, Hu CF, Chen J, et al: LATS2 is de-methylated and overexpressed in nasopharyngeal carcinoma and predicts poor prognosis. *BMC Cancer* 10:538, 2010
27. Kim YT, Seong YW, Jung YJ, et al: The presence of mutations in epidermal growth factor receptor gene is not a prognostic factor for long-term outcome after surgical resection of non-small-cell lung cancer. *J Thorac Oncol* 8:171-8, 2013
28. Liu WS, Zhao LJ, Pang QS, et al: Prognostic value of epidermal growth factor receptor mutations in resected lung adenocarcinomas. *Med Oncol* 31:771, 2014
29. Kosaka T, Yatabe Y, Onozato R, et al: Prognostic implication of EGFR, KRAS, and TP53 gene mutations in a large cohort of Japanese patients with surgically treated lung adenocarcinoma. *J Thorac Oncol* 4:22-9, 2009
30. D'Angelo SP, Janjigian YY, Ahye N, et al: Distinct clinical course of EGFR-mutant resected lung cancers: results of testing of 1118 surgical specimens and effects of adjuvant gefitinib and erlotinib. *J Thorac Oncol* 7:1815-22, 2012
31. Izar B, Sequist L, Lee M, et al: The impact of EGFR mutation status on outcomes in patients with resected stage I non-small cell lung cancers. *Ann Thorac Surg* 96:962-8, 2013
32. Lee YJ, Park IK, Park MS, et al: Activating mutations within the EGFR kinase domain: a molecular predictor of disease-free survival in resected pulmonary adenocarcinoma. *J Cancer Res Clin Oncol* 135:1647-54, 2009



33. Huang JM, Nagatomo I, Suzuki E, et al: YAP modifies cancer cell sensitivity to EGFR and survivin inhibitors and is negatively regulated by the non-receptor type protein tyrosine phosphatase 14. *Oncogene*, 2012
34. Zhang L, Iyer J, Chowdhury A, et al: KIBRA regulates aurora kinase activity and is required for precise chromosome alignment during mitosis. *J Biol Chem* 287:34069-77, 2012
35. Harvey KF, Zhang X, Thomas DM: The Hippo pathway and human cancer. *Nat Rev Cancer* 13:246-57, 2013
36. Yu FX, Guan KL: The Hippo pathway: regulators and regulations. *Genes Dev* 27:355-71, 2013
37. Mo JS, Yu FX, Gong R, et al: Regulation of the Hippo-YAP pathway by protease-activated receptors (PARs). *Genes Dev* 26:2138-43, 2012
38. Yu FX, Zhao B, Panupinthu N, et al: Regulation of the Hippo-YAP pathway by G-protein-coupled receptor signaling. *Cell* 150:780-91, 2012
39. Kili G, Kyriakis JM: Mammalian Ste20-like kinase (Mst2) indirectly supports Raf-1/ERK pathway activity via maintenance of protein phosphatase-2A catalytic subunit levels and consequent suppression of inhibitory Raf-1 phosphorylation. *J Biol Chem* 285:15076-87, 2010
40. Matallanas D, Romano D, Al-Mulla F, et al: Mutant K-Ras activation of the proapoptotic MST2 pathway is antagonized by wild-type K-Ras. *Mol Cell* 44:893-906, 2011
41. Visser S, Yang X: Identification of LATS transcriptional targets in HeLa cells using whole human genome oligonucleotide microarray. *Gene* 449:22-9, 2010
42. Bundschu K, Walter U, Schuh K: Getting a first clue about SPRED functions. *Bioessays* 29:897-907, 2007
43. Lo TL, Fong CW, Yusoff P, et al: Sprouty and cancer: the first terms report. *Cancer Lett* 242:141-50, 2006
44. Sutterluty H, Mayer CE, Setinek U, et al: Down-regulation of Sprouty2 in non-small cell lung cancer contributes to tumor malignancy via extracellular signal-regulated kinase pathway-dependent and -independent mechanisms. *Mol Cancer Res* 5:509-20, 2007
45. Garcia-Dominguez CA, Martinez N, Gragera T, et al: Sprouty2 and Spred1-2 proteins inhibit the activation of the ERK pathway elicited by cyclopentenone prostanoids. *PLoS One* 6:e16787, 2011
46. Andjelkovic M, Jakubowicz T, Cron P, et al: Activation and phosphorylation of a pleckstrin homology domain containing protein kinase (RAC-PK/PKB) promoted by serum and protein phosphatase inhibitors. *Proc Natl Acad Sci U S A* 93:5699-704, 1996
47. Cinar B, Fang PK, Lutchman M, et al: The pro-apoptotic kinase Mst1 and its caspase cleavage products are direct inhibitors of Akt1. *EMBO J* 26:4523-34, 2007
48. Collak FK, Yagiz K, Luthringer DJ, et al: Threonine-120 phosphorylation regulated by phosphoinositide-3-kinase/Akt and mammalian target of rapamycin pathway signaling limits the antitumor activity of mammalian sterile 20-like kinase 1. *J Biol Chem* 287:23698-709, 2012

49. Yuan Z, Kim D, Shu S, et al: Phosphoinositide 3-kinase/Akt inhibits MST1-mediated pro-apoptotic signaling through phosphorylation of threonine 120. *J Biol Chem* 285:3815-24, 2010
50. Chen G, Kronenberger P, Teugels E, et al: Targeting the epidermal growth factor receptor in non-small cell lung cancer cells: the effect of combining RNA interference with tyrosine kinase inhibitors or cetuximab. *BMC Med* 10:28, 2012
51. Aylon Y, Yabuta N, Besserglick H, et al: Silencing of the Lats2 tumor suppressor overrides a p53-dependent oncogenic stress checkpoint and enables mutant H-Ras-driven cell transformation. *Oncogene* 28:4469-79, 2009
52. Lee SH, Lee SJ, Jung YS, et al: Blocking of p53-Snail binding, promoted by oncogenic K-Ras, recovers p53 expression and function. *Neoplasia* 11:22-31, 6p following 31, 2009
53. Zhang K, Rodriguez-Aznar E, Yabuta N, et al: Lats2 kinase potentiates Snail1 activity by promoting nuclear retention upon phosphorylation. *EMBO J* 31:29-43, 2012
54. Kawahara M, Hori T, Chonabayashi K, et al: Kpm/Lats2 is linked to chemosensitivity of leukemic cells through the stabilization of p73. *Blood* 112:3856-66, 2008

## Figure Legends

**Figure 1:** Survival analysis of disease-free survival (DFS) and overall survival (OS).

A) Significant difference in disease-free survival between high and low LATS2 expression groups was present. The p-value was estimated by the log-rank test. B) AD patients with high LATS2 expression had a significantly better overall survival than those expressing low LATS2. . C) Cox regression analysis demonstrated that patients in the wild type EGFR together with low LATS2 expression group exhibited the worst DFS. D) In non-smoker patients, high LATS2 expression is a significant predictor of better DFS.

**Figure 2:** LATS2 protein expression in AD tumors and cell lines. A) Before transfection, LATS2 protein expression was measured in each cell lines. B) Cells were transfected with siRNA for LATS2 (siLATS2), Negative Control siRNA (siCont) or none (Mock Control) for 48hr. Number below each blot indicates fold change to mock control (n=3). C) Immunohistochemistry staining of LATS2 expression in clinical samples of AD. Female cases bearing EGFR mutations had moderately lower expression of LATS2 in the cytoplasm (mean expression value, 77.7 vs. 92.5, p = 0.031).

**Figure 3:** The effects of LATS2 silencing on ERK activation, Akt phosphorylation and p53 level. Cells were transfected with siRNA for LATS2 (siLATS2), Negative Control siRNA (siCont) or none (Mock Control) for 48hr. Cellular lysates were analyzed by Western blotting with the indicated antibodies. Number below each blot indicates fold change to mock control (n=3). One asterisk indicates p<0.05 determined by Student's t test. A) Changes in phosphorylated ERK levels in cell lines after 48-hour transfection with siRNAs described above. B) Analysis of Thr308-phosphorylated Akt levels in cells after transfected with siRNAs described above. C) Levels of Ser473-phosphorylated Akt in HKULC-2, H1648 and H1650 cells with or without LATS2 knockdown. D) The top left picture showed the basal levels of p53 in different cell lines. Modulation of p53 levels in these cells after transfected with siRNAs described above was demonstrated in the lower left picture and in the graph.

**Figure 4:** Modulation of Ras expression by LATS2. A) Cells were transfected with siRNA for LATS2 (siLATS2), Negative Control siRNA (siCont) or none (Mock Control) for 48hr. Cellular lysates were analyzed by Western blotting with the indicated antibodies. Number below each blot indicates fold change to mock control (n=3).

Ms. Ref. No.: LUNGCANCER-D-14-00074R1

Title: Aberrant Large Tumor Suppressor 2 (LATS2) gene expression correlates with EGFR mutation and survival in lung adenocarcinomas

Lung Cancer

### **Acknowledgement**

We would like to thank Professor PC Yang, National Taiwan University, for providing the lung cancer cell line (PC9) used in this study.

This study was supported in part by the Hong Kong SK Yee Medical Grant and the University of Hong Kong Seed Funding.

### **Conflict of Interest Statement**

All the authors in this manuscript declare no conflict of interest that could inappropriately influence or bias this piece of work.

**Table 1:** Patient and tumor characteristics according to *LATS2* mRNA expression (A) and *LATS2* protein expression (B)(A) *LATS2* mRNA expression

		High <i>LATS2</i> Expression	Low <i>LATS2</i> Expression	<i>P</i> value
N (%)		23 (46.9)	26 (53.1)	
<b>Follow-up Duration (Day), mean ± S.D.</b>			1276 ± 404	
<b>Gender, n(%), M:F</b>			23:26	0.907
	Male	11 (22.4)	12 (24.5)	
	Female	12 (24.5)	14 (28.6)	
<b>Age, mean ± S.D.</b>			64.1 ± 9.2	0.336
		66.3 ± 7.6	62.2 ± 10.2	
<b>Smoking, n(%), SM:NS</b>			19:30	0.590
	SM	8 (16.3)	11 (22.4)	
	NS	15 (30.6)	15 (30.6)	
<b>Stage, n(%), I:II:III</b>			29:12:08	0.709
	Stage I	15 (30.6)	14 (28.6)	
	Stage II	5 (10.2)	7 (14.3)	
	Stage III	3 (6.1)	5 (10.2)	
<b>In groups</b>				0.706
	Earlier stage (I & II)	20 (40.8)	21 (42.9)	
	Advanced stage (III)	3 (6.1)	5 (10.2)	
<b>Relapse, n(%)</b>				0.015*
	Yes	7 (14.3)	17 (34.7)	
	No	16 (32.7)	9 (18.4)	
<b>DFS (Day), median</b>			919	
		1303	746	0.006*
<b>Death, n(%)</b>				0.044*
	Yes	3 (6.1)	10 (20.4)	
	No	20 (40.8)	16 (32.7)	
<b>OS (Day), median</b>			1095	
		1364	794	0.059
<b>EGFR, n(%), Mut:WT</b>			21:28	0.283
	Mutation (Mut)	8 (16.3)	13 (26.5)	
	Wild type (WT)	15 (30.6)	13 (26.5)	

B) *LATS2* IHC staining

		Above mean	Below mean	<i>P</i> value
N (%)		32 (50.8)	31 (49.2)	
<b>Gender, n(%), M:F</b>			36:27	0.021*
	Male	18 (36)	18 (36)	
	Female	13 (26)	14 (28)	
<b>EGFR, n(%), Mut:WT</b>			37:22	0.01*
	Mutation (Mut)	18 (36)	19 (38)	
	Wild type (WT)	11 (22)	11 (22)	

\*Statistically significant ( $p < 0.05$ )

C) *EGFR* mutation status of samples included in this study

		Samples for mRNA Measurement	Samples for IHC
EGFR wild-type		28	22
EGFR Mutation	Exon 18	4	3
	Exon 19	8	10
	Exon 20	1	1
	Exon 21	7	6
	L858R+T790M	1	1
	Unknown	0	27

**Table 2:** Multivariate analysis of disease-free survival (DFS) and overall survival (OS)  
(Cox proportional regression model)

Univariate Analysis	DFS			OS		
	HR	95% CI	p-value	HR	95% CI	p-value
<b>Gender (female/male)</b>	1.129	0.497-2.566	0.772	2.364	0.769-7.270	0.133
<b>Age</b>	0.951	0.905-1.000	0.051	1.020	0.956-1.088	0.546
<b>Smoking History (SM/NS)</b>	1.157	0.477-2.807	0.747	0.561	0.187-1.678	0.301
<b>Stage (I - II/III)</b>	3.975	1.569-10.071	0.004*	1.040	0.229-4.719	0.960
<b>EGFR Mutation Status (WT/Mut)</b>	0.577	0.247-1.351	0.205	0.348	0.095-1.268	0.110
<b>LATS2 expression (low/high)</b>	0.269	0.110-0.662	0.004*	0.281	0.077-1.029	0.055

Multivariate Analysis	DFS			OS		
	HR	95% CI	p-value	HR	95% CI	p-value
<b>Gender (female/male)</b>	2.431	0.654-9.039	0.185	2.246	0.448-11.251	0.325
<b>Age</b>	0.997	0.939-1.058	0.913	1.015	0.933-1.103	0.729
<b>Smoking History (SM/NS)</b>	2.119	0.564-7.961	0.266	1.482	0.327-6.724	0.610
<b>Stage (I - II/III)</b>	5.102	1.512-17.208	0.009*	1.832	0.320-10.504	0.497
<b>EGFR Mutation Status (WT/Mut)</b>	0.207	0.068-0.631	0.006*	0.304	0.071-1.306	0.109
<b>LATS2 expression (low/high)</b>	0.221	0.081-0.605	0.003*	0.238	0.062-0.908	0.036*

HR = Hazard Ratio; CI = Confidence interval; SM = Smoker; NS = Non-smoker; WT = Wild type; Mut = Mutant



\*Statistically significant


**Highlights**

- The expression level of Large Tumor Suppressor 2 (*LATS2*) gene associate with survival in patients with resected lung adenocarcinoma (AD).
- *LATS2* kinase may interact with *EGFR* signaling, such as Erk and Akt pathways.
- High *LATS2* expression may suppress p53 expression in lung AD cell lines.


Accepted Manuscript



**A****B**



C



D

**A**

H2023

HKULC-2

H1648

HKULC-4

PC9


H1975

H1650


Short Exposure

**LATS2**


Long Exposure

 **$\beta$ -actin**


B




C




**Negative LATS2 staining**




**LATS2 staining only in the cytoplasm**




**LATS2 staining only in the nucleus**



**LATS2 staining high in the cytoplasm and low in the nucleus**




**LATS2 staining low in the cytoplasm and high in the nucleus**



**High LATS2 staining in both cytoplasm and nucleus**

C (Con't)




A

**Phospho-  
ERK****Total-ERK** **$\beta$ -actin**Mock  
Control    siLATS2    siContMock  
Control    siLATS2    siContMock  
Control    siLATS2    siCont


H2023




1    1.28    0.86



1    1.04    0.97




HKULC-2




1    0.91    1.08




1    0.98    1.16




H1648




1    0.84    1.21




1    1.08    1.14




HKULC-4




1    0.86    0.82



1    0.93    0.99



H1650



1    0.98    1.04




1    1.06    1.13




PC9




1    1.38    1.08



1    0.91    0.93




A (Con't)







B




## B (Con't)



C

**p-Akt(S473)****Total-Akt** **$\beta$ -actin**



## C (Con't)



D

H2023      HKULC-2      H1648      HKULC-4      PC9      H1975      H1650

p53

 $\beta$ -actin

p53

 $\beta$ -actinMock  
Control

siLATS2

siCont

Mock Control

siLATS2

siCont


H2023



1

0.92

0.90



HKULC-2




1

1.17

0.61



D (Con't)



**Ras** **$\beta$ -actin**