



Title	Roles of a novel splice variant of human IFI16 in innate immune response
Author(s)	Wang, P; Siu, KL; Zhang, Y; Yang, W; Kok, KH; Jin, D
Citation	The 2014 Hong Kong Inter-University Biochemistry Postgraduate Symposium, Hong Kong, 14 June 2014. In Program Booklet, 2014, p. 44, poster no. 58
Issued Date	2014
URL	http://hdl.handle.net/10722/203872
Rights	Creative Commons: Attribution 3.0 Hong Kong License

58 **Roles of a novel splice variant of human IFI16 in innate immune response**

PEIHUI WANG,¹ Kam-Lung Siu,¹ Yan Zhang,² Wanling Yang,² Kin-Hang Kok¹ and Dong-Yan Jin¹

¹Department of Biochemistry and ²Department of Pediatrics and Adolescent Medicine, LKS Faculty of Medicine, The University of Hong Kong

DNA from viral or bacterial pathogens activates innate immune response. The recognition of self-DNA would induce autoimmune diseases such as systemic lupus erythematosus (SLE). In human, AIM2 like receptors (ALRs) including AIM2, IFI16, IFIX and MNDA are DNA binding proteins implicated in DNA sensing. Most ALRs contain an N-terminal pyrin domain and C-terminal HIN200 domains. However, mouse SLE susceptibility locus p202 encodes only HIN200 domains. A human homolog of p202 was not found. Here, we identified and characterized a novel splice variant of human IFI16, which has a similar domain structure as mouse p202. We named it as IFI16 β and the original version became IFI16 α . We found that IFI16 β has its own promoter and is regulated independently of IFI16 α . IFI16 β was more abundant in human THP-1 monocytic leukemia cells and it was induced substantially by interferon β in HEK293T and HepG2 cells. Expression of IFI16 β was significantly higher in peripheral blood leukocytes of SLE patients. IFI16 β was also found to interact with AIM2 and STING. Whereas IFI16 α was predominantly found in the nucleus, IFI16 β localized to the cytoplasm. Interestingly, IFI16 β was a potent inhibitor of NLRP3 inflammasomes. Further investigations are required to determine whether IFI16 β exerts its suppressive activity on NLRP3 inflammasomes by counteracting AIM2. Taken together, human IFI16 β is a novel alternatively spliced variant that might have inhibitory activity on inflammasome activation in the cytoplasm.

Supported by Hong Kong Research Grants Council (HKU1/CRF/11G).

59 **Transcriptional activation and biochemical activities of Small Alarmone Synthase (SAS) proteins from *Staphylococcus aureus***

Ning YANG and Rory M. WATT

Oral Biosciences, Faculty of Dentistry, The University of Hong Kong

The stringent response is a key regulatory process exhibited by bacteria in response to amino acid deprivation or other challenging environmental conditions. It is mediated by the nucleotides guanosine 3',5'-bis(diphosphate) (ppGpp) and guanosine 3'-diphosphate, 5'-triphosphate (pppGpp), collectively known as (p)ppGpp. Rel-family proteins (RSH: RelA, SpoT) are responsible for the synthesis and hydrolysis of (p)ppGpp in most bacterial species. However, many bacteria also encode one or two smaller Rel-family proteins, known as small alarmone synthases (SAS), which are also capable of producing (p)ppGpp. In this study, we investigated the transcription and biochemical activities of the Rel-family proteins (SA-Rel, SA-RelP, SA-RelQ) from the notable bacterial pathogen *Staphylococcus aureus*. The respective abilities of the SA-Rel, SA-RelP and SA-RelQ proteins to catalyze the synthesis of alarmones were characterized by analyzing enzymatic reaction mixtures by anion exchange chromatography. Quantitative real-time PCR (qRT-PCR) was used to determine their transcription levels under various 'stress-inducing' conditions. The SA-RelP and SA-RelQ proteins could effectively synthesize (p)ppGpp alarmones, but had no hydrolytic activities. The transcription of SA-Rel, SA-RelP and SA-RelQ was induced by the presence of the antibiotic mupirocin, which is known to induce the stringent response. Transcription patterns for SA-Rel, SA-RelP and SA-RelQ were more variable in response to other stressful environmental conditions. In brief conclusion, our data indicates that all three Rel-family proteins play important roles in helping *S. aureus* alter its cellular physiology to respond to challenging nutritional and environmental conditions.