

The HKU Scholars Hub

## The University of Hong Kong 香港大学学術庫 場 明 物 通

Title	Genomic insights into high exopolysaccharide-producing dairy starter bacterium Streptococcus thermophilus ASCC 1275
Author(s)	Wu, Q; Shah, N
Citation	The 2nd IDF Symposium on Microstructure of Dairy Products and 5th IDF Symposium on Science and Technology of Fermented Milk, Melbourne, Australia, 3-7 March 2014.
Issued Date	2014
URL	http://hdl.handle.net/10722/205074
Rights	Creative Commons: Attribution 3.0 Hong Kong License

## Genomic insights into high exopolysaccharide-producing dairy starter bacterium *Streptococcus thermophilus* ASCC 1275

## Q. Wu<sup>1</sup>, N.P. Shah<sup>1</sup>

## <sup>1</sup>Food and Nutritional Science, School of Biological Sciences, The University of Hong Kong, Pokfulam Road, Hong Kong

Streptococcus thermophilus is an essential dairy starter for the manufacture of yogurt and cheese. Whole-genome sequencing of this organism is expected to provide insights into the genetic basis of metabolic pathways for biotechnological and probiotic applications. Streptococcus thermophilus ASCC 1275, a high EPS-producing dairy starter, has shown texture-enhancing properties for yogurt and cheese. After genomic DNA extraction using CTAB/NaCl method, whole genome sequencing including one shot-gun sequencing, two extra paired-end sequencing and Sanger sequencing was performed for strain Streptococcus thermophilus ASCC 1275 to obtain estimated 10x coverage of the genome. The raw reads were *de novo* assembled using Newbler 2.7. The genome was annotated by NCBI Prokaryotic Genomes Annotation Pipeline (PGAAP) and was compared against two other sequenced strains of S. thermophilus ND03 and MN-ZLW-002. Common features shared by S. thermophilus strains, such as efficient protocooperation with Lactobacillus delbrueckii subsp. bulgaricus, lactate production, well-equipped acid tolerance and proteolytic activity may account for dairy fermentation. Specific physiological properties of ASCC 1275, such as unique EPS synthesis and assembly, folic acid biosynthesis and CRISP/Cas system against phage infection might benefit industrial fermentation processes. This study highlights the specific biosynthesis pathway for EPS production for the strain Streptococcus thermophilus ASCC 1275.