



Title	Gender difference in schizophrenic symptomatology and subjective stress of Chinese adults with schizophrenia in a long-stay residential setting
Author(s)	Tam, SYY; Siu, JCY; Wan, AHY; Chan, CKP; Wong, CPK; Au Yeung, FSW; Ho, RTH
Citation	The 2014 Regional Congress of the World Psychiatry Association (WPA), Hong Kong, China, 12-14 December 2014. In East Asian Archives of Psychiatry, 2014, v. 24 n. 4 suppl., p. 72, abstract OP3.3.1
Issued Date	2014
URL	http://hdl.handle.net/10722/206150
Rights	East Asian Archives of Psychiatry. Copyright © Hong Kong Academy of Medicine Press.

GENDER DIFFERENCE IN SCHIZOPHRENIC SYMPTOMATOLOGY AND SUBJECTIVE STRESS OF CHINESE ADULTS WITH SCHIZOPHRENIA IN A LONG-STAY RESIDENTIAL SETTING

SYT TAM¹, JCY SIU², AHY WAN², CKP CHAN², CPK WONG³, FSW AU-YEUNG³, RTH HO²

1 Department of Psychiatry, Queen Mary Hospital, Hong Kong SAR, China

2 Department of Social Work and Social Administration, Centre on Behavioral Health, The University of Hong Kong, Hong Kong SAR, China

3 Hong Kong Sheng Kung Hui, The Providence Garden for Rehabilitation, Hong Kong SAR, China

OBJECTIVES: Males showed a higher risk of schizophrenia.¹ This study aimed to explore the manifestation of schizophrenia between genders, focusing on schizophrenic symptomatology and perceived stress in Chinese adults with schizophrenia at long-term care residential setting. Results of the present study yield implications for effective health care strategies for this population.

METHODS: A total of 146 Chinese in-patients (80 males and 66 females), aged 24 to 69 (mean \pm standard deviation [SD], 53.97 ± 8.54) years, participated in this cross-sectional study. Independent sample *t*-tests were performed to explore potential gender differences in symptomatology and perceived stress. Schizophrenia symptomatology was indexed by positive and negative symptom, gender psychopathology subscales of the Positive and Negative Symptoms Scale (PANSS), while subjective stress was indicated by the Perceived Stress Scale (PSS).

RESULTS: Results indicated difference on measures of subjective stress ($t(138) = -2.15$, $p \leq 0.05$), negative symptoms ($t(138) = 4.82$, $p \leq 0.01$), and general psychopathology ($t(138) = 3.55$, $p \leq 0.01$) between males and females. Males reported higher level of negative symptoms (mean \pm SD, 21.12 ± 7.64) and general psychopathology (38.43 ± 14.43), but lower level of subjective stress (14.93 ± 6.67) when compared with females (15.38 ± 11.81 ; 38.43 ± 14.34 ; 17.35 ± 6.60).

CONCLUSIONS: The present study replicated existing gender differences findings in schizophrenia.² Males present more negative symptoms, while females reported higher prevalence of perceived stress. The study allows better understanding of illness presentation between the genders, which implies the need for developing gender-specific health care treatments in achieving better functioning outcome.

References

1. McGrath JJ. Variations in the incidence of schizophrenia: data versus dogma. *Schizophr Bull* 2006;32:195-7.
2. Usall J, Suarez D, Haro JM; SOHO Study Group. Gender differences in response to antipsychotic treatment in outpatients with schizophrenia. *Psychiatry Res* 2007;153:225-31.