



<b>Title</b>	<b>A case of Hemiplegia Vegetativa Alterna, Paroxysmal Sympathetic Hyperactivity and Ogilvie's Syndrome: the role of central sympathetic pathways in their pathophysiology</b>
<b>Author(s)</b>	<b>Chan, RSK; Ho, JWK; Chan, STM; Woo, PYM; Leung, KM; Chan, KY</b>
<b>Citation</b>	<b>The 21st Annual Scientific Meeting of the Hong Kong Neurosurgical Society, Hong Kong, 5-6 December 2014. In Program Book, 2014, p. 63</b>
<b>Issued Date</b>	<b>2014</b>
<b>URL</b>	<b><a href="http://hdl.handle.net/10722/207373">http://hdl.handle.net/10722/207373</a></b>
<b>Rights</b>	<b>Creative Commons: Attribution 3.0 Hong Kong License</b>



## Title:

### **A Case of Hemiplegia Vegetativa Alterna, Paroxysmal Sympathetic Hyperactivity and Ogilvie's Syndrome: The Role of Central Sympathetic Pathways in their Pathophysiology**

## Authors and Institution:

Robert SK Chan, Joanna WK Ho, Sharon TM Chan, Peter YM Woo; KM Leung, KY Chan

*Department of Neurosurgery, Kwong Wah Hospital*

## Abstract:

Hemiplegia vegetativa alterna (HVA) is the clinical syndrome of contralateral hemiparesis, hemisensory loss, hemihyperhydrosis and ipsilateral Horner's syndrome<sup>1,2</sup>. The term vegetativa alterna denotes that a single brainstem lesion manifests with ipsilateral and contralateral, i.e. crossed, signs of autonomic ("vegetative") sympathetic nervous system dysfunction. Fewer than five cases have been reported and most were a result of stroke involving the occlusion of posterior cerebral artery (PCA) perforators that supply the anterolateral mesencephalon<sup>1-3</sup>.

We describe a 46 year old male who suffered from aneurysmal subarachnoid hemorrhage and exhibited HVA as a consequence of mesencephalic injury. The patient also experienced paroxysmal sympathetic hyperactivity (PSH) and recurrent colonic pseudo-obstruction, known as Ogilvie's syndrome. PSH is identified as a pathological state of elevated sympathetic activity characterized by episodic tachycardia, hypertension, tachypnea, hyperthermia diaphoresis and decerebrate dystonia occurring in 10% of severe traumatic brain injury patients<sup>4</sup>. Although the pathogenesis has yet to be elucidated the excitatory-inhibitory ratio (EIR) disconnection theory postulates that certain brainstem centers are inhibitory in nature and when severely injured, the spinal cord is released from higher regulation leading to sympathetic hyperactivity<sup>5</sup>. Colonic pseudo-obstruction is a gastro-intestinal emergency with brain injury accounting for 10% of cases<sup>6,7</sup>. It is understood that sympathetic hyperactivity also plays an important role in its pathophysiology<sup>7</sup>.

To our knowledge this is the first documented case of HVA, PSH and colonic pseudo-obstruction in the literature. We propose that the three conditions are related to an interruption of central sympathetic pathways and present clinico-radiological evidence to demonstrate that focal lesions of the anterolateral mesencephalon can account for this unique presentation.