



Title	Doctor-diagnosed Gastro-Oesophageal Reflux Disease in Hong Kong adolescents: prevalence and atypical symptoms
Author(s)	Mok, HY; Ho, DSY; Yung, HYD; Lo, WS; Lam, TH
Citation	The 2014 East-West Alliance Global Symposia, Hong Kong, China, 27-28 October 2014
Issued Date	2014
URL	http://hdl.handle.net/10722/207861
Rights	Creative Commons: Attribution 3.0 Hong Kong License



Doctor-diagnosed Gastro-Oesophageal Reflux Disease in Hong Kong adolescents: prevalence and atypical symptoms

Mok HY, Ho SY, Yung D, Lo WS, Lam TH
School of Public Health, University of Hong Kong

Background

Some atypical symptoms have been suggested to identify subjects with Gastro-Oesophageal Reflux Disease (GORD)

Lower Esophageal Sphincter Open
Allowing Reflex

Abdominal pain



Cough




Halitosis



Sore throat



Vomiting



Objectives

- Investigate the prevalence of GORD in Hong Kong adolescents
- Investigate its association with 5 atypical symptoms

Methods

34678 secondary school students (mean age 14.6, SD 2.0; 44.4% boys) from 42 randomly selected schools

Questionnaire

- Diagnosed with GORD by Western medical practitioners (yes or no)
- Presence of the 5 atypical symptoms in the past 30 days (yes or no)
- Number of atypical symptoms (0 as reference, 1-3 and 4-5)



Logistics regression was used to assess associations, adjusting for confounders

Results


1 The prevalence of GORD was 1.4% (95% CI 1.3%-1.5%)

2 Most adolescents with GORD had 1 or more symptoms

3 Higher prevalence of atypical symptoms in adolescents with GORD



4 Each atypical symptom was significantly associated with GORD



5 Likelihood of GORD increased with the number of atypical symptoms

No. of symptoms	Adjusted odds ratio	95% CI
0 (reference)		
1-3	1.87	1.56-2.28
4-5	15.80	11.40-21.88

Conclusion

Our study supports the use of these atypical symptoms for GORD identification.