



Title	The effectiveness of low-dose desmopressin in improving hypothermia-induced impairment of primary haemostasis under influence of aspirin - a randomized controlled trial.
Author(s)	Tsui, PY; Cheung, CW; Lee, Y; Leung, SWS; Ng, JKF
Citation	BMC Anesthesiol, 2015, v. 15, p. 80
Issued Date	2015
URL	http://hdl.handle.net/10722/211693
Rights	Creative Commons: Attribution 3.0 Hong Kong License

RESEARCH ARTICLE

Open Access



The effectiveness of low-dose desmopressin in improving hypothermia-induced impairment of primary haemostasis under influence of aspirin – a randomized controlled trial

Pui Yee Tsui¹, Chi Wai Cheung^{2*}, Yvonne Lee², Susan Wai Sum Leung³ and Kwok Fu Jacobus Ng²

Abstract

Background: Mild hypothermia (34–35 °C) increases perioperative blood loss. Our previous studies showed that desmopressin could have *in vitro* beneficial effects on hypothermia-induced primary haemostasis impairment. In this study, we investigate the *in vitro* effects of desmopressin on hypothermia-induced primary haemostasis impairment under the influence of aspirin in healthy volunteers.

Methods: Sixty healthy volunteers were randomly allocated to taking aspirin 100 mg or placebo for three days. On the sixth day blood samples were taken before and after the injection of desmopressin (1.5 microgram or 5 microgram) or normal saline subcutaneously. Measurements including Platelet Function Analyzer (PFA-100®) closure times, plasma von Willebrand Factor antigen, haemoglobin and platelet levels were made at 32 °C and 37 °C respectively.

Results: Collagen/epinephrine closure time (EPICT) was significantly prolonged by 21.13 % (95 %CI 2.34–39.74 %, $p = 0.021$) in aspirin group at 37 °C. While hypothermia alone prolonged both collagen/adenosine diphosphate (ADPCT) and EPICT by 17.63 % (95 %CI 13.5–20.85 %, $p < 0.001$) and 8.0 % (95 %CI 6.38–10.04 %, $p = 0.024$) respectively, addition of aspirin only further prolonged EPICT by 19.9 % (95 %CI 3.32–36.49 %, $p = 0.013$). In aspirin group, desmopressin 1.5 microgram and 5 microgram significantly reduced ADPCT to below baseline levels at 37 °C ($p = 0.025$ and < 0.001 respectively), whereas reduction in EPICT was seen with desmopressin 5 microgram ($p = 0.008$). The effect was less pronounced at 32 °C, with a significant reduction in EPICT obtained with a dosage of 5 microgram only ($p = 0.011$).

Conclusion: It was shown that aspirin could further potentiate the hypothermia-induced closure time prolongations. Low dose desmopressin (1.5 microgram) reduced PFA-100® closure times towards baseline. A higher dosage (5 microgram) further reduced the closure times below baseline. Therefore low dose desmopressin (1.5 microgram) might have the potential to correct hypothermia-induced primary haemostasis impairment under the influence of aspirin during the perioperative period.

Trial registration: ClinicalTrials.gov: NCT01382134

Keywords: Aspirin, Blood coagulation, Blood loss, Desmopressin, Hypothermia

* Correspondence: cheucw@hku.hk

²Department of Anaesthesiology, The University of Hong Kong, Room K424, 4th Floor, K Block, Queen Mary Hospital, 102 Pokfulam Road, Pokfulam, Hong Kong, SAR, China

Full list of author information is available at the end of the article

Background

Aspirin is the most commonly taken anti-platelet drug. While it has roles in primary and secondary preventions in a variety of thromboembolic conditions, it can also lead to impairment in primary haemostasis. In particular, pre-operative aspirin intake has been associated with increased post-operative blood loss [1–3]. In the recent POISE-2 study, it was shown that peri-operative intake of aspirin (100–200 mg daily) increased risk of major bleeding in patients, while risk of death or non-fatal myocardial infarction was not reduced when compared to the placebo group [4]. There is always a dilemma on whether cessation of anti-platelet therapy during the peri-operative period would increase the risk of thromboembolic event, especially in patients with increased cardiovascular risks [5, 6]. Given the widespread use of aspirin in primary and secondary cardiovascular disease preventions, it is not uncommon that patients will be under its influence during the peri-operative period. On the other hand, intra-operative hypothermia may also have a negative impact on perioperative blood loss as well. It was previously shown that hypothermia (34–35 °C) increased surgical bleeding and transfusion requirements in both cardiac and non-cardiac surgery [7–10]. Hypothermia also plays a significant role in coagulopathy in trauma patients and those requiring massive transfusion [11, 12]. While hypothermia might be difficult to avoid totally, it was found that low dose desmopressin might have beneficial effect on the primary haemostasis impairment it causes [13, 14]. We therefore conducted this study to evaluate any effect of aspirin on hypothermia-induced primary haemostasis impairment, and whether low dose desmopressin could possibly influence such effect.

Methods

The study was approved by the Hong Kong West Cluster/University of Hong Kong Institutional Review Board (Reference number: UW11-075) and registered at www.ClinicalTrials.gov with Trial Registration Number NCT01382134. The study was carried out according to the Declaration of Helsinki. Sixty healthy volunteers were recruited from August 2011 to June 2012 and informed consents were obtained from all of them. All subjects were over eighteen years old and had no known medical disease nor pregnant. Smokers, regular alcohol users and those with a known history of bleeding disorder were excluded. Subjects were instructed not to take any medication or herbal preparations that would affect haemostasis in the preceding two weeks. They were also instructed not to have red wine or chocolate intake 24 h before blood collection.

Subjects were randomized by computer into aspirin group, taking low dose aspirin 100 mg daily (Cartia 100 mg, Glaxo Smith Kline, Sigma Pharmaceuticals Pty Ltd, Dandenong, Australia) or placebo group, taking

placebo daily for a total of three days. Fifteen males and fifteen females were assigned to the aspirin group, while thirteen males and seventeen females to the placebo group. On the fourth day, morning urine samples were collected for detection of urine 11-dehydro thromboxane B2 (11-dehydro thromboxane B2 EIA Kit Cat. no. 19501, Cayman Chemical Company, Michigan, USA) to check for response to aspirin. On the sixth day, subjects were invited to our laboratory for venous blood sampling and drug testing. After collection of baseline blood samples, subjects were randomized to receive subcutaneous injection of either 0.9 % sodium chloride (placebo), desmopressin 1.5 microgram or 5 microgram made up to 1 ml solution (Octostim 15microgram/ml Desmopressin Acetate, Ferring International Center SA, St Prex, Switzerland). Blood samples were collected two hours later from the contralateral arm for post-treatment analysis.

All blood sampling, drug administration and measurements were performed by investigators who were blinded to the randomization codes. The venous blood samples were stored in citrated samples (BD Vacutainer 0.109 M, 3.2 % Citrate Tube Cat. no. 363047, BD Diagnostics, Dubai, United Arab Emirates). The blood samples were then divided into two groups and subjected to *in-vitro* hypothermia (37 °C versus 32 °C) and haemodilution (undiluted versus dilution with 20 % by volume of normal saline). They were subjected to the following tests: (1) PFA-100° closure times (PFA-100 Platelet Function Analyzer, Dade Behring Inc, West Sacramento, USA), using collagen/adenosine diphosphate (ADP) and collagen/epinephrine (EPI) cartridges as described previously [15] (Dade PFA collagen/EPI Test Cartridge Cat. no. B4170-20; Dade PFA collagen/ADP Test Cartridge Cat. no. B4170-21; Both from Siemens Healthcare Diagnostic Products GmbH, Marburg, Germany). They were incubated at 32 or 37 °C for 15 min before performing any PFA-100° measurements; (2) full blood count (pocH-100i Automated Hematology Analyzer, Sysmex Corporation, Kobe, Japan) and plasma fibrinogen (CA-50 Automated Blood Coagulation Analyzer, Sysmex Corporation, Kobe, Japan); (3) plasma von Willebrand Factor (vWF) antigen (Imubind vWF ELISA plate Cat. no. 828, American Diagnostica Inc, Stamford, US).

Repeated measures analysis of variance (RANOVA) and Chi-square tests were used in analysis of closure times within each group. Inter-group comparisons were analyzed using Mann–Whitney *U* test (for 2 independent groups) and Wilcoxon Signed-rank test (for 2 dependent groups) due to violation of normal distribution. Post-hoc comparisons were performed with paired *t*-test with Bonferroni correction. Statistical analysis was performed using SAS System for Windows Release 9.2, SAS Institute Inc., Cary, NC, USA. A *p* value of <0.05 is considered significant. Using a standard deviation of 16 % of the values

of mean closure times obtained from other healthy volunteers at our centre, our study would have a power of 90 % for detection of a 25 % change in PFA-100® closure times with ten subjects in each group [16].

Results

A total of sixty subjects were recruited in the study and all were included in data analysis (Fig. 1). One subject from the placebo group reported mild facial swelling and stomach discomfort later the day after having desmopressin

injection, which was self-limiting and no serious adverse reactions were reported.

Levels of urinary 11 dehydro thromboxane B2 in aspirin-taking subjects were significantly lower than that of placebo group (Fig. 2). Table 1 showed the baseline closure time measurements of both aspirin and placebo groups. The baseline EPICTs were significantly prolonged in aspirin group by 21.13 % (95 %CI 2.34–39.74 %, *p* = 0.021) when compared with the placebo group. Hypothermia alone prolonged both ADPCT and EPIC T by 17.63 % (95 %CI 13.5–20.85 %, *p* < 0.001) and 8.0 % (95 %CI 6.38–





Fig. 1 Flow diagram of patient recruitment



10.04 %, $p = 0.024$) respectively when compared to normothermic values. The combination of aspirin and hypothermia further prolonged EPIC T by 19.9 % (95 % CI 3.32–36.49 %, $p = 0.013$) when compared with hypothermia alone, but ADPCT was not significantly prolonged.

The administration of desmopressin significantly reduced both ADPCT and EPIC T of aspirin group at 37 °C (Fig. 3). While there was significant reduction of ADPCT at doses 1.5 microgram ($p = 0.025$) and 5 microgram ($p < 0.001$) when comparing to baseline values, an insignificant reduction of EPIC T was seen with 1.5 microgram ($p = 0.65$) and significant reduction at 5 microgram ($p = 0.008$).

Table 1 Baseline closure times of aspirin and placebo groups at 37 °C


	Aspirin group 37 °C (n = 30)	Placebo group 37 °C (n = 30)	p value
ADPCT	120.7 ± 46.6	123.9 ± 42.5	0.728
EPIC T	211.3 ± 68.9	174.6 ± 56.8	0.021 ⁺
	Aspirin group 32 °C (n = 30)	Placebo group 32 °C (n = 30)	p value
ADPCT	142.3 ± 58.2	145.7 ± 62.6	0.690
EPIC T	226.1 ± 67.6	188.6 ± 52.5	0.013 ⁺

Values in mean ± SD


⁺Significant at the 0.05 level

ADPCT: Collagen/adenosine diphosphate closure time

EPIC T: Collagen/adrenaline closure time



At 32 °C under the influence of aspirin, desmopressin produced a less pronounced effect on the closure times (Fig. 4). With desmopressin 1.5 microgram, both ADPCT and EPIC T were not significantly different from their baseline values at 37 °C. At a dose of 5 microgram, desmopressin caused a mild but insignificant reduction of ADPCT when compared to baseline values at 37 °C ($p = 0.431$), while a significant reduction was only seen with EPIC T ($p = 0.011$). The dose response curves of desmopressin did not seem to be temperature-dependent, as the slopes of dose–response curves are nearly parallel as shown in Fig. 5.




The vWF antigen did not show a significant increase after desmopressin 1.5 microgram and 5 microgram (Table 2). Other parameters did not show any significant change, except for haemoglobin level which showed a statistical significant ($p < 0.05$) but probably not clinically significant difference.

While results from hypothermia group were presented in this paper, those from haemodilution group shall be presented in a separated report in the future.

Discussion

The design of this study was based on our previous studies on evaluation of hypothermia-induced primary haemostasis



impairment and the effect of desmopressin on such impairment [14, 17]. In this study, it was shown once again the baseline ADPCT and EPICT were significantly prolonged in presence of hypothermia. The addition of aspirin further prolonged the EPICT, showing a synergistic effect between hypothermia and aspirin on primary haemostasis impairment. As it is not uncommon to see patients taking aspirin peri-operatively, this may have clinical implication on peri-operative blood loss where hypothermia is difficult to avoid totally. In addition, it was previously shown that an EPICT >188 s could predict increased drain output in patients taking non-steroidal anti-inflammatory drugs after total knee arthroplasty with a sensitivity of 89 % [18]. EPICT prolongation in the presence of hypothermia and aspirin could therefore potentially predict an increase in peri-operative blood loss.

Table 2 Aspirin group - concentrations of von Willebrand factor antigen (vWF:Ag), haemoglobin (Hb), fibrinogen (FIB) and platelet count, in subjects receiving saline or desmopressin 1.5 and 5 µg subcutaneously

	Saline (n = 10)		Desmopressin 1.5 (n = 10)		Desmopressin 5 (n = 10)	
	Baseline	Post	Baseline	Post	Baseline	Post
von Willebrand factor antigen; mU.ml ⁻¹	1051.3 ± 320.9	978.8 ± 509.5	918.6 ± 346.0	1028.8 ± 384.2	1025.6 ± 430.1	1266.5 ± 589.7
Hb; g.dl ⁻¹	14.4 ± 1.4	14.3 ± 1.3	13.9 ± 1.1	13.5 ± 1.0 ⁺	14.8 ± 1.4	13.9 ± 1.8 ⁺
Platelets; x 10 ⁹ l ⁻¹	218.0 ± 65.9	232.8 ± 52.6	222.0 ± 54.0	228.2 ± 32.3	258.9 ± 36.6	259.6 ± 41.2
FIB; mg.dl ⁻¹	332.0 ± 72.1	342.0 ± 69.6	337.1 ± 105.9	368.5 ± 75.1	364.4 ± 47.8	355.1 ± 53.4

Values in mean ± SD

⁺p < 0.05 within group post-treatment compared with baseline (Wilcoxon Signed-rank test)

In our previous studies, we have shown that low dose subcutaneous desmopressin could improve hypothermia-induced impairment in primary haemostasis [17]. It was shown that the dosage required could be as low as 1.5 microgram, much lower than the standard dose of 15 microgram used in treating platelet dysfunction associated with uraemia or cirrhosis. In this follow-up study, it was shown that desmopressin at dosage of 1.5 microgram and 5 microgram could correct the ADPCT and EPICT to close to baseline values under the influence of hypothermia and aspirin. Such effect was milder at 32 °C when compared to 37 °C measurements. In fact, there seemed to be a trend for the 5 microgram dose to over-correct closure times to below baseline values (Figs. 2 and 3). Therefore we postulated that at dosage of 1.5 microgram desmopressin might be sufficient to cause improvement in hypothermia induced primary haemostasis impairment under the influence of aspirin.

Being a point-of-care monitoring of coagulation status, it has been suggested that the PFA-100° could be used as a means to monitor the effectiveness of aspirin therapy [19–22]. “Aspirin responders” and “aspirin non-responders” could be identified using population based cut-off values. In one particular study, Coakley et al. showed that PFA-100° could be used to identify “aspirin hyper-responders” peri-operatively [22]. This could allow us to identify a small group of “aspirin hyper-responders” who might be at risk of increased perioperative bleeding in presence of hypothermia and have their aspirin therapy evaluated before major operations. Indiscriminate cessation of aspirin therapy peri-operatively might unnecessarily pose the patients to increased risk of thrombotic events. PFA-100° could also be used as a screening tool for primary haemostasis abnormalities before surgery [23]. Although it is worth noting that studies so far had failed to show any clinical benefit of using PFA-100° (such as prediction of increased peri-operative blood loss) in peri-operative situations [24, 25].

The thrombotic risk of desmopressin could be related to its ability of causing prolonged von Willebrand factor (vWF) release. Our data showed that at the doses used

in this study, there was no significant increase in vWF release. This is in concordance with our previous studies where low dose desmopressin corrects primary haemostasis impairment without causing an increase in vWF levels [17]. In an *in vitro* setting, the absence of endothelial cells means that there will not be vWF release accounting for the improvement in primary haemostasis. These suggested that desmopressin has direct effect on platelets itself, possibly via stimulated expression of adhesion molecules such as glycoprotein Ib and p-selectin [26–30]. Our studied dosages might allow safe and effective use of desmopressin on improving primary haemostasis impairment while minimizing the risk of thrombotic complications.

There are several limitations of our study. Firstly, this is an *in vitro* study based on healthy volunteers. Blood samples were first collected then incubated to 32 °C and 37 °C respectively. The *in vitro* setting may not reveal *in vivo* effects of desmopressin in hypothermic patients. Previous studies on the effectiveness of desmopressin on reduction of peri-operative blood loss had also shown mixed results [31–33]. In addition, whether desmopressin could improve haemostasis impairment due to other causes (e.g., haemodilution, acidosis, noval anti-platelet agents) remained elusive.

Conclusion

In conclusion, aspirin could have a synergistic effect with hypothermia on primary haemostasis impairment. Such impairment could be reversed by subcutaneous low-dose desmopressin (1.5 microgram). This holds a promising and safe way to improve primary haemostasis impairment in aspirin taking patients undergoing major operations, where hypothermia may be difficult to avoid. Large scale clinical trials are warranted to further evaluate the efficacy and safety of desmopressin on this aspect. In addition, the PFA-100° might have a role in identifying “aspirin hyper-responders” who might be at risk of increased surgical bleeding. This could allow selective identification of high risk groups and have their peri-operative anti-thrombotic therapy reviewed, without

risking the majority of thrombotic events with indiscriminate cessation of aspirin therapy.

Abbreviations

ADPCT: Closure Time of Collagen/Adenosine Diphosphate; EPICT: Closure Time of Collagen/Epinephrine; PFA: Platelet Function Analyzer; vWF: von Willebrand Factor; RANOVA: Repeated measures analysis of variance.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

PYT and YL: Study design and protocol, subject recruitment, conduct of study, statistical analysis, draft proof-reading, approval of manuscript. CWC: Study design and protocol, subject recruitment, conduct of study, draft proof-reading, approval of manuscript. SWSL: Study design and protocol, conduct of study, laboratory testing, approval of manuscript. JKFN: Study design and protocol, conduct of study, statistical analysis, draft proof-reading, approval of manuscript.

Funding

This study was solely supported by the departmental funding of Department of Anaesthesiology, The University of Hong Kong.

Author details

¹Department of Anaesthesia, Pamela Youde Nethersole Eastern Hospital, Chai Wan, Hong Kong, SAR, China. ²Department of Anaesthesiology, The University of Hong Kong, Room K424, 4th Floor, K Block, Queen Mary Hospital, 102 Pokfulam Road, Pokfulam, Hong Kong, SAR, China. ³Department of Pharmacology & Pharmacy, The University of Hong Kong, Hong Kong, SAR, China.

Received: 1 October 2014 Accepted: 21 May 2015

Published online: 28 May 2015

References

- Kallis P, Tooze JA, Talbot S, Cowans D, Bevan DH, Treasure T. Pre-operative aspirin decreases platelet aggregation and increases post-operative blood loss—a prospective, randomised, placebo controlled, double-blind clinical trial in 100 patients with chronic stable angina. *Eur J Cardiothorac Surg.* 1994;8:404–9.
- Sun JC, Whitlock R, Cheng J, Eikelboom JW, Thabane L, Crowther MA, et al. The effect of pre-operative aspirin on bleeding, transfusion, myocardial infarction, and mortality in coronary artery bypass surgery: a systematic review of randomized and observational studies. *Eur Heart J.* 2008;29:1057–71.
- Alghamdi AA, Moussa F, Fremes SE. Does the use of preoperative aspirin increase the risk of bleeding in patients undergoing coronary artery bypass grafting surgery? Systematic review and meta-analysis. *J Card Surg.* 2007;22:247–56.
- Devereaux PJ, Mrkobrada M, Sessler DI, Leslie K, Alonso-Coello P, Kurz A, et al. Aspirin in Patients Undergoing Noncardiac Surgery. *N Engl J Med.* 2014;370:1494–503.
- Gerstein NS, Schulman PM, Gerstein WH, Petersen TR, Tawil I. Should more patients continue aspirin therapy perioperatively? Clinical impact of aspirin withdrawal syndrome. *Ann Surg.* 2012;255:811–9.
- Oscarsson A, Gupta A, Fredrikson M, Järhult J, Nyström M, Pettersson E, et al. To continue or discontinue aspirin in the perioperative period: a randomized, controlled clinical trial. *Br J Anaesth.* 2010;104:305–12.
- Cavallini M, Barufaldi Preis FW, Casati A. Effects of mild hypothermia on blood coagulation in patients undergoing elective plastic surgery. *Plast Reconstr Surg.* 2005;116:316–21.
- Hofer CK, Worn M, Tavakoli R, Sander L, Maloigne M, Klaghofer R, et al. Influence of body core temperature on blood loss and transfusion requirements during off-pump coronary artery bypass grafting: a comparison of 3 warming systems. *J Thorac Cardiovasc Surg.* 2005;129:838–43.
- Rajagopalan S, Mascha E, Na J, Sessler DI. The effects of mild perioperative hypothermia on blood loss and transfusion requirement. *Anesthesiology.* 2008;108:71–7.
- Schmied H, Kurz A, Sessler DI, Kozek S, Reiter A. Mild hypothermia increases blood loss and transfusion requirements during total hip arthroplasty. *Lancet.* 1996;347:289–92.
- Holcomb JB, Jenkins D, Rhee P, Johannigman J, Mahoney P, Mehta S, et al. Damage control resuscitation: directly addressing the early coagulopathy of trauma. *J Trauma.* 2007;62:307–10.
- Moor P, Rew D, Midwinter MJ, Doughty H. Transfusion for trauma: civilian lessons from the battlefield? *Anaesthesia.* 2009;64:469–72.
- Ferrara A, MacArthur JD, Wright HK, Modlin IM, McMillen MA. Hypothermia and acidosis worsen coagulopathy in the patient requiring massive transfusion. *Am J Surg.* 1990;160:515–8.
- Ying CLA, Tsang SF, Ng JKF. The potential use of desmopressin to correct hypothermia-induced impairment of primary haemostasis – An in vitro study using PFA-100*. *Resuscitation.* 2008;76:129–33.
- Harrison P, Robinson MS, Mackie IJ, Joseph J, McDonald SJ, Liesner R, et al. Performance of the platelet function analyser PFA-100 in testing abnormalities of primary haemostasis. *Blood Coagul Fibrinolysis.* 1999;10:25–31.
- Ng KF, Lawmin JC, Li CC, Tsang SF, Tang WM, Chiu KY. Comprehensive preoperative evaluation of platelet function in total knee arthroplasty patients taking diclofenac. *J Arthroplasty.* 2008;23:424–30.
- Ng JKF, Cheung CW, Lee Y, Leung SWS. Low-dose desmopressin improves hypothermia-induced impairment of primary haemostasis in healthy volunteers. *Anaesthesia.* 2011;66:999–1005.
- Ng JKF, Lawmin JC, Tsang SF, Tang WM, Chiu KY. Value of a single preoperative PFA-100* measurement in assessing the risk of bleeding in patients taking cyclooxygenase inhibitors and undergoing total knee replacement. *Br J Anaesth.* 2009;102:779–84.
- Homoncik M, Jilma B, Hergovich N, Stohlawetz P, Panzer S, Speiser W. Monitoring of aspirin (ASA) pharmacodynamics with the platelet function analyzer PFA-100. *Thromb Haemost.* 2000;83:316–21.
- Favaloro EJ. Clinical utility of the PFA-100. *Semin Thromb Hemost.* 2008;34:709–33.
- Crescente M, Di Castelnuovo A, Iacoviello L, Vermulen J, Cerletti C, de Gaetano G. Response variability to aspirin as assessed by the platelet function analyzer (PFA)-100. A systematic review. *Thromb Haemost.* 2008;99:14–26.
- Coakley M, Self R, Marchant W, Mackie I, Mallett SV, Mythen M. Use of the platelet function analyser (PFA-100) to quantify the effect of low dose aspirin in patients with ischaemic heart disease. *Anaesthesia.* 2005;60:1173–8.
- Franchini M. The platelet-function analyzer (PFA-100) for evaluating primary hemostasis. *Hematology.* 2005;10:177–81.
- Karger R, Reuter K, Rohlf J, Nimsky C, Sure U, Kretschmer V. The Platelet Function Analyzer (PFA-100*) as a Screening Tool in Neurosurgery. *ISRN Hematol.* 2012;2012:839242.
- Islam N, Fulop T, Zsom L, Miller E, Mire CD, Lebrun CJ, et al. Do platelet function analyzer-100 testing results correlate with bleeding events after percutaneous renal biopsy? *Clin Nephrol.* 2010;73:229–37.
- Sloand EM, Alyono D, Klein HG, Chang P, Yu M, Lightfoot FG, et al. 1-Deamino-8-D-arginine vasopressin (DDAVP) increases platelet membrane expression of glycoprotein Ib in patients with disorders of platelet function and after cardiopulmonary bypass. *Am J Hematol.* 1994;46(3):199–207.
- Wun T, Paglieroni TG, Lachant NA. Desmopressin stimulates the expression of P-selectin on human platelets in vitro. *J Lab Clin Med.* 1995;126(4):401–9.
- Balduini CL, Noris P, Belletti S, Spedini P, Gamba G. In vitro and in vivo effects of desmopressin on platelet function. *Haematologica.* 1999;84:891–6.
- Cattaneo M, Moia M, le Valle P, Castellana P, Mannucci PM. Desmopressin shortens the prolonged bleeding times of patients with severe von Willebrand disease treated with cryoprecipitate. Evidence for a mechanism of action independent of released von Willebrand factor. *Blood.* 1989;74:1972–5.
- Horstman LL, Valle Riestra BJ, Jy W, Wang F, Mao W, Ahn YS. Desmopressin acts on platelets to generate platelet microparticles and enhanced procoagulant activity. *Thromb Res.* 1995;79:163–74.
- Mannucci PM, Levi M. Prevention and treatment of major blood loss. *NEJM.* 2007;356:2301–11.
- Pleym H, Stenseth R, Wahba A, Bjella L, Tromsdal A, Karevold A, et al. Prophylactic treatment with desmopressin does not reduce postoperative bleeding after coronary surgery in patients treated with aspirin before surgery. *Anesth Analg.* 2004;98:578–84.
- Franchini M. The use of desmopressin as a hemostatic agent: a concise review. *Am J Hematol.* 2007;82:731–5.