



Title	High prevalence of Escherichia coli sequence type 131 among antimicrobial-resistant E. coli isolates from geriatric patients
Author(s)	Ho, PL; Chu, YPS; Lo, WU; Chow, KH; Law, PYT; Tse, CWS; Ng, TK; Cheng, VCC; Que, TL
Citation	Journal of Medical Microbiology, 2015, v. 64 n. 3, p. 243-247
Issued Date	2015
URL	http://hdl.handle.net/10722/213730
Rights	Journal of Medical Microbiology. Copyright © Society for General Microbiology.

1 **High prevalence of *Escherichia coli* sequence type 131 among**
2 **antimicrobial-resistant *E. coli* isolates from geriatric patients**

3
4 Pak-Leung Ho^{1*}, Yuki Pui-Shan Chu², Wai-U Lo¹, Kin-Hung Chow¹, Pierra Y. Law¹, Cindy
5 Wing-Sze Tse³, Tak-Keung Ng⁴, Vincent Chi-Chung Cheng¹, Tak-Lun Que²

6
7 ¹Department of Microbiology and Carol Yu Centre for Infection, University of Hong Kong,
8 Hong Kong SAR, PR China

9 ²Department of Clinical Pathology, Tuen Mun Hospital, Hong Kong SAR, PR China

10 ³Department of Clinical Pathology, Kwong Wah Hospital, Hong Kong SAR, PR China

11 ⁴Department of Clinical Pathology, Princess Margaret Hospital, Hong Kong SAR, PR China

12
13 Running title: *Escherichia coli* ST131 subclones from geriatric patients

14 Word counts: Abstract 207 words

15 Key words: Enterobacteriaceae; CTX-M type beta-lactamase; multilocus sequence typing;
16 antimicrobial resistance

17
18 *Correspondence: Pak-Leung Ho, Department of Microbiology, Queen Mary Hospital, The
19 University of Hong Kong, Pokfulam Road, Pokfulam, Hong Kong, China.

20 Tel: +852 22554892; Fax: +852 28551241.

21 E-mail address: plho@hkucc.hku.hk (P.L. Ho)

22

23 **ABSTRACT**

24 Previous work on the subclones within *Escherichia coli* ST131 predominantly involved
25 isolates from Western countries. This study assessed the prevalence and antimicrobial
26 resistance attributed to this clonal group. A total of 340 consecutive, nonduplicated urinary *E.*
27 *coli* isolates originating from four clinical laboratories in Hong Kong in 2013 were tested.
28 ST131 prevalence among the total isolates was 18.5% (63/340) and was higher among
29 inpatients isolates (23.0%) than outpatient isolates (11.8%, $P<0.001$); and higher among
30 isolates from patients aged ≥ 65 years than from patients aged 18-50 years and 51-64 years
31 (25.4% vs. 2.4% and 4.0%, respectively, $P<0.001$). Of the 63 ST131 isolates, 43 (68.3%)
32 isolates belonged to the *H30* subclone, whereas the remaining isolate belonged to *H41* (n=17),
33 *H54* (n=2) and *H22* (n=1). All *H30* isolates were ciprofloxacin-resistant of which 18.6%
34 (8/43) belonged to the *H30-Rx* subclone. Twenty-six (41.3%) ST131 isolates were
35 ESBL-producers of which 19 had *bla*_{CTX-M-14} (12 non-*H30-Rx*, two *H30-Rx* and five *H41*),
36 six had *bla*_{CTX-M-15} (five non-*H30-Rx* and one *H30-Rx*) and one was *bla*_{CTX-M} negative (*H30*).
37 In conclusion, ST131 accounts for a large share of the antimicrobial-resistant *E. coli* isolates
38 from geriatric patients. Unlike previous reports, ESBL-producing ST131 strains mainly
39 belonged to non-*H30-Rx* rather than the *H30-Rx* subclone with *bla*_{CTX-M-14} as the dominant
40 enzyme type.

41

42 **INTRODUCTION**

43 The incidence of infections due to antimicrobial-resistant *Escherichia coli* is increasing
44 worldwide (Barber *et al.*, 2013). Resistance rates for cotrimoxazole, fluoroquinolones and
45 third generation cephalosporins, which are often used for empirical therapy, are now above
46 the 15-20% threshold recommended for choosing first-line antimicrobial agents for empirical
47 treatment in Hong Kong (Ho *et al.*, 2007b; Ho *et al.*, 2010). Discordant therapy may cause
48 treatment failure, persistence and recurrence of infection leading to more patient morbidity
49 and mortality (Barber *et al.*, 2013; Shin *et al.*, 2012). Emerging resistance in *E. coli* involves
50 acquisition of resistance determinant by susceptible strain and the expansion of preexisting
51 resistant clones (Naseer & Sundsfjord, 2011). ST131 is a highly successful *E. coli* clone
52 which has received considerable attentions due to its wide geographic distribution, ability to
53 cause a wide range of extra-intestinal infections and association with CTX-M β -lactamases
54 and multidrug resistance (Nicolas-Chanoine *et al.*, 2014).

55

56 ST131 can be divided into different subclones by other typing methods, of which sequencing
57 the type 1 fimbrial adhesion gene *fimH* is one widely used approach (Weissman *et al.*, 2012).
58 *H30*, designated according to the *fimH30* variant is currently the most prevalent subclone of
59 ST131 (Nicolas-Chanoine *et al.*, 2014). Two studies involving whole genome sequencing of
60 ST131 isolates collected from multiple countries came to the same conclusion that

61 fluroquinolone resistance within ST131 was confined almost entirely to the *H30* subclone and
62 that CTX-M-15 producers clustered within a nested subclone, designated as *H30-Rx* (Petty *et*
63 *al.*, 2014; Price *et al.*, 2013). Reported prevalence of *H30* and *H30-Rx* among ST131 isolates
64 ranged 66.7%-95.8% and 16.9%-66.2%, respectively, depending on the isolate sources and
65 selection criteria (Banerjee *et al.*, 2013b; Peirano *et al.*, 2014; Peirano & Pitout, 2014; Price
66 *et al.*, 2013). Majority of the ST131 isolates that have been tested for the *H30* and *H30-Rx*
67 subclones were collected from North America and Europe; relatively few isolates were from
68 Asia (Banerjee *et al.*, 2013a; Banerjee *et al.*, 2013b; Colpan *et al.*, 2013; Johnson *et al.*, 2013;
69 Johnson *et al.*, 2014; Peirano *et al.*, 2014; Tchesnokova *et al.*, 2013). Additionally, few
70 studies have assessed the association of host factors with the two ST131 subclones (Banerjee
71 & Johnson, 2014). Here, we used an unselected collection of urinary *E. coli* isolates from
72 four laboratories in Hong Kong to evaluate the relationship between patient demographics,
73 antimicrobial-resistant phenotypes, ST131 and its major subclones.

74

75 **METHODS**

76 **Study design.** A total of 340 non-duplicated, urinary *E. coli* isolates were studied. The
77 isolates were consecutive single-patient *E. coli* isolates from four clinical microbiology
78 laboratories in Hong Kong over a two week period, from May to June 2013. The laboratories
79 together served about a quarter of the Hong Kong populations in different geographic

80 districts. The inclusion criteria were: (1) patient age 18 years or above, (2) mid-stream urine
81 specimen, and (3) significant growth at $\geq 10^5$ CFU/ml. Patient identities was kept anonymous.
82 The following information was provided by the submitting laboratories: sex, age, date of
83 collection and patient location (outpatient or inpatient). One isolate per patient was included.

84

85 **Microbiological methods.** The VITEK GNI system (bioMerieux Vitek Inc., Hazelwood, MO)
86 was used for bacterial identification. Antibiotic susceptibilities were tested by the disc
87 diffusion method using Mueller-Hinton agar (Oxoid, Basingstoke, UK) and interpreted
88 according to the Clinical and Laboratory Standard Institute. All antibiotic discs were obtained
89 commercially (BBL, Becton Dickinson, Cockeysville, MD, USA). The double disk synergy
90 test was used for detection of extended-spectrum β -lactamases (ESBL) (Ho *et al.*, 2010). The
91 susceptibility testing of all isolates were performed in a central laboratory at the University of
92 Hong Kong. On each day of testing, standard strains (ATCC 25922 and 35218) were included
93 as quality controls. For each isolate, the resistance score was the number of antimicrobials
94 (including ampicillin, amoxicillin-clavulanate, cefuroxime, ceftriaxone, ertapenem, nalidixic
95 acid, ciprofloxacin, co-trimoxazole, gentamicin, nitrofurantoin and fosfomycin which were
96 chosen to represent 11 classes) for which it exhibited resistance (including both intermediate
97 and resistant categories).

98

99 **Molecular studies.** PCR assays were used to assign the *E. coli* isolates to phylogroups A, B1,
100 B2, C, D, E and F (Clermont *et al.*, 2013). Phylogroup B isolates were investigated for ST131
101 status by PCR assays targeting SNPs in *mdh* and *gyrB* (Johnson *et al.*, 2009), and the O25b
102 variant and SNPs in *pabB* (Clermont *et al.*, 2009). A subset of the isolates were further tested
103 by multilocus sequence typing (MLST) for confirmation (Wirth *et al.*, 2006).
104 ST131-associated O serotype, *fimH* subtype and the H30-Rx subsubclone were determined by
105 established methods (Banerjee *et al.*, 2013b; Clermont *et al.*, 2007; Weissman *et al.*, 2012).
106 The *bla*_{CTX-M} genes were detected by PCR and sequencing using primers with specificity for
107 the CTX-M subgroups (*bla*_{CTX-M1G}, *bla*_{CTX-M2G}, *bla*_{CTX-M8G}, *bla*_{CTX-M9G}, and *bla*_{CTX-M25G}) (Ho *et*
108 *al.*, 2007a; Ho *et al.*, 2012). Alleles were assigned by sequencing the full length of *bla*_{CTX-M} as
109 previously described (Ho *et al.*, 2012).

110

111 **Statistical analysis.** The Chi-square, Fisher's exact test or Student's *t*-test were used for
112 statistical analysis. Univariate and multivariate analyses were used to assess risk factors
113 associated with ST131 subclones. The following parameters were included in the multivariate
114 analysis: age, sex, laboratory source, and patient care location. The values of parameters are
115 given as mean (\pm standard deviation) where appropriate. A two-tailed *P* value of <0.05 was
116 considered significant. All analyses were performed using statistical software (SPSS, version
117 14.0; SPSS Inc; Chicago, IL).

118 **RESULTS**

119 **Patient demographics**

120 A total of 340 urinary isolates were included in the study; 204 (60.0%) from inpatients, 136
121 (40.0%) from outpatients; 259 (76.2%) from females and 81 (23.8%) from males. Each
122 laboratory contributed 82 to 89 isolates. Overall, 50 (14.7%) were obtained from patients
123 aged 18-50 years, 58 (17.1%) from patients aged 51-64 years and 232 (68.2%) from patients
124 aged ≥ 65 years. The patients had mean age of 69.7 ± 17.3 years.

125

126 **Distribution of phylogroups and ST131 by patient sources**

127 Phylogroup B2 predominated among the isolates with similar frequencies among isolates
128 from different age groups (62.1%-66.0%) and among inpatients (63.7%) and outpatients
129 (63.2%) isolates (Table 1). Allele-specific PCR assays targeting *mdh* and *gyrB* identified 63
130 isolates as ST131 of which 45 isolates were also positive for O25b and *pabB*. One isolate was
131 *pabB* positive and *mdh*-negative, *gyrB*-positive. MLST confirmed the isolate as ST131.
132 Another 22 isolates were randomly chosen for MLST and all were confirmed to be ST131.
133 The prevalence of ST131 among all urinary isolates was 18.5% (63/340) overall, but this
134 varied according to isolate sources (Table 1). The prevalence of ST131 was higher among
135 isolates from patients aged ≥ 65 years (25.4%) than the other age groups (3.4%-4.0%), and
136 higher among inpatients (23.0%) than in outpatients (11.8%). Of the 63 ST131 isolates, 45

137 (71.4%) were serogroup O25b, 17 (27.0%) were serogroup O16 and one (1.6%) was
138 O-non-typeable. Forty-three (68.3%) ST131 isolates belonged to the *H30* subclone, whereas
139 the remaining 20 isolates belonged to *H41* (n=17), *H54* (n=2) and *H22* (n=1). All *H30*
140 isolates were ciprofloxacin-resistant and 18.6% (8/43) of *H30* isolates belonged to the
141 *H30-Rx* subclone. In general, serogroup O25b isolates were of *H30* (93.3%, 42/45) subclone
142 and serogroup O16 were of *H41* subclone (100%, 17/17). The frequency of *H30* subclone
143 was higher among patients aged ≥ 65 years and inpatients while those for *H41* and other *fimH*
144 subtypes were similar among the patient subsets. In multivariate analysis, aged ≥ 65 y was the
145 only factor significantly associated with ST131 (OR 8.9, 95% CI 3.1-25.1, $P < 0.001$), *H30*
146 (OR 7.3, 95% CI 2.2-24.1, $P = 0.001$) and *H41* (OR 7.9, 95% CI 1.04-60.6, $P = 0.046$).

147

148 **Distribution of antibiotic resistance by ST131 status**

149 Among ST131 isolates, resistance rates for 8 of the 11 antimicrobials were high, ranging
150 from 33.3% to 87.3% while those for nitrofurantoin (1.6%) and fosfomycin (1.6%) were rare.
151 All isolates including all ST131 subclones were susceptible to ertapenem. ST131 isolates
152 were significantly more likely than non-ST131 isolates to be resistant to ampicillin (87.3% vs.
153 67.9%, respectively), amoxicillin-clavulanate (33.3% vs. 18.8%), cefuroxime (41.3% vs.
154 20.9%), nalidixic acid (100% vs. 69.0%), ciprofloxacin (71.4% vs. 32.5%) and gentamicin
155 (38.1% vs. 25.6%), and to be ESBL-producers (41.3% vs. 18.8%). Resistance rates among

156 *H30* and *H41* isolates were similar except for resistance to ciprofloxacin which is
157 substantially higher among *H30* isolates (100% for *H30* vs. 11.8% for *H41*, $P < 0.001$). The
158 resistance score was highest for *H30* isolates (5.1 ± 2.0), followed by *H41* isolates (3.5 ± 1.4)
159 and non-ST131 isolates (3.0 ± 2.4). Within *H30* isolates, resistance score for *H30*-Rx ($4.9 \pm$
160 2.7) and non-*H30*-Rx (5.1 ± 1.8) isolates were similar ($P = 0.737$).

161

162 Rates of ESBL production, and ciprofloxacin, cotrimoxazole and gentamicin resistance were
163 similar among isolates from different age groups. ST131 accounted for 33.3%, 33.3%, 25.3%
164 of all ESBL-producing, ciprofloxacin-resistant, and gentamicin-resistant *E. coli* populations,
165 respectively. In contrast, prevalence of ST131 among the antimicrobial-sensitive counterparts
166 were significantly lower ($P < 0.05$ for all comparisons), being 14.1% for ESBL-negative
167 isolates, 8.8% for ciprofloxacin-sensitive isolates and 15.9% for gentamicin-sensitive isolates.
168 The prevalence of ST131 among cotrimoxazole-resistant (17.4%) and -sensitive (19.2%)
169 isolates was similar. Stratification by age groups revealed that there were variations in the
170 resistant populations attributed to ST131 (Fig. 1). Among the ST131 isolates, 26 (41.3%)
171 were ESBL-producers. PCR and sequencing showed that 19 had *bla*_{CTX-M-14} (12 non-*H30*-Rx,
172 two *H30*-Rx and five *H41*), six had *bla*_{CTX-M-15} (five non-*H30*-Rx and one *H30*-Rx) and one
173 was *bla*_{CTX-M} negative (*H30*).

174

175 **DISCUSSION**

176 We evaluated 340 *E. coli* urine isolates, collected from four laboratories in 2013 for the
177 ST131 clonal group and its subclones. The prevalence of ST131 among total *E. coli* isolates
178 (18.5%) is concordant with other studies in the United State (17%-27%) and Europe
179 (12%-22%) (Nicolas-Chanoine *et al.*, 2014). Our findings showed that the prevalence of
180 ST131 and its H30 subclone were higher among older age, inpatients and
181 antimicrobial-resistant isolates. These findings indicated that expansion of ST131 is an
182 important mechanism of increased antimicrobial resistance in the geriatric population. The
183 reason for the higher prevalence of ST131 among geriatric patients is not clear but could
184 possibly be related to selection from over-prescription of broad-spectrum antimicrobials
185 (third generation cephalosporins, fluoroquinolones), institutional acquisition from exposure in
186 old age homes and hospitals, and underlying comorbidities (Ho *et al.*, 2014). In previous
187 studies, approximately 25% of hospitalized patients and elderly residents of long-term care
188 facilities were found to carry ST131 in their feces (Banerjee & Johnson, 2014), comparing
189 with <5% among healthy young adults (Kudinha *et al.*, 2013; Leflon-Guibout *et al.*, 2008);
190 suggesting institutions may pose risk for ST131 transmission. However, a recent study found
191 that only 2.1% of 240 residents from 11 nursing homes in Germany had fecal carriage of
192 ST131 (Arvand *et al.*, 2013).

193

194 We found that *H30* comprised 68.3% of all ST131 isolates which is lower than the
195 proportions reported for unselected clinical isolates from the United States (87.3%) and
196 France (86.5%) (Lafolie *et al.*, 2014). The prevalence of *H30*-Rx subclone among our *H30*
197 isolates was 18.6%, which is substantially lower than the >70% among *H30* ST131 isolates
198 preselected by specific resistance phenotypes (Banerjee *et al.*, 2013b; Peirano *et al.*, 2014).
199 *H30*-Rx described previously among isolates from Europe and North America was almost
200 always ESBL-positive and had CTX-M-15 (Peirano *et al.*, 2014; Petty *et al.*, 2014; Price *et*
201 *al.*, 2013). Here, only three of the eight *H30*-Rx isolates were ESBL-producers. Unlike
202 previous reports (Peirano *et al.*, 2014; Petty *et al.*, 2014; Price *et al.*, 2013), ESBL-producing
203 ST131 strains in the present study mainly belonged to non-*H30*-Rx rather than the *H30*-Rx
204 subclone. Among all ST131 subclones, CTX-M-14 was the predominant ESBL found.
205 Plasmid IncF family played a major role in the dissemination of CTX-M-15 in Europe and
206 the United States (Nicolas-Chanoine *et al.*, 2014). In Asia, IncF plasmids were found to more
207 often carry CTX-M-14 instead of CTX-M-15 (Ho *et al.*, 2007a; Nicolas-Chanoine *et al.*,
208 2014). Among ST131 isolates, IncF plasmids carrying CTX-M-14 have been reported from
209 Hong Kong, mainland China and South Korea (Ho *et al.*, 2012; Nicolas-Chanoine *et al.*,
210 2014). In Japan, CTX-M-14 was detected in 44% and 73% of ESBL-producing ST131-O25b
211 and ST131-O16 isolates, respectively, comparing with 18% and 8% for CTX-M-15,
212 respectively. (Matsumura *et al.*, 2012).

213

214 In summary, this study found that antimicrobial-resistant *E. coli* from geriatric patients are
215 substantially more likely to be caused by ST131 than those from younger patients, and that
216 the *H30* and *H41* subclones possess certain resistance traits different from those reported in
217 other locales.

218

219 **Conflict of interest statement**

220 All authors have no competing interests

221

222 **Acknowledgements**

223 This work was supported by grants from the Health and Medical Research Fund (formerly

224 Research Fund for the Control of Infectious Diseases) of the Food and Health Bureau of the

225 Government of the HKSAR and the RGC Collaborative Research Fund Project on Syndromic

226 Surveillance and Modelling for Infectious Diseases

227

228

229

230

231 **REFERENCES**

232 **Arvand, M., Moser, V. & Pfeifer, Y. (2013).** Prevalence of
233 extended-spectrum-beta-lactamase-producing *Escherichia coli* and spread of the epidemic
234 clonal lineage ST131 in nursing homes in Hesse, Germany. *J Antimicrob Chemother* **68**,
235 2686-2688.

236 **Banerjee, R. & Johnson, J. R. (2014).** A New Clone Sweeps Clean: the Enigmatic
237 Emergence of *Escherichia coli* Sequence Type 131. *Antimicrob Agents Chemother* **58**,
238 4997-5004.

239 **Banerjee, R., Johnston, B., Lohse, C., Chattopadhyay, S., Tchesnokova, V., Sokurenko,**
240 **E. V. & Johnson, J. R. (2013a).** The clonal distribution and diversity of extraintestinal
241 *Escherichia coli* isolates vary according to patient characteristics. *Antimicrob Agents*
242 *Chemother* **57**, 5912-5917.

243 **Banerjee, R., Robicsek, A., Kuskowski, M. A., Porter, S., Johnston, B. D., Sokurenko, E.,**
244 **Tchesnokova, V., Price, L. B. & Johnson, J. R. (2013b).** Molecular epidemiology of
245 *Escherichia coli* sequence type 131 and its H30 and H30-Rx subclones among
246 extended-spectrum-beta-lactamase-positive and -negative *E. coli* clinical isolates from the
247 Chicago Region, 2007 to 2010. *Antimicrob Agents Chemother* **57**, 6385-6388.

248 **Barber, A. E., Norton, J. P., Spivak, A. M. & Mulvey, M. A. (2013).** Urinary tract
249 infections: current and emerging management strategies. *Clin Infect Dis* **57**, 719-724.

250 **Clermont, O., Christenson, J. K., Denamur, E. & Gordon, D. M. (2013).** The Clermont
251 *Escherichia coli* phylo-typing method revisited: improvement of specificity and detection of
252 new phylo-groups. *Environ Microbiol Rep* **5**, 58-65.

253 **Clermont, O., Dhanji, H., Upton, M. & other authors (2009).** Rapid detection of the
254 O25b-ST131 clone of *Escherichia coli* encompassing the CTX-M-15-producing strains. *J*
255 *Antimicrob Chemother* **64**, 274-277.

256 **Clermont, O., Johnson, J. R., Menard, M. & Denamur, E. (2007).** Determination of
257 *Escherichia coli* O types by allele-specific polymerase chain reaction: application to the O
258 types involved in human septicemia. *Diagn Microbiol Infect Dis* **57**, 129-136.

259 **Colpan, A., Johnston, B., Porter, S. & other authors (2013).** *Escherichia coli* sequence
260 type 131 (ST131) subclone H30 as an emergent multidrug-resistant pathogen among US
261 veterans. *Clin Infect Dis* **57**, 1256-1265.

262 **Ho, P. L., Chau, P. H., Yan, M. K., Chow, K. H., Chen, J. H., Wong, S. C. & Cheng, V. C.**
263 **(2014).** High burden of extended-spectrum beta-lactamase-positive *Escherichia coli* in
264 geriatric patients. *J Med Microbiol* **63**, 878-883.

265 **Ho, P. L., Poon, W. W., Loke, S. L., Leung, M. S., Chow, K. H., Wong, R. C., Yip, K. S.,**
266 **Lai, E. L. & Tsang, K. W. (2007a).** Community emergence of CTX-M type
267 extended-spectrum beta-lactamases among urinary *Escherichia coli* from women. *J*
268 *Antimicrob Chemother* **60**, 140-144.

269 **Ho, P. L., Wong, R. C., Yip, K. S., Loke, S. L., Leung, M. S., Mak, G. C., Chow, F. K.,**
270 **Tsang, K. W. & Que, T. L. (2007b).** Antimicrobial resistance in *Escherichia coli* outpatient
271 urinary isolates from women: emerging multidrug resistance phenotypes. *Diagn Microbiol*
272 *Infect Dis* **59**, 439-445.

273 **Ho, P. L., Yeung, M. K., Lo, W. U., Tse, H., Li, Z., Lai, E. L., Chow, K. H., To, K. K. &**
274 **Yam, W. C. (2012).** Predominance of pHK01-like incompatibility group FII plasmids
275 encoding CTX-M-14 among extended-spectrum beta-lactamase-producing *Escherichia coli*
276 in Hong Kong, 1996-2008. *Diagn Microbiol Infect Dis* **73**, 182-186.

277 **Ho, P. L., Yip, K. S., Chow, K. H., Lo, J. Y., Que, T. L. & Yuen, K. Y. (2010).**
278 Antimicrobial resistance among uropathogens that cause acute uncomplicated cystitis in
279 women in Hong Kong: a prospective multicenter study in 2006 to 2008. *Diagn Microbiol*
280 *Infect Dis* **66**, 87-93.

281 **Johnson, J. R., Clermont, O., Johnston, B., Clabots, C., Tchesnokova, V., Sokurenko, E.,**
282 **Junka, A. F., Maczynska, B. & Denamur, E. (2014).** Rapid and Specific Detection,
283 Molecular Epidemiology, and Experimental Virulence of the O16 Subgroup within
284 *Escherichia coli* Sequence Type 131. *J Clin Microbiol* **52**, 1358-1365.

285 **Johnson, J. R., Menard, M., Johnston, B., Kuskowski, M. A., Nichol, K. & Zhanel, G. G.**
286 **(2009).** Epidemic clonal groups of *Escherichia coli* as a cause of antimicrobial-resistant
287 urinary tract infections in Canada, 2002 to 2004. *Antimicrob Agents Chemother* **53**,

288 2733-2739.

289 **Johnson, J. R., Tchesnokova, V., Johnston, B. & other authors (2013).** Abrupt emergence
290 of a single dominant multidrug-resistant strain of *Escherichia coli*. *J Infect Dis* **207**, 919-928.

291 **Kudinha, T., Johnson, J. R., Andrew, S. D., Kong, F., Anderson, P. & Gilbert, G. L.**
292 **(2013).** *Escherichia coli* sequence type 131 as a prominent cause of antibiotic resistance
293 among urinary *Escherichia coli* isolates from reproductive-age women. *J Clin Microbiol* **51**,
294 3270-3276.

295 **Lafolie, J., Nicolas-Chanoine, M. H., Grenouillet, F., Hocquet, D. & Bertrand, X. (2014).**
296 Prevalence of *Escherichia coli* sequence type 131 and its H30 subclone among *E. coli*
297 isolates in a French hospital. *Int J Antimicrob Agents*.

298 **Leflon-Guibout, V., Blanco, J., Amaqdouf, K., Mora, A., Guize, L. & Nicolas-Chanoine,**
299 **M. H. (2008).** Absence of CTX-M enzymes but high prevalence of clones, including clone
300 ST131, among fecal *Escherichia coli* isolates from healthy subjects living in the area of Paris,
301 France. *J Clin Microbiol* **46**, 3900-3905.

302 **Matsumura, Y., Yamamoto, M., Nagao, M., Hotta, G., Matsushima, A., Ito, Y., Takakura,**
303 **S. & Ichiyama, S. (2012).** Emergence and spread of B2-ST131-O25b, B2-ST131-O16 and
304 D-ST405 clonal groups among extended-spectrum-beta-lactamase-producing *Escherichia*
305 *coli* in Japan. *J Antimicrob Chemother* **67**, 2612-2620.

306 **Naseer, U. & Sundsfjord, A. (2011).** The CTX-M conundrum: dissemination of plasmids

307 and *Escherichia coli* clones. *Microb Drug Resist* **17**, 83-97.

308 **Nicolas-Chanoine, M. H., Bertrand, X. & Madec, J. Y. (2014).** *Escherichia coli* ST131, an
309 intriguing clonal group. *Clin Microbiol Rev* **27**, 543-574.

310 **Peirano, G. & Pitout, J. D. (2014).** Fluoroquinolone-resistant *Escherichia coli* sequence
311 type 131 isolates causing bloodstream infections in a canadian region with a centralized
312 laboratory system: rapid emergence of the H30-Rx sublineage. *Antimicrob Agents Chemother*
313 **58**, 2699-2703.

314 **Peirano, G., van der Bij, A. K., Freeman, J. L., Poirel, L., Nordmann, P., Costello, M.,
315 Tchesnokova, V. L. & Pitout, J. D. (2014).** Characteristics of *Escherichia coli* sequence type
316 131 isolates that produce extended-spectrum beta-lactamases: global distribution of the
317 H30-Rx sublineage. *Antimicrob Agents Chemother* **58**, 3762-3767.

318 **Petty, N. K., Ben Zakour, N. L., Stanton-Cook, M. & other authors (2014).** Global
319 dissemination of a multidrug resistant *Escherichia coli* clone. *Proc Natl Acad Sci U S A* **111**,
320 5694-5699.

321 **Price, L. B., Johnson, J. R., Aziz, M. & other authors (2013).** The epidemic of
322 extended-spectrum-beta-lactamase-producing *Escherichia coli* ST131 is driven by a single
323 highly pathogenic subclone, H30-Rx. *MBio* **4**, e00377-13.

324 **Shin, J., Kim, J., Wie, S. H. & other authors (2012).** Fluoroquinolone resistance in
325 uncomplicated acute pyelonephritis: epidemiology and clinical impact. *Microb Drug Resist*

326 **18**, 169-175.

327 **Tchesnokova, V., Billig, M., Chattopadhyay, S. & other authors (2013).** Predictive
328 diagnostics for *Escherichia coli* infections based on the clonal association of antimicrobial
329 resistance and clinical outcome. *J Clin Microbiol* **51**, 2991-2999.

330 **Weissman, S. J., Johnson, J. R., Tchesnokova, V. & other authors (2012).** High-resolution
331 two-locus clonal typing of extraintestinal pathogenic *Escherichia coli*. *Appl Environ*
332 *Microbiol* **78**, 1353-1360.

333 **Wirth, T., Falush, D., Lan, R. & other authors (2006).** Sex and virulence in *Escherichia*
334 *coli*: an evolutionary perspective. *Mol Microbiol* **60**, 1136-1151.

335

336

337 **Table 1.** Distribution of phylogroups and ST131 among 340 urinary *E. coli* isolates

Categories	No (column %) by age group				No (column %) by source			
	<i>n</i>	18-50y (n=50)	51-64y (n=58)	≥65y (n=232)	<i>P</i>	Inpatients (n=204)	Outpatients (n=136)	<i>P</i>
ST131								
<i>H30</i> subclone	43	1 (2.0)	2 (3.4)	40 (17.2)	0.001	31 (15.2)	12 (8.8)	0.083
H41 subclone	17	1 (2.0)	0 (0)	16 (6.9)	0.056	13 (6.4)	4 (2.9)	0.155
Others*	3	0 (0)	0 (0)	3 (1.3)	0.494	3 (1.5)	0 (0)	0.155
Subtotal	63	2 (4.0)	2 (3.4)	59 (25.4)	<0.001	47 (23.0)	16 (11.8)	<0.001
<i>H30</i> -Rx	8	0 (0)	1 (1.7)	7 (3.0)	0.417	6 (2.9)	2 (1.5)	0.381

338 * Including H54 (two isolates) and H22 (one isolate).


339

340

341

342 **Fig 1.** Antimicrobial-resistant *E. coli* populations attributed to ST131 according to age groups.

343 The *P* values are indicated for between age groups comparison.



344

345

346

347

348

349