



Title	The protective effects of catechin on palmitic acid-induced cytotoxicity in mouse brain endothelial cell
Author(s)	Wong, KL; Cheung, CW; Chao, CZ; Su, TH; Leung, YM
Citation	The 2015 Annual Scientific Sessions of the International Academy of Cardiology cum 20th World Congress on Heart Disease, Vancouver, BC., Canada, 25-27 July 2015.
Issued Date	2015
URL	http://hdl.handle.net/10722/217815
Rights	Creative Commons: Attribution 3.0 Hong Kong License

THE PROTECTIVE EFFECTS OF CATECHIN ON PALMITIC ACID - INDUCED CYTOTOXICITY IN MOUSE BRAIN ENDOTHELIAL CELL

K.L. Wong^{1,2,3,4}, C.W. Cheung², C.Z. Chao^{3,4}, T.H. Su^{1,5}, Y.M. Leung⁵

1. Department of Anesthesiology, China Medical University & Hospital, Taichung, Taiwan
2. Department of Anesthesiology, LKS Faculty of Medicine, University of Hong Kong, Hong Kong
3. Shandong University, Qilu Hospital-Nanshan Hospital, Shandong, China
4. Department of Anesthesiology, Taishan Medical University, Shandong, China
5. Graduate Institute of Neural and Cognitive Sciences, China Medical University, Taichung, Taiwan

AIMS: The approximate prevalence of the metabolic syndrome (MS) in patients with coronary heart disease (CAD) is 50%, with a prevalence of 37% in patients with premature CAD. Effective prevention or treatment of MS significantly reduces the risk for developing serious complications. Palmitic acid (PA) is a saturated fatty acid, when being excessive, is a significant risk factor for development of MS or cardiovascular accident. Lipotoxicity in endothelial cells (EC) has been well documented but how PA affects EC Ca²⁺-signaling and other functions remain largely unexplored. Catechin has been implicated in benefiting almost every organ system such as cardioprotective and anti-obesity; and also beneficial for blood vessel health. This study aims to investigate the lipotoxic alteration of mouse brain endothelial cell line (bEND.3 cells) function mediated by PA; and how PA modulates EC ion channels, and also to assess the potential protective effects of catechin.

METHODS: Cell apoptosis assessed by TUNEL-Assay. Cytosolic Ca²⁺ in bEND was measured with Fura-2 method. Mitochondria membrane potential (MMP) measured by MMP-Assay Kit. Cell viability was measured By MTT-Assay. The $p < 0.05$ were considered significant (ANOVA).

RESULTS: Exposure of bEND to PA (300 micromolar) for 48 h resulted in apoptosis. PA (100, 300 micromolar) increased expression of CHOP but not phosphorylated eIF2-alpha. PA (300 micromolar) pretreatment diminished (Ca²⁺-agonist) ATP-triggered Ca²⁺ release and Ca²⁺ influx. 300 micromolar PA pretreatment diminished (SR Ca²⁺-pump blocker) CPA-triggered Ca²⁺ release and Ca²⁺ influx. Thus PA at this high concentration reduced the size of Ca²⁺ pool. PA at 100 micromolar, however, did not reduce CPA-induced Ca²⁺ release but suppressed Ca²⁺ influx. Thus PA at this concentration inhibits store-operated Ca²⁺ entry. PA-induced cell death was significantly alleviated by co-treatment of bEND with catechin (300 micromolar).

CONCLUSION: Cell death was apoptotic and related to endoplasmic reticulum(ER) stress and cytosolic Ca²⁺ elevation. Co-treatment of bEND with catechin (300 & micro;M) significantly prevented PA-induced cell death.