



Research Article

Seasonal variations in proximate composition and metallic elements of three cyprinids from a Central Himalayan river Alaknanda in Garhwal Himalaya, India

Sharali Sharma

Freshwater Biodiversity Laboratory, Department of Zoology, H.N.B. Garhwal University, Srinagar- Garhwal -246174 (Uttarakhand), India

Sumit Kumar

Freshwater Biodiversity Laboratory, Department of Zoology, H.N.B. Garhwal University, Srinagar- Garhwal -246174 (Uttarakhand), India

Deepak Singh*

Freshwater Biodiversity Laboratory, Department of Zoology, H.N.B. Garhwal University, Srinagar- Garhwal -246174 (Uttarakhand), India

*Corresponding author. Email: bhandaridrdeepak5@gmail.com

Article Info

<https://doi.org/10.31018/jans.v12i4.2413>

Received: October 28, 2020

Revised: December 7, 2020

Accepted: December 12, 2020

How to Cite

Sharma S. et al. (2020). Seasonal variations in proximate composition and metallic elements of three cyprinids from a Central Himalayan river Alaknanda in Garhwal Himalaya, India. *Journal of Applied and Natural Science*, 12(4):661 - 669. <https://doi.org/10.31018/jans.v12i4.2413>

Abstract

Fish is an irreplaceable, highly nutritious aquatic food in a balanced diet. On the contrary, fish can also be a source of metal contamination when ingested in excess through food and water. The aim of the present study was to determine the seasonal variations in the proximate body composition and levels of Na, K, Ca, Fe, Zn, Cu, and Pb in the muscle tissue of three fish species namely, *Schizothorax plagiostomus*, *Schizothorax progastus* and *Barilius bendelisis* dwelling in Central Himalayan river Alaknanda at Srinagar (Garhwal) in Garhwal Himalaya. The protein (21.01-21.74%) and carbohydrates (2.31-2.69%) were found maximum in summer season in all the studied fishes while lipid was found to be highest during monsoon season in *S. plagiostomus* (2.73%) and *S. progastus* (2.70%) and lowest (1.62-1.68%) in the winter season. *B. bendelisis* showed a different pattern of variation in lipid content being highest (6.32%) during summer and lowest (3.34%) in monsoon season. Significantly ($F=5.69$, $p<0.04$) high content of lipid was found in *B. bendelisis* as compared to *S. plagiostomus* and *S. progastus* during the three seasons. Na, Ca, K, and Fe were detected highest during monsoon, Pb, and Cu during summer and Zn during the winter season. However, Pb was not detected in all the fish samples during monsoon season. *B. bendelisis* was significantly ($p<0.001$) rich in Na and Ca, *S. plagiostomus* in K ($p<0.01$), and *S. progastus* in Fe ($p<0.001$) content during all three seasons. Ca was rich among macroelements while Fe among microelements in all the studied fishes. The amount of Zn (1.48-4.55 mg/100g), Cu (0.429-0.869 mg/100g), and Pb (0.117-0.447 mg/100g) were within the permissible limits of International food safety guidelines. The seasonal variations in the parameters are attributed to food availability, temperature changes, fish metabolism, and prevailing environmental conditions. Thus, these cyprinid species were regarded as safe and useful for human consumption.

Keywords: Alaknanda, Fish, Heavy metals, Lipid, Macroelements, Protein

INTRODUCTION

Fishes act as essential foodstuff due to their high protein content and affordability. Overall production of fishes in the world is about 167.2 million tons, of which about 146.3 million tons is the part of human food and rest is discarded as waste material (FAO, 2016). The fishery sector not only provides nutrition and food security but also improves employment and poverty in developing countries (Moreau and Garaway, 2018). There is a continuous increase in the demand of quality fish and fishery products as they have bountiful val-

uable, healthy substances like protein, lipid, vitamins, mineral elements, etc. The biochemical composition is used in the study of food or feed. The biochemical constituents form 95-98% of the total weight of the tissue (Pal et al., 2018). Therefore, monitoring of changes in the biochemical parameters of fish species is required.

Fishes are also a good source of mineral elements which include calcium, sodium, potassium, zinc, phosphorus, iron etc. These elements have high bioavailability, i.e. they are readily absorbed by the body (Balachandan, 2002). Many essential elements have

their role in metabolic processes and are essential to all living organisms (Durmus *et al.*, 2017). The deficiency in these essential elements induces malfunctioning and causes diseases. Iron is required for the synthesis of haemoglobin so as to transport oxygen throughout the body. Calcium is important for the normal functioning of the muscular and nervous system and also for strong bones. Fish generally occupies the high position in aquatic food chains and are more susceptible to poisoning from chemicals in their diet as well as water. They are also well consumed by the human population. Also, contamination of the river ecosystem by metals has been receiving attention worldwide. Heavy metal accumulation in biota poses a risk to the wildlife and human health (Ali *et al.*, 2017). Therefore, it is utmost important to determine the level of metals in their body tissue.

The river Ganga is one of the major Himalayan rivers having two main parent headstreams, Alaknanda and Bhagirathi which have their origin in the mountainous region of North Indian state Uttarakhand. Alaknanda, being the longer of the two, rises about 50 km north of Himalayan Nanda Devi peak at the confluence of Satopanth and Bhagirath Kharak glaciers near Vasudhara falls (3650m. a.s.l.). After covering a course of 191km, it merges with Bhagirathi river at Devprayag to form the river Ganga (Panwar *et al.*, 2017). Alaknanda harbours a variety of flora and fauna. A majority of fish species belonging to family Cyprinidae flourish in this river ecosystem. The flow in the Alaknanda River is highly regulated and sometimes highly variable due to seasonal flash floods (SANDRP, 2014). A majority of people inhabit along the bank of this river and depend upon the fishes for their nutritional requirements and livelihood. Therefore, the knowledge of seasonal variation in proximate composition and concentrations of essential metallic elements like Na, Ca, K, Fe, Zn, Cu and Pb in the body tissue of the fishes dwelling in the river becomes important. The present study also verifies the interspecific differences in the levels of these elements.

MATERIALS AND METHODS

Collection, identification and measurements: The study was conducted on the fishes dwelling in Central Himalayan river Alaknanda, a parent stream of river Ganga at Srinagar in Garhwal Himalaya. A total of 180 fish samples, 60 of each cyprinid species (*Schizothorax plagiostomus*, *Schizothorax progastus* and *Barilius bendelisis*) were procured directly from local fish market at monthly interval for one year (2017-18). As the fishes were procured from the local fish market and no harm was caused to live animals, there was no issue to take the approval of the Animal Ethic Committee.

Collected samples were transported to the laboratory in fresh condition for identification and body measurements. The fish species were identified using keys of

Day (1878), Jhingran (1982), Talwar and Jhingran (1991) and Jayaram (2002). After necessary measurements, fish gut and skin were removed and samples were then packed in airtight plastic bags and stored in deep freezer at -21°C until further analysis.

Proximate and mineral analysis: 1gm muscle tissue of each selected fish species was homogenized separately in 10 ml phosphate buffer followed by centrifugation at 10,000rpm for 20 minutes. The supernatant/sample solutions collected were subjected to total soluble protein and total carbohydrate determination. Protein was estimated following Lowry *et al.* (1951). Total carbohydrates were estimated following the procedure of Dubois *et al.* (1956) Total lipid was determined according to Folch *et al.* (1954), using olive oil as a standard. Total moisture was evaluated by oven drying 1gm of fish muscle tissue at 105°C for 24 hours. The difference in initial and final weight represents the amount of moisture (AOAC, 2000). The results were expressed in percentage of wet weight of tissue. Minerals, Na, K, Ca, Fe, Zn, Cu and Pb were determined by acid digestion of tissue samples (0.5-0.6 g) in 1:3 HCl and HNO_3 in a closed digestion system at 150°C . Samples were allowed to undergo complete digestion for about 1-2 h until a clear solution was obtained (AOAC, 2000). Samples were then cooled and filtered. The final volume of filtrate was made up to 100 ml by using millipore water. Samples were allowed to run on ICP-OES (Thermo Scientific, iCAP 6000 Series) and the concentration of mineral elements was determined (AOAC, 2000).

Statistical analysis: The data so obtained were analyzed statistically with the help of Data Analysis Tool Pack available on MS-Excel. t-test was performed to determine the significance of the difference between means of two samples, and ANOVA was performed to find out the significance of difference among the means of the seasons.

RESULTS AND DISCUSSION

The proximate composition, macroelements and level of metals in the muscle tissue of *S. plagiostomus*, *S. progastus* and *B. bendelisis* are presented in Table 1 to 3. The proximate composition of the body tissue of these fishes varied significantly among the seasons. Moisture content in the three fish species ranged from 67.40 to 83.16% (Table 1). Highest moisture content was recorded in *S. plagiostomus* (79.72%) and *S. progastus* (79.26%) during monsoon season while lowest was observed during summer (77.81%) and winter season (77.62%). However, a different pattern of moisture content highest (76.92%) during winter and lowest (73.08%) during summer season was reported in *B. bendelisis*. Average moisture content was significantly ($p < 0.05$) higher in *S. plagiostomus* and *S. progastus* than *B. bendelisis* during summer and monsoon. Percent protein values ranged from 10.14 to 27.86 in studied fishes (Table 1). Maximum protein content was

Table 1: Seasonal variations in proximate composition of three cyprinid species.

Proximate composition	Min-Max	Summer	Monsoon	Winter
Moisture (%)				
<i>S. plagiostomus</i>	77.16-81.62	77.81±0.42 ^b	79.72±1.91 ^b	78.70±1.30 ^b
<i>S. progastus</i>	75.54-83.16	78.33±1.02 ^b	79.26±2.66 ^b	77.62±2.70 ^{ab}
<i>B. bendelisis</i>	67.40-80.60	73.08±6.16 ^a	74.02±3.13 ^a	76.92±0.75 ^a
Protein (%)				
<i>S. plagiostomus</i>	11.14-25.92	21.74±2.82 ^a	19.41±3.52 ^{ab}	13.99±3.43 ^b
<i>S. progastus</i>	10.14-23.50	21.01±1.70 ^a	17.80±3.69 ^a	13.08±3.51 ^a
<i>B. bendelisis</i>	10.15-27.86	21.02±2.92 ^a	20.65±5.17 ^b	12.19±1.96 ^a
Lipid (%)				
<i>S. plagiostomus</i>	1.32-3.39	2.18±0.69 ^a	2.73±0.45 ^a	1.62±0.25 ^a
<i>S. progastus</i>	1.41-3.17	2.24±0.71 ^a	2.70±0.37 ^a	1.68±0.19 ^a
<i>B. bendelisis</i>	3.00-7.63	6.32±0.88 ^b	3.34±0.47 ^b	4.06±0.87 ^b
Carbohydrate (%)				
<i>S. plagiostomus</i>	1.14-2.71	2.37±0.48 ^a	2.02±0.5 ^a	1.41±0.20 ^a
<i>S. progastus</i>	1.21-2.84	2.31±0.35 ^a	2.26±0.52 ^b	1.38±0.12 ^a
<i>B. bendelisis</i>	1.12-3.01	2.69±0.45 ^b	2.14±0.68 ^{ab}	1.58±0.34 ^b

Min-Max- Minimum-Maximum. Values are represented as Mean±SD of 5 replicates. Values with different superscript in a row varied significantly (p<0.05).

recorded to be 21.74%, 21.01%, 21.02% during summer season and minimum to be 13.99%, 13.08%, 12.19% during winter season in *S. plagiostomus*, *S. progastus* and *B. bendelisis*, respectively. No significant difference (F=0.157, p=0.856) was observed in muscle protein content of studied fishes. Amount of carbohydrate was found maximum during summer (2.37, 2.31 and 2.69%) followed by monsoon (2.02, 2.26 and 2.14%) and winter season (1.41, 1.38 and 1.58%) in *S. plagiostomus*, *S. progastus* and *B. bendelisis* (Table 1). No significant difference (F=0.388, p=0.682) was observed in the amount of carbohydrate in studied fishes. The highest increment of protein and carbohydrate during summer and lowest during winter season was due to the variations in temperature conditions and availability of food items. During the summer season, the high temperature remains suitable for the growth of fish food like algae, zooplankton, worms and insects and their larvae which causes accumulation of more protein and carbohydrates in their muscle tissue. However, during winter conditions, low temperature results in less growth hence less protein accumulation. Lipid content ranged from 1.32 to 7.63% in studied fish species (Table 1). In *S. plagiostomus* and *S. progastus*, highest lipid accumulation (2.73% and 2.70%) was found during monsoon and lowest (1.62% and 1.68%) during the winter season while in *B. bendelisis*, highest value (6.32%) was recorded during summer and lowest (3.34%) during the monsoon season which is attributed to the variations in their reproduction period and food availability. Lipid content was significantly (p<0.001) higher in *B. bendelisis* than other studied fishes during all the three seasons. Carbohydrate content ranged from 1.12 to 3.01% and it was maximum during summer and minimum during winter

in all the studied fish species in the present study. No significant difference (F=0.388, p=0.682) was observed in the amount of carbohydrate among the studied three species.

The maximum values for the studied macroelements were found during monsoon and minimum during winter season. Na ranged from 60.00-139.18 mg/100g in studied fishes (Table 2). Highest amount of Na was found to be 112.33, 125.46 and 131.55 mg/100g in *S. plagiostomus*, *S. progastus* and *B. bendelisis*, respectively while it was lowest 74.00, 66.67, and 84.67 mg/100g during winter season in all the above three species. A similar range of Na was also recorded by Joshi *et al.* (2017) for five Himalayan species of *Schizothorax* (*S. niger*, *S. curvifrons*, *S. esocinus*, *S. progastus* and *S. plagiostomus*) from three locations of western Himalaya. Na is an essential component and needed for muscle and nerve conduction processes, production of adrenaline hormone, electrolyte and acid-base balancing (Pirestani *et al.*, 2009). Potassium (K), necessary in the regulation of blood pressure, cardiac diseases, kidney function, heart diseases, nerve and muscle conduction, carbohydrate metabolism, and acid-base equilibrium (WHO, 2012) was also found maximum during monsoon and minimum during the winter season in all the three studied fishes. The higher range of K concentration was recorded in the present study than the previous report of Shantosh and Sarojnalini (2018) on cobitid fishes of Manipur. Calcium (Ca) performs the function of blood clotting and bone and skeleton formation. Failure in the maintenance of extracellular calcium can increase the risk of hypertension, obesity, hyperparathyroidism, premenstrual syndrome and polycystic ovary syndrome (Mayanglambam and Chungkham, 2018). The

Table 2: Seasonal variations of macroelements in the muscle tissue of three cyprinid species.

Fishes	Min-Max	Summer	Monsoon	Winter
Na (mg/100g)				
<i>S. plagiostomus</i>	66.00-115.00	87.79±1.55 ^a	112.33±3.79 ^a	74.00±7.21 ^b
<i>S. progastus</i>	60.00-128.89	98.00±2.65 ^b	125.46±4.30 ^{ab}	66.67±5.77 ^a
<i>B. bendelisis</i>	83.00-139.18	118.67±2.31 ^c	131.55±6.74 ^b	84.67±1.53 ^b
K (mg/100g)				
<i>S. plagiostomus</i>	245.00-340.00	309.76±1.01 ^c	330.00±10.00 ^b	251.00±5.29 ^b
<i>S. progastus</i>	235.00-310.05	250.00±5.00 ^b	294.59±6.57 ^b	239.67±5.03 ^b
<i>B. bendelisis</i>	208.00-281.11	216.33±5.51 ^a	276.84±4.04 ^a	212.67±4.51 ^a
Ca (mg/100g)				
<i>S. plagiostomus</i>	310.00-430.00	363.82±9.94 ^b	417.00±11.27 ^b	349.00±9.17 ^b
<i>S. progastus</i>	256.00-356.42	314.00±5.29 ^a	355.41±1.14 ^a	268.67±12.06 ^a
<i>B. bendelisis</i>	443.00-559.32	480.33±19.50 ^c	550.98±10.36 ^c	451.33±8.02 ^c

Min-Max- Minimum-Maximum. Values are Mean±SD of three replicates. Values with different superscript in a row varied significantly ($p < 0.05$).

higher amount of Ca (417.00, 355.41 and 550.98 mg/100g) during monsoon and lower (349.00, 268.67 and 451.33 mg/100g) during winter season were observed in *S. plagiostomus*, *S. progastus* and *B. bendelisis*, respectively. The high abundance of Ca in muscle tissue during monsoon season was also reported by Ahmed and Bat (2015) in *Argyrops spinifer*. Similar findings regarding the seasonal abundance of macroelements were also recorded by Balogun and Talabi (1986) for *Katsuwonus pelamis*, Abdullahi and Abolude (2001) for *Chrysichthys nigrodigitatus*, Bayrus filamentous and *Auchenoglanis occidentals*, Abdullahi (2005) for *Malapterus electricus* and Chakraborty et al. (2014) for *Trichurus lepturus*. The present investigation indicated the presence of highest level of Ca among other studied macroelements during all the three seasons which is in agreement with the study of Jessica et al. (2015) for 55 Bangladeshi fish species, Shantosh and Sarojnalini (2018) for *Lepidocephalichthys guntea*, *Pangio pangia* and *Syncrossus berdmorei*, Mayanglambam and Chungkham (2018) for *Devario acquipinnatus*, *Glossogobius giuris*, *Hypsibarbus myitkyinae*, *Puntius chola* and *Tariqilabeo burmanicus* and Sharma and Singh (2019) for *Schizothorax richardsonii*. The abundant level of macroelements may be attributed to the fact that the body requires them in more amounts for structure and function of the body which was also noticed by Hei and Sarojnalini (2012). Also, there is no exact recommended value for macroelements in muscle tissue of fishes. Swann (2000) also reported that it is unclear that high levels of macroelements in fish tissues are harmful to fish, wildlife species and human consuming such fishes. In the present study, the amount of Na and Ca was significantly ($p < 0.001$) higher in *B. bendelisis* while K was significantly ($p < 0.01$) higher in *S.*

plagiostomus during the three seasons.

Freshwater fishes occupy the top position of the aquatic food chain. High level of metals in fish muscle tissue originates from aquatic resources (Mansour and Sidky, 2002). The accumulation of metals inside the fish body provides evidence against fish exposure to the contaminated aquatic environment. The untreated sewage waste discharged into the river can induce acute toxic effects in fishes (Bhanot and Hundal, 2019).

The muscle tissue of studied fishes showed variation in metal accumulation of Fe, Zn, Cu and Pb in different seasons. Fe was recorded maximum to be 13.26, 22.49 and 14.47mg/100gm during monsoon and minimum to be 9.67, 16.00 and 8.67mg/100g during winter season in *S. plagiostomus*, *S. progastus* and *B. bendelisis*, respectively (Table 3). The range of Fe concentration was recorded higher than those previously reported for sea bass (2.47 mg/100g) by Erkan and Ozden (2007), Nile perch fillet (1.06 mg/100g) by Kabahenda et al. (2011) and *Alestes baremoze* (1.1-3.58 mg/100g) by Kasozi et al., (2014). The present findings indicated a pretty good amount of Fe in muscle tissue of studied cyprinids which could suggest the inclusion of selected fish species in high Fe diet. Also, Fe accumulation was observed highest among the studied metals. The results were in agreement with the study of Njinkoue et al. (2016) for *Pseudotolithus typus* and *Pseudotolithus elongatus*, Jithesh and Radhakrishnan (2017) for *Diplodus annularis*, Khitouni et al. (2018) for *Trichiurus lepturus* and Danabasa et al. (2018) for *Barbus sp.* and *Cyprinus carpio*. The present findings of more Fe followed by Zn were also confirmed by the study of Cross et al. (1973), Hei and Sarojnalini (2012), Kumar et al. (2013), Durmus et al. (2017) and Sarma et al. (2018).

Zn and Cu are the essential elements which are regu-

Table 3: Seasonal variations of different metals present in the muscle tissue of three cyprinid species.

Fishes	Min-Max	Summer	Monsoon	Winter
Fe (mg/100g)				
<i>S. plagiostomus</i>	8.00-14.35	10.33±0.58 ^a	13.26±1.09 ^a	9.67±1.53 ^a
<i>S. progastus</i>	15.00-25.01	18.88±0.07 ^b	22.49±2.35 ^b	16.00±1.00 ^b
<i>B. bendelisis</i>	8.00-15.96	9.67±1.53 ^a	14.47±1.58 ^a	8.67±1.15 ^a
Zn (mg/100g)				
<i>S. plagiostomus</i>	1.50-4.55	1.667±0.21 ^a	1.549±0.01 ^a	4.21±0.50 ^a
<i>S. progastus</i>	1.70-4.11	2.833±0.32 ^a	2.40±0.61 ^a	3.997±0.14 ^a
<i>B. bendelisis</i>	1.48-4.03	2.933±0.90 ^a	1.807±0.57 ^a	3.927±0.10 ^a
Cu (mg/100g)				
<i>S. plagiostomus</i>	0.429-0.683	0.681±0.003 ^b	0.545±0.004 ^a	0.462±0.01 ^a
<i>S. progastus</i>	0.429-0.869	0.867±0.003 ^c	0.593±0.001 ^{ab}	0.447±0.02 ^a
<i>B. bendelisis</i>	0.589-0.665	0.662±0.003 ^a	0.635±0.01 ^b	0.593±0.01 ^a
Pb (mg/100g)				
<i>S. plagiostomus</i>	0.117-0.249	0.248±0.001 ^a	ND	0.118±0.001 ^a
<i>S. progastus</i>	0.311-0.441	0.435±0.01 ^b	ND	0.311±0.001 ^c
<i>B. bendelisis</i>	0.224-0.447	0.443±0.01 ^b	ND	0.226±0.002 ^b

Min-Max- Minimum-Maximum. Values are Mean±SD of three replicates. Values with different superscript in a row varied significantly ($p < 0.05$)

lated by various metabolic processes (Beltcheva *et al.*, 2011). Zinc deficiency leads to the loss of appetite, retardation of growth, skin changes, and abnormalities in immunological functions (Tuzen, 2009). Copper is an essential part of several enzymes and is essential for haemoglobin synthesis, fish growth and reproduction (Sivaperumal *et al.*, 2007) while excessive Cu causes kidney and liver damage (Tuzen, 2009; Ikem and Egiebor, 2005). In the present investigation, Zn contents were observed highest (4.210, 3.997 and 3.927 mg/100g) during winter and lowest (1.549, 2.40 and 1.807 mg/100g) during monsoon season in *S. plagiostomus*, *S. progastus* and *B. bendelisis* (Table 3). The overall amount of Zn was found to be higher in *S. progastus* than *S. plagiostomus* which was also reported by Joshi *et al.* (2017). Cu ranged from 0.429 to 0.869 mg/100g in studied three species of fishes. Cu concentration was found highest during summer (0.681, 0.867 and 0.662 mg/100g) followed by monsoon (0.545, 0.593 and 0.635 mg/100g) and winter season (0.462, 0.447 and 0.593 mg/100g) in the muscle tissue of *S. plagiostomus*, *S. progastus* and *B. bendelisis*. The concentration of Zn and Cu was observed to be significantly higher in the present study than the study of Rahman *et al.* (2018) and Romharsha and Sarojnalini (2018). Pb is non-essential and toxic metal which can be introduced into the ecosystem by natural (erosion of earth's crust and dust transport) and anthropogenic (industrial emissions, use of gasoline and phosphate fertilizers) means (Beltcheva *et al.*, 2011). Pb content was recorded

maximum (0.248, 0.435 and 0.443 mg/100g) in summer and minimum (0.118, 0.311 and 0.226 mg/100g) in winter season in *S. plagiostomus*, *S. progastus* and *B. bendelisis*. However, Pb content was not detected in fish samples collected during the monsoon season (Table 3). The average amount of Fe significantly ($F = 26.91$, $p = 7.41E-07$) varied among the three cyprinids being highest in *S. progastus* however, no significant variation was found in the average amount of Zn, Pb and Cu among the three species in the present study. The overall trend of trace metals accumulation in all the three fish species has been depicted in Table 4. The concentration of lead detected in the present study was comparatively lower than studied other metals. The present range of Pb concentration is in agreement with the study conducted on *Crossocheilus latius* (0.112-0.349 mg/100g) and *Garra lamta* (0.111-0.358 mg/100g) by Sharma and Singh (2020). Low levels of Pb were also detected in *Oreochromis niloticus*, *Oncorhynchus mykiss*, *Cyprinus carpio* and *Pelteobagrus fluvidraco* (Taweel *et al.*, 2013; Reyahikhoram *et al.*, 2016; Rajeshkumar and Li, 2018). Zn, Cu and Pb metals were detected higher in the fish muscle during summer than monsoon, which may be attributed to a higher rate of respiration in fishes during the summer season. Generally, higher temperature conditions affect the level of dissolved oxygen by lowering its concentration in the aquatic environment, which increases the metabolic rate of fishes. Thus, the fish uses more water for higher feeding on plants and grasses, which leads to the uptake of high amount of

Table 4: Season-wise trend of metal accumulation in cyprinids of Alaknanda river.

Trace metal	Abundance order
Fe	Monsoon> Summer>Winter
Zn	Winter>Summer>Monsoon
Cu	Summer>Monsoon>Winter
Pb	Summer>Winter>Monsoon

metals through diffusion or active uptake. However, low content of metals is due to less activity of fish during other seasons (Zayed *et al.*, 1994).

The findings of the present study regarding concentration of metals in the tissue of *S. plagiostomus*, *S. progastus* and *B. bendelisis* were also compared with the previous reports (Table 5). Comparison of our results with other studies demonstrates that our results of Fe concentration are higher than Sinha *et al.* (2002) while similar to Sarkar *et al.* (2017) and Joshi *et al.* (2017). Zn concentration in the present study was higher than Ali *et al.* (2017) but similar to Sinha *et al.* (2002), Sarkar *et al.* (2017), and Joshi *et al.* (2017). The concentration of Cu in the present study was higher than Sinha *et al.* (2002) and Ali and Khan (2018). However, Pb concentration was lower than the study of Ali and Khan (2018). It indicated that the concentration of metallic elements varied in water at different places. The studied metals, Zn, Cu and Pb are toxic metals,

which can cause serious hazards. Therefore, maximum level of heavy metals has been set for human consumption by various authorities of food standards (FAO, 1983; FAO and WHO, 1972; WHO, 1995; FEPA, 2003). The level of Zn, Cu and Pb were below the acceptable limits (100, 30, 2 mg/kg for Zn, Cu and Pb) as set by WHO (1995) for fish. A comparison with FAO (1983) for fish also showed that Zn, Cu and Pb were lower than the guidelines for edible fish. Comparison with FEPA (2003) also showed that Zn and Pb were under the acceptable limits (Zn 75 and Pb 2.0 mg/kg) for human consumption. This study suggests that the studied fishes were able to accumulate metals with different intensity so these may be used to monitor the heavy metal pollution of parent stream of river Ganga and its impact on daily fish consumers.

The level of metal bioconcentration in freshwater fishes may vary which may be attributed to different metal concentration in water and sediments, feeding habits, metabolism, their ecological needs and also the season in which analysis was carried out (Ibrahim and Omar, 2013). Bat (2014) analyzed the level of heavy metals in fish collected from the Black Sea coast and suggested that they are the biomonitors for metal concentration in aquatic ecosystems as they occupy different trophic positions. Thus, fishes can be considered as a valid indicator for the pollution of the aquatic environment (Shakweer, 1998). The accumulation level of

Table 5: Comparative account of trace metal concentration in the fish muscle tissue in various studies.

Species	Region	Fe	Cu	Zn	Pb	Reference
<i>Barilius bendelisis</i>	Parent stream of Ganga river	8.00-15.96	0.589-0.665	1.48-4.03	0.224-0.447	^a Present study
	Hel river, North East India	8.98	1.89	1.06	-	^a Sarkar <i>et al.</i> (2017)
	Kharkai river, Jamshedpur, India	40	2	14	2	^b Sinha <i>et al.</i> (2002)
<i>Schizothorax plagiostomus</i>	Parent stream of Ganga river, India	8.00-14.35	0.429-0.683	1.50-4.55	0.117-0.249	^a Present study
	Kabul river, Pakistan	-	-	-	20.3	^b Ali and Khan (2018)
	Swat river, Pakistan	-	1.07-2.05	7.53-12.80	-	
	Panjhora river, Pakistan	-	0.93-2.72	2.72-10.32	-	^b Ali <i>et al.</i> (2017)
<i>Schizothorax progastus</i>	Barandu river, Pakistan	-	2.02-4.05	10.17-18.00	-	
	Alaknanda river, India	11.83	1.14	2.64	-	^a Joshi <i>et al.</i> (2017)
<i>Schizothorax progastus</i>	Parent stream of Ganga river, India	15.00-25.01	0.429-0.869	1.70-4.11	0.311-0.441	^a Present study
	Alaknanda river, India	12.59	1.81	5.40	-	^a Joshi <i>et al.</i> (2017)
Permissible level of metals in fish for human consumption		-	30	30	0.5	^b FAO (1983)
		-	30	40	0.5	^b FAO/WHO (1972)
		-	30	100	2	^b WHO (1995)
		-	-	75	2.0	^b FEPA (2003)

^aConcentration in mg/100g, ^bConcentration in mg/kg. Blank cells indicate no information available.

heavy metals in the muscle tissue of studied fishes reflects less degree of water pollution by such metal elements. Therefore, it is concluded that the selected fish species of the Alaknanda river were free from pollution load and level of toxicity hence can be considered appropriate for consumption as a source of food.

Conclusion

It was concluded that besides the seasonal variations in nutrient profiling, all the selected fish species, namely, *S. plagiosomus*, *S. progastus* and *B. bendelisis* were found rich in protein and mineral elements. Bioconcentration of heavy metals, Zn (1.48-4.55 mg/100g), Cu (0.429-0.869 mg/100g), and Pb (0.117-0.447 mg/100g) in the muscle tissue of studied fishes collected from a Central Himalayan river Alaknanda at Srinagar (Garhwal) was found below the highest permissible limits as set by International food safety guidelines for fish consumption. Hence, these fishes were nutritious, safe and suitable for inclusion in daily diet. Conservation, management and stock augmentation of these fishes will be instrumental for ameliorating the malnutrition problem of the region.

ACKNOWLEDGEMENTS

Authors are thankful to the Department of Zoology, H.N.B. Garhwal University, Srinagar-Garhwal (Uttarakhand) for providing necessary facilities during the work. Authors (SS) and (SK) thank the University Grants Commission, New Delhi for providing Research Fellowship.

Conflict of interests

There is no conflict of interest regarding the publication of this article.

REFERENCES

1. Abdullahi, S.A. and Abolud, D.S. (2001). Nutrient levels of three species of Bagridae collected in two seasons and two locations of Northern Nigeria. *Bioscience Research Communications*, 13 (3): 231-235.
2. Abdullahi, S.A. (2005). Seasonal study on nutrient variability of the electric catfish (*Malapterurus electricus* Gmelin) obtained from three sites of the Nigerian Savannah. *Journal of Tropical Biosciences*, 5: 62-65.
3. Ahmed, Q. and Bat, L. (2015). Some essential macroelements in the muscles of *Argyrops spinifer* from Karachi Harbour, Pakistan. *Journal of Maritime and Marine Sciences*, 1(1): 8-12.
4. Ali, H. and Khan, E. (2018). Assessment of potentially toxic heavy metals and health risk in water, sediments, and different fish species of River Kabul, Pakistan. *Human Ecological Risk Assessment*, 24 (8): 2101-2118. 10.1080/10807039.2018.1438175
5. Ali, H., Ali, W. and Ullah, K. (2017). Bioaccumulation of Cu and Zn in *Schizothorax plagiosomus* and *Mastacembelus armatus* from river Swat, river Panjkora and river Barandu in Malakand Division, Pakistan. *Pakistan Journal of Zoology*, 49 (5): 1555-1561. 10.17582/journal.pjz/2017.49.5.1555.1561
6. AOAC. (2000). Association of Official Analytical Chemists Official Methods of Analysis (17th ed.). Hortuntzed W. (Ed), Washington. Brogstorm Academic Press, NY and London.
7. Balachandan, K. (2002). Post Harvest Technology of Fish and Fish Products. Daya Publishing House, New Delhi. pp. 1-28.
8. Balogun, A.M. and Talabi, S.O. (1986). Studies on size distribution on skipjack tuna. (*Katsuwonus pelamis*) effect on chemical composition and implications for its utilization. *Journal of Food Technology*, 21: 443-449.
9. Bat, L. (2014). Heavy metal pollution in the Black Sea. In: E. Düzgüneş, B. Öztürk, and M. Zengin (Eds.), *Turkish Fisheries in the Black Sea*. Published by Turkish Marine Research Foundation (TUDAV), Publication number: 40, Istanbul: 71-107.
10. Beltcheva, M., Metcheva, R., Peneva, V. and Marinova, M. (2011). Heavy metals in Antarctic Notothenioid fish from South Bay, Livingston Island, South Shetlands (Antarctica). *Biological Trace Element Research*, 141 (1-3): 150-158. 10.1007/s12011-010-8739-5
11. Bhanot, R. and Hundal, S.S. (2019). Acute toxic effects of untreated sewage water in *Labeo rohita* (Hamilton, 1822). *Journal of Entomology and Zoological Studies*, 7(3): 1351-1355.
12. Chakraborty, K., Joseph, D., Stephy, P.S., Chakkalalal, S.J., Joy, M. and Raola, V.K. (2014). Inter-annual variability and seasonal dynamics of amino acid, vitamin and mineral signatures of ribbon fish, *Trichiurus lepturus* (Linnaeus, 1758). *International Food Research Journal*, 21 (5): 2007-2016.
13. Cross, F.A., Hardy, L.H., Jones, N.Y. and Barber, R.T. (1973). Relation between total body weight and concentrations of manganese, iron, copper, zinc, and mercury in white muscle of bluefish (*Pomatomus saltatrix*) and a bathyl-demersal fish *Antimora rostrata*. *Journal of Fisheries Research Board of Canada*, 30 (9): 1287-1291. 10.1139/f73-208
14. Danabasa, D., Kutluyera, F., Urala, M. and Kocabas, M. (2018). Metal bioaccumulation in selected tissues of barb (*Barbus sp.*) and common carp (*Cyprinus carpio*, Linnaeus 1758) from the Keban Dam Lake, Turkey. *Toxin Reviews*, 39 (1): 78-85. 10.1080/15569543.2018.1479717
15. Day, F. (1878). Fishes of India, Vol. I & II, William Dawson and Sons Ltd., London, XX + 778 & pls.198.
16. Dubois, M., Gilles, K.A., Hamilton, J.K., Rebers, P.A. and Smith, F. (1956). Calorimetric method for determination of sugars and related substances. *Analytical Chemistry*, 28 (3): 350-356. 10.1021/ac60111a017
17. Durmus, M., Kosker, A.R., Ozoglu, Y., Aydin, M., Ucar, Y., Ayas, D. and Ozogul, F. (2017). The effects of sex and season on the metal levels and proximate composition of red mullet (*Mullus barbatus* Linnaeus 1758) caught from the Middle Black Sea. *Human and Ecological Risk Assessment*, 24 (3): 2011-2023. 10.1080/10807039.2017.1398071
18. Erkan, N. and Ozden, O. (2007). Proximate composition and mineral contents in aqua cultured sea bass (*Dicentrarchus labrax*), sea bream (*Sparus aurata*) analyzed by ICPMS. *Food Chemistry*, 102 (3): 721-725. 10.1016/j.foodchem.2006.06.004
19. FAO (2016). The State of World Fisheries and Aquaculture. Food and Agriculture Organization Rome, 200p.
20. WHO and FAO (1972). Evaluation of certain food additives and the contaminants mercury, lead and cadmium,

- Sixteenth Report of the joint FAO/WHO Expert Committee on Food Additives, WHO Technical Report, Series No. 505, FAO Nutrition Meetings Report Series No 51.
21. FEPA. (2003). Guidelines and standards for environmental pollution control in Nigeria. Federal Environmental Protection Agency (FEPA), p. 238.
 22. Folch, J., Lees, M. and Sloane-Stanley, G.H. (1954). A simple method for preparation for total pure lipids extracts from brain. *Federation Proceedings*, 13: 209.
 23. FAO. (1983). Compilation of Legal Limits for Hazardous Substances in Fish and Fishery Products. Food and Agriculture Organization (FAO), Rome, Italy.
 24. Hei, A. and Sarojnalini, C. (2012). Proximate composition, macro and micro mineral elements of some smoke-dried hill stream fishes from Manipur, India. *Nature and Science*, 10 (1): 59-65.
 25. Ibrahim, A.T.A. and Omar, H.M. (2013). Seasonal variation of heavy metals accumulation in muscles of the African Catfish *Clarias gariepinus* and in river Nile water and sediments at Assiut Governorate, Egypt. *Journal of Biology and Earth Sciences*, 3 (2): B236-B248.
 26. Ikem, A. and Egiebor, N.O. (2005). Assessment of trace elements in canned fishes (mackerel, tuna, salmon, sardines and herrings) marketed in Georgia and Alabama (United States of America). *Journal of Food Composition and Analysis*, 18: 771-787. 10.1016/j.jfca.2004.11.002
 27. Jayaram, K.C. (2002). Fundamentals of Fish Taxonomy. Narendra Publishing House, Delhi, 174p.
 28. Jessica, R.B., Shakuntala, H.T., Geoffrey, C.M., Md Abdul, W., Mostafa, A.R.H., Jette, J. and James, S. (2015). Nutrient composition of important fish species in Bangladesh and potential contribution to recommended nutrient intakes. *Journal of Food Composition and Analysis*, 42: 120-133.
 29. Jhingran, V.G. (1982). Fish and Fisheries of India. Hindustan Publishing Corporation, India, 666p.
 30. Jithesh, M. and Radhakrishnan, M.V. (2017). Seasonal variation in accumulation of metals in selected tissues of the Ribbon fish, *Trichiurus lepturus* collected from Chaliyar River, Kerala, India. *Journal of Entomology and Zoology Studies*, 5 (1): 51-56.
 31. Joshi, V., Akhtar, M.S., Sharma, P., Kushwaha, S.S., Baruh, D., Alexander, C., Pande, V. and Sarma, D. (2017). Himalayan fish manifest higher potential of quality nutrients for human health. *Journal of Aquatic Food Product Technology*, 26 (7): 843-855. 10.1080/1049885.0.2017.1340916
 32. Kabahenda, M.K., Amega, R., Okalany, E., Huske, S.M.C. and Heck, S. (2011). Protein and micronutrient composition of low-value fish products commonly marketed in the Lake Victoria Region. *World Journal of Agricultural Sciences*, 7 (5): 521-526.
 33. Kasozi, N., Degu, G.I., Asizua, D., Mukalazi, J. and Kalany, E. (2014). Proximate composition and mineral contents of Pebbly fish, *Alestes baremoze* (Joannis, 1835) filets in relation to fish size. *Uganda Journal of Agricultural Sciences*, 15 (1): 41-50.
 34. Khitouni, I.K., Mihoubi, N.B., Bouain, A. and Rebah, F.B. (2018). Seasonal variation of the chemical composition, fatty acid profiles and mineral elements of *Diplodus annularis* (Linnaeus, 1758) caught in the Tunisian coastal water. *Journal of Food and Nutrition Research*, 2 (6): 1-9. 10.12691/jfnr-2-6-7
 35. Kumar, B., Verma, V.K., Naskar, A.K., Chakraborty, P. and Shah, R. (2013). Human health hazard due to metal uptake via fish consumption from coastal and freshwater waters in Eastern India along the Bay of Bengal. *Journal of Marine Biology and Oceanography*, 2 (3): 1-7. 10.4172/2324-8661.1000115
 36. Lowry, O.H., Rosebrough, N.J., Farr, A.L. and Randall, R.J. (1951). Protein measurement with the folin-phenol reagents. *Journal of Biological Chemistry*, 193: 265-275.
 37. Mansour, S.A. and Sidky, M.M. (2002). Ecotoxicological studies. 3. Heavy metals contaminating water and fish from Fayoum Governorate, Egypt. *Food Chemistry*, 78: 15-22. 10.1016/s0308-8146(01)00197-2
 38. Moreau, M.A. and Garaway, C.J. (2018). Fish rescue us from hunger: the contribution of aquatic resources to household food security on the Rufiji River floodplain, Tanzania, East Africa. *Human Ecology*, 46: 831-848. 10.1007/s10745-018-0030-y
 39. Mayanglambam, S. and Chungkham, S. (2018). Macro and trace mineral elements of five small indigenous fishes of Manipur, India. *Journal of Fisheries and Life Sciences*, 3 (1): 1-8.
 40. Njinkoue, J.M., Gouado, I., Tchoumboungang, F., Yanga Nguegui, J.H., Ndinteh Ntantoh, D., Fomogne-Fodio, C.Y. and Schweigert, F.J. (2016). Proximate composition, mineral content and fatty acid profile of two marine fishes from Cameroonian coast: *Pseudotolithus typus* (Bleeker, 1863) and *Pseudotolithus elongatus* (Bowdich, 1825). *Nutrition and Food Science*, 16 (4): 27-31. 10.1016/j.nfs.2016.07.002
 41. Pal, J., Shukla, B.N., Maurya, A.K., Verma, H.O., Pandey, G. and Amitha. (2018). A review on role of fish in human nutrition with special emphasis to essential fatty acid. *International Journal of Fisheries and Aquatic Studies*, 6 (2): 427-430.
 42. Panwar, S., Gaur, D. and Chakrapani, G.J. (2017). Total organic carbon transport by the Alaknanda river, Garhwal Himalaya, India. *Arabian Journal of Geosciences*, 10 (207): 1-9. 10.1007/s12517-017-3003-3
 43. Pirestani, S., Ali, S.M., Barzegar, M. and Seyfabadi, S.J. (2009). Chemical compositions and minerals of some commercially important fish species from the South Caspian Sea. *International Food Research Journal*, 16: 39-44.
 44. Rahman, R., Chowdhury, M.M., Sultana, N. and Saha, B. (2018). Proximate and major mineral composition of commercially important marine fishes of Bangladesh. *IOSR Journal of Agriculture and Veterinary Science*, 11 (1): 18-25. 10.9790/2380-111021825
 45. Rajeshkumar, S. and Li, X. (2018). Bioaccumulation of heavy metals in fish species from the Meiliang Bay, Taihu Lake, China. *Toxicology Reports*, 5: 288-295. 10.1016/j.toxrep.2018.01.007
 46. Reyahi-Khoram, M., Setayesh-Shiri, F. and Cheraghi, M. (2016). Study of the heavy metals (Cd and Pb) content in the tissues of rainbow trouts from Hamedan coldwater fish farms. *Iranian Journal of Fisheries Sciences*, 15 (2): 858-869.
 47. Romharsha, H. and Sarojnalini, C. (2018). Proximate composition, total amino acids and essential mineral elements of some cyprinid fishes of Manipur, India. *Current Research in Nutrition and Food Science*, 6 (1): 157-164. 10.12944/CRNFSJ.6.1.18
 48. SANDRP (2014). South Asia Network on Dams, Rivers and People: Newsletter- Dams, Rivers and People, 13(3-5): 11.
 49. Sarkar, A.P., Basumatary, S. and Das, S. (2017). Determination of nutritional composition of some selected fish-

- es from Hel river of North-East India. *Asian Journal of Chemistry*, 29 (11): 2493-2496. 10.14233/ajchem.2017.20790
50. Sarma, D., Joshi, V., Akhtar, M.S., Ciji, A., Sharma, P., Kushwaha, S.S., Das, P. and Singh, A.K. (2018). Nutrient composition of six small indigenous fish from NEH region and their contribution potential to human nutrition. *Proceedings of the National Academy of Sciences, India Section B: Biological Sciences*, 89 (B): 1-8.
51. Shakweer, L.M. (1998). Concentration levels of some trace metals in *Oreochromis niloticus* at highly and less polluted areas of Mariut Lake. *Journal of Egyptian-German Society of Zoology*, 25: 125-144.
52. Shantosh, M. and Sarojnalini, Ch. (2018). Nutritional quality of three cobitid fishes of Manipur, India: with special reference to essential mineral elements. *International Journal of Scientific Research in Biological Sciences*, 5 (2): 24-33.
53. Sharma, S. and Singh, D. (2019). Seasonal dynamics of biochemical and mineral components of Himalayan cyprinid *Schizothorax richardsonii* (Gray, 1832). *Research Journal of Biotechnology*, 14 (9): 77-82.
54. Sharma, S. and Singh, D. (2020). Nutritional status of *Crossocheilus latius latius* (Hamilton, 1822) and *Garra lamta* (Hamilton, 1822) from Alaknanda river of Central Himalaya. *Journal of Aquatic Food Product Technology*, 29 (4):1-10. 10.1080/10498850.2020.1737999
55. Sinha, A.K., Dasgupta, P., Chakrabarty, S., Bhattacharyya, G. and Bhattacharjee, S. (2002). Bioaccumulation of heavy metals in different organs of some of the common edible fishes of Kharkai river, Jamshedpur. *Indian Journal of Environmental Health*, 44 (2): 102-107.
56. Sivaperumal, P., Sankar, T.V. and Nair Viswanathan, P.G. (2007). Heavy metals concentrations in fish, Shellfish and fish products from internal markets of India vis-a-vis international standards. *Food Chemistry*, 102: 612-620. 10.1016/j.foodchem.2006.05.041
57. Swann, L. (2000). A fish farmer's guide to understanding water quality. Aquaculture Extension Illinois-Indiana: Sea Grant Program, Purdue University, West Lafayette, IN.
58. Talwar, P.K. and Jhingran, A.G. (1991). Inland Fishes of India and Adjacent Countries. Vol. 1 & 2. Oxford & I.B.H. Publishing Co. Pvt. Ltd., New Delhi, 1158p.
59. Taweel, A., Shuhaimi-Othman, M. and Ahmad, A.K. (2013). Assessment of heavy metals in tilapia fish (*Oreochromis niloticus*) from the Langat river and Engineering lake in Bangi, Malaysia, and evaluation of health risk from tilapia consumption. *Ecotoxicology and Environmental Safety*, 93: 45-51. 10.1016/j.ecoenv.2013.03.031
60. Tuzen, M. (2009). Toxic and essential trace elemental contents in fish species from the Black Sea, Turkey. *Food and Chemical Toxicology*, 7 (8): 1785-1790. 10.1016/j.fct.2009.04.029
61. WHO (1995). Environmental Health Criteria No 165: Inorganic Lead. Geneva (Switzerland): World Health Organization (WHO). <http://www.inchem.org/documents/ehc/ehc/ehc165.htm>
62. WHO (2012). Guideline of potassium intake for adult and children. World Health Organization, 10.
63. Zayed, M.A., Nour El-Dien, F.A. and Rabie, K.A. (1994). Comparative study of seasonal variation in metal concentrations in river Nile sediment, fish and water by atomic absorption spectrophotometry. *Microchemical Journal*, 49: 27-35.