

UDC 519.21

B. ROYNETTE, P. VALLOIS, AND M. YOR

**PENALISATIONS OF BROWNIAN MOTION WITH  
ITS MAXIMUM AND MINIMUM PROCESSES AS  
WEAK FORMS OF SKOROKHOD EMBEDDING**

We develop a Brownian penalisation procedure related to weight processes  $(F_t)$  of the type:  $F_t := f(I_t, S_t)$  where  $f$  is a bounded function with compact support and  $S_t$  (resp.  $I_t$ ) is the one-sided maximum (resp. minimum) of the Brownian motion up to time  $t$ . Two main cases are treated: either  $F_t$  is the indicator function of  $\{I_t \geq \alpha, S_t \leq \beta\}$  or  $F_t$  is null when  $\{S_t - I_t > c\}$  for some  $c > 0$ . Then we apply these results to some kind of asymptotic Skorokhod embedding problem.

1. INTRODUCTION

In a series of papers [18], [17], [16], [23], [24], [21], [22] (see also the surveys [19] and [20] and the monograph [25]) we have introduced a penalisation procedure of Brownian paths and applied it to many settings. To present the aim of this paper, we first introduce a few notations.

Let  $(\Omega = \mathcal{C}(\mathbb{R}_+, \mathbb{R}), (X_t)_{t \geq 0}, (\mathcal{F}_t)_{t \geq 0})$  be the canonical space, where  $(X_t)_{t \geq 0}$  denotes the coordinate maps:  $X_t(\omega) = \omega(t)$ , for any  $t \geq 0$ . Let  $(P_x)_{x \in \mathbb{R}}$  be the family of Wiener probability measures on  $\Omega$ : under  $P_x$ ,  $(X_t)_{t \geq 0}$  is a standard one-dimensional Brownian motion started at  $x$ .

Next, we consider a stochastic process  $(F_t)_{t \geq 0}$  which takes its values in  $[0, \infty[$  and satisfies:

$$(1.1) \quad 0 < E_0(F_t) < \infty \quad \forall t \geq 0.$$

We shall say that the penalisation procedure (associated with the weight process  $(F_t)$ ) holds if:

$$(1.2) \quad Q_0^F(\Lambda_s) := \lim_{t \rightarrow \infty} \frac{E_0[1_{\Lambda_s} F_t]}{E_0[F_t]} \text{ exists for any } s \geq 0 \text{ and } \Lambda_s \in \mathcal{F}_s.$$

We briefly recall (see Theorem 3.6 in [17], [16] and [23]) that the penalisation procedure holds with  $F_t = \varphi(S_t)$  where:

1.  $\varphi : [0, \infty[ \rightarrow [0, \infty[$  is a Borel function such that  $\int_0^\infty \varphi(x) dx = 1$ ;
2.  $(S_t)_{t \geq 0}$  is the one-sided maximum process associated with  $(X_t)_{t \geq 0}$ :

$$(1.3) \quad S_t := \max_{u \leq t} X_u; \quad t \geq 0.$$

According to Theorem 4.6 of [17], under  $Q_0^F$ , the r.v.  $S_\infty$  is finite and admits  $\varphi$  as a density function. Consequently, the procedure (1.2) forces Brownian motion to have a

---

2000 *AMS Mathematics Subject Classification*. Primary 60G17, 60G40, 60G44, 60H10, 60J25, 60J60, 60J65.

*Key words and phrases*. Skorokhod's problem, penalisation, one-sided maximum and minimum, Laplace's method.

finite one-sided total maximum with the given probability density  $\varphi$ . This result presents some analogy with Skorokhod's embedding problem.

**1.1. On Skorokhod's problem for linear Brownian motion.** Let  $\mu$  be a probability measure (p.m.) on  $\mathbb{R}$ , such that:

$$(1.4) \quad \int_{\mathbb{R}} |y| \mu(dy) < \infty \quad \text{and} \quad \int_{\mathbb{R}} y \mu(dy) = 0.$$

A number of constructions of (finite) stopping times  $T$  such that (i)  $T$  is standard, i.e.  $(X_{s \wedge T}; s \geq 0)$  is a uniformly integrable martingale, and (ii) the distribution of  $X_T$ , under  $P_0$  is  $\mu$ , have been made by many authors, see e.g. Oblój's thorough survey [11] of the subject. We briefly recall the particular construction given by Azéma and Yor ([2], [1]): there exists a non-decreasing function  $\phi_\mu : [0, +\infty[ \rightarrow \mathbb{R}$  such that for :

$$(1.5) \quad T_\mu := \inf \{t \geq 0, \quad X_t \leq \phi_\mu(S_t)\},$$

$T_\mu$  is standard and  $X_{T_\mu} \sim \mu$  (under  $P_0$ ).

Precisely,  $\phi_\mu$  is the right-continuous inverse of  $\psi_\mu$  where

$$\psi_\mu(x) = \frac{1}{\mu(]x, \infty[)} \int_{]x, \infty[} y d\mu(y).$$

**1.2. An asymptotic resolution of Skorokhod's problem for diffusions.** Similarly to the Skorokhod embedding problem for Brownian motion, let us start with a target p.m.  $\mu$  on  $\mathbb{R}$ . For simplicity we suppose that  $\mu$  admits a density function  $\mu_0$  which is supposed to be positive and of class  $C_b^1$  :  $\mu(dx) = \mu_0(x)dx$ .

Then, there exists a p.m.  $\mathbb{Q}_0$  on  $(\Omega, \mathcal{F}_\infty)$  such that :

$$(1.6) \quad X_t = B_t + \frac{1}{2} \int_0^t \frac{\mu'_0(X_s)}{\mu_0(X_s)} ds, \quad t \geq 0$$

and  $(B_t)_{t \geq 0}$  is a  $\mathbb{Q}_0$ -Brownian motion started at 0.

Moreover, under  $\mathbb{Q}_0$ ,  $X_t$  converges in distribution to  $\mu$ , as  $t \rightarrow \infty$ . In other words, we have introduced a diffusion process, whose limit distribution is the given p.m.  $\mu$ , which may be considered as an asymptotic kind of resolution of Skorokhod's problem.

**1.3. Solving Skorokhod's problem for  $(S_t, X_t)$ .** The solution of Skorokhod's problem given by Azéma and Yor [2] suggested to consider the embedding problem for the two-dimensional process  $(S_t, X_t)_{t \geq 0}$ . A complete answer has been given by Rogers [13] (see also [15]). Starting with a p.m.  $\nu$  on  $\mathbb{R}_+ \times \mathbb{R}$ , assumed to satisfy:

$$(1.7) \quad \int_{\mathbb{R}_+ \times \mathbb{R}} |y| \nu(dx, dy) < \infty, \quad \int_{\mathbb{R}_+ \times \mathbb{R}} y \nu(dx, dy) = 0$$

$$(1.8) \quad a \nu([a, \infty[ \times \mathbb{R}) = \int_{\mathbb{R}_+ \times \mathbb{R}} \mathbf{1}_{\{x \geq a\}} y \nu(dx, dy), \quad \forall a \geq 0,$$

it is shown that there exists a finite standard stopping time  $T$  such that  $(S_T, X_T) \sim \nu$  (under  $P_0$ ).

Note that (1.7) and (1.8) correspond to

$$(1.9) \quad E[|X_T|] < \infty, \quad E[X_T] = 0,$$

resp.

$$(1.10) \quad aP(S_T \geq a) = E[X_T | S_T \geq a], \quad \forall a \geq 0.$$

**1.4. The  $(S_t, X_t)$  asymptotic resolution of Skorokhod's problem for diffusion processes.** Comparing Subsections 1.2 and 1.3, we may ask the following question : for which class of p.m.'s  $\nu$  on  $\mathbb{R}_+ \times \mathbb{R}$ , does there exist a p.m.  $\mathbb{Q}_1$  on  $(\Omega, \mathcal{F}_\infty)$  under which:

$$(1.11) \quad X_t = B_t + \int_0^t b(u, X_u) du$$

where

$$(1.12) \quad (B_t)_{t \geq 0} \text{ is a } \mathbb{Q}_1\text{-standard Brownian motion with, } B_0 = 0,$$

$$(1.13) \quad (b(t, X_t))_{t \geq 0} \text{ is an } (\mathcal{F}_t)\text{-adapted process,}$$

$$(1.14) \quad \text{and the couple } (S_t, X_t) \text{ converges in distribution towards } \nu, \text{ as } t \rightarrow \infty.$$

Our approach is based on a Brownian penalisation procedure. This method is well fitted for our purpose since it permits to obtain Markov processes whose distributions are locally equivalent to that of Brownian motion and enjoy new path properties (for instance a finite total unilateral maximum, see the beginning of Introduction).

Unfortunately we have not been able to solve completely this question. In Section 6, we only give a class of p.m.'s  $\nu$  verifying (1.11)-(1.14).

**1.5. Organisation of the paper.** Section 2 is a short survey of Brownian penalisations. To show that the penalisation procedure (see (1.2) or Section 2 for more details) associated with a weight process  $(F_t)$  holds, we need to be able to determine the rates of decay of  $E_0[F_t]$  and  $E[F_t | \mathcal{F}_s]$  as  $t \rightarrow \infty$ . In this paper, we consider penalisations with  $F_t = f(I_t, S_t)$  where  $f : ]-\infty, 0] \times [0, \infty[ \rightarrow ]0, \infty[$  is a Borel bounded function with compact support and

$$I_t := \inf_{u \leq t} X_u \quad (t \geq 0).$$

To determine the rate of decay of  $t \mapsto E[f(I_t, S_t)]$  as  $t \rightarrow \infty$ , we have been led to consider two classes of functions  $f$ . These developments are given in Section 3. With these estimates at hand, we shall show in Section 4 that the associated penalisation procedures hold. These schemes give rise to two families of p.m. on the canonical space  $(\Omega, \mathcal{F}_\infty)$ . In Section 5 we determine the law of the process  $(X_t)$  under each of these new p.m.'s. Finally, in Section 6 we apply our results to the question discussed above in subsection 1.4.

## 2. A SURVEY OF BROWNIAN PENALISATIONS

We keep notations from the Introduction.

**2.1. The goal.** Penalisations provide a method to define  $P_0(\cdot | A)$  for  $A$  a negligible event in  $\mathcal{F}_\infty := \bigvee_{t \geq 0} \mathcal{F}_t$ , i.e.  $P_0(A) = 0$ .

This question arises naturally in probability theory and especially in the study of stochastic processes. Let us give a few explicit examples :

$$(2.1) \quad 1) A_1 = \{X_t \geq 0; \forall t \geq 0\}$$

$$(2.2) \quad 2) A_2 = \{\sup_{t \geq 0} X_t \leq a\}, \quad (a > 0)$$

$$(2.3) \quad 3) A_3 = \{\inf_{t \geq 0} X_t \geq \alpha, \sup_{t \geq 0} X_t \leq \beta\} \quad (\alpha < 0, \beta > 0).$$

Conditioning by  $A_1$  may be treated by  $h$ -Doob's transforms (see for instance Section 4, Chap. V in [4]). As for (2.2), it is proved in [9], [10] that for any  $0 < t_1 < \dots < t_n$ , the

conditional distribution of the random vector  $(X_{t_1}, \dots, X_{t_n})$  given  $\{S_t \leq a\}$  converges as  $t \rightarrow \infty$ .

The third case is the subject of our study. It is then demanded that  $(X_t)_{t \geq 0}$  stays in the strip  $[\alpha, \beta]$ .

**2.2. A solution via approximation.** Given a decreasing family  $(A_t)_{t \geq 0}$  of events in  $\mathcal{F}_\infty$  such that  $P_0(A_t) > 0$ ,  $\forall t \geq 0$ , we set  $A = \bigcap_{t \geq 0} A_t$ .

As an example, the set  $A_3$  given by (2.3) satisfies :  $A_3 = \bigcap_{t \geq 0} A_{3,t}$ , with  $A_{3,t} := \{I_t \geq \alpha, S_t \leq \beta\}$ , where  $(I_t)_{t \geq 0}$  is the one-sided minimum process, i.e.

$$(2.4) \quad I_t := \inf_{u \leq t} X_u, \quad t \geq 0.$$

Going back to the the general case of the family  $(A_t)_{t \geq 0}$ , we would like to define :

$$(2.5) \quad P_0^A(\Lambda) := \lim_{t \rightarrow \infty} P_0(\Lambda | A_t),$$

for  $\Lambda \in \mathcal{F}_\infty$  such that the limit exists.

At this stage, three questions arise immediately :

$$(2.6) \quad 1) \text{ for which } \Lambda, \text{ does } P_0^A(\Lambda) \text{ exist ?}$$

$$(2.7) \quad 2) \text{ Can } P_0^A \text{ be extended to a p.m. on } (\Omega, \mathcal{F}_\infty) \text{ ?}$$

$$(2.8) \quad 3) \text{ How does } P_0^A \text{ depend on the family } (A_t)_{t \geq 0} \text{ ?}$$

**2.3. A penalisation procedure.** It is actually easier to generalize the previous approach by replacing  $(A_t)_{t \geq 0}$  by a stochastic process  $(F_t)_{t \geq 0}$  which takes its values in  $[0, \infty[$  and satisfies (1.1).

Our penalisation procedure is the following : the assumptions of the next theorem have been shown to be satisfied for a large number of such weight processes  $(F_t)_{t \geq 0}$ ; see [18], [17], [16], [23], [24] and [22].

**Theorem 2.1.** *Let  $(F_t)_{t \geq 0}$  as above. Assume that:*

$$(2.9) \quad \frac{E_0[F_t | \mathcal{F}_s]}{E_0(F_t)} \xrightarrow[t \rightarrow \infty]{a.s.} M_s^F \quad \forall s \geq 0$$

and

$$(2.10) \quad E_0(M_s^F) = 1 \quad \forall s \geq 0.$$

Then:

1.  $(M_s^F ; s \geq 0)$  is a non-negative  $P_0$ -martingale.
2. For any  $s \geq 0$  and  $\Lambda_s \in \mathcal{F}_s$  :

$$\lim_{t \rightarrow \infty} \frac{E_0[1_{\Lambda_s} F_t]}{E_0[F_t]} = Q_0^F(\Lambda_s)$$

and

$$Q_0^F(\Lambda_s) = E_0[1_{\Lambda_s} M_s^F].$$

3.  $Q_0^F$  extends as a p.m. on  $(\Omega, \mathcal{F}_\infty)$ .

Note that if we choose for the weight process  $(F_t)_{t \geq 0} : F_t = 1_{A_t}$ , where  $(A_t)_{t \geq 0}$  satisfies (??)-(??), then assuming that (2.9) and (2.10) hold in this framework, we get:

$$Q_0^F(\Lambda_s) = \lim_{t \rightarrow \infty} P_0(\Lambda_s | A_t).$$

Consequently  $Q_0^F$  agrees with  $P_0^A$ , as tentatively defined by (2.5). Moreover a solution to the questions (2.6) and (2.7) has been given.

### 3. PRELIMINARY RESULTS

In subsection 7 below we shall study in a general framework the asymptotic behavior of  $t \mapsto E[f(I_t, S_t)]$  as  $t \rightarrow \infty$ , where  $f : ] - \infty, 0] \times [0, \infty[ \rightarrow [0, \infty[$  is a bounded function with compact support. However, in order to obtain an explicit rate of convergence, we will need to impose some restrictions on  $f$ , see subsection 3.2.

**3.1. A general result.** For any Borel function  $f : ] - \infty, 0] \times [0, \infty[ \rightarrow [0, \infty[$  with compact support, let us define :

$$(3.1) \quad K_f = \sup\{\beta - \alpha ; f(\alpha, \beta) > 0\}.$$

This means that the support of  $f$  is included in the triangle with vertices  $(-K_f, 0)$ ,  $(0, 0)$  and  $(0, K_f)$ .

Let us state the main result of this subsection.

**Proposition 3.1.** *Let  $f : ] - \infty, 0] \times [0, \infty[ \rightarrow [0, \infty[$ , bounded with compact support, then*

$$(3.2) \quad E[f(I_t, S_t)] = \Delta_t(fa_0^{(t)}) + t\Delta_t(fa_1^{(t)}) + t^2\Delta_t(fa_2) + R_t(f) ; t \geq 1$$

where

1.  $\Delta_t$  is the linear operator acting on functions  $g : ] - \infty, 0] \times [0, \infty[ \rightarrow [0, \infty[$  :

$$(3.3) \quad \Delta_t(g) := \int_{]-\infty, 0] \times [0, \infty[} \frac{g(\alpha, \beta)}{(\beta - \alpha)^6} \exp\left\{-\frac{\pi^2 t}{2(\beta - \alpha)^2}\right\} d\alpha d\beta$$

2.  $a_0^{(t)}$  and  $a_1^{(t)}$  are two continuous functions defined on  $] - \infty, 0] \times [0, \infty[$  satisfying:

$$(3.4) \quad |a_i^{(t)}(\alpha, \beta)| \leq C(1 + K_f^4) \quad i = 0, 1$$

and

$$(3.5) \quad a_2(\alpha, \beta) := 4\pi^3 \sin\left(\frac{\pi\beta}{\beta - \alpha}\right).$$

3.  $R_t(f)$  is a remainder term, which satisfies :

$$(3.6) \quad |R_t(f)| \leq C(1 + K_f^4) \left(\sup_{\alpha, \beta} f(\alpha, \beta)\right) \exp\left\{-\frac{8\pi^2 t}{K_f^2}\right\} ; t \geq 1.$$

To prove Proposition 3.1, three Lemmas are required.

**Lemma 3.2.** *For any  $t > 0$ ,  $\alpha < 0$  and  $\beta > 0$ , we have :*

$$(3.7) \quad P_0(I_t > \alpha, S_t < \beta) = \frac{4}{\pi} \sum_{k \geq 0} \frac{1}{2k+1} \sin\left(\frac{(2k+1)\pi\beta}{\beta - \alpha}\right) \exp\left\{-\frac{(2k+1)^2\pi^2}{2(\beta - \alpha)^2}t\right\}$$

*Proof.* According to ([3] section 11 of chap 2 ; [12], ex. 3.15, chap. III) we have:

$$(3.8) \quad P_0(I_t > \alpha, S_t < \beta, X_t \in dx) = \int_{[\alpha, \beta]}(x)dx \times \sum_{k \in \mathbb{Z}} p_t(x + 2k(\beta - \alpha)) - p_t(2\beta - x + 2k(\beta - \alpha))$$

where  $p_t(x)$  is the density function of the Gaussian distribution with mean 0 and variance  $t$  :

$$(3.9) \quad p_t(x) = \frac{1}{\sqrt{2\pi t}} \exp\left\{-\frac{x^2}{2t}\right\}.$$

Using the Poisson summation formula (see for instance [7], Chap. XIX, p. 630) we get :

$$\begin{aligned} & \sum_{k \in \mathbb{Z}} p_t(x + 2k(\beta - \alpha)) - p_t(2\beta - x + 2k(\beta - \alpha)) \\ &= \frac{1}{\beta - \alpha} \sum_{k \geq 1} \left[ \cos\left(\frac{k\pi x}{\beta - \alpha}\right) - \cos\left(\frac{k\pi(2\beta - x)}{\beta - \alpha}\right) \right] \exp\left\{-\frac{k^2\pi^2 t}{2(\beta - \alpha)^2}\right\} \end{aligned}$$

Since  $\cos a - \cos b = -2 \sin\left(\frac{a+b}{2}\right) \sin\left(\frac{a-b}{2}\right)$ , we obtain:

$$(3.10) \quad \begin{aligned} P_0(I_t > \alpha, S_t < \beta, X_t \in dx) &= 1_{[\alpha, \beta]}(x) dx \\ &\times \frac{2}{\beta - \alpha} \sum_{k \geq 1} \sin\left(\frac{k\pi\beta}{\beta - \alpha}\right) \sin\left(\frac{k\pi(\beta - x)}{\beta - \alpha}\right) \exp\left\{-\frac{k^2\pi^2 t}{2(\beta - \alpha)^2}\right\}. \end{aligned}$$

Integrating (3.10) over  $[\alpha, \beta]$  we easily obtain the announced result.  $\blacksquare$

**Lemma 3.3.** *Let  $h_1 : [0, \infty[\times] - \infty, 0]\times]0, \infty[ \rightarrow \mathbb{R}$  be the function:*

$$(3.11) \quad h_1(t, \alpha, \beta) := \frac{4}{\pi} \sin\left(\frac{\pi\beta}{\beta - \alpha}\right) \exp\left\{-\frac{\pi^2 t}{2(\beta - \alpha)^2}\right\}.$$

Then

$$(3.12) \quad \frac{\partial^2 h_1}{\partial \alpha \partial \beta}(t, \alpha, \beta) = \left(b_0(t, \alpha, \beta) + b_1(t, \alpha, \beta)t + b_2(\alpha, \beta)t^2\right) \exp\left\{-\frac{\pi^2 t}{2(\beta - \alpha)^2}\right\}$$

where  $b_0(t, \cdot)$  and  $b_1(t, \cdot)$  are of  $C^\infty$  class (in the variables  $\alpha$  and  $\beta$ ) except at 0, and:

$$(3.13) \quad |b_i(t, \alpha, \beta)| \leq \frac{C}{(\beta - \alpha)^6} (1 + (\beta - \alpha)^4), \quad i = 0, 1$$

and

$$(3.14) \quad b_2(\alpha, \beta) := -\frac{4\pi^3}{(\beta - \alpha)^6} \sin\left(\frac{\pi\beta}{\beta - \alpha}\right)$$

*Proof.* We first observe that  $h_1(t, \alpha, \beta)$  is the term which is obtained by taking  $k = 0$  in the series (3.7).

We begin with the  $\alpha$ -partial derivative of  $h_1$ :

$$\frac{\partial h_1}{\partial \alpha}(t, \alpha, \beta) = \frac{4}{\pi} \left[ \frac{\pi\beta}{(\beta - \alpha)^2} \cos\left(\frac{\pi\beta}{\beta - \alpha}\right) - \frac{\pi^2 t}{(\beta - \alpha)^3} \sin\left(\frac{\pi\beta}{\beta - \alpha}\right) \right] \exp\left\{-\frac{\pi^2 t}{2(\beta - \alpha)^2}\right\}.$$

Taking the  $\beta$ -partial derivative in the above expression, it is clear that (3.12) holds. Then (3.14) and (3.13) follow after straightforward calculations and estimates.  $\blacksquare$

**Lemma 3.4.** *Let  $h_2$  be the function:*

$$(3.15) \quad h_2(t, \alpha, \beta) := \frac{4}{\pi} \sum_{k \geq 2} \frac{1}{2k+1} \sin\left(\frac{(2k+1)\pi\beta}{\beta - \alpha}\right) \exp\left\{-\frac{(2k+1)^2\pi^2 t}{2(\beta - \alpha)^2}\right\}.$$

Then

$$(3.16) \quad \left| \frac{\partial^2 h_2}{\partial \alpha \partial \beta}(t, \alpha, \beta) \right| \leq \frac{C(1 + (\beta - \alpha)^4)}{t} \exp\left\{-\frac{8\pi^2 t}{(\beta - \alpha)^2}\right\}.$$

*Proof.* We proceed as in the proof of Lemma 3.3:

$$\left| \frac{\partial^2 h_2}{\partial \alpha \partial \beta}(t, \alpha, \beta) \right| \leq C t^2 (1 + (\beta - \alpha)^4) \left( \sum_{k \geq 2} \frac{(2k+1)^3}{(\beta - \alpha)^6} \exp\left\{-\frac{(2k+1)^2\pi^2 t}{2(\beta - \alpha)^2}\right\} \right)$$

Since  $k \geq 2$ , we have :

$$(2k + 1)^2 = (2k - 3 + 4)^2 \geq (2k - 3)^2 + 16$$

As a result:

$$\left| \frac{\partial^2 h_2}{\partial \alpha \partial \beta}(t, \alpha, \beta) \right| \leq C t^2 (1 + (\beta - \alpha)^4)$$

$$\left( \sum_{k \geq 2} \frac{(2k + 1)^3}{(\beta - \alpha)^6} \exp \left\{ - \frac{(2k - 3)^2 \pi^2 t}{2(\beta - \alpha)^2} \right\} \right) \times \exp \left\{ - \frac{8\pi^2 t}{(\beta - \alpha)^2} \right\}.$$

Let  $C := \sup_{x \geq 0} x^3 e^{-x}$ . Then:

$$x^3 e^{-ax} \leq \frac{C}{a^3}, \quad \forall x \geq 0.$$

Taking  $x = \frac{1}{(\beta - \alpha)^2}$  and  $a = \frac{(2k - 3)^2 \pi^2 t}{2}$  in the above inequality, we get:

$$\frac{1}{(\beta - \alpha)^6} \exp \left\{ - \frac{(2k - 3)^2 \pi^2 t}{2(\beta - \alpha)^2} \right\} \leq \frac{C}{(2k - 3)^6 t^3}.$$

This implies (3.16). ■

*Proof of Proposition 3.1.* It is clear that the definition of  $h_1(t, \alpha, \beta)$  (resp.  $h_2(t, \alpha, \beta)$ ) given by (3.11) (resp. (3.15)) implies that

$$P(I_t > \alpha, S_t < \beta) = h_1(t, \alpha, \beta) + h_2(t, \alpha, \beta), \quad \alpha < 0, \beta > 0.$$

Consequently:

$$E_0[f(I_t, S_t)] = - \int_{]-\infty, 0] \times ]0, \infty[} f(\alpha, \beta) \frac{\partial^2 h_1}{\partial \alpha \partial \beta}(t, \alpha, \beta) d\alpha d\beta + R_t(f)$$

where

$$R_t(f) := - \int_{]-\infty, 0] \times ]0, \infty[} f(\alpha, \beta) \frac{\partial^2 h_2}{\partial \alpha \partial \beta}(t, \alpha, \beta) d\alpha d\beta.$$

It is obvious that (3.16) implies (3.6).

From (3.13) and (3.14), we may deduce:

$$E_0[f(I_t, S_t)] = - \sum_{i=0}^1 t^i \int_{]-\infty, 0] \times ]0, \infty[} b_i(t, \alpha, \beta) (\beta - \alpha)^6 \frac{f(\alpha, \beta)}{(\beta - \alpha)^6} \exp \left\{ - \frac{\pi^2 t}{(\beta - \alpha)^2} \right\} d\alpha d\beta$$

$$+ t^2 \int_{]-\infty, 0] \times ]0, \infty[} b_2(\alpha, \beta) (\beta - \alpha)^6 \frac{f(\alpha, \beta)}{(\beta - \alpha)^6} \exp \left\{ - \frac{\pi^2 t}{2(\beta - \alpha)^2} \right\} d\alpha d\beta + R_t(f).$$

Setting:

$$a_i^{(t)}(\alpha, \beta) = -(\beta - \alpha)^6 b_i(t, \alpha, \beta) \quad (i = 0, 1)$$

leads to (3.2).

Obviously, (3.4) is a consequence of (3.13). □

**3.2. Applications.** Let  $f : ]-\infty, 0] \times [0, \infty[ \rightarrow [0, \infty[$  be a bounded function with compact support. Recall that  $K_f$  has been defined by (3.1). This quantity and the set:

$$\mathcal{S}_f := \{\beta \in ]0, K_f[; f(\beta - K_f, \beta) > 0\}$$

will play an important role in our study.

One aim of our paper is to show that the penalisation procedure holds with the weight process  $F_t := f(I_t, S_t)$ . We briefly detail our approach. Formula (2.9) shows that it is natural to first investigate the asymptotic behavior of  $t \mapsto E[f(I_t, S_t)]$ , as  $t \rightarrow \infty$ . Roughly speaking (3.2) tells us that the dominant term is  $\Delta_t(fa_2)$ . We observe that:

$$(3.17) \quad \sup_{\alpha < 0, \beta > 0, f(\alpha, \beta) > 0} \exp \left\{ -\frac{\pi^2 t}{2(\beta - \alpha)^2} \right\} = \exp \left\{ -\frac{\pi^2 t}{2K_f^2} \right\},$$

and the above maximum is achieved at any point of the type  $(\beta - K_f, \beta)$  where  $\beta \in \mathcal{S}_f$ .

In this general setting, it seems difficult to obtain an equivalent for  $E[f(I_t, S_t)]$  as  $t \rightarrow \infty$ . This led us to consider two extreme cases: either  $\mathcal{S}_f$  reduces to a single point or  $\mathcal{S}_f = ]0, K_f[$ . The two corresponding prototypes of functions  $f$  are either:

$$(3.18) \quad f(\alpha, \beta) = 1_{\{\alpha \geq \alpha_0, \beta \leq \beta_0\}} \quad \text{where } \alpha_0 < 0, \beta_0 > 0$$

or

$$(3.19) \quad f(\alpha, \beta) = 1_{\{\beta - \alpha \leq c\}} \quad \text{where } c > 0.$$

More generally we have been able to deal with the two following cases.

**Case 1.**

$$(3.20) \quad f(\alpha, \beta) = \Phi(\alpha, \beta) 1_{\{\alpha \geq \alpha_0, \beta \leq \beta_0\}}$$

where  $\alpha_0 < 0, \beta_0 > 0$  and  $\Phi : [\alpha_0, 0] \times [0, \beta_0] \rightarrow \mathbb{R}_+$  is continuous and

$$(3.21) \quad \Phi(\alpha_0, \beta_0) > 0.$$

Note that then  $K_f = \beta_0 - \alpha_0, \mathcal{S}_f = \{\beta_0\}$ , and  $(\alpha_0, \beta_0)$  is the unique point which achieves the maximum in (3.17). It seems reasonable to believe that:

$$(3.22) \quad E[f(I_t, S_t)] \underset{t \rightarrow \infty}{\sim} C_{\alpha_0, \beta_0} \Phi(\alpha_0, \beta_0) \exp \left\{ -\frac{\pi^2 t}{2(\beta_0 - \alpha_0)^2} \right\}.$$

**Case 2.**

$$(3.23) \quad f(\alpha, \beta) = \Phi(\alpha, \beta) 1_{\{\beta - \alpha \leq c\}}$$

where  $c > 0, \Phi : \{(\alpha, \beta); \alpha < 0, \beta > 0, \beta - \alpha \leq c\} \rightarrow \mathbb{R}_+$  is continuous and

$$(3.24) \quad \int_0^c \Phi(\beta - c, \beta) d\beta > 0.$$

In this case  $K_f = c$  and  $\mathcal{S}_f = \{\beta \in ]0, c[; \Phi(\beta - c, \beta) > 0\}$ . Therefore the maximum in (3.17) is achieved at any  $(\beta - c, \beta)$ , where  $\beta$  belongs to  $]0, c[$ . The expected behavior would be:

$$(3.25) \quad E[f(I_t, S_t)] \underset{t \rightarrow \infty}{\sim} C(\Phi, t) \exp \left\{ -\frac{\pi^2 t}{2c^2} \right\},$$

where  $t \mapsto C(\Phi, t)$  has a polynomial rate of decay in  $t$  and depends on all the values of  $\Phi$  over the segment in  $\mathbb{R}^2$  with endpoints  $(-c, 0)$  and  $(0, c)$ .

These heuristic arguments will be justified in the remainder of the section.



**Proposition 3.5.** 1. In Case 1, we have:

$$(3.26) \quad E[f(I_t, S_t)] \underset{t \rightarrow \infty}{\sim} \frac{8}{\pi} \Phi(\alpha_0, \beta_0) \sin\left(\frac{\pi\beta_0}{\beta_0 - \alpha_0}\right) \exp\left\{-\frac{\pi^2 t}{2(\beta_0 - \alpha_0)^2}\right\}$$

2. In Case 2, we have:

$$(3.27) \quad E[f(I_t, S_t)] \underset{t \rightarrow \infty}{\sim} \frac{4\pi}{c^2} \left( \int_0^1 \Phi(c(r-1), cr) \sin(\pi r) dr \right) t \exp\left\{\frac{-\pi^2 t}{2c^2}\right\}.$$

*Remark 3.6.* Suppose that  $\varphi : [0, \infty[ \rightarrow [0, \infty[$  is a Borel function such that  $\int_0^\infty \varphi(y) dy < \infty$ . Note that the rate of decay of  $t \mapsto E[\varphi(S_t)]$  as  $t \rightarrow \infty$  is very different from that of  $E[f(I_t, S_t)]$ . Indeed, (Lemma 3.8 of [17]) it is easy to prove that:

$$E[\varphi(S_t)] \underset{t \rightarrow \infty}{\sim} \sqrt{\frac{2}{\pi}} \int_0^\infty \varphi(y) dy \frac{1}{\sqrt{t}}.$$

Our proof of Proposition 3.5 requires 3 steps. First, we specify the  $\Delta_t^i(f)$  introduced in Proposition 3.1, in Cases 1 and 2.

**Lemma 3.7.** Let  $a : ]-\infty, 0] \times [0, \infty[ \rightarrow \mathbb{R}$  be a continuous function. Assume that  $f$  is a function which satisfies either (3.20) or (3.23). Then

$$(3.28) \quad \Delta_t(fa) = \frac{1}{2} \int_{1/K_f^2}^\infty \Lambda(\Phi a)(x) x \exp\left\{-\frac{\pi^2 t x}{2}\right\} dx$$

where

$$(3.29) \quad \Lambda(g)(x) = \begin{cases} \int_{(1+\alpha_0\sqrt{x})_+}^{\inf\{1, \beta_0\sqrt{x}\}} g\left(\frac{r-1}{\sqrt{x}}, \frac{r}{\sqrt{x}}\right) & \text{in Case 1} \\ \int_0^1 g\left(\frac{r-1}{\sqrt{x}}, \frac{r}{\sqrt{x}}\right) dr & \text{in Case 2} \end{cases}$$

*Proof.* We only consider Case 2. According to (3.3) we have

$$\Delta_t(fa) = \int_{]-\infty, 0] \times [0, \infty[} \Phi(\alpha, \beta) \frac{a(\alpha, \beta)}{(\beta - \alpha)^6} \exp\left\{-\frac{\pi^2 t}{2(\beta - \alpha)^2}\right\} 1_{\{\beta - \alpha \leq c\}} d\alpha d\beta$$

Setting  $x = \frac{1}{(\beta - \alpha)^2}$  ( $\beta$  fixed), we get:

$$\Delta_t(fa) = \frac{1}{2} \int_{1/c^2}^\infty x^{3/2} \exp\left\{-\frac{\pi^2 t x}{2}\right\} \left( \int_0^{1/\sqrt{x}} \Phi\left(\beta - \frac{1}{\sqrt{x}}, \beta\right) a\left(\beta - \frac{1}{\sqrt{x}}, \beta\right) d\beta \right) dx.$$

The change of variable  $\beta = r/\sqrt{x}$  leads directly to (3.29).  $\blacksquare$

The proof of (3.26) and (3.27) is based on Laplace's method, whose main result we briefly recall (see for instance [6], chap. IV). We consider:

$$I(t) = \int_0^\infty g(x) e^{t h(x)} dx,$$

where  $g, h : ]0, \infty[ \rightarrow \mathbb{R}$  are continuous and satisfy the two following properties:

$$(3.30) \quad \int_0^\infty |g(x)| e^{t h(x)} dx < \infty, \quad \forall t > 0$$

$$(3.31) \quad \exists \delta_0 > 0, \text{ such that } h(x) \leq h(\delta) \text{ for any } x \geq \delta \text{ and } 0 < \delta < \delta_0.$$

**Proposition 3.8.** *Suppose that the functions  $g$  and  $h$  satisfy:*

$$(3.32) \quad g(x) \underset{x \rightarrow 0_+}{\sim} g_0 x^\rho$$

$$(3.33) \quad h(x) = h_0 - h_1 x^\tau + o(x^\tau), \quad x \rightarrow 0$$

for some:

$$(3.34) \quad g_0 \neq 0, \rho > -1, h_1 \geq 0, \tau > 0.$$

Then:

$$(3.35) \quad I(t) \underset{t \rightarrow \infty}{\sim} \frac{g_0}{\tau} \Gamma\left(\frac{\rho+1}{\tau}\right) (h_1 t)^{-\frac{\rho+1}{\tau}} e^{h_0 t}.$$

As an application of the previous instance of Laplace's method, we obtain the following asymptotics.

**Lemma 3.9.** *Let  $a : ]-\infty, 0] \times [0, \infty[ \rightarrow \mathbb{R}$  be a continuous function.*

1. *In Case 1, we have:*

$$(3.36) \quad \Delta_t(fa) \underset{t \rightarrow \infty}{\sim} \frac{2}{\pi^4} (\Phi a)(\alpha_0, \beta_0) \frac{1}{t^2} \exp\left\{-\frac{\pi^2 t}{2K_f^2}\right\}$$

when  $a(\alpha_0, \beta_0) \neq 0$ .

2. *In Case 2, we have*

$$(3.37) \quad \Delta_t(fa) \underset{t \rightarrow \infty}{\sim} \frac{1}{\pi^2 K_f^2} \left( \int_0^1 (\Phi a)((r-1)K_f, rK_f) dr \right) \frac{1}{t} \exp\left\{-\frac{\pi^2 t}{2K_f^2}\right\}$$

where it is assumed that  $\int_0^1 (\Phi a)((r-1)K_f, rK_f) dr \neq 0$ .

*Proof.* According to (3.28), we have

$$\Delta_t(fa) = \frac{1}{2} \int_{1/K_f^2}^\infty \Lambda(\Phi a)(y) y \exp\left\{-\frac{\pi^2 t y}{2}\right\} dy.$$

Setting  $y = x + 1/K_f^2$  we get:

$$(3.38) \quad \Delta_t(fa) = \frac{1}{2} \left( \int_0^\infty \Lambda(\Phi a)\left(x + \frac{1}{K_f^2}\right) \left(x + \frac{1}{K_f^2}\right) \exp\left\{-\frac{\pi^2 t x}{2}\right\} dx \right) \exp\left\{-\frac{\pi^2 t}{2K_f^2}\right\}.$$

**a)** We begin with Case 2 which is easier.

From (3.29) we deduce:

$$\Lambda(\Phi a)\left(x + \frac{1}{K_f^2}\right) = \int_0^1 (\Phi a)\left(\frac{r-1}{\sqrt{x + 1/K_f^2}}, \frac{r}{\sqrt{x + 1/K_f^2}}\right) dr.$$

Since  $\Phi$  and  $a$  are continuous, we obtain:

$$(3.39) \quad \lim_{x \rightarrow 0} \Lambda(\Phi a)\left(x + \frac{1}{K_f^2}\right) = \int_0^1 (\Phi a)((r-1)K_f, rK_f) dr.$$

**b)** Next, we deal with Case 1. Due to (3.29) we have:

$$\Lambda(\Phi a)\left(x + \frac{1}{K_f^2}\right) = \int_{\tau_0(x)}^{\tau_1(x)} (\Phi a)\left(\frac{r-1}{\sqrt{x + 1/K_f^2}}, \frac{r}{\sqrt{x + 1/K_f^2}}\right) dr$$

with

$$\tau_0(x) = \left(1 + \alpha_0 \sqrt{x + 1/K_f^2}\right)_+, \quad \tau_1(x) = \inf \left\{1, \beta_0 \sqrt{x + 1/K_f^2}\right\}.$$

We observe that

$$\begin{aligned} \lim_{x \rightarrow 0_+} \tau_0(x) &= \left(1 + \frac{\alpha_0}{K_f}\right)_+ = \left(1 + \frac{\alpha_0}{\beta_0 - \alpha_0}\right)_+ = \frac{\beta_0}{\beta_0 - \alpha_0} \\ \lim_{x \rightarrow 0_+} \tau_1(x) &= \inf \left\{1, \frac{\beta_0}{K_f}\right\} = \inf \left\{1, \frac{\beta_0}{\beta_0 - \alpha_0}\right\} = \frac{\beta_0}{\beta_0 - \alpha_0}. \end{aligned}$$

This implies that:

$$\Lambda(\Phi a)\left(x + \frac{1}{K_f^2}\right) \underset{x \rightarrow 0_+}{\sim} (\tau_1(x) - \tau_0(x))(\Phi a)(\alpha_0, \beta_0).$$

When  $x$  is small, we have

$$\tau_1(x) - \tau_0(x) = \beta_0 \sqrt{x + 1/K_f^2} - 1 - \alpha_0 \sqrt{x + 1/K_f^2} = \sqrt{x K_f^2 + 1} - 1.$$

As a result:

$$(3.40) \quad \Lambda(\Phi a)\left(x + \frac{1}{K_f^2}\right) \underset{x \rightarrow 0_+}{\sim} \frac{K_f^2}{2} (\Phi a)(\alpha_0, \beta_0) x.$$

c) We apply Laplace's method with

$$g(x) = \left(x + \frac{1}{K_f^2}\right) \Lambda(\Phi a)\left(x + \frac{1}{K_f^2}\right), \quad \text{and} \quad h(x) = -\frac{\pi^2 x}{2}.$$

It is clear that (3.30), (3.31) and (3.33) hold with

$$h_0 = 0, \quad h_1 = \frac{\pi^2}{2} \quad \text{and} \quad \tau = 1.$$

In Case 2 (resp. Case 1), relation (3.39) (resp. (3.40)) implies that (3.32) is satisfied with  $g_0 = \frac{1}{K_f^2} \int_0^1 (\Phi a)((r-1)K_f, rK_f) dr$  and  $\rho = 0$  (resp.  $g_0 = \frac{1}{2} (\Phi a)(\alpha_0, \beta_0)$  and  $\rho = 1$ ).

Lemma 3.9 is a direct consequence of (3.35) and (3.38).  $\blacksquare$

*Proof of Proposition 3.5.* **1)** We begin with Case 2. It is clear that (3.37) and (3.5) imply:

$$(3.41) \quad \Delta_t(f a_2) \underset{t \rightarrow \infty}{\sim} \frac{4\pi}{K_f^2} \left( \int_0^1 \Phi((r-1)K_f, rK_f) \sin(\pi r) dr \right) \frac{1}{t} \exp \left\{ -\frac{\pi^2 t}{2K_f^2} \right\}.$$

Let  $i = 0, 1$ . From (3.4) we have:

$$(3.42) \quad |\Delta_t(f a_i^{(t)})| \leq C(1 + K_f^4) \Delta_t(f).$$

Using (3.37), we get:

$$(3.43) \quad \Delta_t(f) \underset{t \rightarrow \infty}{\sim} \frac{\mu}{t} \exp \left\{ -\frac{\pi^2 t}{2K_f^2} \right\} \quad (\text{for some } \mu > 0).$$

Applying (3.2), (3.6) and (3.41)–(3.43) shows (3.27).

**2)** Similarly to Case 2, in Case 1, the main term of  $E[f(I_t, S_t)]$  is  $\Delta_t(f a)$ . As a result, (3.26) follows from (3.36).  $\square$

## 4. PENALISATION WITH THE MAXIMUM AND THE MINIMUM

**4.1. Penalisation with  $f(I_t, S_t)$ , Case 1.** In this subsection we suppose that  $f$  satisfies (3.20).

**Theorem 4.1.** *The Brownian penalisation procedure holds with the weight process  $F_t = f(I_t, S_t)$ , i.e.*

1. *Property (2.9) holds:*

$$(4.1) \quad \lim_{t \rightarrow \infty} \frac{E_0[f(I_t, S_t)|\mathcal{F}_u]}{E_0[f(I_t, S_t)]} = M_u^{\alpha_0, \beta_0}, \quad \text{for any } u \geq 0$$

where

$$(4.2) \quad M_u^{\alpha_0, \beta_0} = N^{\alpha_0, \beta_0}(u \wedge T_{\alpha_0} \wedge T_{\beta_0}); \quad u \geq 0$$

$$(4.3) \quad N_u^{\alpha_0, \beta_0} = \frac{1}{\sin\left(\frac{\pi\beta_0}{\beta_0 - \alpha_0}\right)} \sin\left(\frac{\pi(\beta_0 - X_u)}{\beta_0 - \alpha_0}\right) \exp\left\{\frac{\pi^2 u}{2(\beta_0 - \alpha_0)^2}\right\} \quad u \geq 0.$$

$$(4.4) \quad T_x = \inf\{t \geq 0, X_t = x\}.$$

2. *Moreover (2.10) is satisfied:*

$$(4.5) \quad E_0[M_u^{\alpha_0, \beta_0}] = 1 \quad \forall u \geq 0.$$

*Remark 4.2.* 1. According to Theorem 2.1:

(a)  $(M_u^{\alpha_0, \beta_0}; u \geq 0)$  is a non-negative  $P_0$ -martingale, which converges to 0.

(b) For any  $u \geq 0$  and  $\Lambda_u \in \mathcal{F}_u$ ,

$$(4.6) \quad Q_0^{\alpha_0, \beta_0}(\Lambda_u) := \lim_{t \rightarrow \infty} \frac{E_0[1_{\Lambda_u} \Phi(I_t, S_t) 1_{\{I_t \geq \alpha_0, S_t \leq \beta_0\}}]}{E_0[\Phi(I_t, S_t) 1_{\{I_t \geq \alpha_0, S_t \leq \beta_0\}}]}$$

exists and  $Q_0^{\alpha_0, \beta_0}$  is a p.m. on  $(\Omega, \mathcal{F}_\infty)$  which satisfies:

$$(4.7) \quad Q_0^{\alpha_0, \beta_0}(\Lambda_u) = E_0[1_{\Lambda_u} M_u^{\alpha_0, \beta_0}] \quad u \geq 0, \Lambda_u \in \mathcal{F}_u.$$

2. In the particular case:  $f = 1$  and  $\beta_0 = -\alpha_0$ , then

$$F_t = 1_{\{I_t \geq -\beta_0, S_t \leq \beta_0\}} = 1_{\{X_t^* \leq \beta_0\}},$$

where  $X_t^* = S_t \vee (-I_t) = \max_{u \leq t} |X_u|$ . Moreover:

$$M_u^{-\beta_0, \beta_0} = \cos\left(\frac{\pi X(u \wedge T_{\beta_0}^*)}{2\beta_0}\right) \exp\left\{\frac{\pi^2(u \wedge T_{\beta_0}^*)}{2\beta_0^2}\right\},$$

where  $T_{\beta_0}^* = \inf\{t \geq 0, |X_t| = \beta_0\}$ .

3. Note that in [17], a penalisation procedure has been considered with weight processes:

$$F_t := \int_{]-\infty, 0] \times [0, \infty]} 1_{\{I_t \geq \alpha, S_t \leq \beta\}} \exp\left\{\frac{1}{2}\left(\frac{1}{\beta} - \frac{1}{\alpha}\right)L_t^0\right\} \nu(d\alpha, d\beta)$$

where  $\nu$  is a p.m. on  $] -\infty, 0] \times [0, \infty]$  and  $(L_t^0)$  is the local time process at 0 associated with  $(X_t)$ .

It has been proved (cf Theorem 3.18) that the martingales generated by this penalisation are functions of the triplet  $(I_t, S_t, L_t^0)$ . Consequently, they are not of the form  $(M_t^{-\beta_0, \beta_0})$ .

*Proof of Theorem 4.1.* To show that the penalisation procedure holds with  $F_t = f(I_t, S_t)$  we need to prove (4.1) and (4.6). Observe that relation (3.26) in Proposition 3.5 gives the

rate of decay of  $t \rightarrow E_0[f(I_t, S_t)]$ ,  $t \rightarrow \infty$ . Next, we need to determine the asymptotic behavior of  $t \rightarrow E[f(I_t, S_t)|\mathcal{F}_u]$ . We will prove in step 1 below that the rate of decay of  $E[f(I_t, S_t)|\mathcal{F}_u]$  may be deduced from (3.26). In step 2 we will prove (4.4).

**1) Proof of (4.1)**

Let  $u > 0$  be fixed. We introduce  $X'_t = X_{t+u} - X_u$ ,  $t \geq 0$ .

Under  $P_0$ ,  $(X'_t)_{t \geq 0}$  is a Brownian motion started at 0 and independent from  $\mathcal{F}_u$ . Moreover:

$$S_{u+h} = S_u \vee (X_u + S'_h), \quad I_{u+h} = I_u \wedge (X_u + I'_h), \quad h > 0$$

where  $S'_h = \max_{v \leq h} X'_v$  and  $I'_h = \min_{v \leq h} X'_v$ .

This implies that

$$(4.8) \quad E_0[f(I_{u+v}, S_{u+v})|\mathcal{F}_u] = H(I_u, S_u, X_u, v); \quad v \geq 0$$

where

$$(4.9) \quad H(a, b, x, v) := E_0[f(a \wedge (x + I'_v), b \vee (x + S'_v))].$$

Suppose that  $a, b, x$  are fixed,  $a \leq x$ ,  $b \geq x$ . We introduce:

$$(4.10) \quad \tilde{f}(\alpha, \beta) = f(a \wedge (x + \alpha), b \vee (x + \beta)) \quad \alpha \leq 0, \beta \geq 0.$$

Since:

$$a \wedge (x + \alpha) \geq \alpha_0 \quad \Leftrightarrow \quad a \geq \alpha_0 \quad \text{and} \quad \alpha \geq \alpha_0 - x$$

and

$$b \vee (x + \beta) \leq \beta_0 \quad \Leftrightarrow \quad b \leq \beta_0 \quad \text{and} \quad \beta \leq \beta_0 - x$$

then (3.20) implies that

$$H(a, b, x, v) = 1_{\{a \geq \alpha_0, \beta \leq \beta_0\}} E_0[\Phi(a \wedge (x + I_v), b \vee (x + S_v)) 1_{\{I_v \geq \alpha_0 - x, S_v \leq \beta_0 - x\}}].$$

We may apply (3.26):

$$H(a, b, x, v) \underset{v \rightarrow \infty}{\sim} \frac{8}{\pi} \Phi(\alpha_0, \beta_0) \sin\left(\frac{\pi(\beta_0 - x)}{\beta_0 - \alpha_0}\right) \exp\left\{-\frac{\pi^2 v}{2(\beta_0 - \alpha_0)^2}\right\} 1_{\{a \geq \alpha_0, b \leq \beta_0\}}.$$

Consequently:

$$(4.11) \quad E_0[f(I_{u+v}, S_{u+v})|\mathcal{F}_u] \underset{v \rightarrow \infty}{\sim} \frac{8}{\pi} \Phi(\alpha_0, \beta_0) \sin\left(\frac{\pi(\beta_0 - X_u)}{\beta_0 - \alpha_0}\right) \exp\left\{-\frac{\pi^2 v}{2(\beta_0 - \alpha_0)^2}\right\} \\ \times 1_{\{I_u \geq \alpha_0, S_u \leq \beta_0\}}.$$

Recall that

$$E_0[f(I_{u+v}, S_{u+v})] \underset{v \rightarrow \infty}{\sim} \frac{8}{\pi} \Phi(\alpha_0, \beta_0) \sin\left(\frac{\pi\beta_0}{\beta_0 - \alpha_0}\right) \exp\left\{-\frac{\pi^2(u+v)}{2(\beta_0 - \alpha_0)^2}\right\}.$$

This proves (4.1) because

$$N^{\alpha_0, \beta_0}(T_{\alpha_0}) = N^{\alpha_0, \beta_0}(T_{\beta_0}) = 0$$

and

$$\{I_u \geq \alpha_0\} = \{u \leq T_{\alpha_0}\}, \quad \{S_u \leq \beta_0\} = \{u \leq T_{\beta_0}\}.$$

**2) Proof of (4.5)**

$(N_u^{\alpha_0, \beta_0}, u \geq 0)$  is a continuous local martingale as combination of exponential martingales. Itô's formula confirms this:

$$(4.12) \quad dN_u^{\alpha_0, \beta_0} = -\frac{\pi}{\beta_0 - \alpha_0} \frac{1}{\sin\left(\frac{\pi\beta_0}{\beta_0 - \alpha_0}\right)} \cos\left(\frac{\pi(\beta_0 - X_u)}{\beta_0 - \alpha_0}\right) \exp\left\{\frac{\pi^2 u}{2(\beta_0 - \alpha_0)^2}\right\} dX_u.$$

It is clear that  $N_u^{\alpha_0, \beta_0}$  is uniformly bounded on any interval  $[0, T]$ ,  $T$  fixed. This shows that  $(N_u^{\alpha_0, \beta_0}, u \geq 0)$  is a martingale. Applying the Doob's optional stopping theorem we get:

$$E_0[M_u^{\alpha_0, \beta_0}] = E[N^{\alpha_0, \beta_0}(u \wedge T_{\alpha_0} \wedge T_{\beta_0})] = N^{\alpha_0, \beta_0}(0) = 1. \quad \blacksquare$$

**4.2. Penalisation with  $f(I_t, S_t)$ , Case 2 .** We study penalisation in Case 2. Namely  $f$  satisfies (3.23):

$$f(\alpha, \beta) = \Phi(\alpha, \beta) 1_{\{\beta - \alpha \leq c\}}.$$

**Theorem 4.3.** *The Brownian penalisation holds with  $F_t = f(I_t, S_t)$ ,  $f$  satisfying (3.23).*

1. Property (2.9) holds:

$$(4.13) \quad \lim_{t \rightarrow \infty} \frac{E_0[f(I_t, S_t) | \mathcal{F}_u]}{E_0[f(I_t, S_t)]} = M_u^{\Phi, c}, \quad \forall u \geq 0$$

where

$$(4.14) \quad M_u^{\Phi, c} = N^{\Phi, c}(u \wedge \theta(c)); \quad u \geq 0$$

$$(4.15) \quad N^{\Phi, c}(u) = \frac{c + I_u - S_u}{c\rho(F)} \left\{ \int_0^1 \left[ \Phi(S_u - c + r(c + I_u - S_u), S_u + r(c + I_u - S_u)) \right. \right. \\ \left. \left. \times \sin\left(\frac{\pi}{c}(S_u - X_u + r(c + I_u - S_u))\right) \right] dr \right\} \exp\left\{\frac{\pi^2 u}{2c^2}\right\}$$

$$(4.16) \quad = \frac{1}{\rho(\Phi)} \left( \int_{(S_u - X_u)/c}^{(I_u - X_u + c)/c} \Phi(X_u + c(r - 1), X_u + rc) \sin(\pi r) dr \right) \\ \times \exp\left\{\frac{\pi^2 u}{2c^2}\right\}$$

$$(4.17) \quad \rho(\Phi) = \int_0^1 \Phi(c(r - 1), cr) \sin(\pi r) dr$$

$$(4.18) \quad \theta(c) = \inf\{t \geq 0; S_t - I_t = c\}$$

2. Moreover (2.10) holds:

$$(4.19) \quad E_0[M^{\Phi, c}(u)] = 1.$$

*Remark 4.4.* 1. Applying Theorem 2.1, we may deduce that:

- (a)  $(M_u^{\Phi, c}; u \geq 0)$  is a non-negative  $P_0$ -martingale, which converges to 0.
- (b) For any  $u \geq 0$  and  $\Lambda_u \in \mathcal{F}_u$ ,

$$Q_0^{\Phi, c}(\Lambda_u) := \lim_{t \rightarrow \infty} \frac{E_0[1_{\Lambda_u} \Phi(I_t, S_t) 1_{\{S_t - I_t \leq c\}}]}{E_0[\Phi(I_t, S_t) 1_{\{S_t - I_t \leq c\}}]}$$

exists.

Moreover  $Q_0^{\Phi, c}$  is a p.m. on  $(\Omega, \mathcal{F}_\infty)$  and

$$(4.20) \quad Q_0^{\Phi, c}(\Lambda_u) = E_0[1_{\Lambda_u} M_u^{\Phi, c}] \quad u \geq 0, \Lambda_u \in \mathcal{F}_u.$$

2. When  $\Phi(\alpha, \beta) = \Phi_0(\beta - \alpha)$  then  $f(\alpha, \beta) = \Phi_0(\beta - \alpha)1_{\{\beta - \alpha \leq c\}}$  and

$$(4.21) \quad N_u^{\Phi, c} = \frac{1}{2} \left[ \cos \left( \frac{\pi(S_u - X_u)}{c} \right) + \cos \left( \frac{\pi(I_u - X_u)}{c} \right) \right] \exp \left\{ \frac{\pi^2 u}{2c^2} \right\}.$$

Recall that  $M_u^{\Phi, c} = N^{\Phi, c}(u \wedge \theta(c))$   $u \geq 0$ .

*Proof of Theorem 4.3.* Our proof of Theorem 4.3 is close to that of Theorem 4.1. The details are left to the reader. However we would like to explain how the martingale  $(M_t^{\Phi, c})_{t \geq 0}$  appears. We go back to the proof of Theorem 4.1. Obviously (4.8)-(4.10) are still valid. We have to make  $\tilde{f}$  explicit in Case 2. From (3.23):

$$\tilde{f}(\alpha, \beta) = \Phi(a \wedge (x + \alpha), b \vee (x + \beta))1_{\{b \vee (x + \beta) - a \wedge (x + \alpha) \leq c\}}$$

Since:

$$b \vee (x + \beta) - a \wedge (x + \alpha) \leq c \quad \Leftrightarrow \quad b - a \leq c, \quad b - x - \alpha \leq c, \quad x + \beta - a \leq c, \quad \beta - \alpha \leq c$$

then

$$\tilde{f}(\alpha, \beta) = 1_{\{b - a \leq c\}} \tilde{\Phi}(\alpha, \beta) 1_{\{\beta - \alpha \leq c\}}$$

where

$$\tilde{\Phi}(\alpha, \beta) = \Phi(a \wedge (x + \alpha), b \vee (x + \beta)) 1_{\{b - x - \alpha \leq c, x + \beta - a \leq c\}}.$$

Consequently,  $K_{\tilde{f}} = K_f = c$  and

$$\begin{aligned} \tilde{\Phi}(c(r-1), cr) &= \Phi(a \wedge (x + cr - c), b \vee (x + cr)) 1_{\{b - x \leq cr \leq c + a - x\}} \\ &= \Phi(x + cr - c, x + cr) 1_{\{b - x \leq cr \leq c + a - x\}}. \end{aligned}$$

Applying (3.27) leads to (4.13) and (4.16). ■

## 5. THE LAW OF $(X_t)$ UNDER $Q_0^{\alpha_0, \beta_0}$ AND $Q_0^{\Phi, c}$

We first consider the distribution of the canonical process  $(X_t)$  under  $Q_0^{\alpha_0, \beta_0}$ . Let  $\alpha_0 < 0$  and  $\beta_0 > 0$  be two fixed real numbers. Recall the definition of the p.m.  $Q_0^{\alpha_0, \beta_0}$  on  $(\Omega, \mathcal{F}_\infty)$ :

$$(5.1) \quad Q_0^{\alpha_0, \beta_0}(\Lambda_u) = E_0[1_{\Lambda_u} M_u^{\alpha_0, \beta_0}] \quad \Lambda_u \in \mathcal{F}_u$$

where  $(M_u^{\alpha_0, \beta_0})_{u \geq 0}$  is the  $P_0$ -martingale defined by (4.2).

**Theorem 5.1.** *Under  $Q_0^{\alpha_0, \beta_0}$ :*

1.  $(X_t)$  is a diffusion process solving:

$$(5.2) \quad X_t = B_t - \frac{\pi}{\beta_0 - \alpha_0} \int_0^t \cot \left( \frac{\pi(\beta_0 - X_u)}{\beta_0 - \alpha_0} \right) du; \quad t \geq 0,$$

where  $(B_t)_{t \geq 0}$  is a  $Q_0^{\alpha_0, \beta_0}$ -Brownian motion started at 0.

2.  $(X_t)$  has the following path properties:

$$(5.3) \quad \alpha_0 < X_t < \beta_0 \quad \forall t \geq 0$$

$$(5.4) \quad S_\infty = \sup_{t \geq 0} X_t = \beta_0, \quad I_\infty = \inf_{t \geq 0} X_t = \alpha_0.$$

3.  $X_t$  converges in distribution, as  $t \rightarrow \infty$  to the p.m.  $p^{\alpha_0, \beta_0}(x)dx$  on  $\mathbb{R}$ , with:

$$(5.5) \quad p^{\alpha_0, \beta_0}(x) := \frac{2}{\beta_0 - \alpha_0} \sin^2 \left( \frac{\pi(\beta_0 - x)}{\beta_0 - \alpha_0} \right) 1_{(\alpha_0, \beta_0)}(x).$$

*Remark 5.2.* 1. Property (5.3) follows intuitively from our penalisation procedure and the fact that the support of the p.m. on  $\mathcal{F}_u$ :

$$\Lambda_u \mapsto \frac{E_0[1_{\Lambda_u} \Phi(I_t, S_t) 1_{\{I_t \geq \alpha_0, S_t \leq \beta_0\}}]}{E_0[\Phi(I_t, S_t) 1_{\{I_t \geq \alpha_0, S_t \leq \beta_0\}}]} \quad (t > u)$$

is included in  $\{I_u \geq \alpha_0, S_u \leq \beta_0\}$ .

2. Note that  $Q_0^{\alpha_0, \beta_0}$  do not depend on  $\Phi$ .

*Proof of Theorem 5.1.*

**a)**  $(M_t^{\alpha_0, \beta_0}; t \geq 0)$  is a non-negative  $Q_0^{\alpha_0, \beta_0}$ -martingale and can be written as:

$$M_t^{\alpha_0, \beta_0} = \mathcal{E}(J)_t := \exp \left\{ \int_0^t J_u dX_u - \frac{1}{2} \int_0^t J_u^2 du \right\}$$

for any  $t < T_{\alpha_0} \wedge T_{\beta_0}$ . Indeed, from (4.12) and (4.2) we have:

$$J_u = -\frac{\pi}{\beta_0 - \alpha_0} \cot \left( \frac{\pi(\beta_0 - X_u)}{\beta_0 - \alpha_0} \right), \quad 0 \leq u < T_{\alpha_0} \wedge T_{\beta_0}.$$

Then Girsanov's theorem implies (5.2).

**b)** To investigate the path properties of  $(X_t)$ , we define  $Y_t = \beta_0 - X_t$ , then

$$(5.6) \quad Y_t = \beta_0 - B_t + \frac{\pi}{\beta_0 - \alpha_0} \int_0^t \cot \left( \frac{\pi Y_u}{\beta_0 - \alpha_0} \right) du$$

Obviously  $(Y_t)_{t \geq 0}$  is a one-dimensional diffusion. Let  $S$  (resp.  $m(dy)$ ) denote its scale function (resp. speed measure), see for instance: (Section 1, chap. 4 of [8]), (Section 1, chap II of [4]). Using standard calculations (i.e. [14] (Section 52, chap V), [5] (Section 12, chap 16))) we easily get:

$$S(y) = -\cot \left( \frac{\pi y}{\beta_0 - \alpha_0} \right), \quad y \in ]0, \beta_0 - \alpha_0[$$

$$m([0, y]) = \frac{(\beta_0 - \alpha_0)}{\pi} y - \frac{(\beta_0 - \alpha_0)^2}{2\pi^2} \sin \left( \frac{2\pi y}{\beta_0 - \alpha_0} \right), \quad y \in ]0, \beta_0 - \alpha_0[.$$

According to the classification of boundary points of a linear diffusion (cf Section 1, chap II of [4]), we have:

*i)* 0 is not an exit point since :

$$(5.7) \quad \int_{(0,x)} m([y, z]) S'(y) dy = \infty \quad (z \in ]0, \beta_0 - \alpha_0[).$$

*ii)* 0 is an entrance point since:

$$(5.8) \quad \int_{(0,z)} (S(z) - S(y)) m(dy) < \infty \quad (z \in ]0, \beta_0 - \alpha_0[).$$

Similarly  $\beta_0 - \alpha_0$  is not an exit point and is an entrance one. This shows (5.3).

**c)** The diffusion  $(Y_t)_{t \geq 0}$  which takes its values in  $(0, \beta_0 - \alpha_0)$  is recurrent. This implies (5.4). Let  $p$  be the density of its invariant p.m. From the Fokker Planck equation,  $p$  solves:

$$(5.9) \quad \frac{1}{2} p''(y) - \frac{\pi}{\beta_0 - \alpha_0} \left( \cot \left( \frac{\pi y}{\beta_0 - \alpha_0} \right) p(y) \right)' = 0.$$

It is easy to verify that

$$p(y) = \frac{2}{\beta_0 - \alpha_0} \sin^2 \left( \frac{\pi y}{\beta_0 - \alpha_0} \right) \quad y \in (0, \beta_0 - \alpha_0)$$



is the unique density function solving (5.9).  $\blacksquare$

To study the law of  $(X_t)_{t \geq 0}$  under  $Q_0^{\Phi, c}$  it is more convenient to express the p.m.  $Q_0^{\Phi, c}$  via the family of p.m.'s  $\{Q_0^{\beta-c, \beta}; 0 < \beta < c\}$  (see Theorem 5.3 below). This result will allow to determine easily the distribution of  $(X_t)$  under  $Q_0^{\Phi, c}$ .

**Theorem 5.3.** *Let  $c > 0$ , and  $\Phi : \{(\alpha, \beta); \alpha < 0, \beta > 0, \beta - \alpha \leq c\} \rightarrow \mathbb{R}_+$  be a continuous function satisfying (3.24). Then:*

$$(5.10) \quad Q_0^{\Phi, c}(\cdot) = \frac{1}{c\rho(\Phi)} \int_0^c \Phi(\beta - c, \beta) \sin\left(\frac{\pi\beta}{c}\right) Q_0^{\beta-c, \beta}(\cdot) d\beta.$$

*Proof.* **a)** Assume for a while that the followings holds:

$$(5.11) \quad E_0^{\Phi, c}[1_{\Lambda_u} h(S_u, S_\infty)] = \frac{1}{c\rho(\Phi)} \int_0^c \Phi(v - c, v) \sin\left(\frac{\pi v}{c}\right) E_0^{v-c, v}[1_{\Lambda_u} h(S_u, v)] dv$$

for any  $h : \mathbb{R}_+^2 \rightarrow \mathbb{R}_+$  Borel,  $u > 0$ ,  $\Lambda_u \in \mathcal{F}_u$ , and that  $E_0^{\Phi, c}$  (resp.  $E_0^{\alpha_0, \beta_0}$ ) stands for the expectation under  $Q_0^{\Phi, c}$  (resp.  $Q_0^{\alpha_0, \beta_0}$ ).

Taking  $h = 1$  in (5.1) implies that (5.10) holds on  $\mathcal{F}_u$ . Since the p.m.'s in (5.10) are defined on  $\mathcal{F}_\infty$  they coincide on  $\mathcal{F}_\infty$ .

**b)** The remainder is devoted to the proof of (5.11).

*i)* We claim that:

$$(5.12) \quad Q_0^{\Phi, c}(S_\infty - I_\infty \leq c) = 1.$$

Indeed, from the definition of  $Q_0^{\Phi, c}$  we have:

$$Q_0^{\Phi, c}(S_t - I_t > c) = E_0^{\Phi, c}[1_{\{S_t - I_t > c\}} M_t^{\Phi, c}] = E_0^{\Phi, c}[1_{\{t > \theta(c)\}} M_t^{\Phi, c}]$$

From Doob's optional stopping theorem and the fact that  $M_{\theta(c)}^{\Phi, c} = 0$ , we get:

$$Q_0^{\Phi, c}(S_t - I_t > c) = E_0^{\Phi, c}[1_{\{t > \theta(c)\}} M_{\theta(c)}^{\Phi, c}] = 0.$$

Taking  $t \rightarrow \infty$ , we obtain (5.12).

*ii)* Due to the monotone class theorem and (5.12), it is sufficient to show (5.11) with

$$h(x, y) := 1_{[0, \beta]}(x) 1_{[\beta', c]}(y) \quad (0 < \beta < \beta' < c)$$

In this case, (5.11) reduces to:

$$(5.13) \quad A = \frac{1}{c\rho(\Phi)} \int_{\beta'}^c \Phi(v - c, v) \sin\left(\frac{\pi v}{c}\right) P_0^{v-c, v}(\Lambda_u \cap \{S_u \leq \beta\}) dv,$$

where

$$(5.14) \quad A := P_0^{\Phi, c}(\Lambda_u \cap \{S_u \leq \beta, S_\infty > \beta'\}).$$

It is clear that

$$A = \lim_{t \rightarrow \infty} A(t), \quad \text{with } A(t) := P_0^{\Phi, c}(\Lambda_u \cap \{S_u \leq \beta, S_t > \beta'\}).$$

Then, for  $t > u$ , we have:

$$\begin{aligned} A(t) &= E_0[1_{\{\Lambda_u \cap \{S_u \leq \beta\}\}} 1_{\{S_t > \beta'\}} M_t^{\Phi, c}] \\ &= E_0[1_{\{\Lambda_u \cap \{S_u \leq \beta\}\}} 1_{\{T_{\beta'} < t\}} M_t^{\Phi, c}] \end{aligned}$$

Due to Doob's optional stopping theorem we get:

$$A(t) = E_0[1_{\{\Lambda_u \cap \{S_u \leq \beta\}\}} 1_{\{T_{\beta'} < t\}} M^{\Phi, c}(T_{\beta'})]$$

Taking the limit  $t \rightarrow \infty$ , we obtain:

$$A = E_0 \left[ 1_{\Lambda_u \cap \{S_u \leq \beta\}} M^{\Phi, c}(T_{\beta'}) \right].$$

Using (4.14) and (4.16) we have:

$$A = \frac{1}{\rho(\Phi)} E_0 \left[ 1_{\Lambda_u \cap \{S_u \leq \beta\}} \left( \int_0^1 1_{\{rc < I(T_{\beta'}) - \beta' + c\}} \sin(\pi r) \Phi(\beta' + c(r-1), \beta' + rc) dr \right) \right. \\ \left. \times 1_{\{\beta' - I(T_{\beta'}) < c\}} \exp \left\{ \frac{\pi^2 T_{\beta'}}{2c^2} \right\} \right].$$

Observe that  $0 > I(T_{\beta'}) > rc + \beta' - c$  implies that  $I(T_{\beta'}) > \beta' - c$  and  $\beta' + rc < c$ . Making the change of variable  $v = \beta' + rc$  leads to

$$(5.15) \quad A = \frac{1}{c\rho(\Phi)} \int_{\beta'}^c \Phi(v-c, v) \sin \left( \frac{\pi(v-\beta')}{c} \right) A_1(v) dv$$

where

$$A_1(v) := E_0 \left[ 1_{\Lambda_u \cap \{S_u \leq \beta\}} 1_{\{I(T_{\beta'}) > v-c\}} \exp \left\{ \frac{\pi^2 T_{\beta'}}{2c^2} \right\} \right].$$

Since  $u \leq T_\beta < T_{\beta'}$ , we get, by using the Markov property at time  $u$ :

$$E_0 \left[ 1_{\{I(T_{\beta'}) > v-c\}} \exp \left\{ \frac{\pi^2 T_{\beta'}}{2c^2} \right\} \middle| \mathcal{F}_u \right] = 1_{\{I_u > v-c\}} A_2(X_u) \exp \left\{ \frac{\pi^2 u}{2c^2} \right\}$$

with:

$$A_2(x) := E \left[ 1_{\{I(T_{\beta'-x}) > v-c-x\}} \exp \left\{ \frac{\pi^2 T_{\beta'-x}}{2c^2} \right\} \right].$$

Let  $\alpha_1 < 0 < \beta_1$  and assume that  $\beta_1 - \alpha_1 < c$ . Then, using the fact that

$$Z_t := \sin \left( \frac{\pi(X_t - \alpha_1)}{c} \right) \exp \left\{ \frac{\pi^2 t}{2c^2} \right\}, \quad t \geq 0$$

is a martingale, and Doob's optional stopping theorem at  $T_{\beta_1} \wedge T_{\alpha_1}$  lead to:

$$(5.16) \quad E_0 \left[ 1_{\{I(T_{\beta_1}) > \alpha_1\}} \exp \left\{ \frac{\pi^2 T_{\beta_1}}{2c^2} \right\} \right] = - \frac{\sin \left( \frac{\pi \alpha_1}{c} \right)}{\sin \left( \frac{\pi}{c} (\beta_1 - \alpha_1) \right)}.$$

Consequently, we obtain successively:

$$A_2(x) = \frac{\sin \left( \frac{\pi(v-x)}{c} \right)}{\sin \left( \frac{\pi(v-\beta')}{c} \right)}$$

$$A = \frac{1}{c\rho(\Phi)} \int_{\beta'}^c \Phi(v-c, v) A_3(v) dv,$$

with

$$A_3(v) := E_0 \left[ 1_{\Lambda_u \cap \{S_u \leq \beta, I_u > v-c\}} \sin \left( \frac{\pi(v-X_u)}{c} \right) \exp \left\{ \frac{\pi^2 u}{2c^2} \right\} \right].$$

Note that  $S_u \leq \beta < \beta' < v$ ; then according to (4.2) and (4.3) we have:

$$A_2(v) = \sin \left( \frac{\pi v}{c} \right) E_0 \left[ 1_{\Lambda_u \cap \{S_u \leq \beta\}} M_u^{v-c, v} \right].$$

Finally

$$A = \frac{1}{c\rho(\Phi)} \int_{\beta'}^c \Phi(v-c, v) \sin \left( \frac{\pi v}{c} \right) E_0 \left[ 1_{\Lambda_u} M_u^{v-c, v} 1_{\{S_u \leq \beta\}} \right] dv. \\ = \frac{1}{c\rho(\Phi)} \int_{\beta'}^c \Phi(v-c, v) \sin \left( \frac{\pi v}{c} \right) P_0^{v-c, v} (\Lambda_u \cap \{S_u \leq \beta\}) dv.$$

This shows (5.13). ■

We will deduce from Theorem 5.3 two main consequences (see Theorems 5.4 and 5.5 below). We first interpret the identity (5.10) in a more probabilistic way.

**Theorem 5.4.** 1. *Conditionally on  $S_\infty = v$ ,  $(I_t, S_t, X_t)_{t \geq 0}$  is distributed under  $Q_0^{\Phi, c}$  as the three dimensional process  $(I_t, S_t, X_t)_{t \geq 0}$  under  $Q_0^{v-c, v}$ .*

2. *The density function of  $S_\infty$  under  $Q_0^{\Phi, c}$  is*

$$\frac{1}{c\rho(\Phi)} \Phi(v-c, v) \sin\left(\frac{\pi v}{c}\right) 1_{(0, c)}(v).$$

*Proof.* Let  $u > 0$  and  $\Lambda_u \in \mathcal{F}_u$ . Let us apply (5.11) with  $h(x, y) = H(y)$ , for  $H : [0, \infty[ \rightarrow [0, \infty[$ :

$$E_0^{\Phi, c}[1_{\Lambda_u} H(S_\infty)] = \frac{1}{c\rho(\Phi)} \int_0^c \Phi(v-c, v) \sin\left(\frac{\pi v}{c}\right) H(v) Q_0^{v-c, v}(\Lambda_u) dv$$

If we take in particular  $\Lambda_u = \Omega$ , we get

$$E_0^{\Phi, c}[H(S_\infty)] = \frac{1}{c\rho(\Phi)} \int_0^c H(v) \Phi(v-c, v) \sin\left(\frac{\pi v}{c}\right) dv.$$

This shows *ii*).

Then *i*) follows from

$$\begin{aligned} E_0^{\Phi, c}[1_{\Lambda_u} H(S_\infty)] &= E_0^{\Phi, c}[H(S_\infty) P_0^{\Phi, c}(\Lambda_u | S_\infty)] \\ &= \frac{1}{c\rho(\Phi)} \int_0^c H(v) P_0^{\Phi, c}(\Lambda_u | S_\infty = v) \Phi(v-c, v) \sin\left(\frac{\pi v}{c}\right) dv. \end{aligned}$$

■

We are now able to present a few path properties of  $(X_t)$  under  $Q_0^{\Phi, c}$ .

**Theorem 5.5.** *Under  $Q_0^{\Phi, c}$  :*

1. *We have:*

$$(5.17) \quad I_t > I_\infty \quad \text{and} \quad S_t < S_\infty \quad \text{for any } t \geq 0$$

$$(5.18) \quad S_\infty - I_\infty = c.$$

2. *When  $t$  goes to infinity, the couple  $(S_t, X_t)$  converges in distribution to the p.m. on  $\mathbb{R}_+ \times \mathbb{R}$ :*

$$\frac{2}{c^2 \rho(\Phi)} \sin^2\left(\frac{\pi(x-y)}{c}\right) \sin\left(\frac{\pi x}{c}\right) \Phi(x-c, x) 1_{\{0 < x < c, x-c < y < x\}} dx dy.$$

*Remark 5.6.* 1. Property (5.18) can be deduced intuitively from our penalisation procedure and the fact that the support of the p.m. on  $\mathcal{F}_u$ :

$$\Lambda_u \mapsto \frac{E_0[1_{\Lambda_u} \Phi(I_t, S_t) 1_{\{S_t - I_t \leq c\}}]}{E_0[\Phi(I_t, S_t) 1_{\{S_t - I_t \leq c\}}]} \quad (t > u)$$

is included in  $\{S_u - I_u \leq c\}$ .

2. Recall that the p.m.  $Q_0^{\alpha_0, \beta_0}$  which arises from penalisation with

$$\Phi(I_t, S_t) 1_{\{I_t \geq \alpha_0, S_t \leq \beta_0\}}$$

does not depend on the values of  $\Phi$ . Therefore penalisation associated with

$$\Phi(I_t, S_t) 1_{\{S_t - I_t \leq c\}}$$

is very different since the p.m.  $Q_0^{\Phi, c}$  depends on the values of  $\Phi$  over the segment  $\{(\beta - c, \beta); \beta \in [0, c]\}$ .

*Proof of Theorem 5.5.*

It is easy to deduce (5.17) (resp. (5.18)) from (5.3) (resp. (5.4)) and Theorem 5.4. The details are left to the reader. Let  $g : \mathbb{R}_+ \times \mathbb{R}$  be a continuous and bounded function. According to Theorem 5.4 we have:

$$E_0^{\Phi,c}[g(S_t, X_t)] = \frac{1}{c\rho(\Phi)} \int_0^c \Phi(v-c, v) \sin\left(\frac{\pi v}{c}\right) E_0^{v-c,v}[g(S_t, X_t)] dv.$$

Applying Theorem 5.1, we get:

$$\lim_{t \rightarrow \infty} E_0^{v-c,v}[g(S_t, X_t)] = \frac{2}{c} \int_{v-c}^v g(v, y) \sin^2\left(\frac{\pi(v-y)}{c}\right) dy.$$

Point 2) of Theorem 5.5 is a direct consequence of the dominated convergence theorem. ■

We formulate differently item 2) of Theorem 5.5.

**Corollary 5.7.** *The pair  $\left(\frac{S_t + I_t}{2}, X_t - \frac{S_t + I_t}{2}\right)$  converges in distribution as  $t \rightarrow \infty$  to:*

$$\left(\frac{1}{c\rho(\Phi)} \Phi\left(x - \frac{c}{2}, x + \frac{c}{2}\right) \cos\left(\frac{\pi x}{c}\right) 1_{[-\frac{c}{2}, \frac{c}{2}]}(x) dx\right) \times \left(\frac{2}{c} \cos^2\left(\frac{\pi y}{c}\right) 1_{[-\frac{c}{2}, \frac{c}{2}]}(y) dy\right)$$

hence, its two components are asymptotically independent.

*Proof.* This is a direct consequence of 2) of Theorem 5.4, (5.18) and simple changes of variables. ■

*Remark 5.8.* Using Girsanov's theorem it may be proved that  $(X_t)$  solves:

$$(5.19) \quad X_t = B_t + \int_0^t \frac{\partial \Gamma^{\Phi,c}}{\Gamma^{\Phi,c}}(I_u, S_u, X_u) 1_{\{S_u - I_u < c\}} du$$

where  $(B_t)_{t \geq 0}$  is a  $Q_0^{\Phi,c}$  Brownian motion, and

$$(5.20) \quad \Gamma^{\Phi,c}(a, b, x) = \int_{(b-x)/c}^{(a-x+c)/c} \Phi(x + c(r-1), x + rc) \sin(\pi r) dr$$

$$\frac{\partial \Gamma^{\Phi,c}}{\partial x}(a, b, x) = -\frac{\pi}{c} \int_{(b-x)/c}^{(a-x+c)/c} \Phi(x + c(r-1), x + rc) \cos(\pi r) dr.$$

In the particular case  $\Phi = 1$  (i.e. penalisation with  $S_t - I_t \leq c$ ) then:

$$\Gamma^{1,c}(a, b, x) = \frac{1}{\pi} \left( \cos\left(\frac{\pi(b-x)}{c}\right) + \cos\left(\frac{\pi(a-x)}{c}\right) \right)$$

This implies that  $(X_t)$  solves:

$$(5.21) \quad X_t = B_t + \frac{\pi}{c} \int_0^t \tan\left(\frac{\pi}{c} \left(\frac{S_u + I_u}{2} - X_u\right)\right) 1_{\{S_u - I_u < c\}} du.$$

## 6. APPLICATION TO DIFFUSIONS

Our approach is based on Theorem 5.1. Let  $\mu$  be a p.m. on  $] - \infty, 0] \times [0, \infty[$  which does not charge  $(0, 0)$ . Let us consider the associated p.m.  $Q_0^\mu$  on  $(\Omega, \mathcal{F}_\infty)$  :

$$(6.1) \quad Q_0^\mu(\cdot) = \int_{]-\infty, 0] \times [0, \infty[} Q_0^{\alpha, \beta}(\cdot) \mu(d\alpha, d\beta)$$

where  $Q_0^{\alpha, \beta}$  has been defined by (5.1). Note that from Theorem 5.3, the p.m.  $Q_0^{\Phi, c}$  is equal to  $Q_0^\mu$  where

$$(6.2) \quad \mu(d\alpha, d\beta) = \mu^{\Phi, c}(d\alpha, d\beta) := \frac{1}{c\rho(\Phi)} \Phi(\beta - c, \beta) \sin\left(\frac{\pi\beta}{c}\right) \delta_{\beta-c}(d\alpha) 1_{[0, c]}(\beta) d\beta.$$

**Proposition 6.1.** *Under  $Q_0^\mu$ :*

1.  $(I_t, S_t)$  converges a.s. as  $t \rightarrow \infty$  to  $(I_\infty, S_\infty)$  and the distribution of  $(I_\infty, S_\infty)$  is  $\mu$ .
2.  $(I_t, S_t, X_t)$  converges in distribution, as  $t \rightarrow \infty$  to the p.m. on  $] - \infty, 0] \times [0, \infty[ \times \mathbb{R}$ :

$$(6.3) \quad \lambda(d\alpha, d\beta, dx) = p^{\alpha, \beta}(x) \mu(d\alpha, d\beta) dx,$$

where the density function  $p^{\alpha, \beta}(x)$  has been defined by (5.5).

In particular,  $(S_t, X_t)$  converges in law, as  $t \rightarrow \infty$ , to

$$(6.4) \quad \nu(d\beta, dx) := \left( \int_{-\infty}^0 p^{\alpha, \beta}(x) \mu(d\alpha, d\beta) \right) dx.$$

*Proof.*

Proposition 6.1 is a direct consequence of Theorem 5.1. ■

*Remark 6.2.* 1. It seems difficult to characterize all possible p.m.'s  $\nu$  obtained by this randomization procedure, i.e. to describe the set of p.m.'s  $\nu$  which are defined by (6.4),  $\mu$  varying in the set of p.m.'s on  $] - \infty, 0] \times [0, \infty[$ .

2. It can be proved that if  $\mu(d\alpha, d\beta)$  satisfies:

$$(6.5) \quad \int_{]-\infty, 0] \times [0, \infty[} (\beta - \alpha) \mu(d\alpha, d\beta) < \infty,$$

$$(6.6) \quad \int_{]-\infty, 0] \times [0, \infty[} (\beta + \alpha) \mu(d\alpha, d\beta) = 0,$$

$$(6.7) \quad 2a \mu(] - \infty, 0] \times [a, \infty[) = \int_{]-\infty, 0] \times [0, \infty[} (\alpha + \beta) \mu(d\alpha, d\beta).$$

Then, the Rogers conditions (1.7) and (1.8) hold.

3. The p.m.  $Q_0^\mu$  is locally absolutely continuous with respect to the Wiener measure  $P_0$ , namely:

$$(6.8) \quad Q_0^\mu(\Lambda_t) = E_0[1_{\Lambda_t} M_t^\mu], \quad \Lambda_t \in \mathcal{F}_t$$

where

$$(6.9) \quad M_t^\mu = \int_{]-\infty, 0] \times [0, \infty[} M_t^{\alpha, \beta} \mu(d\alpha, d\beta),$$

and  $(M_t^{\alpha, \beta})_{t \geq 0}$  is the  $P_0$ -martingale introduced in Theorem 4.1. Moreover

$$(6.10) \quad M_t^\mu = \Gamma^\mu(I_t, S_t, X_t, t)$$

where

$$(6.11) \quad \Gamma^\mu(a, b, x, t) := \int_{]-\infty, 0] \times [0, \infty[} \frac{1}{\sin\left(\frac{\pi\beta}{\beta-\alpha}\right)} \sin\left(\frac{\pi(\beta-x)}{\beta-\alpha}\right) \exp\left\{\frac{\pi^2 t}{2(\beta-\alpha)^2}\right\} \\ \times 1_{\{\alpha \geq a, \beta \leq b\}} \mu(d\alpha, d\beta).$$

Due to Girsanov's theorem, it may be inferred that  $X_t$  solves:

$$(6.12) \quad X_t = B_t + \int_0^t \frac{\frac{\partial}{\partial x} \Gamma^\mu}{\Gamma^\mu}(I_u, S_u, X_u, u) du$$

where  $(B_t)_{t \geq 0}$  is a  $Q_0^\mu$ -Brownian motion started at 0 and

$$(6.13) \quad \frac{\partial}{\partial x} \Gamma^\mu(a, b, x, t) = -\pi \int_{]-\infty, 0] \times [0, \infty[} \frac{1}{(\beta-\alpha) \sin\left(\frac{\pi\beta}{\beta-\alpha}\right)} \cos\left(\frac{\pi(\beta-x)}{\beta-\alpha}\right)$$

$$(6.14) \quad \times \exp\left\{\frac{\pi^2 t}{2(\beta-\alpha)^2}\right\} 1_{\{\alpha \geq a, \beta \leq b\}} \mu(d\alpha, d\beta)$$

Consequently,  $(I_t, S_t, X_t)$  is a non-homogeneous Markov process.

4. There exists a p.m.  $\mu$  satisfying (6.5)-(6.7). It is easy to show that these conditions hold with

$$(6.15) \quad \mu(d\alpha, d\beta) = \left(\frac{2c^2}{(\beta+c)^3} 1_{\{\beta > 0\}} d\beta\right) \mu_1(d\alpha)$$

where  $\mu_1(d\alpha)$  is a p.m. on  $] -\infty, 0]$ , such that  $-\int_{-\infty}^0 \alpha \mu_1(d\alpha) = c \in ]0, \infty[$ .

#### BIBLIOGRAPHY

1. J. Azéma and M. Yor, *Le problème de Skorokhod: compléments à "Une solution simple au problème de Skorokhod"*, In Séminaire de Probabilités, XIII (Univ. Strasbourg, Strasbourg, 1977/78), vol. 721 of Lecture Notes in Math., Springer, Berlin, 1979, pp. 625–633.
2. J. Azéma and M. Yor, *Une solution simple au problème de Skorokhod*, In Séminaire de Probabilités, XIII (Univ. Strasbourg, Strasbourg, 1977/78), vol. 721 of Lecture Notes in Math., Springer, Berlin, 1979, pp. 90–115.
3. P. Billingsley, *Convergence of probability measures*, John Wiley & Sons Inc., New York, 1968.
4. A. N. Borodin and P. Salminen, *Handbook of Brownian motion—facts and formulae*, Probability and its Applications (Second, eds.), Birkhäuser Verlag, Basel, 2002.
5. L. Breiman, *Probability*, Addison-Wesley Publishing Company, Reading, Mass., 1968.
6. J. Dieudonné, *Calcul infinitésimal*, Hermann, Paris, 1968.
7. W. Feller, *An introduction to probability theory and its applications. Vol. II*, John Wiley & Sons Inc., New York, 1966.
8. K. Itô and H. P. McKean, Jr., *Diffusion processes and their sample paths*, Second printing, corrected, Die Grundlehren der mathematischen Wissenschaften, Band 125, Springer-Verlag, Berlin, 1974.
9. F. B. Knight, *Brownian local times and taboo processes*, Trans. Amer. Math. Soc. **143** (1969), 173–185.
10. F. B. Knight, *Essentials of Brownian motion and diffusion*, vol. 18 of Mathematical Surveys, American Mathematical Society, Providence, R.I., 1981.
11. Jan Oblój, *The Skorokhod embedding problem and its offspring*, Probab. Surv. **1** (2004), 321–390 (electronic).
12. D. Revuz and M. Yor, *Continuous martingales and Brownian motion*, vol. 293 of Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences] (Third, eds.), Springer-Verlag, Berlin, 1999.
13. L. C. G. Rogers, *The joint law of the maximum and terminal value of a martingale*, Probab. Theory Related Fields **95** (1993), no. 4, 451–466.

14. L. C. G. Rogers and David Williams, *Diffusions, Markov processes, and martingales. Vol. 2, Itô calculus*, Reprint of the second (1994) edition, Cambridge Mathematical Library. Cambridge University Press, Cambridge, 2000.
15. B. Roynette, P. Vallois and M. Yor, *A solution to Skorokhod's embedding for linear Brownian motion and its local time*, *Studia Sci. Math. Hungar.* **39** (2002), no. 1-2, 97–127.
16. B. Roynette, P. Vallois and M. Yor, *Limiting laws for long Brownian bridges perturbed by their one-sided maximum, III*, *Period. Math. Hungar.* **50** (2005), no. 1-2, 247–280.
17. B. Roynette, P. Vallois and M. Yor, *Limiting laws associated with Brownian motion perturbed by its maximum, minimum and local time, II*, *Studia Sci. Math. Hungar.* **43** (2006), no. 3, 295–360.
18. B. Roynette, P. Vallois and M. Yor, *Limiting laws associated with Brownian motion perturbed by normalized exponential weights I*, *Studia Sci. Math. Hungar.* **43** (2006), no. 2, 171–346.
19. B. Roynette, P. Vallois and M. Yor, *Pénalisations et extensions du théorème de Pitman, relatives au mouvement brownien et à son maximum unilatère*, In *Séminaire de Probabilités, XXXIX (P.A. Meyer, in memoriam)*, vol. 1874 of *Lecture Notes in Math.*, Springer, Berlin, 2006, pp. 305–336.
20. B. Roynette, P. Vallois and M. Yor, *Some penalisations of the Wiener measure*, *Japan. J. Math.* **1** (2006), 263–290.
21. B. Roynette, P. Vallois and M. Yor, *Penalisations of multidimensional Brownian motion, VI*, Submitted (2007).
22. B. Roynette, P. Vallois and M. Yor, *Penalizing a Brownian motion with a function of the lengths of its excursions, VII*, In preparation (2007).
23. B. Roynette, P. Vallois and M. Yor, *Some extensions of Pitman's and Ray-Knight's theorems for penalized Brownian motions and their local times, IV*, *Studia Sci. Math. Hungar.* **44** (2007), no. 4, 469–516.
24. B. Roynette, P. Vallois and M. Yor, *Penalizing a  $Bes(d)$  process ( $0 < d < 2$ ) with a function of its local time at 0, V*, To appear in *Studia Sci. Math. Hungar.* **45** (2008).
25. B. Roynette and M. Yor, *Penalising Brownian paths: rigorous results and meta-theorems*, Monograph, 2008.

INSTITUT ELIE CARTAN, UNIVERSITÉ HENRI POINCARÉ, B.P. 239, 54506 VANDOEUVRE LES NANCY CEDEX

INSTITUT ELIE CARTAN, UNIVERSITÉ HENRI POINCARÉ, B.P. 239, 54506 VANDOEUVRE LES NANCY CEDEX

LABORATOIRE DE PROBABILITÉS ET MODÈLES ALÉATOIRES, UNIVERSITÉ PARIS VI ET VII, 4 PLACE JUSSIEU – CASE 188; F – 75252 PARIS CEDEX 05; INSTITUT UNIVERSITAIRE DE FRANCE