

(.), « »

PROTEOLYSIS SYSTEM IN PATIENTS WITH DYSPLASTIC NEVI

D. V. Prokhorov

SUMMARY

The state of proteinase-inhibitory capacity of serum skin 31 patients with dysplastic nevi. Found that malignant neoplastic process leads to a unidirectional increase in the activity of proteinases and their inhibitors in the serum (the skin).

31

()

:

5-

[3,10,12].

« » ()

W. H. Clark [1,2,4,6].

42 83 (13 18 54,16±3,02)

1 6).

20

() N- -L- N- -L-

()

[7,11,13].

t- -n- ()

(1-)

()

[5].

[8].

P<0,05.

(m),

(M)

1

	(.) /	(.) /	/	,/
n=20	0,196±0,2	0,22±0,03	7,8±0,2	29,3±1,6
n=31	0,193±0,02	0,225±0,02	7,5±0,2	29,3±1,1

2

	(.) /	(.) /	/	,/
n=20	22,8±1,9	10,7±0,9	0,116±0,01	0,09±0,09
n=31	25,4±0,9	12,93±1,1	0,129±0,01	0,105±0,01

11,52%; +11,28%; +10,79%; +12,03%).

5.- .22-26.

5.

6.

7.

8.

[. . .]- .: , 1985. - 358 .

9. Nomura T. Involvement of cathepsins in the invasion, metastasis and proliferation of cancer cells / T. Nomura, N. Katunuma // J. Med. Invest. - 2005. - V. 52 (1-2). - P. 1-9.

10. Proteolytic-antiproteolytic balance and its regulation in carcinogenesis / E. Skrzydlewska, M. Sulowska, M. Koda, S. Sulowski // World J. Gastroenterol. - 2005. - 11 (9). - P. 1251-1266.

11. Rubin H. Systemic effects of cancer: Role of multiple proteases and their toxic peptide products / H. Rubin // Med. Sci. Monit. - 2005. - 11 (7). - P. 221-228.

12. Sondak V. K., Taylor J. M., Sabel M. S. et al. Mitotic rate and younger age are predictors of sentinel lymph node positivity: lessons learned from the generation of a probabilistic model // Ann. Surg. Oncol. - 2004. - Vol. 11. - P. 247-258.

1.

2.

3.

4.

13. Sun Z. Role of imbalance between neutrophil elastase and alpha 1-antitrypsin in cancer development and progression / Z. Sun, P. Yang // The lancet oncology. 2004. – 5. – P. 182–190.