

1 Improved Respiratory Characteristics in Non-specific Low Back Pain: Comparison of 2 Feldenkrais Method versus Routine Physiotherapy

3 Abstract

4 **Purpose:** Abnormal breathing patterns, decrease in respiratory muscle strength and
5 endurance are some of the alterations, which are observed in non-specific low back pain (NS-
6 LBP). The purpose of this study was to determine the efficacy of the Feldenkrais method
7 (FM) on respiratory muscle strength, Maximum Voluntary Ventilation (MVV), Total Faulty
8 Breathing Scale (TFBS), Cloth Tape Measure (CTM) and core stability among NS-LBP
9 participants. **Methods:** Participants were recruited from a rehabilitation clinic and
10 randomized either to experimental group (EG) or the control group (CG). For the EG (FM
11 and routine physiotherapy), and for the CG routine physiotherapy alone were carried out
12 three days per week over a period of 8 weeks. Outcome measures including Respiratory
13 Muscle Strength, MVV, TFBS, Numeric Rating Scale (NRS), CTM, and Pressure
14 biofeedback device (PBU) were evaluated at baseline and 8 weeks. **Results:** Forty
15 participants were assigned to an EG (n=20) and CG (n=20) based on the study criteria. There
16 was a significant increase in inspiratory muscle strength (MIP) ($p=0.004$) for the EG, but no
17 significant change in the CG ($p=0.455$). There was also a significant increase in the
18 expiratory muscle strength (MEP) for the EG ($p=0.001$), but no changes in the CG ($p=0.574$).
19 In addition, decrease in pain, increase in xiphoid process chest expansion and improvement in
20 core stability were observed in EG and improvement in MVV was observed in CG.
21 **Conclusions:** FM is a potential training program that can improve respiratory variables
22 among NS-LBP.

23 1. Introduction

24 Feldenkrais is an educational approach whereby people correct their faulty movement
25 patterns through self-exploration of their own bodily movement [1]. The Feldenkrais method
26 (FM) is recommended as an alternate therapy in the field of musculoskeletal practice and is
27 increasingly being used in current practice [2,3]. The FM approach is directed through two
28 methods which are Awareness Through Movement (ATM) and Functional Integrations (FI).
29 The fundamental principles related to efficient use of the neuro-musculoskeletal system in
30 FM are reduction of effort, attending body' parts, speed of movement, coordinated well-
31 learnt action, co-contraction of muscles and respiratory mechanic principles [1]. A key aspect
32 of FM is to pay attention to and develop awareness of breathing to maximize movement
33 patterns, which eases the aggravating symptoms [4]. The FM breathing mechanic principles
34 focus mainly on movement of the diaphragm and movement of the rib cage [1].

35 Recently, there has been renewed interest regarding the involvement of respiratory
36 characteristics in NS-LBP [5,6,7]. A case-control study of 18 participants with Chronic LBP
37 and 29 healthy subjects examined the function of the diaphragm during postural limb
38 activities in performing isometric flexion of upper and lower limbs. The study concluded that
39 participants with chronic LBP had an abnormal diaphragm position and the steeper slope of
40 diaphragm using Magnetic Resonance Imaging [5]. An earlier study hypothesized that the
41 increased respiratory demand compromises spinal control, especially in individuals with LBP

42 [6]. The study was carried out comparing healthy controls to participants with LBP using
43 trans diaphragmatic pressure; findings suggested that the individuals with LBP exhibit greater
44 diaphragm fatiguability compared to healthy controls [6]. Additionally, a recently published
45 study suggested that eight weeks of IMT showed an increased reliance on back
46 proprioceptive signals during postural control, increased in inspiratory muscle strength, and
47 reported a deficit associated with LBP severity [7]. **In addition, it was projected that the**
48 **models such as multifactorial model, a model of movement dysfunction, and ‘Puzzle’ model**
49 **theorized that there existed a relationship between LBP and respiratory variables [8,9,10].**
50 These studies suggest a relationship between LBP and respiratory characteristics. Therefore,
51 the exercises that are related to the respiratory component of FM will be advantageous to
52 LBP population, and there is a clear need to explore this area of research.

53 The existing body of research on FM suggests that FM helps to manage pain for people with
54 LBP following a single session of ATM which was implemented through pre-recorded tape
55 for visualization and breathing sequences [11]. Recently, investigators have examined the
56 efficacy of FM for relieving pain in people with LBP and investigated the improvement of
57 interoceptive awareness, which is the ability to detect internally generated bodily signals
58 involved in maintaining the homeostasis [12]. The intervention used in the study was based
59 on ATM lessons for a period of five weeks. It has been observed that FM was more effective
60 in improving visual analogue scale (VAS) and McGill Pain Questionnaire, Present Pain
61 Intensity scores. [12]. In light of recent evidence in FM, it is becoming extremely difficult to
62 ignore the potential impact of FM on LBP, as it is known that no single intervention is
63 superior to the other for management of LBP. The main challenge faced by these two
64 experiments is the implementation of ATM. However, research has consistently shown that
65 there is improvement following FM irrespective of different ATM approaches. Although,
66 research has been carried out regarding FM and LBP and musculoskeletal disorders, no single
67 study explored the potential impact of respiratory characteristics on NS-LBP [13,14,15].

68 The present study looked at the potential of ATM sessions to influence respiratory
69 characteristics among participants with NS-LBP as FM has a respiratory mechanism as one
70 of the principles related to efficient use of the neuromusculoskeletal system. Hence, the study
71 hypothesized that inclusion of FM would be advantageous to the LBP participants in
72 ameliorating respiratory parameters.

73

74 **2. Materials and methods**

75 2.1 Design

76 The trial was a prospective design with pre-test and post-test evaluation and followed the
77 Consolidated Standards of Reporting Trial statement for Non-pharmacologic treatment [16].
78 This study received ethics approval from local Research Ethics Committee [600-IRMI
79 (5/1/6)/ REC/256/16], and all participants provided informed consent before entering the
80 study.

81

82 2.2 Participants

83 Eligible participants were male or female aged between 18-55 years, diagnosed by the
84 physicians with chronic LBP [17,18] with the pain intensity of LBP in the range of a minimal

85 pain intensity (2/10 – 5/10) by the numeric rating scale (NRS). Participants were excluded if
86 they had any respiratory disease, pregnancy or a history of surgeries to the lumbar spine [7].
87 The study criteria were based on a recent study used by Mohan et al. (2018) [19]. The study
88 was conducted in a Centre of Physiotherapy at a public university. **Initially, leaflets were**
89 **displayed in the rehabilitation clinic of the university hospital. Potential patients who**
90 **approached the researcher were recruited and allocated consequently.**

91

92 2.3 Randomization-sequence generation

93 Two research assistants **that were final year Physiotherapy students who are trained in the**
94 **protocol** were randomly assigned and delivered the protocol; either for the experimental
95 group (EG) or for the control group (CG). Participants were randomly assigned to EG or CG
96 by block randomization using computer randomization method and drawing lots from the
97 concealed envelopes. The assessors remained blinded to the treatment conditions throughout
98 the study.

99

100 2.4 Interventions

101 **The CG received routine physiotherapy using modalities such as infrared rays or**
102 **interferential therapy or shortwave diathermy, spinal flexion or extension exercises whereas**
103 **the EG received a predesigned exercise protocol along with routine physiotherapy (Appendix**
104 **1).**

105 **Both groups received treatment for a period of 8 weeks. The participants in both groups were**
106 **instructed to carry out the exercises 3 days per week. Once a week, the training was**
107 **supervised by a research assistant, and the exercises were progressed based on the patient's**
108 **level of pain. If the level of pain remained the same or reduced, then the exercise was**
109 **progressed. If the patient was unable to maintain the lumbar stability with a pressure of +/- 10**
110 **mmHg using a pressure biofeedback device (PBU), the exercise was not progressed.**

111

112 2.5 Outcomes

113 The primary outcomes were the respiratory muscle strength variables: maximal inspiratory
114 pressure (MIP) and maximal expiratory pressure (MEP), maximum voluntary ventilation
115 (MVV) for measuring respiratory muscle endurance [19]. Secondary outcome measures were
116 **Total Faulty Breathing Scale (TFBS) for assessing faulty breathing pattern [20], Cloth Tape**
117 **Measure (CTM) for measuring chest expansion at the level of axilla, 4th Intercostal space and**
118 **xiphoid [21], NRS for measuring pain level and PBU for core stability [19,22]. The stability**
119 **was tested using 7 levels (level 1 – level 7) with the participant in supine lying with knees**
120 **bent and feet flat on the floor, and the levels of testing were described in previous literature**
121 **[22]. The measurement procedures for all the outcome measures were based on the**
122 **procedures used by Mohan et al. 2018 [19]. The reliability measures of TFBS and CTM were**
123 **established in earlier studies [20,21]. All the outcome measures were evaluated at baseline**
124 **and after 8 weeks of treatment by a blinded assessor.**

125

126 2.6 Sample size

127 MIP which is considered as one of the primary outcomes in the study was used to calculate
128 the sample size using the G*power program 3.1.0 for two tails, paired test. The mean and

129 standard deviation (SD) of MIP were taken from an earlier study for sample size estimation
130 [7]. The estimated sample to obtain a power of minimum 80% at a significant alpha level of
131 95% required a total of 34 participants. Therefore, at least 17 participants with NS-LBP were
132 in both EG and CG to identify a difference between the two interventions. However, to
133 account for the possibility of drop-out during the therapeutic treatment program, 10% of the
134 sample size was added, therefore at least 20 participants per group were included in this
135 study.

136

137 2.7 Statistical methods

138 The data was analysed using SPSS statistical software, version 20.0. The measurement
139 variables were subjected to descriptive and inferential analysis. Description of demographic
140 variables and study variables are presented as mean, standard deviation, frequency and
141 percentage. Results were tested for normal distribution using the Shaapiro-wilk test.
142 Demographic details between the groups were tested using Mann-Whitney U-test. Based on
143 the assumption of normality, Wilcoxon signed rank test were used to compare **baseline** and
144 post **intervention** of the EG and CG.

145 3. Results

146 **A total of 40 participants (n=40; 8 males, 32 females) were recruited and randomized. EG**
147 **(n=20) aged with mean±SD 22.85±2.10 years and CG (n=20) aged with mean±SD**
148 **24.00±2.57 years.** The demographic characteristics showed that there were no significant
149 differences in participants details between EG and CG at baseline. This indicates that the
150 participants in both groups had similar characteristics with regard to age, gender and body
151 mass index (BMI) at the start of the study. The clinical background and the results of the
152 baseline and post values were presented in **Table 1 - Table 4** for primary and secondary
153 variables **‘Insert Table 1, 2, 3 & 4 here’**. **Three participants from each group dropped out**
154 **during the training as they are unable to meet the required follow-ups (Figure 1).** MVV
155 values were lower in both **baseline** and **post intervention** values in CG as compared to EG.

156

157 3.1 Primary outcome variables

158 There was a significant increase in MIP values from **baseline to post intervention** (p=0.004)
159 in the EG. Similarly, with regard to MEP values, there was significant increase in the values
160 (p=0.001) for the EG. On the other hand, there were no significant changes for the MIP and
161 MEP in the CG. There was no significant increase in MVV scores in the EG from **baseline** to
162 post **intervention** (p=0.367). There was a significant increase in respiratory muscle endurance
163 score in CG (p=0.005).

164

165 3.2 Secondary outcome variable

166 In relation to chest expansion the participants in the EG showed improvement at the level of
167 xiphoid process (p=0.004) but did not show improvement at the **level** of the axilla and 4th ICS
168 (p=0.582, and 0.084, respectively). With regard to the CG, the participants did not show
169 improvement in chest expansion for axilla, 4th ICS and xiphoid (p=0.480, 0.679, 0.317,
170 respectively).

171 In relation to NRS values, there was significant reduction in the pain ($p=0.004$) for the EG,
172 but there was no reduction in pain for the CG ($p=0.746$). TFBS scores did not change for the
173 either EG or the CG ($p>0.05$). The scores for the core stability component for the EG
174 ($p=0.001$) and for the CG ($p=0.414$) showed that there was improvement in lumbo-pelvic
175 stability in the EG alone.

176

177 **4. Discussion**

178 This study achieved its aim by improving certain respiratory variables and reducing pain in
179 people with NS-LBP following FM training in EG. Similarly, there were effects on
180 respiratory muscle endurance and on pain among CG exercise training protocols.
181 Specifically, the FM was effective in respiratory muscle strength components, pain and in
182 promoting breathing pattern components. These results corroborate the findings of a great
183 deal of the multifactorial model, a model of movement dysfunction and system-based
184 classification of 'Puzzle' model proposed for the relationship between respiratory variables
185 and LBP [8,9,10]. Therefore, the hypothesis of improving respiratory variables and reducing
186 pain following a pre-designed FM was supported.

187

188 With regard to respiratory muscle strength, there was improvement in both MIP and MEP
189 following FM exercise sessions as compared to CG exercise sessions. The results of the study
190 cannot be compared with other studies related to FM as this is the first study to use these
191 outcome measures in this manner. Most of the studies are qualitative in nature and the
192 outcome measures used are mostly related to pain and interoceptive awareness [11,12]. There
193 was also significant improvement in respiratory muscle endurance following CG, that might
194 be due to the type of exercises which was interspersed from the initial exercise session
195 onwards. It is known that the FM promotes respiratory mechanics rather than respiratory
196 muscle endurance [1].

197

198 Even though, the results of the study cannot be compared directly with earlier research, the
199 results could be compared with relation to respiratory muscle strength. Firstly, trunk
200 stabilizing functions of diaphragm which could have been achieved by promoting symmetry
201 through FM sessions. Secondly, it is assumed that suboptimal position of diaphragm would
202 have been improved because of FM. Potential future studies could explore if there is an
203 association between diaphragm position and the development and recurrence of LBP.

204 The reason behind including pain and lumbo-pelvic instability as one of the outcome
205 measures is, pain can alter an individual's breathing pattern and lumbo-pelvic instability
206 leading to low back pain. The EG reported a greater decrease in pain score compared to the
207 CG. This indicates that the present study results with relation to pain score was supportive of
208 the hypothesis that FM could alter pain through increased body awareness and symmetrical
209 postural alignment [1]. Physiologically, FM is believed to stimulate the neuro-plastic
210 properties of the nervous system. This could have reduced pain through exploration of normal
211 movement, improving a person's neuro-muscular self-image through sensory-motor
212 awareness [1]. In addition, it could be argued, FM might have an impact on descending pain

213 control pathways, may utilise several neurotransmitters in their interaction with the dorsal
214 horn cell pain transmission neurons contributing to a reduction in pain. Fear avoidance that
215 could reduce movement because of an emotional component of pain would have been
216 mitigated through mindful learning of FM [12]. These skills might have helped in organizing
217 the body to transfer to other forms of mental activity there by reducing pain.

218

219 There were changes in xiphoid level chest expansion following FM lessons, but there were no
220 changes in any of the levels of chest expansion in the CG. There was also improvement in
221 breathing pattern from moderate to mild following **EG interventions** as measured by the
222 **TFBS**.

223

224 The changes in breathing pattern and chest expansion would have happened because of
225 emphasis on the body through mindfulness, which is not being considered in their image of
226 movement [12]. In addition, the respiratory mechanics, which are promoted through efficient
227 use neuro-musculoskeletal system would have facilitated an appropriate breathing pattern and
228 improved chest expansion [4]. The brain becomes aware of using a symmetrical breathing
229 pattern through neuroplasticity as a result of mindfulness and body awareness following FM.
230 The significant changes in lumbo-pelvic core stability were observed in FM lesson group
231 alone, and this was not observed in the routine physiotherapy exercise group. A total of three
232 participants achieved level 5 which can be compared with the base line in which none of the
233 participants achieved level 5 among EG. This signifies lumbo-pelvic stability improved
234 through proper positioning and alignment following FM training sessions.

235 **4.1 Limitations**

236 The findings of the study could be viewed in light of a few limitations. First, no **long-term**
237 follow-up tests was conducted which could establish longer-terms effects of the intervention.
238 Second, most of the participants were younger females which could limit external validity of
239 the findings. Thirdly, the participants had mild-moderate pain intensity, and this data might
240 not be applicable for those participants with severe pain. In addition, the study did not
241 consider data imputation technique for the dropped-out participants, and there was a
242 significant difference in baseline value between the group which need to be interpreted
243 carefully while interpreting the study results.

244 **5. Conclusions**

245 FM technique is suggested to be a potential additional exercise for participants with LBP
246 which could improve respiratory, pain and lumbo-pelvic stability components. **Further**
247 **research is needed to compare FM with other forms of physiotherapy exercises in order to**
248 **clarify their effects, and the potential of combination of exercises with FM in treating LBP.**

249 **Acknowledgments** The authors would like to thank the study participants who participated in
250 this study and Dr. Kathryn Collins, Bournemouth University who spent her valuable time in
251 editing and proofreading the manuscript.

252

253 **References**

- 254 [1] Lyttle T. The Feldenkrais method: application, practice and principles. *J Bodyw Mov*
255 *Ther.* 1997;1:262-269.
- 256 [2] Henry LJ, Paungmali A, Mohan V, Ramli A. Feldenkrais method and movement
257 education – An alternate therapy in musculoskeletal rehabilitation. *Polish Ann Med.*
258 2016;23:68-74.
- 259 [3] Mohan V, Paungmali A, Silitertpisan P, Henry LJ, Mohamad NB, Kharami NNB.
260 Feldenkrais method on neck and low back pain to the type of exercises and outcome
261 measurement tools: A systematic review. *Polish Ann Med.* 2017;24:77-83.
- 262 [4] Mohan V, Paungmali A, Silitertpisan P. Letter to Editor: Application of neuroplasticity
263 theory through the use of the Feldenkrais Method with a runner with scoliosis and
264 lower quarter pain: Additional respiratory mechanics principle, implication of the
265 Feldenkrais method for clinical practice. *J Bodyw Mov Ther.* 2017;21:470-471.
- 266 [5] Kolar P, Sulc J, Kyncl M, Sanda J, Cakrt O, Andel R, Kumagai K, Kobesova A. Postural
267 function of the diaphragm in persons with and without chronic low back pain. *J Orthop*
268 *Sports Phys Ther.* 2012; 42: 352-362.
- 269 [6] Janssens L, Brumagne S, McConnell AK, Hermans G, Troosters T, Gayan-Ramirez G.
270 Greater diaphragm fatigability in individuals with recurrent low back pain. *Respir*
271 *Physiol Neurobiol.* 2013;188:119-123.
- 272 [7] Janssens L, McConnell AK, Pijnenburg M, et al. Inspiratory muscle training affects
273 proprioceptive use and low back pain. *Med Sci Sports Exerc.* 2015;47:12-19.
- 274 [8] Richmond J. Multi-factorial causative model for back pain management; relating
275 causative factors and mechanisms to injury presentations and designing time- and cost
276 effective treatment thereof. *Med Hypotheses.* 2012;79:232-240.
- 277 [9] Key J, Clift A, Condie F, Harley C. A model of movement dysfunction provides a
278 classification system guiding diagnosis and therapeutic care in spinal pain and related
279 musculoskeletal syndromes: a paradigm shift-Part 1. *J Bodyw Mov Ther.* 2008;12:7-
280 21.
- 281 [10] Lee DG, Lee LJ, McLaughlin L. Stability, continence and breathing: the role of fascia
282 following pregnancy and delivery. *J Bodyw Mov Ther.* 2008;12(4):333-348.
- 283 [11] Smith A, Kolt G, McConville J. The Effect Of The Feldenkrais Method On Pain And
284 Anxiety In People Experiencing Chronic Low Back Pain. *New Zeal J Physiother.*
285 2001;29:6-14.
- 286 [12] Paolucci T, Zangrando F, Iosa M, et al. Improved interoceptive awareness in chronic low
287 back pain: a comparison of Back school versus Feldenkrais method. *Disabil Rehabil.*
288 2017;39:994-1001.
- 289 [13] Malmgren-Olsson E-B, Armelius B-A, Armelius K. A comparative outcome study of
290 body awareness therapy, feldenkrais, and conventional physiotherapy for patients
291 with nonspecific musculoskeletal disorders: changes in psychological symptoms,
292 pain, and self-image. *Physiother Theory Pract.* 2001;17:77–95.
293 doi:10.1080/095939801750334
- 294 [14] Pugh JD, Williams AM. Feldenkrais method empowers adults with chronic back pain.
295 *Holist Nurs Pract.* 2014;28:171–83. doi:10.1097/HNP.000000000000026.

296 [15] Myers LK. Application of neuroplasticity theory through the use of the Feldenkrais
297 Method® with a runner with scoliosis and hip and lumbar pain: A case report. *J*
298 *Bodyw Mov Ther.* 2016;20:300–309. doi:10.1016/j.jbmt.2015.06.003.

299 [16] Boutron I, Moher D, Altman DG, Schulz KF, Ravaud P. Extending the CONSORT
300 statement to randomized trials of nonpharmacologic treatment: Explanation and
301 elaboration. *Ann Intern Med.* 2008:295-309.

302 [17] Brumagne S, Janssens L, Janssens E, Goddyn L. Altered postural control in anticipation
303 of postural instability in persons with recurrent low back pain. *Gait Posture.*
304 2008;28:657-662.

305 [18] Lawand P, Lombardi Júnior I, Jones A, Sardim C, Ribeiro LH, Natour J. Effect of a
306 muscle stretching program using the global postural reeducation method for patients
307 with chronic low back pain: A randomized controlled trial. *Jt Bone Spine.*
308 2015;82:272-277.

309 [19] Mohan V, Paungmali A, Sitilerpisan P, Hashim UF, Mazlan MB, Nasuha TN.
310 Respiratory characteristics of individuals with non-specific low back pain: A cross-
311 sectional study. *Nurs Health Sci.* 2018;1-7.

312 [20] Mohan V, Perri M, Paungmali A, et al. Intra-rater and inter-rater reliability of total faulty
313 breathing scale using visual observation and videogrammetry methods. *J Bodyw Mov*
314 *Ther.* 2017;21:694-698.

315 [21] Mohan V, Dzulkifli NH, Justine M, Haron R, Leonard Joseph H, Rathinam C. Intrarater
316 reliability of chest expansion using cloth tape measure technique. *Bangladesh J Med*
317 *Sci.* 2012;11:307-311.

318 [22] Hagins M, Adler K, Cash M, Daugherty J, Mitrani G. Effects of Practice on the Ability to
319 Perform Lumbar Stabilization Exercises. *J Orthop Sport Phys Ther.* 1999;29:546–55.
320 doi:10.2519/jospt.1999.29.9.546.

321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339

340 **Table 1**
 341 Demographic Details of Participants between Experimental and Control Groups [mean +/-
 342 SD; number (%)]

Characteristics	Experimental (n=20)	Control (n=20)
Age (Years)	22.85±2.10	24.00±2.57
BMI (Kg/m ²)	23.99±4.20	25.25±5.64
Gender (%)	F- 16 (80%) M- 4 (20%)	F- 16 (70%) M- 4 (30%)

343 **Note:** No significant differences in participants' demographics between groups (p>0.05)

344
 345
 346 **Table 2**
 347 Comparison of the Primary Outcome variables (MVV, MIP, MEP) between Experimental and Control
 348 Groups [data represented as mean (95% CI)]

Parameters	Groups	Before [Experimental: n=17, Control :n=17]	After [Experimental: n=17, Control :n=17]
MVV (l/min)	Experimental Control	95.27 (86.18 to 104.36) 75.47 (63.60 to 87.33)	93.61 (85.13 to 102.09) 87.49 (76.28 to 98.71) ^a
MIP (cm H ₂ O)	Experimental Control	61.47 (52.80 to 70.13) 76.64 (66.46 to 86.83)	70.88 (63.23 to 78.53) ^a 75.23 (66.61 to 83.85)
MEP (cm H ₂ O)	Experimental Control	52.17 (46.56 to 57.78) 61.23 (53.63 to 68.83)	62.94 (56.92 to 68.95) ^a 62.05 (54.26 to 69.85)

349 **Note:** ^aSignificant change within group (p<0.05) from pre- to post

350
 351
 352
 353
 354
 355
 356
 357
 358
 359
 360
 361
 362
 363
 364
 365
 366
 367
 368
 369
 370
 371

372 **Table 3**
 373 Comparison of the Cloth Tape Measure (CTM) at different levels and Numerical Rating Scale (NRS)
 374 between Experimental and Control Groups [data represented as mean (95% CI)]

Parameters	Groups	Pre- Values (Experimental: n=17, Control: n=17)	Post Values (Experimental: n=17, Control: n=17)
Axilla (cm)	Experimental	1.62 (1.39 to 1.84)	1.53 (1.12 to 1.94)
	Control	1.41 (1.15 to 1.67)	1.29 (1.05 to 1.53)
4 th ICS (cm)	Experimental	1.31 (1.08 to 1.55)	1.57 (1.26 to 1.89)
	Control	1.55 (1.32 to 1.79)	1.52 (1.22 to 1.93)
Xiphoid (cm)	Experimental	1.33 (1.06 to 1.60)	1.81 (1.44 to 2.17) ^a
	Control	2.11 (1.72 to 2.50)	2.17 (1.80 to 2.55)
Numerical Rating Scale (10)	Experimental	3.58 (2.51 to 4.66)	1.23 (.567 to 1.90) ^a
	Control	2.88 (2.34 to 3.42)	2.41 (1.86 to 2.95) ^a

375 **Note:** ^aSignificant change within group (p<0.05) from pre- to post


376 **Table 4**
 377 Comparison of Total Faulty Breathing Scale (TFBS) and lumbo-pelvic core stability using pressure
 378 biofeedback device between Experimental and Control Groups [represented as number (%)]
 379

Parameters	Groups	Pre-Values (Experimental: n=17, Control: n=17)	Post Values (Experimental: n=17, Control: n=17)
Total Faulty Breathing Scale (TFBS)	Experimental	Mild- 16(94%) Moderate – 1(6%)	Mild- 17(100%)
	Control	Mild- 17(100%)	Mild- 17(100%)
Pressure biofeedback device (mmHg)	Experimental	Level 0- 2(12%)	Level 0- 1(6%) ^a
		Level 1- 8(47%)	Level 1- 1(6%)
	Control	Level 2- 2(12%)	Level 2 - 5(29%)
		Level 3- 4(23%)	Level 3 - 4(23%)
Experimental	Level 4- 1(6%)	Level 4 - 3(18%)	
	Control	Level 1- 2(12%)	Level 5 - 3(18%)
Control	Level 2- 2(12%)	Level 1- 2(12%)	
	Level 3- 8(47%)	Level 2- 1(6%)	
Experimental	Level 4- 5(29%)	Level 3- 9(53%)	
	Control	Level 4- 4(23%)	Level 4- 4(23%)
Experimental	Level 5- 1(6%)	Level 5- 1(6%)	
	Control	Level 5- 1(6%)	Level 5- 1(6%)

380
 381
 382
 383
 384
 385
 386
 387

388 **Figure 1- Flow of participants**

389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414



Appendix 1 - Feldenkrais Method Training Protocols

415

416 Week 1

417 1. Tilting legs:

418 **Patient position:** Initially, the participant were asked to lie on their back, with the knees bent
419 and the soles of the feet in contact with the floor.

420 **Instruction for Movements:** Then gently, they were asked to let the knees tilt a little bit to
421 the left, and then smoothly move to tilt them to the right. Make each repetition a little bit
422 different – smoother, softer, easier, more comfortable. Try slowing down the breath so that
423 when inhaling tilt the knees and while exhaling bring them back to the middle.

424 **Variation 1:** Movements are tried in knees close together and knees apart to know which
425 position is comfortable.

426 **Variation 2:** Cross the right knee over the left. Reposition the knees on the floor if the
427 subjects are fully comfortable

428 **Duration:** 1 hour **Rest period:** 3 minutes between each set of educational program

429 Week 2

430 2. Pelvic tilt:

431 **Patient position:** Lie on the back, with knees bent and soles of the feet in contact with the
432 floor.

433 **Instruction for movements:** The participants are instructed to feel the flat, low back or
434 slowly they are asked to flatten the back to feel the roll on the back of the pelvis. This
435 reminded the spine that it can change the shape.

436 **Duration:** 1 hour **Rest period:** 3 minutes between each set of educational program.

437 Week 3

438 3. Spine like a chain:

439 **Patient position:** Same position as above.

440 **Instruction for movements:** Same as above exercises the participant should feel the lower
441 back to flatten into the floor. Then they are instructed to go little farther in that direction and
442 feel the tailbone peak out into the room. Roll back down, take an easy breath and then roll
443 again, but a bit farther this movement in order to feel the sacrum.

444 **Duration:** 1 hour **Rest period:** 3 minutes between each set of educational program

445 Week 4

446 4. Prone kneeling:

447 **Patient position:** The arms need to be at right angle to the torso and the knees can be directly
448 below your hip joints.

449 **Instruction for movements:** Instruction was given such that belly is relaxed and hand down
450 toward the floor. Then, gently pull the belly in. keep the movement small enough and gentle
451 enough so that entirely the participants felt comfortable.

452 **Duration:** 1 hour **Rest period:** 3 minutes between each set of educational program

453 **Week 5**

454 **5. Prone lying:**

455 **Patient position:** Lie on the front and rest the arms on the floor on either side of the head.
456 Let the legs be long and extended, comfortably apart, with the feet resting so that toenails are
457 on the floor.

458 **Instruction for movements:** Comments were given such that to turn the heels to the left and
459 then to the right. At the same time, the pupils should notice turning the heels rolls the pelvis,
460 as rolling across the tummy from one hip-bone to the other. Then, keep rolling across the
461 tummy to roll the pelvis and see how the heel follows.

462 When the heels are pointing to the left and the right leg needs to roll onto its inner edge, and
463 draw up the knee towards the abdomen. Then let it straighten again. Do the exercise for
464 several times and then rest.

465 For each and every exercise the participants are supposed to stand up easily, walk around a
466 bit, and feel comfortable.

467 **Duration:** 1 hour **Rest period:** 3 minutes between each set of educational program

468 **Week 6 -8**

469 All the above mastered techniques were carried out together for a period of 1 hour with rest
470 periods in between the exercise program.

471 There was one session per week, which were supervised for 1 hour for 8 consecutive weeks
472 and the subjects were instructed to perform the exercises 3 days in a week. Each exercise was
473 progressed until 5 weeks and for the last three weeks the whole set of exercises was given.

474

475

476

477

478

Specific Comments	Answers to Reviewers Comments
Can insert information here on where participants were recruited from. E.g. participants were recruited from xxxxx and randomised to either the experimetna group (EG) or the control group (CG)	We have included the sentence as recommended as ‘Participants were recruited from a rehabilitation clinic and randomized either to experimental group (EG) or the control group (CG)’
This is a result move to the results section.	We have moved this to the results section as ‘Forty participants were assigned to an EG (n=20) and CG (n=20) based on the study criteria’
How often? Daily?	We have rephrased this sentence as ‘For the EG (FM and routine physiotherapy), and for the CG routine physiotherapy alone were carried out three days per week over a period of 8 weeks’
Than what?	Thanks for asking this. We would like to inform that FM was effective in improving musculoskeletal parameters and no single study explored the potential impact of respiratory characteristics on NS-LBP. We have mentioned this part in the last line of the paragraph.
Was this not a randomised controlled trial?	This is not a randomised controlled trial
You could provide more information here, were their adverts circulated? Were they patients? How were the participants identified?	We have modified the sentence as ‘Initially, leaflets were displayed in the rehabilitation clinic of the university hospital. Potential patients who approached the researchers were recruited and allocated consequently’ .
What qualifications/training did they have?	We have refrained the contents as ‘Two research assistants that were final year Physiotherapy students who are trained in the protocol were randomly assigned and delivered the protocol;’
It might be helpful to explain both the intervention and the control group separately as it is confusing here when you report both groups and then later on go on to distinguish the control group saying they received spinal flexion or extension exercises was this in addition to the other exercises.	We have revised the whole contents to make it clear. First, we presented both the groups and then presented the level of progression of exercises.

<p>Two research assistants supervised the session is this clear?</p>	<p>We have made this clear by mentioning ‘Once a week, the training was supervised by a research assistant, and the exercises were progressed based on the patient’s level of pain. If the level of pain remained the same or reduced, then the exercise was progressed’.</p>
<p>Please insert the references beside the outcome measure they correspond to.</p>	<p>I have inserted the reference as suggested. Reference 19 corresponds to all outcome measures as these outcome measures are referred in the same literature. References 20 & 21 are referenced beside the outcome measures as they correspond to.</p>
<p>Measured at both the level of the xiphoid and the axilla</p>	<p>We have rephrased the whole sentence as ‘Secondary outcome measures were Total Faulty Breathing Scale (TFBS) for assessing faulty breathing pattern [20], Cloth Tape Measure (CTM) for measuring chest expansion at the level of axilla, 4th Intercostal space and xiphoid [21], NRS for measuring pain level and PBU for core stability [19,22]’.</p>
<p>What position was the participant in for this? As described in level 2</p> <p>This is not explained very clearly. I wonder would it be more beneficial to explain each level clearly with a diagram in an appendices or simply reference where these can be found</p>	<p>The stability was tested using 7 levels (level 1 – level 7) with the participant in supine lying with knees bent and feet flat on the floor, and the levels of testing were described in previous literature [22].</p> <p>We have revised the sentences and referenced as 22 as suggested for its clarity.</p>
<p>For which outcome measures?</p>	<p>We have rephrased the sentence as ‘The measurement procedures for all the outcome measures were based on the procedures used by Mohan et al. 2018’.</p>
<p>Introduce here N=40 participants were recruited and randomised. And some demographic details here: age/gender of the</p>	<p>We have detailed the required details are suggested.</p>

<p>two groups</p> <p>Do you have a CONSORT flow diagram?</p> <p>How many were ineligible and to details reasons for dropout?</p>	<p>‘A total of 40 participants (n=40; 8 males, 32 females) were recruited and randomized. EG (n=20) aged with mean±SD 22.85±2.10 years and CG (n=20) aged with mean±SD 24.00±2.57 years’</p> <p>Yes, we have attached along with the revised script as Figure 1.</p> <p>‘Three participants from each group dropped out during the training because as they are unable to meet the required follow-ups’</p>
<p>Be consistent in use of language i.e. baseline and post intervention</p>	<p>We have refrained the language as recommended</p>
<p>Think about what was your most important finding? This this study achieve its aim. This is all a repetition of the background, consider the This is the first time mentioning “the puzzle” model. I think that this should have been brought in in the background.</p>	<p>We have rephrased the whole contents of the discussion as ‘This study achieved its aim by improving certain respiratory variables and reducing pain in people with NS-LBP following FM training in EG. Similarly, there were effects on respiratory muscle endurance and on pain among CG exercise training protocols. Specifically, the FM was effective in respiratory muscle strength components, pain and in promoting breathing pattern components. These results corroborate the findings of a great deal of the multifactorial model, a model of movement dysfunction and system-based classification of ‘Puzzle’ model proposed for the relationship between respiratory variables and LBP’</p> <p>In addition, we have included those three models in the background.</p>
<p>This is important if this is the first study to use these outcome measures in this manner, then you need to highlight this.</p>	<p>We have rephrased the sentence as ‘The results of the study cannot be compared with other studies related to FM as this is the first study to use these outcome measures in this manner’.</p>
<p>Is it subjective? Qualitative insinuates that they conducted qualitative research....</p>	<p>Yes, most of the studies are qualitative in nature and they are subjective.</p>
<p>I don’t think these sub headings are necessary</p>	<p>We have removed all the sub-heading from the discussion as recommended</p>
<p>Recommend deleting this, it is a repeat of</p>	<p>We have deleted the repeat of results as</p>

the results.	suggested.
Is there any reference for this?	We do not have direct reference to FM. Its our inference.
I was not aware that any group did FM alone?	We have rephrased the sentence as ‘There was also improvement in breathing pattern from moderate to mild following EG interventions as measured by the TFBS ’.
Is further research needed to explore this?	Yes, we have added a sentence to the conclusion as ‘ Further research is needed to compare FM with other forms of physiotherapy exercises in order to clarify their effects, and the potential of combination of exercises with FM in treating LBP ’.

480

481