

Acute Respiratory Distress Syndrome and Prone Position in Covid-19

Abstract

Patients who develop severe Covid-19 disease can develop respiratory failure and subsequently Acute Respiratory Distress Syndrome (ARDS). However, it has to be noted that these patients may not follow the typical ARDS disease trajectory. The causes of this paradox are complex and not yet fully understood, with the result that varying pathophysiological hypotheses have been proposed. This article describes ARDS in Covid-19 patients, the use of the conscious and unconscious prone position as an intervention to improve oxygenation.

Key Points

- Covid-19 can lead to respiratory failure and Acute Respiratory Distress Syndrome.
- The cause of ARDS in Covid-19 is complex and not completely understood.
- The high incidence of ARDS in Covid-19 patients has resulted in the use of the prone position being undertaken early because for both conscious and unconscious patients it can improve oxygenation.

Key Words

ARDS, Prone Position, Covid-19, SARS-CoV-2, Pandemic, Critical Care

Biographies

Rosie Recce-Anthony BSc (Hons) RN(A) is Critical Care Research Nurse at University Hospital Lewisham, London. She is a qualified and experienced critical care nurse.

Competing interests: none declared.

Grace Lao RN(A) is Critical Care Assistant Research Nurse at University Hospital Lewisham, London. She is an experienced nurse who has specialised in critical care.

Competing interests: none declared.

Chris Carter MEd BSc (Hons) DiPHE RN(A) is Senior Lecturer at Birmingham City University and Visiting Professor at Foshan Hospital of Traditional Chinese Medicine and the Sixth Affiliated

Hospital of Guangzhou Medical University, China. A specialist critical care nurse expert in resource-limited service delivery, he was deployed to Afghanistan and has worked in Oman and Zambia. He supports projects in Vietnam, Somaliland and Jamaica. He is a member of the Royal College of Nursing International Committee. Competing interests: none declared.

Joy Notter PhD MSc SRN RHV CPT HVT PGCEA is Professor of Community Health Care at Birmingham City University, and Professor Honoris Causa, Hanoi Medical University, Viet Nam. She is a fellow of the Royal Society of Medicine and is Past President of the European Association for Cancer Education. Competing interests: none declared.

Covid-19 causes acute respiratory failure, with an estimated 10% of patients developing acute respiratory distress syndrome (ARDS), which is associated with a high mortality, of approximately 30- 40% despite advanced treatment [1]. In consequence, the rapid increase in critical care need, resulted in many hospitals, regions and countries coming close to, or being overwhelmed, by the unprecedented number of patients. As knowledge and understanding of this new Coronavirus advances, effective treatments measures have been increasingly identified. One of these, prone positioning, was not commonly used in critical care units prior to the Covid-19 pandemic [2-4], has revolutionised the treatment of both ventilated and non-ventilated patients. However, its re-introduction has confirmed that it should not be seen as a stand-alone measure, to be effective it needs to be a core component of a series of structured interventions. This chapter, explores the use of the prone position in ARDS arising from Covid-19.

Respiratory Failure

The Covid-19 disease progression may range from mild to severe [5]. High numbers of hospitalised patients develop respiratory symptoms, with reported incidence of over 80% of patients needing oxygen therapy [6-7]. Patients with increasingly severe COVID-19 symptoms may go on to develop acute respiratory failure and subsequently Acute Respiratory Distress Syndrome (ARDS). It has to be noted that these patients may not follow the typical ARDS disease trajectory [8]. Within the literature, this has been described by the term 'happy hypoxemia' or 'silent hypoxia', when an individual has profound hypoxia caused by Covid-19, but does not have the proportional signs of respiratory distress [9-11]. In consequence, there has been a suggestion that the term CARDS (Covid-19 with ARDS) should be used instead of the traditional ARDS definition [8]. The causes of this paradox is complex and not yet fully understood, with the result that varying pathophysiological hypothesis have been proposed [8, 12].

The phenotype theory hypothesis is that in severe COVID-19 disease, there is a systemic impact on the vascular endothelium, causing lung injury. Marini and Gattinoni [8] propose that in Covid-19, Type L and Type H phenotypes cause different variants of respiratory failure. Type L patients have a scattered ground glass appearance on chest X-ray, with good lung

compliance and tend not to be Positive End Expiratory Pressure (PEEP) responsive. Contrastingly, individuals with Type H respiratory failure have extensive infiltrates with both atelectasis and oedema on chest X-ray, adding further complexity to the disease management. These patients have a lower lung compliance and are PEEP responsive. However, Marini and Gattinoni [8] describe the type L and H phenotypes as a continuum, with some stages and characteristics overlapping. They argue that during the early phases of respiratory failure, a complex process of pulmonary vascular dysregulation occurs, which instead of causing alveolar oedema, leading to hypoxemia, and a high minute volume ventilation, but they exhibit no signs of respiratory failure. As the pulmonary vascular dysregulation continues and extends, causing vasoplegia, the lungs become increasingly unable to regulate perfusion and maintain adequate ventilation. This exacerbates hypoxemia, causes dead space ventilation and hypercapnia, leading to ARDS.

Jain and Doyle [12] dispute the above phenotype hypothesis and argue that the Type L phenotype relates to stage 2 or 3 of COVID-19 pneumonia. They propose a different pathophysiological process for the cause of severe hypoxia. They suggest that the SARS-CoV-2 enters the type II alveolar epithelial cells binding to the spike protein of the ACE-2 receptor. This in turn leads to downregulation of the alveolar epithelium, allowing for ACE-1 to have an unregulated effect on the pulmonary capillary endothelial cells. The level of protective ACE2-Ang-1-7-mas-R production is reduced, resulting in a harmful increase in the level of ACE 1_AngII-ATI-R. Vasoconstriction in the pulmonary epithelium is caused by endothelin-1. This causes a complex cascade effect, which results in endothelial nitric oxide being inhibited. The associated severe pulmonary vasoconstriction is unevenly distributed within the lungs, as the shunt fraction increases with the increasing hypoxia, the alveolar-capillary barrier is disrupted. The flooding of proteins, fibrin, cells and fluid into the alveolar space causes bilateral patchy ground glass opacities noted on CT scan or Chest X-Ray. The development of pulmonary symptoms and rapid disease progression associated with COVID-19 may be linked to the 'endothelial-epithelial' interaction. Following alveolar-capillary membrane disruption, SARS-CoV-2 is able to enter the pulmonary capillary membrane via the pulmonary capillaries. The pulmonary endothelial cells become infected via the ACE-2 protein on the luminal surfaces, to assume a 'proinflammatory / procoagulant' phenotype. This accelerates apoptosis of alveolar epithelial and endothelial cells and causes a cytokine storm [13].

The decision on when to intubate patients, remains largely subjective and based on practitioners experience and patient's condition [14]. Mohlenkamp et al [15] found that 5-15% of patients with Covid-19 require critical care and ventilatory support, with 17% patients developing ARDS [1]. ARDS was first described in critically ill patients in 1967 by Ashbaugh et al. [16]. However, it was not until 1994, then the first clinical definition was agreed by the International American European Consensus Conference (AECC) [17]. The Berlin Definition aimed to classify the severity of ARDS and to publish treatments and ventilatory strategies depending on the degree of hypoxemia [18]. ARDS is defined using the Berlin Criteria and is based on timing, imaging, evidence of oedema and oxygenation. ARDS, this is defined [19] as:

‘an acute diffuse, inflammatory lung injury, leading to increased pulmonary vascular permeability, increased lung weight, and loss of aerated lung tissue...[with] hypoxemia and bilateral radiographic opacities, associated with increased venous admixture, increased physiological dead space and decreased lung compliance.’

ARDS is described as respiratory failure with an acute onset which affects both lungs and occurs within one week of either a clinical insult or deterioration in respiratory symptoms. Chest imaging reveals bilateral opacities not fully explained by effusions, lung collapse, and nodules. Respiratory failure not explained by cardiac failure and fluid overload and oxygenation:

- Mild $\text{PaO}_2/\text{FiO}_2 \leq 39.9\text{kPa}$ ($\leq 300\text{mmHg}$) with PEEP or CPAP $\geq 5\text{cmH}_2\text{O}$
- Moderate $\text{PaO}_2/\text{FiO}_2 \leq 26.6\text{kPa}$ ($\leq 200\text{mmHg}$) with PEEP $\geq 5\text{cmH}_2\text{O}$
- Severe $\text{PaO}_2/\text{FiO}_2 \leq 13.3\text{kPa}$ ($\leq 100\text{mmHg}$) with PEEP $\geq 5\text{cmH}_2\text{O}$

[19, 20, 21].

Limitations with the current ARDS definition, include that severity can be assessed on a single blood gas without prior standardisation of ventilator settings including PEEP which may affect oxygenation. In consequence, it is recommended that ventilator settings are optimised using tidal volumes of 6ml/kg of predicted body weight and a high PEEP level [22]. Furthermore, the time from optimising ventilator settings and assessing $\text{PaO}_2/\text{FiO}_2$ is deemed more clinical relevant in ARDS classification when measured 24 hours after ARDS onset [22].

ARDS is an acute inflammatory lung condition not a disease, with multi-factorial causes and no proven drug treatments. Therefore, ARDS is always caused by an underlying pulmonary or extra-pulmonary condition. Pulmonary ARDS occurs when there is a direct insult to the lung

damaging the alveolar epithelium, while extra-pulmonary ARDS is caused by an indirect lung injury due to inflammatory mediators damaging the vascular endothelium [23, 24]. Pulmonary ARDS can be caused by bacterial, viral or fungal pneumonia, aspiration of gastric contents, inhalation contusion, pulmonary contusion, and pulmonary vasculitis or near drowning. Extra-pulmonary ARDS can be triggered by non-pulmonary sepsis, non-cardiogenic shock, pancreatitis, major trauma, multiple transfusion or transfusion-related acute lung injury, severe burns or drug overdose [25-26].

The majority of patients with Covid-19 who develop ARDS meet the Berlin Criteria [27]. In patients who develop ARDS, management includes using lung protection strategies: (low volume, low pressure ventilation). Initial ventilation strategies may include pressure controlled modes, with tidal volumes aimed at 6ml/kg using predicted body weight and plateau airway pressure <30cmH₂O. Initial ventilator settings may include a higher respiratory rate (20/min), with positive end expiratory pressure (PEEP). Early evidence suggests pressures may need to be lower than previously recommended [4]. There needs to be pre-oxygenation of patients prior to any intervention e.g. suctioning to prevent prolonged periods of desaturation. Moderate hypoxemia with SpO₂ targets of >90% and PaO₂ of >8kPa with permissive hypercapnia.

The use of nitric oxide or nebulised prostacyclin if using a wet circuit has been noted to improve vasodilation and thus improve oxygenation and reducing airway pressures [4]. PEEP may be used to recruit collapsed alveoli; however, high PEEP should be avoided. Recruitment manoeuvres are used to improve oxygenation by providing brief inspiratory flow cycles to maximum plateau pressure and inflating collapsed alveoli. Recruitment manoeuvres tend to be used as a rescue therapy in severe refractory hypoxia or following accidental disconnection from the ventilator. The procedure remains controversial in routine care, as it provides a temporary increase in oxygenation which is not sustained, and therefore must only be performed by an experienced practitioner [28].

For patients who do not respond and become increasingly difficult to ventilate and oxygenate, the prone position may be considered. With Extra Corporeal Oxygenation Membrane (ECMO) services being limited in many settings during a pandemic, the prone position, may be used as an alternative in an attempt to improve oxygenation and optimise lung compliance [29].

Conservative use of intravenous fluids and careful fluid balance monitoring, with the use of diuretics to remove excess fluid should be considered provided it is not detrimental to other organs.

Considerations for Resource Limited Environments

It is accepted that identifying and applying the internationally agreed ARDS definition in a resource limited setting may be difficult, due to limited availability of resources [30]. In these situations, it may be appropriate to use the Kigali modified ARDS definition and criteria (table xxx). The main difference relates to assessment of deteriorating respiratory deterioration, using the SpO₂/FiO₂ calculation [30].

Table 1: Similarities and differences between Kigali modified ARDS Definition and Berlin Definition of ARDS

Kigali modified ARDS definition	Berlin Definition of ARDS
Acute onset affecting both lungs occurring within one week of: <ul style="list-style-type: none"> • Clinical insult #Deterioration in respiratory symptoms 	Acute onset affecting both lungs occurring within one week of: <ul style="list-style-type: none"> • Clinical insult • Deterioration in respiratory symptoms
Chest X-ray or ultrasound showing: <ul style="list-style-type: none"> • Bilateral opacities not fully explained by effusions, • Lobar/lung collapse • Nodules. 	Chest imaging showing: <ul style="list-style-type: none"> • Bilateral opacities not fully explained by effusions • Lung collapse • Nodules.
Respiratory failure not fully explained by cardiac failure or fluid overload	Respiratory failure not explained by cardiac failure and fluid overload
SpO ₂ /FiO ₂ <315 No Positive End Expiratory Pressure (PEEP) requirement	Mild PaO ₂ /FiO ₂ ≤39.9kPa (≤300mmHg) with PEEP or CPAP ≥5cmH ₂ O Moderate PaO ₂ /FiO ₂ ≤26.6kPa (≤200mmHg) with PEEP ≥5cmH ₂ O Severe PaO ₂ /FiO ₂ ≤13.3kPa (≤100mmHg) with PEEP ≥ 5cm ₂ O
[30]	[19-21]

Prone Position

The prone position involves repositioning the patient from the supine position onto their abdomen. It redistributes perfusion and improves ventilation/perfusion (VQ) matching, through maximising dorsal ventilation. This results in recruitment of the posterior lung segments reverse atelectasis and improved secretion clearance [28]. The prone position was first described in the literature in 1974, as a way to improve oxygenation [31]. Since that date, research has consistently shown that oxygenation can be improved in ventilated patients with ARDS by turning them into the prone position [32-35]. However, more recent studies have shown that prone position improves mortality in moderate to severe ARDS if undertaken early. The position needs to be maintained for 16-18 hours or more while using protective lung strategies [36]. This is a change from traditional practice, in which the prone position was solely used for ventilated patients, however, more recently, experience has shown a beneficial response to prone position by COVID-19 patients not yet requiring invasive ventilation [28]. Early recommendations are that proning patient on admission to ICU during the early phase of their disease may be beneficial and avoid more aggressive ventilation strategies. It can be used irrespective of the PF ratio [4]. Both conscious and unconscious prone position methods are described here.

Conscious Prone Position

For conscious patients with suspected or confirmed Covid-19 who requiring oxygen of >28% or basic respiratory support to achieve SaO₂ 92-96% (88-92% if high risk of hypercapnic respiratory failure) may gain benefit from the conscious prone position. If tolerated, this position can improve oxygenation, reducing the need for non-invasive ventilation (NIV). It has been found that it can delay and/or avert the need for intubation and mechanical ventilation [7, 28]. The current pandemic has also revealed that this is a simple and safe intervention that is suitable for use on general wards.

If there is an improvement in SaO₂ 92-96% (88-92% if risk of hypercapnic respiratory failure) and no obvious distress, the prone position should continue, with a view of changing the patients position every 1-2 hours or longer if possible. When not in the prone position, the patient should be nursed in a 30-60 degree upright position. Vital signs including oxygen saturations and early warning scores should be monitored after every position change and oxygen titrated accordingly. If tolerated continued timed position changes can be used. A

proposed regimen includes 30 minutes to 2 hours in the following positions lying fully prone with the bed flat, lying on their right side with the bed flat, sitting upright 30-60 degrees, lying on left side with the bed flat, prone position again and then repeated. The position should be discontinued if there is no improvement, the patient is unable to tolerate the position, the respiratory rate is ≥ 35 , there is evidence of tiring and / or the use of accessory muscles [28]. If appropriate, the patient should be reviewed by critical care and assessment made regarding transfer into critical care.

Absolute contraindications include respiratory distress ($RR \geq 35$, $PaCO_2 \geq 6.5$, and / or accessory muscle use), the immediate need for intubation, haemodynamic instability (systolic blood pressure < 90 mmHg) or arrhythmia, agitation or altered mental status, unstable spine/thoracic injury/recent abdominal surgery. Relative contraindications include facial injuries, neurological issues, morbid obesity, pregnancy (2/3rd trimesters) and pressure sores / ulcers [28].

To turn a conscious patient into the prone position, it is important to explain the importance of the procedure to the patient, to provide reassurance, to improve oxygenation and reduce their chance of requiring invasive ventilation. Patients should be encouraged to remain in the prone position for as long as possible, ideally up to 18 hours per 24 hours. Patients must be assisted into and out of the prone position, and patients should not do this without assistance in case their oxygen levels drop during the turn. There are two ways the patient can position themselves in the prone position either from a sitting or lying position.

Prone Position (Unconscious)

During the pandemic, the increasing numbers of patients requiring prone positioning has led to the establishment of 'proning teams'. Led by an Anaesthetist (or suitably trained advanced airway provider) the team is pre-trained and when necessary are able to turn a patient into the prone or supine position. Local and national guidelines and checklists are being developed to guide and support these teams [4].

Careful monitoring of patients is essential as there is some evidence that the prone position has been associated with a higher incidence of adverse effects than identified in patients

placed in the supine position [37]. Potential complications include airway obstruction from a kinked or displaced ETT. Once in the prone position, frequent oral suctioning and mouth care is required as secretions may reduce the integrity of ETT securing devices. In addition, proning can lead to facial swelling causing retinal nerve compression, ETT ties becoming too tight and pressure ulcers. Electrodes for cardiac monitoring need to be applied posteriorly on the patients back and in the event of cardiac arrest the anterior/posterior placement of defibrillator pads/paddles should be used [38]. Enteral feeding can continue in the prone position; however, the procedure is high risk for vomiting and/or increased gastric residual vomiting [39-41]. Absolute contra-indications for prone position include spinal instability, unstable fractures, burns, open wounds, pregnancy, and recent tracheal surgery and raised intra-cranial pressure. Relative contra-indications include haemodynamic instability (including the use of vasopressors), cardiac pacemakers and abdominal surgery [37].

Neuromuscular blocking agents may be required to maintain gaseous exchange, this reduces extrapulmonary resistance and ventilatory desynchrony which in turn results in improved oxygenation. Paralysis of the diaphragm allows for metabolic rest, reduced oxygen consumption and invasive control of breathing mechanic.²⁶ Neuromuscular blockage may be required in patients with ARDS as this allows for less PEEP to maintain oxygenation, and reduced mortality.

Once turned into the prone position patients may remain in this position for 12 to 16 hours per day [42]. In addition, patients may need to be placed into this position several times. Patients in the prone position should be nursed on a pressure relieving mattress to reduce pressure damage and periodically it will be necessary to change the position of the head and arms every 2-4 hours. To maintain the prone position, ventilated patients must be adequately sedated with the use of neuromuscular blockade [37, 43].

It is important to note that critically ill patients are at high risk of developing malnutrition and sarcopenia [44]. Therefore, is accepted practice that early enteral feeding should be established unless contra-indicated, in addition, the enteral route is preferred over the parental route (e.g. Total Parental Nutrition (TPN) [41, 45]. However, there is an increased risk of gastric aspiration during repositioning, therefore, interruptions in the enteral feeding regimens have to occur when positioning a patient from supine to prone and vice-versa.

Enteral feeding should be resumed once the patient has been re-positioning, NG tube position confirmed and vital signs recorded.

Research into the impact of prone position on tolerance and gastrointestinal complications is still ongoing. De la Fuente et al's [40] small scale study of enteral feeding tolerance in ventilated prone patients, concluded enteral feeding did not increase the risk of gastrointestinal problems. In contrast, Malhotra et al ([37] found patients in the prone position developed a higher incidence of vomiting and/ or increased in gastric residual volumes. As a result of their findings, Malhotra et al [37] recommend patients' heads should be elevated at least 25° while receiving enteral feeding and that prokinetic drugs such as erythromycin may be appropriate.

Conclusion:

Nurses need understanding of the complex pathophysiological processes that arise from Covid-19 infection. However, as this chapter has indicated the pathophysiology associated with the development Covid-19 related ARDS is still being investigate. Nevertheless, the high incidence of ARDS in Covid-19 patients has resulted in recognition that use of the prone position undertaken early for both conscious and unconscious patients can improve oxygenation. It has been identified as an intervention that for some patients may reduce or avoid the need for invasive ventilation. Nevertheless, it has to be accepted that it poses potential risks and complications. Protocols, guidelines and training are essential to minimize the risk of adverse events during or after prone positioning. The Covid-19 pandemic is relatively new therefore, further research is needed before definitive guidelines and recommendations can be made.

Reference List

1. Ghelichkhani P. Esmaeili M. (2020). Prone Position in Management of COVID-19 Patients; a Commentary. Archives of academic emergency medicine. 8. 1. e48
2. Seaton-Mills D. (2000). Prone positioning in ARDS: a nursing perspective. Clinical Intensive Care. 11. 4. 203-208
3. Telias I. Katira BH. Brochard L. (2020). Is the prone position helpful during spontaneous breathing in patients with COVID-19. JAMA. 323. 22. 2265-67
4. Intensive Care Society. (2019). Guidance For: Prone Positioning in Adult Critical Care. www.ics.ac.uk

5. Centre for Disease Control and Prevention. (2020). Interim considerations for infection prevention and control of coronavirus diseases 2019 (COVID-19) in obstetric healthcare settings. <https://www.cdc.gov/coronavirus/2019-ncov/hcp/disposition-hospitalized-patients.html>
6. Yang D. Leibowitz JL. (2015). The structure and functions of coronavirus genomic 3' and 5' ends. *Virus Res.* 206. 120–133. doi: 10.1016/j.virusres.2015.02.025
7. Chad T. Sampson C. (2020). Prone positioning in conscious patients on medical wards: A review of the evidence and its relevance to patients with COVID-19 infection. *Clinical Medicine.* 20. 4. e97–103
8. Marini JJ. Gattinoni L. (2020). Management of COVID-19 respiratory distress. *JAMA.* 323. 22. 2329-2330.
9. Dhont S et al. (2020). The pathophysiology of 'happy' hypoxemia in Covid-19. *Resp Research.* 21. 198.
10. Tobin MJ. Laghi F. Jubran A. (2020). Why COVID-19 Silent Hypoxemia Is Baffling to Physicians. *Am J Respir Crit Care Med.* 202. 3. 356-360. doi: 10.1164/rccm.202006-2157CP. PMID: 32539537; PMCID: PMC7397783.
11. Xie J. Covassin N. Fan Z. Singh P. Gao W. Li G. Kara T. Somers VK et al (2020). Association between hypoxemia and mortality in patients with COVID-19. *Mayo Clin Proc.* 95. 6. 1138-1147
12. Jain A. Doyle DJ. (2020). Stages or phenotypes? A critical look at COVID-19 pathophysiology. *Intensive Care Medicine.*
13. Coperchia F. Chiovato L. Crocea L. Magria F. Rotondi M. (2020). The cytokine storm in COVID-19: An overview of the involvement of the chemokine/chemokine-receptor system. *Cytokine & Growth Factor Reviews.* 53. 25-32
14. Wilcox S (2020). Management of respiratory failure due to covid-19. *BMJ.* 329: m1786.
15. Mohlenkamp. S. Thiele, H. (2020). Ventilation of COVID-19 patients in intensive care units. *Herz.* 45. 4. 329-331. doi:10.1007/s00059-020-04923-1
16. Ashbaugh DG. Bigelow DB. Petty TL. Levine BE. (1967). Acute respiratory distress in adults. *Lancet.* 12. 2. 7511. 319-23
17. Bernard GR. Artigas A. Brigham KL. Carlet J. Falke K. Hudson L. Lamy M. Legall JO. Morris A. Spragg R. (1994). The American-European Consensus Conference on ARDS. Definitions, mechanisms, relevant outcomes and clinical trial co-ordination. *Am J Respir Crit Care Med.* 149. 3. Part 1. 818-24
18. Ferguson ND. Fan E. Camporota L. Antonelli M. Anzueto A. Beale R. Brochard L. Brower R. Esteban A. Gattinoni L. Rhodes A. Slutsky AS. Vincent JL. Rubenfeld GD. Thompson BT. Tanieri VM. (2012). The Berlin Definition of ARDS: an expanded rationale, justification and supplementary material. *Intensive Care Med.* 38. 10. 1573-82
19. Ranieri VM. Rubenfeld G. Thompson BT. (2012). Acute Respiratory Distress Syndrome: The Berlin Definition. *JAMA.* 307. 23. 2526-33
20. Fanelli V. Vlachou A. Ghannadian S. Simonetti U. Slutsky AS. Zhang H. (2013). Acute respiratory distress syndrome: new definition, current and future therapeutic options. *Journal of thoracic disease.* 5. 3. 326–334. <https://doi.org/10.3978/j.issn.2072-1439.2013.04.05>
21. ARDS Definition Task Force. (2012). Acute respiratory distress syndrome: the Berlin definition. *JAMA* 307. 2526–33.
22. Villar J. Perez-Mendez L. Blanco J. Anon JM. Blanch L. Belda J. Santos-Bouza A. Fenandez RL. Kacmarek RM. (2013). Spanish initiative for epidemiology, stratification, and therapies for ARDS (SIESTA) Network. A universal definition of ARDS: the PaO₂/FiO₂ ratio under a standard ventilator setting- a prospective, multicentre validation study. *Intensive Care Medicine.* 39. 583-592

23. Pelosi P. D'Onofrio D. Chiumello D. Paolo S. Chiara G. Capellozzi VL. Barbas CS. Chiaranda M. Gattinoni L. (2003). Pulmonary and extra pulmonary acute respiratory distress syndrome are different. *European Respiratory Journal*. 22. 48s-56s
24. Tremblay L. Valenza F. Ribeiro SP. Slutsky AS. (1997). Injurious ventilator strategies increase cytokines and c-fos mRNA expression in an isolated rat lung model. *J Clin Invest*. 99. 5. 944–952
25. European Society of Intensive Care Medicine. (2017). *Acute Respiratory Distress Syndrome*. 2nd Edition. www.esicm.org
26. Baid H. (2016). Patient safety: Identifying and managing complications of mechanical ventilation. *Crit Care Nurs Clin N Am*. 28. 451-462
27. Pooni RS. (2020). Research in brief: Prone positioning in COVID-19: What's the evidence? Royal College of Physicians London.
28. Bamford P. Bentley A. Dean J. Whitmore D. Wilson-Baig N. (2020). ICS Guidance for Prone Positioning of the Conscious COVID Patient 2020. www.ics.ac.org
29. Rimmer A. Wilkinson E. (2020). What's happening in covid-19 ICUs? An intensive care doctor answers some common questions. *BMJ*. 369: m1552.
30. Riviello ED. Kiviri W. Twagirumugabe T. Mueller A. Banner-Goodspeed VM. Officer L, Novack V. Mutumwinka M. Talmor DS. Fowler RA. (2020). Hospital Incidence and Outcomes of the Acute Respiratory Distress Syndrome Using the Kigali Modification of the Berlin Definition. *Am J Respir Crit Care Med*. 193. 1. 52-9. doi: 10.1164/rccm.201503-0584OC.
31. Bryan AC. (1974). Comments of a devil's advocate. *Am Rev Respir Dis* 110 [Suppl]:143–144
32. Geurin C. Gaillard S. Lemasson S. (2004). Effects of Systematic Prone Positioning in Hypoxaemic Acute Respiratory Failure. *JAMA*. 292. 2379-2387.
33. Mancebo J. Fernández R. Blanch L. et al. (2006). A Multicenter Trial of Prolonged Prone Ventilation in Severe Acute Respiratory Distress Syndrome. *Am J Respir Crit Care Med*. 173. 1233–1239.
34. Taccone P. Pesenti A. Latini R. et al. (2009). Prone–Supine II Study Group. Prone Positioning in patients with moderate and severe Acute Respiratory Distress Syndrome: A Randomized Controlled Trial. *JAMA*. 302. 1977–1984.
35. Gattinoni L. Tognoni G., Pesenti A. et al. (2001). Effect of Prone Positioning on the Survival of Patients with Acute Respiratory Failure. *N Engl J Med*. 345. 8. 568-573.
36. Sud S. Friedrich J. Adhikari N. et al. (2014). Effect of prone positioning during mechanical ventilation on mortality among patients with acute respiratory distress syndrome: a systematic review and meta-analysis. *CMAJ*. 186. 10. 381-390.
37. Malhotra A. Kacmarek RM. Parsons PE. Finlay G. (2020). Prone ventilation for adults patients with acute respiratory distress syndrome. <https://www.uptodate.com/contents/prone-ventilation-for-adult-patients-with-acute-respiratory-distress-syndrome#H2655777180>
38. Poston JT. Patel BK. Davis AM. (2020). Management of Critically Ill Adults with COVID-19. *JAMA*. 323. 18. 1839–1841. doi:10.1001/jama.2020.4914