

REVIEW

Mass spectrometry analysis of tau and amyloid-beta in iPSC-derived models of Alzheimer's disease and dementia

Charles Arber¹  | Argyro Alatzas¹  | Claire A. Leckey^{1,2,3}  | Ross W. Paterson¹  |
Henrik Zetterberg^{1,3,4,5}  | Selina Wray¹ 

¹Department of Neurodegenerative Disease, UCL Queen Square Institute of Neurology, University College London, London, UK

²Translational Mass Spectrometry Research Group, UCL Great Ormond Street Institute of Child Health, University College London, London, UK

³UK Dementia Research Institute at UCL, London, UK

⁴Department of Psychiatry and Neurochemistry, Institute of Neuroscience and Physiology, Sahlgrenska Academy at the University of Gothenburg, Mölndal, Sweden

⁵Clinical Neurochemistry Laboratory, Sahlgrenska University Hospital, Mölndal, Sweden

Correspondence

Selina Wray, Department of Neurodegenerative Disease, UCL Queen Square Institute of Neurology, University College London, London, WC1N 1PJ., UK.
Email: selina.wray@ucl.ac.uk

Funding information

This work is supported by the NIHR UCL Hospitals Biomedical Research Centre. SW is funded by an Alzheimer's Research UK Senior Research Fellowship (ARUKSRF-2016B). CA is funded by an Alzheimer's Society junior fellowship (AS-JF-18-008). RWP is supported by an Alzheimer's Association Clinician Scientist fellowship and the UK Dementia Research Institute. HZ is a Wallenberg Scholar supported by grants from the Swedish Research Council (#2018-02532), the European Research Council (#681712), Swedish State Support for Clinical Research (#ALFGBG-720931), the Alzheimer Drug Discovery Foundation (ADDF), USA (#201809-2016862), the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 860197 (MIRIAD) and the UK Dementia Research Institute at UCL.

This is a Review for the special issue "Mass Spectrometry in Alzheimer Disease"

Abstract

Induced pluripotent stem cell (iPSC) technology enables the generation of human neurons in vitro, which contain the precise genome of the cell donor, therefore permitting the generation of disease models from individuals with a disease-associated genotype of interest. This approach has been extensively used to model inherited forms of Alzheimer's disease and frontotemporal dementia. The combination of iPSC-derived neuronal models with targeted mass spectrometry analysis has provided unprecedented insights into the regulation of specific proteins in human neuronal physiology and pathology. For example enabling investigations into tau and APP/A β , specifically: protein isoform expression, relative levels of cleavage fragments, aggregated species and functionally critical post-translational modifications. The use of mass spectrometry has enabled a determination of how closely iPSC-derived models recapitulate disease profiles observed in the human brain. This review will highlight the progress to date in studies using iPSCs and mass spectrometry to model Alzheimer's disease and dementia. We go on to convey our optimism, as studies in the near future will make use of this precedent, together with novel techniques such as genome editing and stable isotope labelling, to provide real progress towards an in depth understanding of early neurodegenerative processes and development of novel therapeutic agents.

KEYWORDS

Alzheimer's disease, amyloid-beta, induced pluripotent stem cells, mass spectrometry, tau

Abbreviations: AD, Alzheimer's disease; AGD, argyrophilic grain disease; APP, amyloid precursor protein; A β , amyloid β ; CBD, corticobasal degeneration; CSF, cerebrospinal fluid; DS, down's syndrome; FTD, frontotemporal dementia; GGT, globular glial tauopathy; hESC, human embryonic stem cell; IP, immunoprecipitation; iPSC, induced pluripotent stem cell; LC-MS/MS, liquid chromatography – tandem mass spectrometry; MALDI, TOF, matrix-assisted laser desorption/ionization with time-of-flight; MAPT, microtubule-associated protein tau gene; MS, mass spectrometry; PART, primary age-related tauopathy; PiD, Pick's disease; PRM, parallel reaction monitoring; PSEN1, presenilin-1; PSEN2, presenilin-2; PSP, progressive supranuclear palsy; PTM, post-translational modification; qPCR, quantitative polymerase chain reaction; RT-PCR, reverse transcription polymerase chain reaction; SELDI-TOF, surface-enhanced laser desorption/ionization–time-of-flight; SILK, stable isotope labelling kinetics.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2021 The Authors. *Journal of Neurochemistry* published by John Wiley & Sons Ltd on behalf of International Society for Neurochemistry.

1 | ALZHEIMER'S DISEASE

Alzheimer's disease (AD) is characterized by two main pathologies: the extracellular deposition of A β peptides in amyloid plaques and the intracellular deposition of the tau protein in neurofibrillary tangles. Studies of autosomal-dominant AD have placed amyloid upstream of tau, with causative mutations in the genes encoding the amyloid precursor protein (APP), presenilin-1 (PSEN1) and presenilin-2 (PSEN2) (Hardy, 2017). However, there is substantial evidence that tau plays a central role in AD pathogenesis. The extent of tau pathology correlates with symptom severity and the extent of neurodegeneration (Braak et al. 2003); tau knockout or reduction is protective against amyloid toxicity in pre-clinical models (Gheyara et al. 2014; Ittner et al. 2010; Roberson et al. 2007; Vossel et al. 2010); and mutations in the tau gene, (MAPT) cause frontotemporal dementia (FTD) (Hutton et al. 1998; Poorkaj et al. 1998), confirming a causative link between tau dysfunction and neurodegeneration. Thus, understanding the two pathologies of AD and the molecular mechanisms linking them to neurodegeneration is an area of intense research. Pre-clinical models recapitulating the main pathologies are key to this endeavour.

2 | INDUCED PLURIPOTENT STEM CELL MODELS OF ALZHEIMER'S DISEASE

Although multiple in vitro and in vivo models of AD and FTD exist, it is only recently that induced pluripotent stem cell (iPSC) technology has enabled the generation of unlimited numbers of human neurons in the lab. Pluripotency can be induced in terminally differentiated cells such as fibroblasts by transduction with the four reprogramming factors cMYC, SOX2, OCT4 and KLF4. The resulting iPSC are indistinguishable from human embryonic stem cells (hESC) and can be differentiated into all three germ lineages (Takahashi et al. 2007).

Thus, by generating iPSC from a patient with a genotype and/or phenotype of interest and differentiating those cells into a disease-relevant cell type, patient-specific in vitro models containing the participant's precise genome can be used for pre-clinical studies of disease mechanisms. This approach has revolutionized the in vitro modelling of neurodegenerative disease, by permitting a limitless in vitro supply of human neurons and glia with disease-causing genes expressed at endogenous levels. The ability to generate distinct neuronal subtypes together with astrocytes and microglia from iPSC is particularly advantageous for diseases such as the tauopathies, where the pathophysiological characteristics are not confined to one specific cell type. In the tauopathies, both neurons and/or glial cells can display abnormal tau deposition, aggregates and /or hyperphosphorylated tau. For example primary age-related tauopathy (PART) and ageing-related tau astrogliaopathy (ARTAG) are pure neuronal and astroglial tauopathies, respectively, whereas globular glial tauopathy (GGT), Pick's disease (PiD), progressive supranuclear palsy (PSP), corticobasal degeneration (CBD), argyrophilic grain disease (AGD) and Frontotemporal dementia with parkinsonism-17 (FTDP-17) present

both neuronal and glial cell involvement (Crary et al. 2014; Kovacs et al. 2016; Ahmed et al. 2013; Buée & Delacourte, 1999; Sergeant et al. 2008; Sergeant et al. 1999; Yoshida, 2014; Togo et al. 2002; BRAAK & BRAAK, 1989; Yamada et al. 1992; Ferrer et al. 2019; Mackenzie & Neumann, 2016). Furthermore, the neuronal tau aggregates can be found in different brain regions depending on the disease: PSP and CBD display prominent tau pathology in the brain stem, for example while predominantly cortical tau pathology is present in AD. iPSC technology, therefore offers the opportunity for the specific cell type affected in disease to be generated in vitro in order to investigate the selective vulnerability of specific cell populations to pathology.

Although animal models have been widely used to gain mechanistic insights into AD, it has been particularly challenging to generate models that develop plaques, tangles and neurodegeneration within a single system, often requiring exogenous overexpression of multiple mutant genes (Oddo et al. 2003). This raises the possibility that human neurons are selectively vulnerable to AD-associated pathologies. The use of rodent models has also been hampered by species-specific features of the protein of interest, for example alternative splicing of the tau protein, which is complex, developmentally regulated and differs between rodents and humans (Goedert et al. 1989; Kosik et al. 1989; Takuma et al. 2003). Importantly, the lack of translation of A β and tau-targeted therapeutics from pre-clinical studies to clinical success supports the need for refined models. iPSC therefore provide an attractive approach to address this gap in our toolkit, acting as a human, physiologically relevant model to study the mechanism of a disease in the specific cell type which is selectively vulnerable in disease, and in the absence of exogenous gene expression.

Despite the advantages offered by iPSC, there are still some limitations associated with this model, particularly concerns around the maturity of differentiated cell types. iPSC are 'rejuvenated', that is the reprogramming process erases the cellular epigenetic signatures associated with donor age (Lee et al. 2020). Furthermore, multiple studies have shown that iPSC-neurons closely resemble neurons, for example using transcriptomic comparisons (Patani et al. 2012). This must be taken into consideration when studying diseases such as AD where age is a major risk factor. This can now be overcome in part by using transdifferentiation to directly convert fibroblasts into neurons, bypassing an iPSC intermediate, which allows cells to retain epigenetic signatures of ageing (Mertens et al. 2015). Nonetheless, iPSC are the only means to obtain an unlimited supply of patient-specific neurons and glia for the study of disease mechanisms, and permit the direct cellular consequences of disease-associated mutations to be studied.

Progress made using iPSC technology to model AD and FTD linked to MAPT mutations is reviewed here (Arber et al. 2017; Lines et al. 2020). In this review, we focus specifically on insights gained from the use of mass spectrometry in conjunction with these models, and how this has enabled the precise measurement of peptide isoforms, cleavage fragments, multimeric species and post-translational modifications.

3 | MASS SPECTROMETRY IN ALZHEIMER'S DISEASE RESEARCH

Over the last few decades, mass spectrometry-based analysis of A β has focused primarily on identification and quantification in CSF, plasma and brain tissue (Portelius, Bogdanovic, *et al.* 2010; Portelius *et al.* 2012; Wildburger *et al.* 2017; Nakamura *et al.* 2018). Mass spectrometry (MS) analytical strategies can vary considerably across studies, however, the most common approaches used for the analysis of A β so far have been matrix-assisted laser desorption/ionization (MALDI) or surface enhancement laser desorption/ionization (SELDI) coupled to a time-of-flight (TOF) mass spectrometer as well as liquid chromatography-tandem mass spectrometry (LC-MS/MS). A more

detailed overview and technical comparison of these techniques for A β detection and quantification can be found here (Bros *et al.* 2015). MALDI-TOF MS provides a sensitive, accurate and rapid method for the relative quantification of A β species. Targeted LC-MS/MS approaches have commonly used electrospray ionization (ESI) in setups using high-performance liquid chromatography (HPLC) or ultra performance liquid chromatography (UPLC) coupled to a triple quadrupole mass spectrometer, which offer high specificity, absolute quantification and multiplexing capabilities. A common feature between these approaches has been the need for enrichment of A β peptides during sample preparation, likely because of the low abundance of A β peptides in complex sample matrices. This has most commonly been achieved by immunoprecipitation (IP) using antibodies targeting the mid domain (4G8) and N-terminus (6E10) of the A β peptide before MS analysis. Although IP-MALDI-TOF and SELDI-TOF MS approaches offer relative quantification, they have played a pivotal role in identifying novel and truncated A β species, and with less system complexity and at lower costs than LC-MS/MS (Lewczuk *et al.* 2004; Maddalena *et al.* 2004; Portelius *et al.* 2007).

Similar to A β , MS-based approaches have played a critical role in the characterization of tau in AD research. In the human central nervous system there are six different tau isoforms that are produced and these are subject to extensive post-translational modifications in both normal and disease conditions (Kametani *et al.* 2020; Morris *et al.* 2015). It is this vast heterogeneity of tau species that has posed a particular analytical challenge to mass spectrometrists. Furthermore, thorough characterization and quantification of tau proteoforms by MS has been further complicated by their low concentration in CSF. Therefore, profiling of tau species in CSF has previously required the use of well-established tau antibodies to enrich for specific isoforms/phosphorylation sites by IP before subsequent MS-based analysis (Portelius *et al.* 2008). In the last decade the introduction of quadrupole-Orbitrap hybrid mass spectrometers has allowed for parallel reaction monitoring (PRM) strategies to be developed; utilizing the high resolution and high mass accuracy capabilities of these instruments these methods offer a targeted approach that has high selectivity, high sensitivity and multiplexing capabilities (Gallien *et al.* 2012; Peterson *et al.* 2012). It is these PRM-based strategies in particular that have most recently advanced our understanding and interpretation of tau truncation and metabolism in AD both in vivo and in vitro (Barthélemy *et al.* 2016, 2019, 2020; Sato *et al.* 2018).

4 | A β GENERATION FROM APP

Canonical A β is produced when APP is sequentially cleaved by β -secretase and γ -secretase. However, there exists a spectrum of A β species that are produced when APP is cleaved by alternate enzymes, leading to truncations at both N and C termini (Figure 1). As such, Takami *et al.* used a targeted LC-MS/MS approach to show that γ -secretase has C-terminal truncating activity as the longest forms




FIGURE 1 Amyloid-beta sequence, cleavage sites and insights from mass spectrometry. (a) The amino acid sequence of A β , with residues labile to post-translational modifications highlighted: Green—phosphorylation and dityrosine/nitrotyrosine/nitration of serine and tyrosine residues. Gold—pyroglutamate modification of glutamic acid residues. Red—racemization/isomerization of aspartic acid residues. Arrows show the sites of secretase cleavage of A β . (b) Information learned from iPSC models subject to mass spectrometry analysis. Predominant forms of A β in different stages of neuronal differentiation (Bergström *et al.* 2016). The finding that N terminal cleavage can direct C terminal endoproteolysis (Siegel *et al.* 2017). The anti-amyloidogenic role of BACE2 in reducing relative amounts of amyloidogenic species in Down's syndrome neurons (Alić *et al.* 2020). The specificity of avermectins, increasing the processivity of γ -secretase while leaving short species of A β unchanged (Brownjohn *et al.* 2017)

of A β (A β 1-49 and A β 1-48) are successively processed through parallel tripeptide cleavage pathways (A β 1-49 > 46>43 > 40 versus A β 1-48 > 45>42) to produce the most abundant fragments of A β (Takami et al. 2009).

In CSF, A β 1-40 is the most abundant species (Portelius et al. 2010). The relative levels of other C-terminally truncated A β species such as A β 1-42 and A β 1-43 are prognostic for familial AD and diagnostic for amyloid plaque pathology in both familial and sporadic forms of the disease. These longer forms are a major constituent of pathological amyloid plaques, presumably because of increased hydrophobicity and enhanced aggregation. As a result the A β 1-42:A β 1-40 ratio in the CSF is used as a diagnostic test of AD (Janelidze et al. 2016).

The second most abundant form of A β is A β 1-17, which is also potentially β - and γ -secretase-dependent (Pérez-Grijalba et al. 2014; Portelius et al. 2011). Shorter, non-amyloidogenic isoforms ranging from A β 1-13 to A β 1-16 have been proposed to be α - and β -secretase dependent (Portelius et al. 2011). Additionally, several A β species are produced by β -secretase cleavage alone (BACE1 and BACE2 cleavage), producing species such as A β 1-19, A β 1-20 and A β 1-34 (Shi et al. 2003; Yan et al. 2001). Other enzymes (e.g. MMP2, MMP9 and Caspases) have been shown to be implicated in the generation of further A β species (e.g. A β 1-30) but are less well studied (Baranello et al. 2015).

N-terminal truncation of A β includes cleavage at residues 2, 4, 5 and 11, each thought to be β -secretase dependent (Vassar et al. 1999). Importantly, these N-terminally truncated A β forms, such as A β 3-40/42 and A β 11-40/42, have been shown to be targets of post-translational modifications such as pyroglutamate modification at glutamic acid 3 or 11; generating key amyloidogenic species (Mori et al. 1992; Saido et al. 1996). These modified A β forms are not secreted by neurons but appear to form in the tissue and may well be A β plaque-specific.

Therefore, there is a huge diversity of A β species outside the well-known A β 1-40 and A β 1-42. Mass spectrometry has been critical in detecting and distinguishing between various A β species in human brain tissue, CSF and plasma, and these approaches have recently been applied to iPSC models of AD.

5 | A β DETECTION IN IPSC-DERIVED NEURONS

To date, only a handful of studies, all using IP-MALDI-TOF MS, have investigated the entire spectrum of A β species produced in iPSC-derived neurons (Bergström et al. 2016; Arber et al. 2020). Bergström and colleagues analysed the A β spectrum produced throughout iPSC differentiation; from the stem cell stage, through neural specification, to mature neuronal cultures (Bergström et al. 2016). APP expression and protein levels were found to be relatively consistent throughout differentiation, however, the processing of APP and the resultant A β species was tightly controlled in a cell stage-specific manner. Short A β peptides, A β 1-15, A β 1-16

and A β 1-17, were primarily produced from days 10 to 25, at time points before the specification of neurons. This is consistent with predominant α -secretase cleavage. From day 60 of differentiation onwards, after the generation of functional glutamatergic neurons (Kirwan et al. 2015; Shi et al. 2012), the majority of A β species produced were β - and γ -secretase-dependent, represented by A β 1-40, A β 1-42, A β 1-38 and A β 1-34. This study elegantly shows that A β is produced in a cell type and cell stage-dependent manner, giving an insight into human developmental regulation of A β processing. β -secretase expression (BACE1 and BACE2) and activity appear to be neuronally enriched, supporting the view that amyloidogenic A β is primarily generated within the neuronal population.

It has been shown that the A β spectrum produced from iPSC-derived neurons is a representative model of the in vivo setting. Arber and colleagues showed that the most abundant species are A β 1-40, A β 1-17, A β 1-34, A β 1-38 and A β 1-19 (Arber et al. 2020), which is similar to the profiles in CSF. In addition to the expected C-terminally processed A β species, representative N terminally truncated species were also detectable, including A β 2-x, A β 4-x, A β 5-x and A β 11-x (Arber et al. 2020).

iPSC-derived neuronal models contributed to a detailed analysis of the N-terminal truncation of A β and the downstream consequence on C-terminal processing. Siegel and colleagues show that when A β is cleaved by α -secretase at residue 17, γ -secretase effectively processes the C-terminus to A β 40 (A β 17-40). However, when the substrate for γ -secretase is β -secretase dependent (A β 1-x or A β 11-x), proportionally longer A β is produced (i.e. A β 1-42 or A β 11-42) (Siegel et al. 2017). Enrichment for the APP intracellular domain (C terminal to A β) by IP followed by MALDI-TOF MS analysis showed that this effect was in part driven by different endoproteolytic cleavage sites of γ -secretase on APP. α -secretase cleaved substrates predisposed A β 17-49 (and the A β 49 > 46>43 > 40 processing pathway) whereas β -secretase cleaved substrates predisposed A β 1-48 or A β 11-48 (and the A β 48 > 45>42 processing pathway), thereby helping to explain the differences in relative A β 1-42 production.

In sum, these studies help to show the value of iPSC models in recapitulating the in vivo spectrum of A β generation (Figure 1). Additionally, mechanistic insights can be gained, such as the work by Siegel et al, explaining how the balance of A β 1-40 and A β 1-42 is achieved.

6 | MODELLING FAMILIAL ALZHEIMER'S DISEASE

Stem cell models of A β generation have primarily focused on the ratio of A β 1-42:A β 1-40 (for review see Arber et al., 2017). It is widely accepted that mutations in *PSEN1* reduce the processivity of γ -secretase (Chávez-Gutiérrez et al. 2012; Szaruga et al. 2017), increasing the relative production of A β 1-42 compared to A β 1-40. Indeed, this has been corroborated using mutant *PSEN1* iPSC models (Yagi et al. 2011; Woodruff et al. 2013; Duan et al. 2014; Mahairaki et al. 2014; Sproul

et al. 2014; Koch et al. 2012; Moore et al. 2015; Arber et al. 2020; Kwart et al. 2019; Raja et al. 2016; Liu et al. 2014; Arber, Villegas-Llerena, et al., 2019). Additionally, mutations around the γ -secretase cleavage site in APP have been shown to have a similar effect in the A β 1-42:A β 1-40 ratio, although this is achieved via favouring of the A β 48 > 45>42 tripeptide cleavage pathway (Arber et al. 2020; Kondo et al. 2013; Kwart et al. 2019; Moore et al. 2015; Muratore et al. 2014). Mutations around the β -secretase cleavage site or additional copies of APP because of local genomic duplications or Down's syndrome increase the total production of A β (Alić et al. 2020; Chang et al. 2015; Israel et al. 2012; Kondo et al. 2013; Kwart et al. 2019; Moore et al. 2015; Raja et al. 2016; Shi, Kirwan, Smith, MacLean, et al., 2012).

These studies help to reinforce mechanistic studies of γ -secretase in a human model of familial AD expressing APP and secretase enzymes at physiological levels. However, studies using MALDI-TOF MS have been able to advance these mechanistic insights. Mutations affecting the extracellular domain of PSEN1 (Y115H, Y115C and the splicing mutation int4del) display a relative increase in production of short species A β 1-14, A β 1-15 and A β 1-16 (Moore et al. 2015; Arber et al. 2020). This effect is not shared by PSEN1 mutations in other domains of the protein and so may be a direct consequence of mutations to this substrate-binding domain (Takagi-Niidome et al. 2015). The relative increase in short A β species suggests an overall reduction in γ -secretase activity and a concomitant increase in β - and/or α -secretase activity.

In the last decade, iPSC technology has enabled the generation of organoids (organ-like three-dimensional tissues). Brain organoids provide an in vitro model that recapitulates the development of the human brain in which multilayers of several cell lineages self-assemble and create an architecture of embryonic human brain-like compartmentalization (e.g. forebrain, midbrain and hindbrain) with complex neural networks (Lancaster et al. 2013; Lancaster and Knoblich 2014; Lancaster et al. 2017; Qian et al. 2016, 2018; Jo et al. 2016). Therefore, the investigation of tissue development and disease state could be studied in a more 'physiological' multicellular model rather than into a single-dimensional cellular level. Gonzalez and colleagues successfully generated cerebral organoids from people affected with fAD (PSEN1 mutation), Down syndrome and Creutzfeldt-Jakob disease. Over time, pathological features of AD were observed in the cerebral organoids, including A β and tau aggregates (Gonzalez et al. 2018). Furthermore, using a different organoid generation protocol, Raja and colleagues were able to reduce amyloid and tau pathology using β - and γ -secretase inhibitors treatment (Raja et al. 2016). Recently, the validity of cerebral organoids as a model for investigating AD disease was further confirmed when AD patient-derived organoids with either APOE ϵ 3/ ϵ 3 or APOE ϵ 4/ ϵ 4 mutations recapitulated increased levels of A β and phosphorylated tau (Zhao et al. 2020).

Relative A β generation was further investigated in iPSC-derived cerebral organoid models of Down's syndrome (DS) (Alić et al. 2020). The presence of three copies of APP because of duplication mutations predispose familial Alzheimer's disease with complete penetrance. APP is located on chromosome 21, and so individuals with DS also carry three copies of the APP gene. However, in contrast

to APP duplication carriers, people with DS have only a 70% likelihood of developing Alzheimer's disease, suggesting the presence of protective genes on chromosome 21. Alić and colleagues presented data to support an anti-amyloidogenic role for the β -secretase gene BACE2, a gene also present in three copies in DS. When analysing A β isoforms generated by cerebral organoids, there was an increase in A β 1-19 and A β 1-34 relative to amyloidogenic species (A β 1-40 and A β 1-42) in Down's syndrome organoids when compared to isogenic diploid cells and APP duplication organoids. This increased BACE2-associated β -secretase activity, therefore, reduces the relative amount of amyloidogenic A β species. The authors witnessed a concomitant reduction in disease-like signatures in DS organoids compared to APP duplication lines, such as amyloid-like immunocytochemical staining. Genetic reduction of BACE2 to two copies in trisomic cells exacerbated these disease-like phenotypes, supporting a protective role for BACE2. Together these data demonstrate the delicate balance of cleavage activity and the importance of investigating the entire A β spectrum in a physiological model.

Finally, mass spectrometry analyses of iPSC-derived A β species can have a central role in drug discovery. Brownjohn and colleagues screened compounds capable of increasing the processivity of γ -secretase in Down's syndrome and familial AD iPSCs, i.e. molecules capable of increasing the A β 1-38:A β 1-42 ratio (Brownjohn et al. 2017). Importantly, mass spectrometry was used to support the finding that the lead compounds, avermectins, increased the generation of A β 1-37 and A β 1-38 at the expense of A β 1-40 and A β 1-42 without affecting smaller, γ -secretase-independent A β species. This selectivity of lead compounds is highly desirable in putative therapeutic agents.

Together, these studies demonstrate the importance of a representative appreciation for the entire A β spectrum (Figure 1). Mass spectrometry analysis, together with iPSC technology, highlight the interdependence and the delicate balance of different enzyme cleavages of A β . An understanding of the A β spectrum and this enzymatic balance is crucial when analysing the effect of therapeutic drug candidates or inherited Alzheimer's disease-linked mutations.

7 | FUTURE DIRECTIONS

Future work to analyse A β species without immunoprecipitation will be informative, as new truncated species, unknown post-translational modifications and multimeric/aggregated species may be defined in this way. The use of immunoprecipitation can lead to bias in the species analysed, for example certain post-translational modifications can alter the binding of antibodies used in immunoprecipitation. Of particular interest, examples of modified A β include pyroglutamate-modified species at glutamic acid residues 3 and 11 (Perez-Garmendia & Gevorkian, 2013), phosphorylation at serine 8 and serine 26 (Kumar et al. 2011, 2013), nitration at tyrosine 10 (Kummer et al. 2011), racemization/isomerization at aspartic acids 1, 7, 23 and 26 (Warmack et al. 2019) and dityrosine/nitrotyrosine modifications also at tyrosine 10 (Al-Hilaly et al. 2014) (Figure 1). To date, little information

exists about the post-translational modification of A β species derived from iPSC-neurons and these investigations will be highly informative. For example pyroglutamate modifications might be co-culture or cell state dependent, as transglutaminase activity may be glial cell derived and dependent on inflammatory conditions (Wilhelmus et al. 2016). Furthermore, A β modifications may emerge in a temporal manner that could be linked to disease severity, as recently shown for tau (Dujardin et al. 2020; Wesseling et al. 2020).

8 | TAU

Hyperphosphorylated aggregates of the microtubule-associated protein tau are a pathological hallmark of a diverse group of neurodegenerative diseases collectively termed as tauopathies. Alzheimer's disease (AD) is the most common tauopathy, although it is in fact a secondary tauopathy; tau is downstream of altered A β production as evidence by the genetics of fAD as described above. The most common primary tauopathies (where tau is the defining pathological feature) include progressive supranuclear palsy (PSP), corticobasal degeneration (CBD) and frontotemporal dementia linked to mutations in the *MAPT* gene that encodes tau (Guo et al. 2017). It was the discovery of *MAPT* mutations that confirmed a causative link between tau dysfunction and neurodegenerative disease, and highlighted the importance of understanding the mechanisms linking tau to neuronal demise (Hutton et al. 1998; Poorkaj et al. 1998).

Alternative splicing of *MAPT* results in six protein isoforms of tau in the human central nervous system, differing by the presence of 0, 1 or 2 repeats at the N-terminus of the protein (0N, 1N or 2N), and 3 or 4 microtubule-binding repeats at the C-terminus (3R or 4R). The expression of precise ratios of tau isoforms is regulated developmentally and dysregulated in disease, which will be discussed further below. Tau is further complicated by its extensive post-translational modification. The most heavily studied of these is phosphorylation (Hanger et al. 2009). The phosphorylation of tau plays an important role in regulating its function, with phosphorylation at specific residues favouring the detachment of tau from the microtubules. In disease, tau is hyperphosphorylated, and the aberrant phosphorylation of tau may drive its aggregation.

There are 84 serine, threonine and tyrosine residues in tau, and mass spectrometry has enabled the direct identification of at least 51 specific phosphorylation sites in both in vitro and in vivo pre-clinical models, as well as directly in control and disease post-mortem brain tissue (Hanger et al. 2007; Wray et al. 2008). Additionally, in vitro studies have enabled candidate kinases for many phosphorylation sites to be identified, including GSK3 β , CDK5 and CK1 (Hanger et al. 2009). A comprehensive list of tau phosphorylation sites identified in different diseases together with the kinases able to mediate phosphorylation at specific sites can be found here: https://docs.google.com/spreadsheets/d/1hGYs1ZcupmTnbB7n6qs1r_WVtXHt1O7NBLyKBN7EOUQ/edit#gid=0

Beyond phosphorylation, tau is also the target of numerous other post-translational modifications (PTMs) including ubiquitination,

acetylation, nitration, glycosylation and cleavage (Morris et al. 2015). A thorough catalogue of 170 distinct tau PTMs has recently been described in control and disease tissue (Kametani et al. 2020), and tau PTM state can correlate with clinical severity and heterogeneity (Barthélemy et al. 2020; Dujardin et al. 2020; Wesseling et al. 2020). Thus, there is a diverse range of tau species existing within a neuron at any one time, and it is important to understand the range of tau species, their disease specificity and the temporal manner in which they appear in disease.

There are a limited number of studies that have used mass spectrometry to analyse tau in iPSC-neurons, these are discussed below.

9 | MASS SPECTROMETRY ANALYSIS OF TAU IN IPSC

Tau splicing is developmentally regulated, with only the smallest tau isoform (0N3R) expressed at foetal stages (Goedert et al. 1989; Hefti et al. 2018). Additionally, splicing is dysregulated in disease, particularly in a subgroup of tauopathies where excess 4R is observed, including PSP and CBD. Intronic and splice-site mutations in and around exon 10 of *MAPT* cause an increase in 4R tau and are causative of FTD, confirming a causal link between disrupted tau splicing and neurodegeneration (Hutton et al. 1998). Tau splicing is also species-specific, and rodent models do not recapitulate the human pattern of tau isoforms (Yu et al. 2009). There has therefore been a great deal of interest in studying tau splicing in human neurons generated from iPSC. A number of studies have now shown that 0N3R is the predominant isoform expressed by iPSC neurons. Variable levels of 4R tau isoforms appear at varying culture times, and dependent on differentiation protocol, as detected by techniques including RT-PCR, qPCR and western blot (Beevers et al. 2017; Iovino et al. 2010, 2015; Sposito et al. 2015). Mass spectrometry provides the opportunity for the unambiguous detection of tau isoforms by confirming the presence of isoform-specific peptides at exon-exon junctions.

Disruption of the nuclear membrane and nucleocytoplasmic transport was observed in iPSC-neurons with the *MAPT* 10 + 16 and P301L mutations (Paonessa et al. 2019). As P301L resides within exon 10, which is only expressed in 4R isoforms, there is a requirement for 4R expression in order to have mutant protein present in the model. This was confirmed by MALDI-TOF/TOF mass spectrometry, which identified 4R-specific peptides from both the wild-type (HVPGGGSVQIVYKPVDSLK) and mutant (HVLGGGSVQIVYKPVDSLK) alleles (Paonessa et al. 2019). This confirms the presence of 4R tau, but does not provide quantitative information on the stoichiometry of the different tau isoforms. Sato et al used quantitative proteomics to assess the levels of 4R tau in iPSC-neurons and post-mortem brain tissue via the detection of two 4R-specific peptides, LDLSNVQSK (amino acids 282-290) and HVPGGSVQIVYK (amino acids 299-311) (Sato et al. 2018). As expected, the 4R tau signal in the brain was around 50% of the total tau signal, corresponding to equimolar amounts of 3R:4R. In contrast, very low levels of 4R peptide could be detected in iPSC-neurons,

suggesting that at 6 weeks of culture, levels of total tau are around 100-fold less than in the brain. A similar approach was unable to identify peptides corresponding to 1N or 2N tau isoforms in iPSC-neurons at 5 weeks of differentiation. Low levels of the 4R-specific peptide were detected in comparison to tau peptides ubiquitously present in all isoforms, further confirming the predominance of ON3R in iPSC-neurons (Silva et al. 2016).

Further studies of the detailed quantification of altered splicing in neurons with FTD-linked splicing mutations will be of great interest, to help decipher developmental regulation and disease-associated dysfunction in naïve human neurons.

10 | TAU POST-TRANSLATIONAL MODIFICATION IN IPSC-NEURONS

Tau phosphorylation is also developmentally regulated, with high phosphorylation in early development thought to be related to the requirement for dynamic remodelling of the microtubule network (Brion et al. 1993). Hyperphosphorylation of tau is observed across the tauopathies, and accurate determination of the sites of phosphorylation and the relative stoichiometry in control versus disease is important in inferring the presence of “pathological” tau in pre-clinical models. The majority of studies using iPSCs to model tauopathy have examined phosphorylation at specific sites using phospho-specific antibodies, although a few studies have directly identified sites of phosphorylation using mass spectrometry.

Although the direct identification of phosphorylation sites on tau by mass spectrometry is informative, it is important to note that many of the same phosphorylation sites occur in development, control adult brain tissue and disease states (Kametani et al. 2020). Multiple quantitative approaches to allow the stoichiometry of phosphorylation at particular sites have been developed and will enable future work to identify disease-associated changes. FLEXI-tau is a mass spectrometry-based assay that quantifies the ratio of modified/unmodified peptides in a culture system, with a decrease in the levels of unmodified peptides indicative of PTMs within the peptide (Mair et al. 2016). Caterina-Silva *et al.* used the FLEXI-tau assay to investigate post-translational-modifications of tau in iPSC-neurons with the A152T variant compared with controls after 5 weeks in culture (Mair et al. 2016; Silva et al. 2016). There is an inherent assumption in many studies that equal expression from both the mutant and the unaffected allele is occurring within cells. A152T neurons showed increased levels of total tau overall, as well as increased levels of mutant tau, with A152T tau accounting for ~56.6% of total tau. Modification at several known phosphorylation epitopes was increased in A152T neurons (S202/205, T231/S235 and S396/S404), suggesting an early increase in phosphorylation at disease-associated epitopes in tau mutation neurons. The A152T variant creates a potential novel phosphorylation site in tau through the addition of an extra threonine residue, however, no modifications were observed on the peptide covering this region suggesting it is unlikely to be phosphorylated in iPSC-neurons (Silva et al. 2016). Using an immunoprecipitation-mass

spectrometry (IP-MS) approach, an independent study identified multiple phosphorylated tau peptides in control neurons, including pT212, pS214 and pT217, pS262 and pS356 (Sato et al. 2018).

11 | TAU TURNOVER IN IPSC-NEURONS

Stable isotope labelling kinetics (SILK) allows for the measurement of protein production and turnover rates using labelling with heavy essential amino acids (typically $^{13}\text{C}_6$ -leucine), followed by mass spectrometry to distinguish labelled from unlabelled peptides. Heavy leucine will be incorporated into newly synthesized proteins during the labelling phase, and the relative amount of labelled protein will reduce over time because of degradation. A thorough review of SILK in neurodegeneration is provided here (Paterson et al. 2019). Sato and colleagues used this approach to understand tau turnover in iPSC-neurons (Sato et al. 2018).

Interestingly, disease-associated tau species appear to have an increased turnover. For example 4R tau had a shorter half-life than 3R tau, and several tau peptides containing phosphorylation residues (T111/S113/T123, T212/214, T217, S262/T263 and S356/T361) were shown to have a faster turnover than their non-phosphorylated counterparts (Sato et al. 2018), suggesting differential proteostasis of specific tau species. It is also intriguing to note that the half-life of tau in iPSC-neurons is much shorter than in human participants (6.74 ± 0.45 days vs. 23 ± 6.4 days respectively) (Sato et al. 2018). This indicates developmental changes in tau proteostasis and is concordant with the idea that protective mechanisms against protein aggregation may become less efficient during ageing.

Further insights into the regulation of tau have been obtained by the identification of secreted tau fragments in the conditioned media from iPSC-neurons previously labelled with $^{13}\text{C}_6$ -leucine. Specifically, N-terminal fragments of tau were observed in cell culture media, but no peptides covering the C-terminal region (incorporating the microtubule-binding repeats) were observed (Sato et al. 2018). Together with antibody epitope mapping, the cleavage site was suggested to be between residues 210–230 of tau. The levels of labelled fragments increased in the media 3 days after labelling stopped, suggesting the cleavage and secretion of this fragment is a regulated, physiological process (Sato et al. 2018). The work by Sato *et al.* helped to change our understanding of tau in AD; the increase in total-tau and phosphorylated-tau seen extracellularly in AD represents an active secretion of phosphorylated and non-phosphorylated N-terminal tau fragments from live neurons exposed to A β , not passive release from dying neurons (Zetterberg, 2018).

12 | PROTEOMICS TO UNCOVER MECHANISM OF TAU-LINKED NEURONAL DYSFUNCTION

In addition to targeted proteomics, untargeted proteomics can provide a global and unbiased insight into dysregulated cellular

pathways, an approach commonly utilized for biomarker discovery and one that has successfully identified early metabolic changes in post-mortem AD brain tissue (Johnson et al. 2020). Tau pathology progresses throughout the brain in a predictable and defined manner, and recent research has focussed on mechanisms by which tau may be transferred from cell to cell (Braak et al. 2003). The propagation of tau via exosomes has been implicated in this process. Podvin and colleagues used iPSC-neurons overexpressing the repeat region of tau with the FTD-linked *MAPT* mutations P301L and V337 M to determine whether the presence of mutant tau would affect the composition of exosomes (Podvin et al. 2020). Widespread alterations in the proteome of exosomes from neurons expressing mutant tau were observed, including almost 245 proteins in control exosomes that were absent in those isolated from tau mutant overexpressing neurons. Several proteins uniquely present in tau exosomes have previously been linked to AD, including ANP32A, a potential modulator of tau phosphorylation, and PEN2, which is a subunit of γ -secretase. It would be interesting to further extend this work to iPSC-neurons with tau mutations; examining the impact of full length, mutant tau at the endogenous level.

13 | FUTURE DIRECTIONS

Although the number of studies using mass spectrometry to analyse tau in iPSC-neurons is relatively small, they have revealed the potential to gain insight into the physiological regulation of tau in healthy neurons and disease models. Future studies will likely enable a comprehensive characterization of the full range of tau PTMs, as previously described for mouse and human tau (Kametani et al. 2020; Morris et al. 2015). The importance of mapping tau PTMs has been highlighted by recent studies: cryoEM has enabled the detailed visualization of disease-specific tau structures, which may be influenced by PTM profile (Arakhamia et al. 2020). Furthermore, specific signatures of tau PTMs identified by mass spectrometry in brain tissue and CSF have recently been shown to correlate with clinical heterogeneity and disease severity in AD (Barthélemy et al. 2020; Dujardin et al. 2020; Wesseling et al. 2020). Thus, tau PTMs are directly related to clinical outcomes, and detailed profiling in iPSC-neurons will be essential to determine the extent to which they recapitulate disease pathologies.

14 | CONCLUSIONS

Induced pluripotent stem cell technology is now over a decade old. This technology has enabled 'disease in a dish' studies that provide a human model of the cell type affected by disease, expressing mutant genes at physiological levels. Indeed, the number of iPSC-related studies into Alzheimer's disease and dementia has steadily risen over the last decade, highlighting the relevance of this tool.

Mass spectrometry used in concert with iPSC technology has been instrumental in showing that these pre-clinical models do indeed effectively model the in vivo setting. For example A β and tau

are processed in a developmentally regulated manner and mature neuronal cultures are able to generate appropriate species of tau and A β spectra. The stage is now set to further develop these models to gain a deeper understanding of molecular mechanisms of neurodegeneration and screen for novel therapeutic agents.

iPSC-derived cultures are reductionist, providing simplified models of the brain. This enables a cell-type-specific analysis into the consequences of gene mutations. Additionally, these models are foetal, because of cellular reprogramming and epigenetic erasure. This allows underlying, constitutive effects of a mutation to be distinguished from late stage, complex neurodegenerative processes. What is more, models are continually evolving and the advent of 3D cerebral organoid techniques (Lancaster et al. 2013) allow cell behaviours to be investigated in a model more akin to the developing brain. Additionally, CRISPR/Cas9 genome editing enables complex genetic manipulations to be performed, such as combinations that are not typically present in nature (Cong et al. 2013; Mali et al. 2013).

Going forward, mass spectrometry will become instrumental in these studies (Figure 1). As well as employing global proteomic approaches that can effectively describe the consequences of inherited forms of dementia, detailed analysis of post-translational modifications will further our understanding of the earliest events of neurodegeneration. Adaptations of these techniques, such as SILK provide further crucial information, for example into the kinetics of protein turnover.

In sum, the combination of iPSC and mass spectrometry techniques have already proven the value of modelling Alzheimer's disease in vitro. The time is now for detailed mechanistic insights into early neurodegenerative processes and real progress to be made with drug discovery platforms towards novel therapeutic agents.

CONFLICTS OF INTEREST

HZ has served at scientific advisory boards for Denali, Roche Diagnostics, Wave, Samumed, Siemens Healthineers, Pinteon Therapeutics and CogRx, has given lectures in symposia sponsored by Fujirebio, Alzecure and Biogen, and is a co-founder of Brain Biomarker Solutions in Gothenburg AB (BBS), which is a part of the GU Ventures Incubator Program. The other authors have no conflicts of interest to declare.

ORCID

Charles Arber  <https://orcid.org/0000-0002-9743-8134>

Argyro Alatza  <https://orcid.org/0000-0002-3342-6804>

Claire A. Leckey  <https://orcid.org/0000-0001-5268-3681>

Ross W. Paterson  <https://orcid.org/0000-0001-9372-3635>

Henrik Zetterberg  <https://orcid.org/0000-0003-3930-4354>

Selina Wray  <https://orcid.org/0000-0003-3062-7050>

REFERENCES

Ahmed, Z., Bigio, E. H., Budka, H., Dickson, D. W., Ferrer, I., Ghetti, B., Giaccone, G., Hatanpaa, K. J., Holton, J. L., Josephs, K. A., Powers, J., Spina, S., Takahashi, H., White, C. L., Revesz, T., & Kovacs, G. G.

- (2013). Globular glial tauopathies (GGT): Consensus recommendations. *Acta Neuropathologica*, 126, 537–544.
- Al-Hilaly, Y. K., Williams, T. L., Stewart-Parker, M., Ford, L., Skaria, E., Cole, M., Bucher, W. G., Morris, K. L., Sada, A. A., Thorpe, J. R., & Serpell, L. C. (2014). A central role for dityrosine crosslinking of Amyloid- β in Alzheimer's disease. *Acta Neuropathologica Communications*, 2, 1–17.
- Alić, I., Goh, P. A., Murray, A., Portelius, E., Gkanatsiou, E., Gough, G., Mok, K. Y., Koschut, D., Brunmeir, R., Yeap, Y. J., O'Brien, N. L., Groet, J., Shao, X., Steven, H., Dunn, N. R., Kvartsberg, H., Brinkmalm, G., Hithersay, R., Startin, C., & Hamburg, S. ... Nižetić, D. (2020). Patient-specific Alzheimer-like pathology in trisomy 21 cerebral organoids reveals BACE2 as a gene dose-sensitive AD suppressor in human brain. *Molecular Psychiatry*, <https://doi.org/10.1038/s41380-020-0806-5>
- Arakhamia, T., Lee, C. E., Carlomagno, Y., Duong, D. M., Kundinger, S. R., Wang, K., Williams, D., DeTure, M., Dickson, D. W., Cook, C. N., Seyfried, N. T., Petrucelli, L., & Fitzpatrick, A. W. P. (2020). Posttranslational modifications mediate the structural diversity of Tauopathy strains. *Cell*, 180, 633–644.e12.
- Arber, C., Lovejoy, C., & Wray, S. (2017). Stem cell models of Alzheimer's disease: Progress and challenges. *Alzheimer's Research & Therapy*, 9, 42.
- Arber, C., Toombs, J., Lovejoy, C., Ryan, N. S., Paterson, R. W., Willumsen, N., Gkanatsiou, E., Portelius, E., Blennow, K., Heslegrave, A., Schott, J. M., Hardy, J., Lashley, T., Fox, N. C., Zetterberg, H., & Wray, S. (2020). Familial Alzheimer's disease patient-derived neurons reveal distinct mutation-specific effects on amyloid beta. *Molecular Psychiatry*, 25, 2919–2931.
- Arber, C., Villegas-Llerena, C., Toombs, J., Pocock, J. M., Ryan, N. S., Fox, N. C., Zetterberg, H., Hardy, J., & Wray, S. (2019). Amyloid precursor protein processing in human neurons with an allelic series of the PSEN1 intron 4 deletion mutation and total presenilin-1 knockout. *Brain Communications* 1, 1–10.
- Baranello, R., Bharani, K., Padmaraju, V., Chopra, N., Lahiri, D., Greig, N., Pappolla, M., & Sambamurti, K. (2015). Amyloid-beta protein clearance and degradation (ABCD) pathways and their role in Alzheimer's disease. *Current Alzheimer Research*, 12, 32–46.
- Barthélemy, N. R., Fenaille, F., Hirtz, C., Sergeant, N., Schraen-Maschke, S., Vialaret, J., Buée, L., Gabelle, A., Junot, C., Lehmann, S., & Becher, F. (2016). Tau protein quantification in human cerebrospinal fluid by targeted mass spectrometry at high sequence coverage provides insights into its primary structure heterogeneity. *Journal of Proteome Research*, 15, 667–676.
- Barthélemy, N. R., Li, Y., Joseph-Mathurin, N., Gordon, B. A., Hassenstab, J., Benzinger, T. L. S., Buckles, V., Fagan, A. M., Perrin, R. J., Goate, A. M., Morris, J. C., Karch, C. M., Xiong, C., Allegri, R., Mendez, P. C., Berman, S. B., Ikeuchi, T., Mori, H., Shimada, H., ... Xiong, X. (2020). A soluble phosphorylated tau signature links tau, amyloid and the evolution of stages of dominantly inherited Alzheimer's disease. *Nature Medicine*, 26, 398–407.
- Barthélemy, N. R., Mallipeddi, N., Moiseyev, P., Sato, C., & Bateman, R. J. (2019). Tau phosphorylation rates measured by mass spectrometry differ in the intracellular brain vs. Extracellular cerebrospinal fluid compartments and are differentially affected by Alzheimer's disease. *Frontiers in Aging Neuroscience*, 11, 1–18. <https://doi.org/10.3389/fnagi.2019.00121>.
- Beevers, J. E., Lai, M. C., Collins, E., Booth, H. D. E., Zambon, F., Parkkinen, L., Vowles, J., Cowley, S. A., Wade-Martins, R., & Caffrey, T. M. (2017). MAPT genetic variation and neuronal maturity alter isoform expression affecting axonal transport in iPSC-derived dopamine neurons. *Stem Cell Reports*, 9, 587–599. <https://doi.org/10.1016/j.stemcr.2017.06.005>.
- Bergström, P., Agholme, L., Nazir, F. H., Satir, T. M., Toombs, J., Wellington, H., Strandberg, J., Bontell, T. O., Kvartsberg, H., Holmström, M., Boreström, C., Simonsson, S., Kunath, T., Lindahl, A., Blennow, K., Hanse, E., Portelius, E., Wray, S., & Zetterberg, H. (2016). Amyloid precursor protein expression and processing are differentially regulated during cortical neuron differentiation. *Scientific Reports*, 6, 29200.
- Braak, H., & Braak, E. (1989). Cortical and subcortical argyrophilic grains characterize a disease associated with adult onset dementia. *Neuropathology and Applied Neurobiology*, 15(1), 13–26.
- Braak, H., Tredici, K., Del, R. U., de Vos, R. A., Jansen, S. E. N. H., & Braak, E. (2003). Staging of brain pathology related to sporadic Parkinson's disease. *Neurobiology of Aging*, 24, 197–211.
- Brion, J. P., Smith, C., Couck, A. M., Gallo, J. M., & Anderton, B. H. (1993). Developmental changes in tau phosphorylation: Fetal tau is transiently phosphorylated in a manner similar to paired helical filament-tau characteristic of Alzheimer's disease. *Journal of Neurochemistry*, 61, 2071–2080.
- Bros, P., Delatour, V., Vialaret, J., Lalere, B., Barthelemy, N., Gabelle, A., Lehmann, S., & Hirtz, C. (2015). Quantitative detection of amyloid- β peptides by mass spectrometry: State of the art and clinical applications. *Clinical Chemistry and Laboratory Medicine*, 53, 1483–1493.
- Brownjohn, P. W., Smith, J., Portelius, E., Serneels, L., Kvartsberg, H., Strooper, B., De, B. K., Zetterberg, H., & Livesey, F. J. (2017). Phenotypic screening identifies modulators of amyloid precursor protein processing in human stem cell models of Alzheimer's disease. *Stem Cell Reports*, 8, 870–882. <https://doi.org/10.1016/j.stemcr.2017.02.006>.
- Buée, L. & Delacourte, A. (1999). Comparative biochemistry of tau in progressive Supranuclear palsy, Corticobasal degeneration, FTDP-17 and Pick's disease. *Brain Pathology*, 9(4), 681–693.
- Chang, C. Y., Chen, S. M., Lu, H. E., Lai, S. M., Lai, P. S., Shen, P. W., Chen, P. Y., Shen, C. I., Harn, H. J., Lin, S. Z., Hwang, S. M., & Su, H. L. (2015). N-butylidenephthalide attenuates Alzheimer's disease-like cytopathy in down syndrome induced pluripotent stem cell-derived neurons. *Scientific Reports*, 5, 8744.
- Chávez-Gutiérrez, L., Bammens, L., Benilova, I., Vandersteen, A., Benurwar, M., Borgers, M., Lismont, S., Zhou, L., Van Cleynenbreugel, S., Esselmann, H., Wiltfang, J., Serneels, L., Karran, E., Gijsen, H., Schymkowitz, J., Rousseau, F., Broersen, K., & De Strooper, B. (2012). The mechanism of γ -Secretase dysfunction in familial Alzheimer disease. *EMBO Journal*, 31, 2261–2274. <https://doi.org/10.1038/emboj.2012.79>.
- Cong, L., Ran, F. A., Cox, D., Lin, S., Barretto, R., Habib, N., Hsu, P. D., Wu, X., Jiang, W., Marraffini, L. A., & Zhang, F. (2013). Multiplex genome engineering using CRISPR/Cas systems. *Science*, 339, 819–823. <https://doi.org/10.1126/science.1231143>.
- Crary, J. F., Trojanowski, J. Q., Schneider, J. A., Abisambra, J. F., Abner, E. L., Alafuzoff, I., Arnold, S. E., Attems, J., Beach, T. G., Bigio, E. H., Cairns, N. J., Dickson, D. W., Gearing, M., Grinberg, L. T., Hof, P. R., Hyman, B. T., Jellinger, K., Jicha, G. A., Kovacs, G. G., ... Nelson, P. T. (2014). Primary age-related tauopathy (PART): A common pathology associated with human aging. *Acta Neuropathologica*, 128, 755–766. <https://doi.org/10.1007/s00401-014-1349-0>.
- Duan, L., Bhattacharyya, B. J., Belmadani, A., Pan, L., Miller, R. J., & Kessler, J. A. (2014). Stem cell derived basal forebrain cholinergic neurons from Alzheimer's disease patients are more susceptible to cell death. *Molecular Neurodegeneration*, 9, 3–<https://doi.org/10.1186/1750-1326-9-3>.
- Dujardin, S., Commins, C., Lathuiliere, A., Beerepoot, P., Fernandes, A. R., Kamath, T. V., Santos, D. L., Mark, B., & Klickstein, N., Corjuc, D. L., Corjuc, B. T., Dooley, P. M., Viode, A., Oakley, D. H., Moore, B. D., Mullin, K., & Jean-Gilles, D. ... Bradley, T. (2020). Tau molecular diversity contributes to clinical heterogeneity in Alzheimer's disease. *Nature Medicine*, 26, 1256–1263.
- Ferrer, I., Aguiló, G. M., Carmona, M., Andrés-Benito, P., Torrejón-Escribano, B., Garcia-Esparcia, P., & del Rio, J. A. (2019). Involvement



- of Oligodendrocytes in Tau seeding and spreading in tauopathies. *Frontiers in Aging Neuroscience* 11.
- Gallien, S., Duriez, E., Crone, C., Kellmann, M., Moehring, T., & Domon, B. (2012). Targeted proteomic quantification on quadrupole-orbitrap mass spectrometer. *Molecular & Cellular Proteomics: MCP*, 11, 1709–1723.
- Gheyara, A. L., Ponnusamy, R., Djukic, B., Craft, R. J., Ho, K., Guo, W., Finucane, M. M., Sanchez, P. E., & Mucke, L. (2014). Tau reduction prevents disease in a mouse model of Dravet syndrome. *Annals of Neurology*, 76, 443–456.
- Goedert, M., Spillantini, M. G., Jakes, R., Rutherford, D., & Crowther, R. A. (1989). Multiple isoforms of human microtubule-associated protein tau: Sequences and localization in neurofibrillary tangles of Alzheimer's disease. *Neuron*, 3, 519–526. [https://doi.org/10.1016/0896-6273\(89\)90210-9](https://doi.org/10.1016/0896-6273(89)90210-9).
- Gonzalez, C., Armijo, E., Bravo-Alegria, J., Becerra-Calixto, A., Mays, C. E., & Soto, C. (2018). Modeling amyloid beta and tau pathology in human cerebral organoids. *Molecular Psychiatry*, 23, 2363–2374.
- Guo, T., Noble, W., & Hanger, D. P. (2017). Roles of tau protein in health and disease. *Acta Neuropathologica*, 133(5), 665–704. <https://doi.org/10.1007/s00401-017-1707-9>.
- Hanger, D. P., Anderton, B. H., & Noble, W. (2009). Tau phosphorylation: The therapeutic challenge for neurodegenerative disease. *Trends in Molecular Medicine*, 15, 112–119.
- Hanger, D. P., Byers, H. L., Wray, S., Leung, K.-Y., Saxton, M. J., Seereeram, A., Reynolds, C. H., Ward, M. A., & Anderton, B. H. (2007). Novel phosphorylation sites in tau from alzheimer brain support a role for casein kinase 1 in disease pathogenesis. *Journal of Biological Chemistry*, 282, 23645–23654.
- Hardy, J. (2017). The discovery of Alzheimer-causing mutations in the APP gene and the formulation of the “amyloid cascade hypothesis”. *FEBS Journal*, 284, 1040–1044.
- Hefti, M. M., Farrell, K., Kim, S. H., Bowles, K. R., Fowkes, M. E., Raj, T., & Crary, J. F. (2018). High-resolution temporal and regional mapping of MAPT expression and splicing in human brain development. *PLoS One*, 13(4), e0195771. <https://doi.org/10.1371/journal.pone.0195771>.
- Hutton, M., Lendon, C. L., Rizzu, P., Baker, M., Froelich, S., Houlden, H., Pickering-Brown, S., Chakraverty, S., Isaacs, A., Grover, A., Hackett, J., Adamson, J., Lincoln, S., Dickson, D., Davies, P., Petersen, R. C., Stevens, M., de Graaff, E., Wauters, E., ... Heutink, P. (1998). Association of missense and 5'-splice-site mutations in tau with the inherited dementia FTDP-17. *Nature*, 393, 702–705. <https://doi.org/10.1038/31508>.
- Iovino, M., Agathou, S., González-Rueda, A., Velasco-Herrera, D. C., Martin, B., Barbara, A., Antonella, L., Timothy, O' D., Sean, G., Imbisaat, G., Daniel, V., Ludovic, P., Ole, K., Thóra, R., & Spillantini, M. G. (2015). Early maturation and distinct tau pathology in induced pluripotent stem cell-derived neurons from patients with MAPT mutations. *Brain*, 138, 3345–3359.
- Iovino, M., Patani, R., Watts, C., Chandran, S., & Spillantini, M. G. (2010). Human stem cell-derived neurons: A system to study human tau function and dysfunction. *PLoS One*, 5. <https://doi.org/10.1371/journal.pone.0013947>.
- Israel, M. A., Yuan, S. H., Bardy, C., Reyna, S. M., Mu, Y., Herrera, C., Hefferan, M. P., Van Gorp, S., Nazor, K. L., Boscolo, F. S., Carson, C. T., Laurent, L. C., Marsala, M., Gage, F. H., Remes, A. M., Koo, E. H., & Goldstein, L. S. B. (2012). Probing sporadic and familial Alzheimer's disease using induced pluripotent stem cells. *Nature*, 482, 216–220. <https://doi.org/10.1038/nature10821>.
- Ittner, L. M., Ke, Y. D., Delerue, F., Bi, M., Gladbach, A., van Eersel, J., Wölfing, H., Chieng, B. C., Christie, M. D. J., Napier, I. A., Eckert, A., Staufenbiel, M., Hardeman, E., & Götz, J. (2010). Dendritic function of tau mediates amyloid-beta toxicity in Alzheimer's disease mouse models. *Cell*, 142, 387–397.
- Janelidze, S., Zetterberg, H., Mattsson, N., Palmqvist, S., Vanderstichele, H., Lindberg, O., van Westen, D., Stomrud, E., Minthon, L., Blennow, K., & Hansson, O. (2016). CSF A β 42/A β 40 and A β 42/A β 38 ratios: Better diagnostic markers of Alzheimer disease. *Annals of Clinical and Translational Neurology*, 3, 154–165.
- Jo, J., Xiao, Y., Sun, A. X., Cukuroglu, E., Tran, H.-D., Göke, J., Tan, Z. Y., Saw, T. Y., Tan, C.-P., Lokman, H., Lee, Y., Kim, D., Ko, H. S., Kim, S.-O., Park, J. H., Cho, N.-J., Hyde, T. M., Kleinman, J. E., Shin, J. H., ... Ng, H.-H. (2016). Midbrain-like organoids from human pluripotent stem cells contain functional dopaminergic and neuromelanin-producing neurons. *Cell Stem Cell*, 19, 248–257.
- Johnson, E. C. B., Dammer, E. B., Duong, D. M., Ping, L., Zhou, M., Yin, L., Higginbotham, L. A., Guajardo, A., White, B., Troncoso, J. C., Thambisetty, M., Montine, T. J., Lee, E. B., Trojanowski, J. Q., Beach, T. G., Reiman, E. M., Haroutunian, V., Wang, M., Schadt, E., ... Seyfried, N. T. (2020). Large-scale proteomic analysis of Alzheimer's disease brain and cerebrospinal fluid reveals early changes in energy metabolism associated with microglia and astrocyte activation. *Nature Medicine*, 26, 769–780.
- Kametani, F., Yoshida, M., Matsubara, T., Murayama, S., Saito, Y., Kawakami, I., Onaya, M., Tanaka, H., Kakita, A., Robinson, A. C., Mann, D. M. A., & Hasegawa, M. (2020). Comparison of common and disease-specific post-translational modifications of pathological tau associated with a wide range of tauopathies. *Frontiers in Neuroscience*, 14, 1110.
- Kirwan, P., Turner-Bridger, B., Peter, M., Momoh, A., Arambepola, D., Robinson, H. P. C., & Livesey, F. J. (2015). Development and function of human cerebral cortex neural networks from pluripotent stem cells in vitro. *Development*, 142, 3178–3187. <https://doi.org/10.1242/dev.123851>.
- Koch, P., Tamboli, I. Y., Mertens, J., Wunderlich, P., Ladewig, J., Stüber, K., Esselmann, H., Wiltfang, J., Brüstle, O., & Walter, J. (2012). Presenilin-1 L166P mutant human pluripotent stem cell-derived neurons exhibit partial loss of γ -secretase activity in endogenous amyloid- β generation. *American Journal of Pathology*, 180, 2404–2416. <https://doi.org/10.1016/j.ajpath.2012.02.012>.
- Kondo, T., Asai, M., Tsukita, K., Kutoku, Y., Ohsawa, Y., Sunada, Y., Imamura, K., Egawa, N., Yahata, N., Okita, K., Takahashi, K., Asaka, I., Aoi, T., Watanabe, A., Watanabe, K., Kadoya, C., Nakano, R., Watanabe, D., Maruyama, K., ... Inoue, H. (2013). Modeling Alzheimer's disease with iPSCs reveals stress phenotypes associated with intracellular A β and differential drug responsiveness. *Cell Stem Cell*, 12, 487–496. <https://doi.org/10.1016/j.stem.2013.01.009>.
- Kosik, K. S., Orecchio, L. D., Bakalis, S., & Neve, R. L. (1989). Developmentally regulated expression of specific tau sequences. *Neuron*, 2, 1389–1397. [https://doi.org/10.1016/0896-6273\(89\)90077-9](https://doi.org/10.1016/0896-6273(89)90077-9).
- Kovacs, G. G., Ferrer, I., Grinberg, L. T., Alafuzoff, I., Attems, J., Budka, H., Cairns, N. J., Crary, J. F., Duyckaerts, C., Ghetti, B., Halliday, G. M., Ironside, J. W., Love, S., Mackenzie, I. R., Munoz, D. G., Murray, M. E., Nelson, P. T., Takahashi, H., Trojanowski, J. Q., ... Dickson, D. W. (2016). Aging-related tau astrogliopathy (ARTAG): Harmonized evaluation strategy. *Acta Neuropathologica*, 131, 87–102. <https://doi.org/10.1007/s00401-015-1509-x>.
- Kumar, S., Rezaei-Ghaleh, N., Terwel, D., Thal, D. R., Richard, M., Hoch, M., Donald, M. C., Jessica, M., Wüllner, U., Glebov, K., Heneka, M. T., Walsh, D. M., Zweckstetter, M., & Walter, J. (2011). Extracellular phosphorylation of the amyloid β 2-peptide promotes formation of toxic aggregates during the pathogenesis of Alzheimer's disease. *EMBO Journal*, 30, 2255–2265.
- Kumar, S., Wirths, O., Theil, S., Gerth, J., Bayer, T. A., & Walter, J. (2013). Early intraneuronal accumulation and increased aggregation of phosphorylated A β in a mouse model of Alzheimer's disease. *Acta Neuropathologica*, 125, 699–709. <https://doi.org/10.1007/s00401-013-1107-8>.



- Kummer, M. P., Hermes, M., Delekarte, A., Hammerschmidt, T., Kumar, S., Terwel, D., Walter, J., Pape, H.-C., König, S., Roeber, S., Jessen, F., Klockgether, T., Korte, M., & Heneka, M. T. (2011). Nitration of tyrosine 10 critically enhances amyloid β aggregation and plaque formation. *Neuron*, 71(5), 833–844. <https://doi.org/10.1016/j.neuron.2011.07.001>.
- Kwart, D., Gregg, A., Scheckel, C., Murphy, E. A., Paquet, D., Duffield, M., Fak, J., Olsen, O., Darnell, R. B., & Tessier-Lavigne, M. (2019). A large panel of isogenic APP and PSEN1 mutant human iPSC neurons reveals shared endosomal abnormalities mediated by APP β -CTFs, not A β . *Neuron*, 104, 256–270.e5. <https://doi.org/10.1016/j.neuron.2019.07.010>.
- Lancaster, M. A., Corsini, N. S., Wolfinger, S., Gustafson, E. H., Phillips, A. W., Burkard, T. R., Otani, T., Livesey, F. J., & Knoblich, J. A. (2017). Guided self-organization and cortical plate formation in human brain organoids. *Nature Biotechnology*, 35, 659–666.
- Lancaster, M. A., & Knoblich, J. A. (2014). Generation of cerebral organoids from human pluripotent stem cells. *Nature Protocols*, 9, 2329–2340.
- Lancaster, M. A., Renner, M., Martin, C. A., Wenzel, D., Bicknell, L. S., Hurler, M. E., Homfray, T., Penninger, J. M., Jackson, A. P., & Knoblich, J. A. (2013). Cerebral organoids model human brain development and microcephaly. *Nature*, 501, 373–379. <https://doi.org/10.1038/nature12517>.
- Lee, J., Bignone, P. A., Coles, L. S., Liu, Y., Snyder, E., & Larocca, D. (2020). Induced pluripotency and spontaneous reversal of cellular aging in supercentenarian donor cells. *Biochemical and Biophysical Research Communications*, 525(3), 563–569.
- Lewczuk, P., Esselmann, H., Groemer, T. W., Bibl, M., Maler, J. M., Steinacker, P., Otto, M., Kornhuber, J., & Wiltfang, J. (2004). Amyloid β peptides in cerebrospinal fluid as profiled with surface enhanced laser desorption/ionization time-of-flight mass spectrometry: Evidence of novel biomarkers in Alzheimer's disease. *Biological Psychiatry*, 55, 524–530.
- Lines, G., Casey, J. M., Preza, E., & Wray, S. (2020). Modelling frontotemporal dementia using patient-derived induced pluripotent stem cells. *Molecular and Cellular Neuroscience*, 109, 103553.
- Liu, Q., Waltz, S., Woodruff, G., Ouyang, J., Israel, M. A., Herrera, C., Sarsoza, F., Tanzi, R. E., Koo, E. H., Ringman, J. M., Goldstein, L. S. B., Wagner, S. L., & Yuan, S. H. (2014). Effect of potent γ -secretase modulator in human neurons derived from multiple Presenilin 1-induced pluripotent stem cell mutant carriers. *JAMA Neurol.*, 71, 1481–1482. <https://doi.org/10.1001/jamaneurol.2014.2482>.
- Mackenzie, I. R. A., & Neumann, M. (2016). Molecular neuropathology of frontotemporal dementia: insights into disease mechanisms from postmortem studies. *Journal of Neurochemistry*, 138, 54–70.
- Maddalena, A. S., Papassotiropoulos, A., Gonzalez-Agosti, C., Signorell, A., Hegi, T., Pasch, T., Nitsch, R. M., & Hock, C. (2004). Cerebrospinal fluid profile of amyloid β peptides in patients with Alzheimer's disease determined by protein biochip technology. *Neuro-Degenerative Diseases*, 1, 231–235.
- Mahairaki, V., Ryu, J., Peters, A., Chang, Q., Li, T., Park, T. S., BurrIDGE, P. W., Talbot, C. C., Asnaghi, L., Martin, L. J., Zambidis, E. T., & Koliatsos, V. E. (2014). Induced pluripotent stem cells from familial Alzheimer's disease patients differentiate into mature neurons with amyloidogenic properties. *Stem Cells and Development*, 23, 2996–3010. <https://doi.org/10.1089/scd.2013.0511>.
- Mair, W., Muntel, J., Tepper, K., Tang, S., Biernat, J., Seeley, W. W., Kosik, K. S., Mandelkow, E., Steen, H., & Steen, J. A. (2016). FLEXITau: Quantifying post-translational modifications of tau protein in vitro and in human disease. *Analytical Chemistry*
- Mali, P., Yang, L., Esvelt, K. M., Aach, J., Guell, M., DiCarlo, J. E., Norville, J. E., & Church, G. M. (2013). RNA-guided human genome engineering via cas9. *Science* (80-.). 339, 823–826.
- Mertens, J., Paquola, A. C. M., Ku, M., Hatch, E., Böhnke, L., Ladjevardi, S., McGrath, S., Campbell, B., Lee, H., Herdy, J. R., Gonçalves, J. T., Toda, T., Kim, Y., Winkler, J., Yao, J., Hetzer, M. W., & Gage, F. H. (2015). Directly reprogrammed human neurons retain aging-associated transcriptomic signatures and reveal age-related nucleocytoplasmic defects. *Cell Stem Cell*, 17, 705–718. <https://doi.org/10.1016/j.stem.2015.09.001>.
- Moore, S., Evans, L. D. B., Andersson, T., Portelius, E., Smith, J., Dias, T. B., Saurat, N., McGlade, A., Kirwan, P., Blennow, K., Hardy, J., Zetterberg, H., & Livesey, F. J. (2015). APP metabolism regulates tau proteostasis in human cerebral cortex neurons. *Cell Reports*, 11, 689–696. <https://doi.org/10.1016/j.celrep.2015.03.068>.
- Mori, H., Takio, K., Ogawara, M., & Selkoe, D. J. (1992). Mass spectrometry of purified amyloid beta protein in Alzheimer's disease. *Journal of Biological Chemistry*, 267, 17082–17086.
- Morris, M., Knudsen, G. M., Maeda, S., Trinidad, J. C., Ioanoviciu, A., Burlingame, A. L., & Mucke, L. (2015). Tau post-translational modifications in wild-type and human amyloid precursor protein transgenic mice. *Nature Neuroscience*, 18, 1183–1189.
- Muratore, C. R., Rice, H. C., Srikanth, P., Callahan, D. G., Shin, T., Benjamin, L. N. P., Walsh, D. M., Selkoe, D. J., & Young-Pearse, T. L. (2014). The familial Alzheimer's disease APPV717I mutation alters APP processing and Tau expression in iPSC-derived neurons. *Human Molecular Genetics*, 23, 3523–3536.
- Nakamura, A., Kaneko, N., Villemagne, V. L., Kato, T., Doecke, J., Doré, V., Fowler, C., Li, Q.-X., Martins, R., Rowe, C., Tomita, T., Matsuzaki, K., Ishii, K., Ishii, K., Arahata, Y., Iwamoto, S., Ito, K., Tanaka, K., Masters, C. L., & Yanagisawa, K. (2018). High performance plasma amyloid- β biomarkers for Alzheimer's disease. *Nature*, 554, 249–254.
- Oddo, S., Caccamo, A., Shepherd, J. D., Murphy, M. P., Golde, T. E., Kaye, D., Metherate, R., Mattson, M. P., Akbari, Y., & LaFerla, F. M. (2003). Triple-transgenic model of Alzheimer's disease with plaques and tangles. *Neuron*, 39, 409–421. [https://doi.org/10.1016/S0896-6273\(03\)00434-3](https://doi.org/10.1016/S0896-6273(03)00434-3).
- Paonessa, F., Evans, L. D., Solanki, R., Larrieu, D., Wray, S., Hardy, J., Jackson, S. P., & Livesey, F. J. (2019). Microtubules deform the nuclear membrane and disrupt nucleocytoplasmic transport in tau-mediated frontotemporal dementia. *Cell Reports*, 26, 582–593.e5.
- Patani, R., Lewis, P. A., Trabzuni, D., Puddifoot, C. A., Wyllie, D. J. A., Walker, R., Smith, C., Hardingham, G. E., Weale, M., Hardy, J., Chandran, S., & Ryten, M. (2012). Investigating the utility of human embryonic stem cell-derived neurons to model ageing and neurodegenerative disease using whole-genome gene expression and splicing analysis. *Journal of Neurochemistry*, 122, 738–751. <https://doi.org/10.1111/j.1471-4159.2012.07825.x>.
- Paterson, R. W., Gabelle, A., Lucey, B. P., Barthélemy, N. R., Leckey, C. A., Hirtz, C., Lehmann, S., Sato, C., Patterson, B. W., West, T., Yarasheski, K., Rohrer, J. D., Wildburger, N. C., Schott, J. M., Karch, C. M., Wray, S., Miller, T. M., Elbert, D. L., Zetterberg, H., ... Bateman, R. J. (2019). SILK studies – capturing the turnover of proteins linked to neurodegenerative diseases. *Nature Reviews. Neurology*, 15, 419–427.
- Perez-Garmendia, R., & Gevorkian, G. (2013). Pyroglutamate-modified amyloid beta peptides: emerging targets for Alzheimer's disease immunotherapy. *Current Neuropharmacology*, 11, 491–498.
- Pérez-Grijalba, V., Pesini, P., Allué, J. A., Sarasa, L., Montañés, M., Lacosta, A.-M., Casabona, D., San-José, I., Boada, M., Tárraga, L., Ruiz, A., & Sarasa, M. (2014). A β 1-17 is a major amyloid- β fragment isoform in cerebrospinal fluid and blood with possible diagnostic value in Alzheimer's disease. *J. Alzheimer's Dis.* <https://doi.org/10.3233/JAD-140156>.
- Peterson, A. C., Russell, J. D., Bailey, D. J., Westphall, M. S., & Coon, J. J. (2012). Parallel reaction monitoring for high resolution and high mass accuracy quantitative, targeted proteomics. *Molecular & Cellular Proteomics: MCP*, 11, 1475–1488.



- Podvin, S., Jones, A., Liu, Q., Aulston, B., Ransom, L., Ames, J., Shen, G., Lietz, C. B., Jiang, Z., O'Donoghue, A. J., Winston, C., Ikezu, T., Rissman, R. A., Yuan, S., & Hook, V. (2020). Dysregulation of exosome cargo by mutant tau expressed in human-induced pluripotent stem cell (iPSC) neurons revealed by proteomics analyses. *Molecular & Cellular Proteomics: MCP*, *19*(6), 1017–1034. <https://doi.org/10.1074/mcp.RA120.002079>.
- Poorkaj, P., Bird, T. D., Wijsman, E., Nemens, E., Garruto, R. M., Anderson, L., Andreadis, A., Wiederholt, W. C., Raskind, M., & Schellenberg, G. D. (1998). Tau is a candidate gene for chromosome 17 frontotemporal dementia. *Annals of Neurology*, *43*, 815–825. <https://doi.org/10.1002/ana.410430617>.
- Portelius, E., Bogdanovic, N., Gustavsson, M. K., Volkman, I., Brinkmalm, G., Zetterberg, H., Winblad, B., & Blennow, K. (2010). Mass spectrometric characterization of brain amyloid beta isoform signatures in familial and sporadic Alzheimer's disease. *Acta Neuropathologica*, *120*, 185–193.
- Portelius, E., Dean, R. A., Gustavsson, M. K., Andreasson, U., Zetterberg H., Siemers E., Blennow K. (2010). A novel Abeta isoform pattern in CSF reflects gamma-secretase inhibition in Alzheimer disease. *Alzheimer's Research & Therapy*, *2*, 7.
- Portelius, E., Gustavsson, M. K., Zetterberg, H., Andreasson, U., & Blennow, K. (2012). Evaluation of the performance of novel A β isoforms as theragnostic markers in Alzheimer's disease: From the cell to the patient. *Neuro-Degenerative Diseases*, *10*, 138–140.
- Portelius, E., Hansson, S. F., Tran, A. J., Zetterberg, H., Grognet, P., Vanmechelen, E., Höglund, K., Brinkmalm, G., Westman-Brinkmalm, A., Nordhoff, E., Blennow, K., & Gobom, J. (2008). Characterization of tau in cerebrospinal fluid using mass spectrometry. *Journal of Proteome Research*, *7*, 2114–2120. <https://doi.org/10.1021/pr7008669>.
- Portelius, E., Price, E., Brinkmalm, G., Stiteler, M., Olsson, M., Persson, R., Westman-Brinkmalm, A., Zetterberg, H., Simon, A. J., & Blennow, K. (2011). A novel pathway for amyloid precursor protein processing. *Neurobiology of Aging*, *32*, 1090–1098.
- Portelius, E., Tran, A. J., Andreasson, U., Persson, R., Brinkmann, G., Zetterberg, H., Blennow, K., & Westman-Brinkmalm, A. (2007). Characterization of amyloid β peptides in cerebrospinal fluid by an automated immunoprecipitation procedure followed by mass spectrometry. *Journal of Proteome Research*, *6*, 4433–4439. <https://doi.org/10.1021/pr0703627>.
- Qian, X., Jacob, F., Song, M. M., Nguyen, H. N., Song, H., & Ming, G. (2018). Generation of human brain region-specific organoids using a miniaturized spinning bioreactor. *Nature Protocols*, *13*, 565–580.
- Qian, X., Nguyen, H. N., Song, M. M., Hadiono, C., Ogden, S. C., Hammack, C., Yao, B., Hamersky, G. R., Jacob, F., Zhong, C., Yoon, K.-J., Jeang, W., Lin, L. I., Li, Y., Thakor, J., Berg, D. A., Zhang, C. E., Kang, E., Chickering, M., ... Ming, G.-L. (2016). Brain-region-specific Organoids using mini-bioreactors for modeling ZIKV exposure. *Cell*, *165*, 1238–1254.
- Raja, W. K., Mungenast, A. E., Lin, Y. T., Ko, T., Abdurrob, F., Seo, J., & Tsai, L. H. (2016). Self-organizing 3D human neural tissue derived from induced pluripotent stem cells recapitulate Alzheimer's disease phenotypes. *PLoS One*, *11*, e0161969. <https://doi.org/10.1371/journal.pone.0161969>.
- Roberson, E. D., Scarce-Levie, K., Palop, J. J., Yan, F., Cheng, I. H., Wu, T., Gerstein, H., Yu, G.-Q., & Mucke, L. (2007). Reducing endogenous tau ameliorates amyloid beta-induced deficits in an Alzheimer's disease mouse model. *Science*, *316*, 750–754.
- Saido, T. C., Yamao-Harigaya, W., Iwatsubo, T., & Kawashima, S. (1996). Amino- and carboxyl-terminal heterogeneity of β -amyloid peptides deposited in human brain. *Neuroscience Letters*, *215*, 173–176.
- Sato, C., Barthélemy, N. R., Mawuenyega, K. G., Patterson, B. W., Gordon, B. A., Jockel-Balsarotti, J., Sullivan, M., Crisp, M. J., Kasten, T., Kirmess, K. M., Kanaan, N. M., Yarasheski, K. E., Baker-Nigh, A., Benzinger, T. L. S., Miller, T. M., Karch, C. M., & Bateman, R. J. (2018). Tau kinetics in neurons and the human central nervous system. *Neuron*, *98*, 861–864.
- Sergeant, N., Bretteville, A., Hamdane, M., Caillet-Boudin, M. -L., Grognet, P., Bombois, S., Blum, D., Delacourte, A., Pasquier, F., Vanmechelen, E., Schraen-Maschke, S., & Bué, L. (2008). Biochemistry of Tau in Alzheimer's disease and related neurological disorders. *Expert Rev. Proteomics*, *5*, 207–224.
- Sergeant, N., Watzet, A., & Delacourte, A. (1999). Neurofibrillary degeneration in progressive supranuclear palsy and corticobasal degeneration. Tau pathologies with exclusively "Exon 10" isoforms. *Journal of Neurochemistry*, *72*, 1243–1249.
- Shi, X. P., Tugusheva, K., Bruce, J. E., Lucka, A., Wu, G. X., Chen-Dodson, E., Price, E., Li, Y., Xu, M., Huang, Q., Sardana, M. K., & Hazuda, D. J. (2003). β -secretase cleavage at amino acid residue 34 in the amyloid β peptide is dependent upon γ -secretase activity. *Journal of Biological Chemistry*, *278*, 21286–21294.
- Shi, Y., Kirwan, P., Smith, J., MacLean, G., Orkin, S. H., & Livesey, F. J. (2012). A human stem cell model of early Alzheimer's disease pathology in Down syndrome. *Science Translational Medicine*, *4*, 124ra29.
- Shi, Y., Kirwan, P., Smith, J., Robinson, H. P. C., & Livesey, F. J. (2012). Human cerebral cortex development from pluripotent stem cells to functional excitatory synapses. *Nature Neuroscience*, *15*(477–86), S1.
- Siegel, G., Gerber, H., Koch, P., Bruestle, O., Fraering, P. C., & Rajendran, L. (2017). The Alzheimer's disease γ -secretase generates higher 42:40 ratios for β -amyloid than for p3 peptides. *Cell Reports*, *19*, 1967–1976.
- Silva, M. C., Cheng, C., Mair, W., Almeida, S., Fong, H., Biswas, M. H. U., Zhang, Z., Huang, Y., Temple, S., Coppola, G., Geschwind, D. H., Karydas, A., Miller, B. L., Kosik, K. S., Gao, F.-B., Steen, J. A., & Haggarty, S. J. (2016). Human iPSC-derived neuronal model of tau-A152T frontotemporal dementia reveals tau-mediated mechanisms of neuronal vulnerability. *Stem Cell Reports*, *7*, 325–340.
- Sposito, T., Preza, E., Mahoney, C. J., Setó-Salvia, N., Ryan, N. S., Morris, H. R., Arber, C., Devine, M. J., Houlden, H., Warner, T. T., Bushell, T. J., Zagnoni, M., Kunath, T., Livesey, F. J., Fox, N. C., Rossor, M. N., Hardy, J., & Wray, S. (2015). Developmental regulation of tau splicing is disrupted in stem cell-derived neurons from frontotemporal dementia patients with the 10 + 16 splice-site mutation in MAPT. *Human Molecular Genetics*, *24*(18), 5260–5269.
- Sproul, A. A., Jacob, S., Pre, D., Kim, S. H., Nestor, M. W., Navarro-Sobrinho, M., Santa-Maria, I., Zimmer, M., Aubry, S., Steele, J. W., Kahler, D. J., Dranovsky, A., Arancio, O., Crary, J. F., Gandy, S., & Nogle, S. A. (2014). Characterization and molecular profiling of PSEN1 familial Alzheimer's disease iPSC-derived neural progenitors. *PLoS One*, *9*, e84547. <https://doi.org/10.1371/journal.pone.0084547>.
- Szaruga, M., Munteanu, B., Lismont, S., Veugelen, S., Horréc, K., Mercken, M., Saido, T. C., Ryan, N. S., De Vos, T., Savvides, S. N., Gallardo, R., Schymkowitz, J., Rousseau, F., Fox, N. C., Hopf, C., De Strooper, B., & Chávez-Gutiérrez, L. (2017). Alzheimer's-causing mutations shift A β length by destabilizing γ -secretase-A β n interactions. *Cell*, *170*, 443–456.e14. <https://doi.org/10.1016/j.cell.2017.07.004>.
- Takagi-Niidome, S., Sasaki, T., Osawa, S., Sato, T., Morishima, K., Cai, T., Iwatsubo, T., & Tomita, T. (2015). cooperative roles of hydrophilic loop 1 and the C-terminus of presenilin 1 in the substrate-gating mechanism of γ -secretase. *Journal of Neuroscience*, *35*, 2646–2656.
- Takahashi, K., Tanabe, K., Ohnuki, M., Narita, M., Ichisaka, T., Tomoda, K., & Yamanaka, S. (2007). Induction of pluripotent stem cells from adult human fibroblasts by defined factors. *Cell*, *131*, 861–872. <https://doi.org/10.1016/j.cell.2007.11.019>.
- Takami, M., Nagashima, Y., Sano, Y., Ishihara, S., Morishima-Kawashima, M., Funamoto, S., & Ihara, Y. (2009). Gamma-secretase: Successive tripeptide and tetrapeptide release from the transmembrane domain of -carboxyl terminal fragment. *Journal of Neuroscience*, *29*, 13042–13052.



- Takuma, H., Arawaka, S., & Mori, H. (2003). Isoforms changes of tau protein during development in various species. *Developmental Brain Research*, 142(2), 121–127. [https://doi.org/10.1016/S0165-3806\(03\)00056-7](https://doi.org/10.1016/S0165-3806(03)00056-7).
- Togo, T., Sahara, N., Yen, S. H., Cookson, N., Ishizawa, T., Hutton, M., Silva, R., De, L. A., & Dickson, D. W. (2002). Argrophilic grain disease is a sporadic 4-repeat tauopathy. *Journal of Neuropathology & Experimental Neurology*, 61(6), 547–556.
- Vassar, R., Bennett, B. D., Babu-Khan, S., Kahn, S., Mendiaz, E. A., Denis, P., Teplow, D. B., Ross, S., Amarante, P., Loeloff, R., Luo, Y. I., Fisher, S., Fuller, J., Edenson, S., Lile, J., Jarosinski, M. A., Biere, A. L., Curran, E., Burgess, T., ... Citron, M. (1999). β -Secretase cleavage of Alzheimer's amyloid precursor protein by the transmembrane aspartic protease BACE. *Science*, 286, 735–741.
- Vossel, K. A., Zhang, K., Brodbeck, J., Daub, A. C., Sharma, P., Finkbeiner, S., Cui, B., & Mucke, L. (2010). Tau reduction prevents Abeta-induced defects in axonal transport. *Science*, 330, 198.
- Warmack, R. A., Boyer, D. R., Zee, C., Te, R. L. S., Sawaya, M. R., Cascio, D., Gonen, T., Eisenberg, D. S., & Clarke, S. G. (2019). Structure of amyloid- β (20–34) with Alzheimer's-associated isomerization at Asp23 reveals a distinct protofilament interface. *Nature Communications*, 10, 1–12.
- Wesseling, H., Mair, W., Kumar, M., Schlaffner, C. N., Tang, S., Beerepoot, P., Fatou, B., Guise, A. J., Cheng, L., Takeda, S., Muntel, J., Rotunno, M. S., Dujardin, S., Davies, P., Kosik, K. S., Miller, B. L., Berretta, S., Hedreen, J. C., Grinberg, L. T., ... Steen, J. A. (2020). Tau PTM profiles identify patient heterogeneity and stages of Alzheimer's disease. *Cell*, 183(6), 1699–1713.e13. <https://doi.org/10.1016/j.cell.2020.10.029>.
- Wildburger, N. C., Esparza, T. J., LeDuc, R. D., Fellers, R. T., Thomas, P. M., Cairns, N. J., Kelleher, N. L., Bateman, R. J., & Brody, D. L. (2017). Diversity of amyloid-beta proteoforms in the Alzheimer's disease brain. *Scientific Reports*, 7, 9520.
- Wilhelmus, M. M. M., Jager, M., De, S. A. B., Van Der, L. R. J., & Drukarch, B. (2016). Catalytically active tissue transglutaminase colocalises with A β pathology in Alzheimer's disease mouse models. *Scientific Reports*, 6, 1–12.
- Woodruff, G., Young, J. E., Martinez, F. J., Buen, F., Gore, A., Kinaga, J., Li, Z., Yuan, S. H., Zhang, K., & Goldstein, L. S. B. (2013). The presenilin-1 Δ E9 mutation results in reduced γ -secretase activity, but not total loss of PS1 function, in isogenic human stem cells. *Cell Reports*, 5, 974–985. <https://doi.org/10.1016/j.celrep.2013.10.018>.
- Wray, S., Saxton, M., Anderton, B. H., & Hanger, D. P. (2008). Direct analysis of tau from PSP brain identifies new phosphorylation sites and a major fragment of N-terminally cleaved tau containing four microtubule-binding repeats. *Journal of Neurochemistry*, 105, 2343–2352. <https://doi.org/10.1111/j.1471-4159.2008.05321.x>.
- Yagi, T., Ito, D., Okada, Y., Akamatsu, W., Nihei, Y., Yoshizaki, T., Yamanaka, S., Okano, H., & Suzuki, N. (2011). Modeling familial Alzheimer's disease with induced pluripotent stem cells. *Human Molecular Genetics*, 20, 4530–4539.
- Yamada, T., McGeer, P. L., & McGeer, E. G. (1992). Appearance of paired nucleated, Tau-positive glia in patients with progressive supranuclear palsy brain tissue. *Neuroscience Letters*, 135(1), 99–102.
- Yan, R., Munzner, J. B., Shuck, M. E., & Bienkowski, M. J. (2001). BACE2 functions as an alternative α -Secretase in cells. *Journal of Biological Chemistry*, 276, 34019–34027.
- Yoshida, M. (2014). Astrocytic inclusions in progressive supranuclear palsy and corticobasal degeneration. *Neuropathology*, 34(6), 555–570. <https://doi.org/10.1111/neup.12143>.
- Yu, Y., Run, X., Liang, Z., Li, Y., Liu, F., Liu, Y., Iqbal, K., Grundke-Iqbal, I., & Gong, C.-X. (2009). Developmental regulation of tau phosphorylation, tau kinases, and tau phosphatases. *Journal of Neurochemistry*, 108, 1480–1494. <https://doi.org/10.1111/j.1471-4159.2009.05882.x>.
- Zetterberg, H. (2018). Tauomics and kinetics in human neurons and biological fluids. *Neuron*, 97, 1202–1205. <https://doi.org/10.1016/j.neuron.2018.02.030>.
- Zhao, J., Fu, Y., Yamazaki, Y., Ren, Y., Davis, M. D., Liu, C.-C., Lu, W. et al (2020). APOE4 exacerbates synapse loss and neurodegeneration in Alzheimer's disease patient iPSC-derived cerebral organoids. *Nature Communications*, 11, 5540.

How to cite this article: Arber C, Alatza A, Leckey CA, Paterson RW, Zetterberg H, Wray S. Mass spectrometry analysis of tau and amyloid-beta in iPSC-derived models of Alzheimer's disease and dementia. *J Neurochem*. 2021;00:1–13. <https://doi.org/10.1111/jnc.15315>