



# Minimization of Back-Electron Transfer Enables the Elusive $sp^3$ C–H Functionalization of Secondary Anilines

DOI:

[10.1002/anie.202100051](https://doi.org/10.1002/anie.202100051)

## Document Version

Accepted author manuscript

[Link to publication record in Manchester Research Explorer](#)

## Citation for published version (APA):

Zhao, H., & Leonori, D. (2021). Minimization of Back-Electron Transfer Enables the Elusive  $sp^3$  C–H Functionalization of Secondary Anilines. *Angewandte Chemie. International Edition*. <https://doi.org/10.1002/anie.202100051>

## Published in:

Angewandte Chemie. International Edition

## Citing this paper

Please note that where the full-text provided on Manchester Research Explorer is the Author Accepted Manuscript or Proof version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version.

## General rights

Copyright and moral rights for the publications made accessible in the Research Explorer are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

## Takedown policy

If you believe that this document breaches copyright please refer to the University of Manchester's Takedown Procedures [<http://man.ac.uk/04Y6Bo>] or contact [uml.scholarlycommunications@manchester.ac.uk](mailto:uml.scholarlycommunications@manchester.ac.uk) providing relevant details, so we can investigate your claim.



# Minimization of Back-Electron Transfer Enables the Elusive $sp^3$ C–H Functionalization of Secondary Anilines


Huaibo Zhao<sup>[a]</sup> and Daniele Leonori<sup>\*[a]</sup>

**Abstract:** Anilines are some of the most used class of substrates for application in photoinduced electron transfer. *N,N*-Dialkyl-derivatives enable radical generation  $\alpha$  to the *N*-atom by oxidation followed by deprotonation. This approach is however elusive to monosubstituted anilines owing to fast back-electron transfer (BET). Here we demonstrate that BET can be effectively minimised by using photoredox catalysis in the presence of an exogenous alkylamine. This approach synergistically aids aniline SET oxidation and then accelerates the following deprotonation. In this way, the generation of  $\alpha$ -anilinoalkyl radicals is now possible and these species can be used in a general sense to achieve divergent  $sp^3$  C–H functionalization.

Anilines are one of the most important building blocks in organic chemistry. They are widespread in the core structure of high-value materials like drugs and agrochemicals, and also have a rich chemistry for utilization in both polar and radical strategies.<sup>[1]</sup> Indeed, the electron rich nature of the *N*-atom, combined with the stabilisation offered by the neighbouring aromatic ring, makes anilines powerful electron donors and this has been pivotal for their exploitation in redox-based processes.<sup>[2]</sup> However, *N*-substitution is also known to exert a large difference in the way these substrates behave upon single-electron transfer (SET). Taking *N,N*-Me<sub>2</sub>-aniline **1** as an example, it is remarkable that while  $\alpha$ -anilinoalkyl radical **B** generation by SET oxidation followed by deprotonation (**1** → **A** → **B**) is a standard process in photocatalysis, the translation of the same elementary steps to *N*-Me-aniline **2** (**2** → **C** → **D**) is unprecedented (Scheme 1A).<sup>[3]</sup> This lack of reactivity can be exemplified considering the photooxidative demethylation of anilines, one of the most studied and applied reaction in the field of photoinduced SET.<sup>[2c, 4]</sup> This transformation can be triggered by a broad range of photocatalysts but it always stops upon the removal of the first Me-group (**1** → **2**) and **2** never engages in a following demethylation to the free aniline **3**. In the broader sense it is remarkable that the direct conversion of **2** to **3** is currently beyond the scope not just of photoredox catalysis but also of any redox, thermal and transition metal-mediated approach.<sup>[5]</sup>

Since the redox properties of **1** and **2** are relatively similar and well within the range accessible by excited photocatalysts, the ability of **2** to resist photooxidative demethylation is generally rationalised on the basis of back electron transfer (BET).<sup>[6]</sup> Indeed, pioneering studies from Dinnocenzo and Mariano demonstrated that, upon oxidation, the deprotonation of aromatic aminium radicals **C** is slow ( $k_{\text{obs}} \sim 10^5 \text{ M}^{-1} \text{ s}^{-1}$ ) which leads to re-generation of **2** by BET (Scheme 1B).<sup>[7]</sup> The kinetics of deprotonation can be

favoured by installing electron withdrawing groups onto the aniline aromatic ring but this has the detrimental effect of making the initial SET more difficult therefore accelerating BET.<sup>[8]</sup> Overall, the subtle interplay of the elementary steps of electron transfer and deprotonation has, so far, thwarted the development of methodologies for direct  $sp^3$  C–H functionalization of secondary *N*-alkyl-anilines. Currently, the only feasible way to access and exploit the reactivity of radical **D** is to use either *N*-Ph-glycine **2a** or the  $\alpha$ -silylated derivative **2b**. These two systems have the intrinsic advantage the once SET oxidation occurs, subsequent and diffusion-controlled decarboxylation (**E** → **D**) or desilylation (**F** → **D**) take place ( $k_{\text{obs}} \sim 10^{10} \text{ M}^{-1} \text{ s}^{-1}$ ).<sup>[9]</sup>



Here we demonstrate that BET can be minimised by simply using basic alkylamines to synergistically facilitate aniline SET oxidation and the following deprotonation (Scheme 1C). This approach has enabled effective access to  $\alpha$ -anilinoalkyl radicals which can now be used for the elusive  $\alpha$ -*N*  $sp^3$  C–H functionalisation of monosubstituted anilines. Most notably, this strategy has also

[a] Dr Huaibo Zhao and Prof Daniele Leonori  
Department of Chemistry, University of Manchester  
Oxford Road, Manchester, M13 9PL, UK  
E-mail: [daniele.leonori@manchester.ac.uk](mailto:daniele.leonori@manchester.ac.uk)  
Website: <https://leonoriresearchgroup.com>  
Supporting information for this article is given via a link at the end of the document.

## COMMUNICATION

provided the first catalytic approach to achieve unprecedented *N*-dealkylation.

This work originated by a serendipitous observation made during routine cyclic voltammetry experiments. Specifically, we noticed that upon addition of Et<sub>3</sub>N or piperidine to **2**, new peaks appeared in the voltammogram with, crucially, oxidation potentials lower than then the ones of the isolated species (Scheme 2A).<sup>[10]</sup> A similar outcome, albeit less dramatic, was also observed upon performing Stern-Volmer luminescence quenching studies using the photoredox catalyst Ir(dtbbpy)(ppy)<sub>2</sub>PF<sub>6</sub>.<sup>[10]</sup> These unforeseen results led us to consider the formation of a H-bonded complex between **2** and the two amines (e.g. Ph(Me)N–H⋯NEt<sub>3</sub>) which should increase electron density on the aniline *N*-atom. This interaction, would lower the aniline oxidation potential and also increase the rate of quenching of the \*[Ir] photocatalyst. At this stage, we propose this effect might be perceived as an attenuated version of what usually observed in proton-coupled electron transfers (PCETs) between amides and phosphate bases.<sup>[11]</sup> We were therefore intrigued by the possibility that this pre-association upon facilitating SET oxidation could also be used to synergistically aid a following deprotonation that, circumventing BET, would provide access to the  $\alpha$ -anilinoalkyl radical **D**.


In approaching this blueprint for  $\alpha$ -anilinoalkyl radical generation, we envisaged its utilization to target the still elusive demethylation of *N*-methyl anilines (Scheme 2B). Specifically, we proposed the development of a dual photoredox-cobalt<sup>[12]</sup> catalytic system where a visible light-excited [Ir(III)] photocatalyst would, helped by the amine, generate **D** by oxidation and deprotonation of **2** while minimizing BET.<sup>[13]</sup> As **D** is a nucleophilic radical (local electrophilicity index,<sup>[14]</sup>  $\omega^+_{rc} = 0.36$ ), we speculated it could react with a [Co(II)] co-catalyst and undergo dehydrogenation to give a [Co(III)]–H species and imine **4**. A following hydrolysis would provide the desired demethylated product **3**. The pathway for dehydrogenation can be rationalised considering the following two mechanistic options. (1) Radical capture of **D** by [Co(II)] could generate an alkyl-[Co(III)] species followed by  $\beta$ -hydride-type elimination<sup>[15]</sup> across the N–H bond. (2) Since **D** is also an electron rich radical (IP = 6.1 V), an alternative process would involve direct SET between **D** and [Co(II)] to give, via a [Co(I)],<sup>[16]</sup> **4** and [Co(III)]–H. While it is difficult at this stage to distinguish between the two mechanistic options, both of them ultimately result in the formation of the [Co(III)]–H. This species would then evolve H<sub>2</sub> by reaction with a protic source,<sup>[17]</sup> with the resulting electron poor [Co(III)] closing both photoredox and cobalt cycles by SET with [Ir(II)].<sup>[18]</sup>

To put this proposal into practice, we first evaluated the demethylation of **2** using both Et<sub>3</sub>N and piperidine (Scheme 2C). While in their absence no reactivity could be observed,<sup>[10]</sup> a survey of reaction conditions revealed an optimal process that involved the use of the Ir(dtbbpy)(ppy)<sub>2</sub>PF<sub>6</sub><sup>[19]</sup> photocatalyst and the cobaloxime co-catalyst Co(dmgH)(dmgH<sub>2</sub>)Cl<sub>2</sub> in CH<sub>3</sub>CN solvent under blue light irradiation.<sup>[20]</sup> Under these mild conditions **3** was formed in high yields. Crude reaction analysis enabled the detection of formaldehyde, indicative of the generation of **4**, as well as H<sub>2</sub>, which supports the interplay of a [Co(II)/(III)] cycle.<sup>[10]</sup> The scope of the demethylation process was initially evaluated by using mono-substituted *N*-methyl anilines. *para*-Substituted aryl groups displaying electron-rich (**5**, **6**), electron-deficient (**7**, **8**) and

synthetically useful functionalities (**9**–**11**) provided the desired products in good to high yields. The ability to engage substrates containing electron poor arenes is remarkable since BET from their corresponding aminium radicals is accelerated.<sup>[8, 21]</sup> *meta*-Substitution was evaluated next and electronically different functionalities were tolerated albeit in lower yield in the case of electron poor systems (**12**–**15**). We then trialled *ortho*-substituted aromatics and also in this case very high yields were obtained across a broad range of derivatives (**16**–**22**). In this case, the steric hinderance provided by the substituent decreases the stabilisation of the aminium radical and therefore enhances the rate of deprotonation which cooperates to minimise BET.<sup>[8]</sup> This effect is also operating in the demethylations leading to **23**–**25** which were quantitative. The chemistry was amenable to a less activated dichloro aniline (**26**) and found compatible to other aromatics like *N*-methylnaphthalen-1-amine (**27**), several aminopyridines (**28**–**30**) and 5-amino-benzothiophene (**31**). In terms of limitations, we did not succeed in engaging heteroaromatic anilines located on electron rich 5-membered ring systems (e.g. **32**).

Having developed the first catalytic approach for the demethylation of secondary anilines, we were keen to understand if this dual photoredox-cobalt approach could be used in a general sense as a *N*-dealkylation platform. In this case, to evaluate the impact of the alkyl chain we initially looked at a series of *N*-Ph derivatives. Pleasingly, the process displayed broad scope enabling the removal of a benzyl group (**33**) as well as challenging primary (**34** and **35**) and secondary (**36** and **37**) alkyl chains. As an  $\alpha$ -anilinoalkyl radical cannot be formed in the case of a *t*-Bu substituent, **38** resulted in complete starting material recovery which agrees with our mechanistic analysis. The chemistry was also expanded to the removal of functionalised alkyl chains containing inductively electron withdrawing substituents (**39** and **40**) and could also be used to fully dealkylate an ethane-1,2-diamine derivative (**41**). In line with our mechanistic analysis, dealkylation of *N*-Me-*N*-Bn-amine **42** was not possible and this substrate was recovered at the end of the reaction. In this case, Stern-Volmer and cyclic voltammetry analysis revealed no interaction with Et<sub>3</sub>N or piperidine, which does not enable to overcome BET. The approach was also demonstrated on a series of more complex derivatives (**43**–**45**) which included the blockbuster drug primaquine. We then benchmarked the strategy on the demethylation of **1**. While all previous approaches stop after the removal of the first methyl group,<sup>[3,4]</sup> ours enables, for the first time, complete dealkylation to **3** in good yield by simply extending the reaction time. Finally, we were able to demonstrate the orthogonality that this strategy might offer with respect to standard hydrogenation protocols. Taking *N*-Me-*N*-Bn-aniline **46**, all reported deprotections enable debenzilation (**46** → **2**) while this approach provides a switch in selectivity and targets the removal of the Me-group (**46** → **33**).<sup>[22]</sup>

The chemistry described here demonstrates that  $\alpha$ -anilinoalkyl radicals can be conveniently accessed under mild conditions from the corresponding secondary anilines. Since previous exploitation of these open shell intermediates in synthetic chemistry required the preparation and use of *N*-aryl glycines or  $\alpha$ -silyl derivatives (see also Scheme 1B),<sup>[23]</sup> this approach has the potential to facilitate divergent sp<sup>3</sup> C–H functionalization.



**Scheme 2.** (A) Cyclic voltammetry and Stern-Volmer studies. (B) Proposed mechanism for aniline demethylation. (C) Scope of the process. <sup>a</sup> The cyclic voltammetry data for Et<sub>3</sub>N is in the SI.  $\beta$ -HE =  $\beta$ -hydride elimination-type process.


A relevant application would involve application in radical–polar crossover<sup>[24]</sup> reactions involving acrylates for the construction of poly-substituted pyrrolidones (Scheme 3A).<sup>[25]</sup> This would exploit the intrinsic vicinal radical–ionic di-nucleophilic nature of **D** and constitute a novel [3+2]-like retrosynthetic disconnection alternative to more classical approaches based on azomethine ylide chemistry.<sup>[26]</sup> Pleasingly, the realization of this cascade process was achieved by employing conditions similar to the ones presented above but, in this case, quinuclidine gave slightly increased yields over Et<sub>3</sub>N and piperidine (Scheme 3B).<sup>[10, 27]</sup> Scope investigation started by using **2** as the aniline and a range of different acrylates. This enabled the preparation of several C3- (**48–51**) and C4-substituted derivatives (**52–54**) as well as products containing a 4-*gem*-dimethyl group (**55**). The process tolerated valuable functionalities like free alcohol (**49** and **54**), fluorine (**50**) and acetamide (**51**). We also demonstrated reactivity

with tri-substituted acrylates that led to challenging 3,4-disubstituted pyrrolidones (**56–59**) in good yields. We then evaluated the range of anilines compatible as modification of the *N*-substitution would enable introduction of alkyl/aryl substituents at the 5-position – a key feature of many bioactive pyrrolidone- and pyrrolidine-based natural products. Pleasingly, the use of *N*-Et- and *N*-Bn-anilines resulted in the high-yielding formation of 5-Me and -Ph derivatives (**60** and **61**). Other *N*-alkylanilines containing several functionalities were tolerated and these include nitrile (**65**), free and TBS-protected hydroxyl (**66** and **67**), alkene (**68**) and, by using a protected  $\beta$ -alanine, also ester (**66**). At the moment, this reactivity is only feasible on primary *N*-alkyl chains as efforts towards the inclusion of *N*-*i*-Pr-aniline (**70**) proved unsuccessful.





## COMMUNICATION

## A) Radical-to-polar cross-over approach for pyrrolidinone assembly



## B) Scope of the process

C) Further application in photoredox catalysis for divergent sp<sup>3</sup> C–H functionalization

**Scheme 3.** (A) Utilization of  $\alpha$ -anilinoalkyl radicals in radical–polar cross-over reactions for pyrrolidinone assembly. (B) Scope of the process. (C) Extension of  $\alpha$ -anilinoalkyl radical reactivity in other photoredox mediated transformations and current limitations. <sup>a</sup> Yield using benzyl acrylate. EWG = electron-withdrawing group.

The scope of anilines amenable to this cascade reactivity was further extended by *N*-aryllating the neurotransmitters histamine and tryptamine and the alkaloid (–)-*cis*-myrtylamine that provided 71–73 in good yields. These examples demonstrate that this strategy can be used to install a pyrrolidinone system across the linear structure of bio-active molecules. This can be used to

prepare structurally rigid analogues with potential application in bio-organic and medicinal chemistry.

As a finale effort, we question if this approach for  $\alpha$ -anilinoalkyl radical generation could be used in a general sense within the remit of photoredox catalysis. Scheme 3C shows successful application in  $\alpha$ -alkylation of anilines via Giese addition to  $\alpha,\beta$ -unsaturated nitrile (74), amide (75), 3-vinylpyridine (76) and

ketones (**77** and **78**).  $\alpha$ -Aryl amines are valuable synthetic intermediates but standard procedures for  $\alpha$ -N  $sp^3$  C–H arylation generally require the use of carbamates/amides or tertiary amines and H-atom abstraction as radical-generating step.<sup>[28]</sup> In this case, treatment of several *N*-alkyl anilines with 1,4-dicyanobenzene resulted in the formation of **79–84** in good yield. This demonstrates that these conditions are compatible for radical-radical couplings with persistent aromatic radical anions.<sup>[29]</sup> Finally, we also succeed in performing a reaction with the *N*-Ph-imine of benzaldehyde, which furnished 1,2-diamine **85** where both amines can be further functionalised.

The results reported here demonstrate that secondary *N*-alkyl anilines can be converted into the corresponding  $\alpha$ -anilinoalkyl radicals using photoredox catalysis. This reactivity is possible through the minimization of back-electron transfer and by-passes the current requirement for  $\alpha$ -N pre-functionalized aniline substrates. The ability to engage the  $\alpha$ -anilinoalkyl radicals in a broad array of C–C bond-forming processes has enabled divergent  $sp^3$  C–H functionalization while, dual catalysis with cobalt catalysis has provided the first catalytic approach for secondary aniline dealkylation.

## Acknowledgements

D.L. thanks EPSRC for a Fellowship (EP/P004997/1), and the European Research Council for a research grant (758427). HB thanks the Marie Curie actions for a Fellowship (840318).

**Keywords:** photoredox catalysis • C–H functionalization • electron transfer • late-stage modification

[1] (a) A. Ricci, **2008**, Wiley-VCH, Weinheim, Germany; (b) in *Science of Synthesis: Houben–Weyl Methods of Molecular Transformations*, Vol. 31b, 1st Edition ed. (Eds.: C. A. Ramsden, D. Bellus), Georg Thieme Verlag, Stuttgart, **2007**.

[2] Selected examples: (a) S. Andreades, E. W. Zahnow, *J. Am. Chem. Soc.* **1969**, *91*, 4181–4190; (b) M. Ohashi, S. Suwa, K. Tsujimoto, *J. Chem. Soc., Chem. Commun.* **1977**, 348–349; (c) G. Pandey, K. Sudha Rani, U. T. Bhalerao, *Tetrahedron Lett.* **1990**, *31*, 1199–1202; (d) G. Pandey, K. S. Rani, G. Lakshmaiah, *Tetrahedron Lett.* **1992**, *33*, 5107–5110; (e) S. Araneo, F. Fontana, F. Minisci, F. Recupero, A. Serri, *Tetrahedron Lett.* **1995**, *36*, 4307–4310; (f) S. Bertrand, N. Hoffmann, J.-P. Pete, V. Bulach, *Chem. Commun.* **1999**, 2291–2292; (g) Z. Li, C.-J. Li, *J. Am. Chem. Soc.* **2004**, *126*, 11810–11811; (h) Z. Li, R. Yu, H. Li, *Angew. Chem., Int. Ed.* **2008**, *47*, 7497–7500; (i) C. M. R. Volla, P. Vogel, *Org. Lett.* **2009**, *11*, 1701–1704; (j) A. G. Condie, J. C. González-Gómez, C. R. J. Stephenson, *J. Am. Chem. Soc.* **2010**, *132*, 1464–1465; (k) A. McNally, C. K. Prier, D. W. C. MacMillan, *Science* **2011**, *334*, 1114 (l) M. Nishino, K. Hirano, T. Satoh, M. Miura, *J. Org. Chem.* **2011**, *76*, 6447–6451; (m) J. W. Beatty, C. R. J. Stephenson, *Acc. Chem. Res.* **2015**, *48*, 1474–1484; (n) K. Nakajima, Y. Miyake, Y. Nishibayashi, *Acc. Chem. Res.* **2016**, *49*, 1946–1956; (o) R. A. Aycock, C. J. Pratt, N. T. Jui, *ACS Catalysis* **2018**, *8*, 9115–9119.

[3] For a recent example of reactivity of a NHMe on guanidine residues in DNA using photoredox catalysis, see: M. Nappi, A. Hofer, S. Balasubramanian, M. J. Gaunt, *J. Am. Chem. Soc.* **2020**, *142*, 21484–21492.

[4] Selected examples: (a) Q.-Y. Chen, Z.-T. Li, *J. Fluor. Chem.* **1994**, *66*, 59–62; (b) M. Rueping, C. Vila, A. Szadkowska, R. M. Koenigs, J. Fronert, *ACS Catal.* **2012**, *2*, 2810–2815.

[5] (a) D. H. Hunter, D. H. R. Barton, W. J. Motherwell, *Tetrahedron Lett.* **1984**, *25*, 603–606; (b) M. Sako, K. Shimada, K. Hirota, Y. Maki, *J. Am. Chem. Soc.* **1986**, *108*, 6039–6041; (c) A. Sobkowiak, D. T. Sawyer, *J. Am. Chem. Soc.* **1991**,

*113*, 9520–9523; (d) T. Rosenau, A. Hofinger, A. Potthast, P. Kosma, *Org. Lett.* **2004**, *6*, 541–544.

[6] (a) S. Tazuke, N. Kitamura, Y. Kawanishi, *J. Photochem.* **1985**, *29*, 123–138; (b) N. Kitamura, R. Obata, H. B. Kim, S. Tazuke, *J. Phys. Chem.* **1987**, *91*, 2033–2035; (c) C. Zou, J. B. Miers, R. M. Ballew, D. D. Dlott, G. B. Schuster, *J. Am. Chem. Soc.* **1991**, *113*, 7823–7825; (d) S. Narra, Y. Nishimura, H. A. Witek, S. Shigeto, *ChemPhysChem* **2014**, *15*, 2945–2950.

[7] J. P. Dinnocenzo, T. E. Banach, *J. Am. Chem. Soc.* **1989**, *111*, 8646–8653.

[8] G. W. Dombrowski, J. P. Dinnocenzo, P. A. Zielinski, S. Farid, Z. M. Wosinska, I. R. Gould, *J. Org. Chem.* **2005**, *70*, 3791–3800.

[9] (a) X. Zhang, S.-R. Yeh, S. Hong, M. Freccero, A. Albini, D. E. Falvey, P. S. Mariano, *J. Am. Chem. Soc.* **1994**, *116*, 4211–4220; (b) E. R. Gaillard, D. G. Whitten, *Acc. Chem. Res.* **1996**, *29*, 292–297; (c) Z. Su, D. E. Falvey, U. C. Yoon, P. S. Mariano, *J. Am. Chem. Soc.* **1997**, *119*, 5261–5262; (d) Z. Su, P. S. Mariano, D. E. Falvey, U. C. Yoon, S. W. Oh, *J. Am. Chem. Soc.* **1998**, *120*, 10676–10686.

[10] See SI for more information.

[11] L. Q. Nguyen, R. R. Knowles, *ACS Catal.* **2016**, *6*, 2894–2903.

[12] K. C. Cartwright, A. M. Davies, J. A. Tunge, *Eur. J. Org. Chem.* **2020**, 1245–1258.

[13] Upon SET oxidation of **2**, the alkylamine could also deprotonate the N–H to give an aminyl radical. This species can then be converted back to **2** by polarity matched HAT on CH<sub>3</sub>CN solvent.

[14] F. D. Vleschouwer, V. V. Speybroeck, M. Waroquier, P. Geerlings, F. D. Proft, *Org. Lett.* **2007**, *9*, 2721–2724.

[15] A. A. Gridnev, S. D. Ittel, *Chem. Rev.* **2001**, *101*, 3611–3660.

[16] S. M. Chemaly, J. M. Pratt, *J. Chem. Soc., Dalton Trans.* **1984**, 595–599.

[17] J. L. Dempsey, B. S. Brunschwig, J. R. Winkler, H. B. Gray, *Acc. Chem. Res.* **2009**, *42*, 1995–2004.

[18] (a) D. Dodd, M. D. Johnson, *J. Organomet. Chem.* **1973**, *52*, 1–232; (b) X. Sun, J. Chen, T. Ritter, *Nat. Chem.* **2018**, *10*, 1229–1233; (c) S. U. Dighe, F. Juliá, A. Luridiana, J. J. Douglas, D. Leonori, *Nature* **2020**, *584*, 75–81.

[19] BET minimization is mostly possible on triplet excited state photocatalysts as it would require a change in spin. Indeed, attempts to extend this chemistry to photoredox catalysts active through the singlet excited state like Fukuzumi's acridinium, has proven not possible despite its stronger oxidation power.<sup>[10]</sup>

[20] Inorganic bases have also been evaluated and they provided the desired product in generally much lower yield.<sup>[10]</sup> In this case, cyclic voltammetry analysis was hampered by their low solubility.

[21] The intensity of peak resulting from the interaction of the alkylamine with the *N*-Me anilines increases with the presence of electron withdrawing groups onto the aromatic ring as this enhances the N–H acidity.<sup>[10]</sup>

[22] In this case, careful monitoring of the reaction was required to stop it after the demethylation. When the process was left for longer time, the following debenzoylation took place.

[23] (a) V. Corcé, L.-M. Chamoreau, E. Derat, J.-P. Goddard, C. Ollivier, L. Fensterbank, *Angew. Chem., Int. Ed.* **2015**, *54*, 11414–11418; (b) C. Lévêque, L. Chenneberg, V. Corcé, C. Ollivier, L. Fensterbank, *Chem. Commun.* **2016**, 52, 9877–9880.

[24] L. Pitzer, J. L. Schwarz, F. Glorius, *Chem. Sci.* **2019**, *10*, 8285–8291.

[25] For a recent radical–polar crossover method for pyrrolidone assembly using HAT on free tertiary alkylamines (alkyl–NH<sub>2</sub>), see: A. S. H. Ryder, W. B. Cunningham, G. Ballantyne, T. Mules, A. G. Kinsella, J. Turner-Dore, C. M. Alder, L. J. Edwards, B. S. J. McKay, M. N. Grayson, A. J. Cresswell, *Angew. Chem., Int. Ed.* **2020**, *59*, 14986–14991.

[26] I. Coldham, R. Hufton, *Chemical Reviews* **2005**, *105*, 2765–2810.

[27] A mechanism based on SET oxidation of quinuclidine followed by HAT on **2** should not be feasible as the oxidation of 2•••quinuclidine is much faster as determined by luminescence quenching (Stern–Volmer) studies.<sup>[10]</sup>


[28] A. McNally, C. K. Prier, D. W. C. MacMillan, *Science* **2011**, *334*, 1114–1117.

[29] A. Albini, E. Fasani, M. Mella, in *Photoinduced Electron Transfer V* (Ed.: J. Mattay), Springer Berlin Heidelberg, Berlin, Heidelberg, **1993**, pp. 143–173.

Entry for the Table of Contents (Please choose one layout)

Layout 2:

## COMMUNICATION



Huaibo Zhao and Daniele Leonori\*

Page No. – Page No.

**Minimization of Back-Electron Transfer Enables the Elusive  $sp^3$  C–H Functionalization of Secondary Anilines**

Divergent radical  $sp^3$  C–H functionalization of *N*-alkyl anilines has been achieved using a photoredox strategy. This approach enables to overcome back-electron transfer and therefore enable direct access to previously elusive  $\alpha$ -anilinoalkyl radicals.