



# The proficiency of the original host species determines community-level plasmid dynamics

**DOI:**

[10.1093/femsec/fiab026](https://doi.org/10.1093/femsec/fiab026)

**Document Version**

Accepted author manuscript

[Link to publication record in Manchester Research Explorer](#)

**Citation for published version (APA):**

Kottara, A., Hall, J. P. J., & Brockhurst, M. A. (2021). The proficiency of the original host species determines community-level plasmid dynamics. *FEMS Microbiology Ecology*, 97(4), [fiab026].  
<https://doi.org/10.1093/femsec/fiab026>

**Published in:**

FEMS Microbiology Ecology

**Citing this paper**

Please note that where the full-text provided on Manchester Research Explorer is the Author Accepted Manuscript or Proof version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version.

**General rights**

Copyright and moral rights for the publications made accessible in the Research Explorer are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

**Takedown policy**

If you believe that this document breaches copyright please refer to the University of Manchester's Takedown Procedures [<http://man.ac.uk/04Y6Bo>] or contact [uml.scholarlycommunications@manchester.ac.uk](mailto:uml.scholarlycommunications@manchester.ac.uk) providing relevant details, so we can investigate your claim.



1 **The proficiency of the original host species determines community-level plasmid dynamics**

2 **Anastasia Kottara<sup>1‡</sup>, James P.J. Hall<sup>2</sup>, Michael A. Brockhurst<sup>3\*</sup>**

3 <sup>1</sup>Department of Animal and Plant Sciences, University of Sheffield, Sheffield, S10 2TN, UK

4 <sup>2</sup>Department of Evolution, Ecology and Behaviour, Institute of Infection, Veterinary and Ecological  
5 Sciences, University of Liverpool, Liverpool, L69 7ZB, UK

6 <sup>3</sup>Faculty of Biology, Medicine and Health, University of Manchester, Manchester, M13 9PT, UK.

7 <sup>‡</sup>Current address: Instituto Gulbenkian de Ciência, Rua Quinta Grande 6, Oeiras 2780-156, Portugal

8

9

10 Corresponding author: Michael Brockhurst, Division of Evolution and Genomic Sciences, Faculty of  
11 Biology, Medicine and Health, University of Manchester, Manchester, M13 9PT, UK. Email-

12 [michael.brockhurst@manchester.ac.uk](mailto:michael.brockhurst@manchester.ac.uk)

13

14 Keywords: horizontal gene transfer, mobile genetic elements, conjugative plasmids, bacterial  
15 communities, plasmid transfer.

16

17

18

19

20

21

22

23

24

25

26 **ABSTRACT**

27 Plasmids are common in natural bacterial communities, facilitating bacterial evolution via horizontal  
28 gene transfer. Bacterial species vary in their proficiency to host plasmids: Whereas plasmids are stably  
29 maintained in some species regardless of selection for plasmid-encoded genes, in other species, even  
30 beneficial plasmids are rapidly lost. It is, however, unclear how this variation in host proficiency affects  
31 plasmid persistence in communities. Here, we test this using multispecies bacterial soil communities  
32 comprising species varying in their proficiency to host a large conjugative mercury resistance plasmid,  
33 pQBR103. The plasmid reached higher community-level abundance where beneficial and when  
34 introduced to the community in a more proficient host species. Proficient plasmid host species were  
35 also better able to disseminate the plasmid to a wider diversity of host species. These findings suggest  
36 that the dynamics of plasmids in natural bacterial communities depend not only upon the plasmid's  
37 attributes and the selective environment, but also upon the proficiency of their host species.

38

39

40

41

42

43

44

45

46

47

48

49

50

51

## 52 INTRODUCTION

53 Mobile genetic elements (MGEs) like plasmids, temperate bacteriophages, and transposons, are  
54 important agents of horizontal gene transfer (HGT) driving the diversification of bacterial genomes  
55 (Frost *et al.* 2005; Hall, Brockhurst and Harrison 2017a; Brockhurst *et al.* 2019). Conjugative plasmids  
56 contain genes encoding core plasmid functions – including their own propagation, replication, stability  
57 and transfer – along with accessory genes that encode traits like antibiotic and metal resistance  
58 (Norman, Hansen and Sørensen 2009). While the plasmid’s accessory genes may directly benefit the  
59 host cell by providing them with new ecological functions, plasmid acquisition can impose a heavy  
60 fitness burden on the host cell (Baltrus 2013; San Millan and Maclean 2017). Mathematical models of  
61 plasmid population dynamics suggest that the plasmid cost, conjugation rate, segregation rate, and the  
62 strength of positive selection are key parameters determining plasmid survival in bacterial populations  
63 (Stewart and Levin 1977; Levin, Stewart and Rice 1979; Simonsen *et al.* 1990; Bergstrom, Lipsitch and  
64 Levin 2000).

65 Plasmids are expected to spread under positive selection for their encoded accessory genes (San  
66 Millan *et al.* 2014; Harrison *et al.* 2015), however, because accessory genes can be captured by the  
67 bacterial chromosome rendering the plasmid redundant, positive selection does not guarantee the long-  
68 term survival of plasmids (Bergstrom, Lipsitch and Levin 2000). Meanwhile, in the absence of positive  
69 selection, plasmids are expected to decline in frequency due to purifying selection because the benefits  
70 of accessory genes do not outweigh the costs of plasmid carriage (Bergstrom, Lipsitch and Levin 2000).  
71 Since rates of conjugation appear to often be too low for plasmids to persist as infectious elements  
72 (although see: Lopatkin *et al.* (2017) and Stevenson *et al.* (2017)), it has been argued that the widespread  
73 distribution of plasmids is paradoxical (the plasmid paradox: Harrison and Brockhurst (2012)). Yet,  
74 plasmids have been found to stably persist in natural bacterial communities in the absence of measurable  
75 positive selection, where the factors allowing plasmid stability are puzzling (Heuer and Smalla 2012).

76 Most studies of plasmid dynamics focus on a single-host species, whereas, in natural bacterial  
77 communities, many potential host species co-exist, potentially broadening the range of conditions under  
78 which plasmids can survive. This limitation of current understanding is particularly interesting

79 considering that several studies have shown that plasmids are not equally stable across host species (De  
80 Gelder *et al.* 2007; Kottara *et al.* 2018; Sakuda *et al.* 2018). For example, while the mercury resistance  
81 plasmid pQBR103 was highly stable for >400 generations with or without mercury selection in *P.*  
82 *fluorescens* and *P. savastanoi*, it was unstable to varying degrees in *P. stutzeri* (generally lost within  
83 ~100-400 generations), *P. aeruginosa* and *P. putida* (<6 generations) even with strong mercury  
84 selection (Kottara *et al.* 2018). Here, we define host proficiency as the ability of a bacterial host species  
85 to stably maintain a conjugative plasmid within its population.

86 Hall *et al.* (2016) showed, by tracking the dynamics of the mercury resistance plasmid pQBR57  
87 in a two-species soil community of *P. fluorescens* and *P. putida*, that between-species transfer of the  
88 plasmid from a proficient host, *P. fluorescens*, to an unstable host, *P. putida*, allowed the plasmid to  
89 persist in *P. putida* both with and without mercury selection. This finding suggests that the dynamics  
90 of a plasmid in a bacterial community is likely to depend on the proficiency of the plasmid host species  
91 to stably maintain the plasmid. This leads to the prediction that, at the community-level, plasmid  
92 abundance will be higher in communities where it is carried by a proficient original plasmid host, since  
93 this species will both be able to maintain the plasmid in its own population, and then disseminate the  
94 plasmid to other species in the community.

95 To test this prediction, we tracked the dynamics of pQBR103 in a three-species community of  
96 *P. fluorescens*, *P. stutzeri* and *P. putida* with and without mercury selection. We varied which of the  
97 species carried the plasmid at the start of the experiment. We hypothesised that the community-level  
98 plasmid abundance would vary according to the proficiency of the original plasmid host species to act  
99 as hosts to pQBR103, which varies hierarchically – *P. fluorescens* > *P. stutzeri* > *P. putida* (Kottara *et*  
100 *al.* 2018). Replicate communities were propagated in effectively sterile potting soil (artificial soil  
101 microcosms), which provide spatial structure and a low resource environment that more closely  
102 resemble the natural physical and chemical conditions in soil and promote the stable co-existence of  
103 multiple bacterial species (Gómez and Buckling 2011; Heuer and Smalla 2012; Hall *et al.* 2015; Hall  
104 *et al.* 2016).

105

## 106 MATERIALS AND METHODS

### 107 Bacterial strains and plasmid

108 Three *Pseudomonas* species – *P. fluorescens* SBW25 (Rainey, Bailey and Thompson 1994), *P. stutzeri*  
109 JM300 (DSM 10701) (Busquets *et al.* 2012) and *P. putida* KT2440 (Bagdasarian *et al.* 1981) – were  
110 utilised in this study. *Pseudomonas* species were labelled by directed insertion of either streptomycin  
111 ( $\text{Sm}^{\text{R}}$ ) or gentamicin resistance ( $\text{Gm}^{\text{R}}$ ) marker using the mini-Tn7 transposon system (Lambertsen,  
112 Sternberg and Molin 2004). The plasmid used in this study, pQBR103 is a large conjugative plasmid  
113 (425 kb) that confers mercury resistance via a *mer* operon encoded on a Tn5042 transposon (Lilley *et*  
114 *al.* 1996; Tett *et al.* 2007). pQBR103 plasmid is part of a group of 136 plasmids that were isolated from  
115 the bacterial community inhabiting the sugar beet rhizosphere and phyllosphere during a long-term field  
116 experiment (Lilley *et al.* 1996). pQBR103 was exogenously acquired by conjugation into labelled strain  
117 of *P. fluorescens* that was introduced onto the naturally occurring bacterial community colonising the  
118 sugar beet rhizosphere with the primary plasmid-host remaining unknown (Lilley *et al.* 1996). To obtain  
119 the initial plasmid-bearing clones of each host species to start the selection experiment, pQBR103  
120 plasmid was conjugated into *P. stutzeri*  $\text{Gm}^{\text{R}}$ , *P. putida*  $\text{Sm}^{\text{R}}$  and *P. fluorescens*  $\text{Sm}^{\text{R}}\text{lacZ}$  from the  
121 plasmid-bearing *P. fluorescens* SBW25  $\text{Sm}^{\text{R}}$  or  $\text{Gm}^{\text{R}}$  stocks. Plasmid conjugation was performed by  
122 mixing 1:1 each of the plasmid-free with the plasmid-bearing strains, incubating for 48 h and spreading  
123 on King's B growth (KB) agar plates containing  $5 \mu\text{g mL}^{-1}$  gentamicin or  $50 \mu\text{g mL}^{-1}$  streptomycin ( $50$   
124  $\mu\text{g mL}^{-1}$  X-Gal) and  $20 \mu\text{M}$  of mercury(II) chloride to select for transconjugant colonies (Simonsen *et*  
125 *al.* 1990). The conjugation assays were conducted in 6 mL KB medium in 30 mL universal vials  
126 ('microcosms') at  $28^{\circ}\text{C}$  in shaking conditions (180 rpm).

127

### 128 Selection experiment

129 To account for the high segregation rate of the plasmid in *P. putida* KT2440 (Kottara *et al.* 2018) and  
130 to ensure high starting frequencies of plasmid carriage across all the tested bacterial strains, single  
131 colonies of each plasmid-bearing species were reconditioned overnight and then transferred in fresh  
132 media containing mercury. Specifically, individual colonies ( $n=12$ ) of each plasmid-bearing

133 *Pseudomonas* species were picked into separate 6 mL KB microcosms and incubated overnight at 28°C  
134 with shaking 180 rpm after which time 1% of each population was transferred to grow for 24 h in fresh  
135 KB microcosms containing 50 µM of mercury(II) chloride at same temperature and shaking conditions;  
136 this concentration of mercury was used to select for the pQBR103 plasmid based on previous findings  
137 (Kottara *et al.* 2018). Similarly, 24 colonies of each plasmid-free *Pseudomonas* species were each  
138 grown overnight in KB 6 mL microcosms and transferred to grow for 24 h in fresh KB microcosms at  
139 same temperature and shaking conditions.

140

#### 141 *Bacterial communities*

142 We used artificial soil microcosms to evolve three different bacterial communities differing by which  
143 species carried the plasmid at the beginning of the experiment (original plasmid host). To prepare the  
144 artificial soil microcosms, we added 10 g of John Innes No. 2 compost soil in 30 mL universal vials  
145 which we autoclaved twice. By autoclaving the compost soil two times, we established an effectively  
146 sterile micro-environment with the physical and chemical properties of soil which did not contain other  
147 culturable bacteria than our inoculum (Gómez and Buckling 2010; Hall *et al.* 2015; Hall *et al.* 2016).  
148 Three different bacterial communities were then constructed: *P. fluorescens* (carrying pQBR103) with  
149 *P. stutzeri* and *P. putida*; *P. fluorescens* with *P. stutzeri* (pQBR103) and *P. putida*; *P. fluorescens* with  
150 *P. stutzeri* and *P. putida* (pQBR103). Six replicates of each community were grown either without  
151 mercury or with mercury (16 µg g<sup>-1</sup> Hg(II)); this concentration of mercury was used to select for the  
152 pQBR103 plasmid while could allow the survival of the plasmid-free species based on previous findings  
153 (Hall *et al.* 2015). After the addition of Hg(II), each artificial soil microcosm was briefly vortexed to  
154 ensure homogeneous distribution of mercury in soil. Each community had a starting ratio of 1:1:1 of  
155 each *Pseudomonas* species such that the starting frequency of pQBR103 in the community was  
156 approximately 33%. To remove spent media and residual mercury from overnight cultures each  
157 inoculum was briefly vortexed, then centrifuged for 1 min at 10,000 rpm and resuspended in 1 mL M9  
158 salt solution (Sambrook, Fritsch and Maniatis 1989). 100 µL was then inoculated into artificial soil  
159 microcosms, briefly vortexed to ensure spread homogeneity in artificial soil microcosms and incubated  
160 at 28°C at 80% humidity (Hall *et al.* 2016).

161

162 *Serial transfers and bacterial counts*

163 Every 4 days, 10 mL of M9 buffer and 20 glass beads were added to each artificial soil microcosm and  
164 mixed by vortexing for 1 min, and 100  $\mu\text{L}$  of soil wash was transferred to a fresh artificial soil  
165 microcosm as previously described by Hall *et al.* (2016). Bacterial counts for each species were  
166 estimated by plating onto selective media: 50  $\mu\text{g mL}^{-1}$  streptomycin + 50  $\mu\text{g mL}^{-1}$  X-Gal KB agar plates  
167 and 5  $\mu\text{g mL}^{-1}$  gentamicin KB agar plates, each of which was then replica plated onto mercury KB agar  
168 plates (100  $\mu\text{M}$  mercury(II) chloride). The bacterial communities were evolved for 10 transfers (~40  
169 days, estimated to be approximately 70 bacterial generations).

170

171 *Plasmid and mercury resistance transposon screening*

172 Twenty-four mercury-resistant colonies of each *Pseudomonas* species were sampled every 2 transfers  
173 from the mercury containing plates and tested for the presence of the plasmid and mercury resistance  
174 transposon by PCR screening. The PCR used the same sets of primers as previously described [*mer*  
175 operon on the Tn5042 transposon – forward primer: 5'-TGCAAGACACCCCCTATTGGAC-3', reverse  
176 primer: 5'-TTCGGCGACCAGCTTGATGAAC-3' and origin of replication of the plasmid (*oriV*) –  
177 forward primer: 5'-TGCCTAATCGTGTGTAATGTC-3', reverse primer: 5'-  
178 ACTCTGGCCTGCAAGTTTC-3'] (Harrison *et al.* 2015; Kottara *et al.* 2018).

179

180 **Statistics**

181 Statistical analyses were performed using RStudio version 3.2.3 (R Core Team 2013). Shapiro-Wilk  
182 test, normal Q-Q plots, histograms and box-plots were used to examine the normality of the data. We  
183 found that in most cases the data were not normally distributed, and in such cases used a non-parametric  
184 test. Cumulative plasmid abundance in each community over time was estimated as the area under the  
185 curve using the function *auc* of the package 'flux' (Jurasinski, Koebisch and Hagemann 2012).  
186 Community-level plasmid abundances in the plasmid host treatments were compared by using the  
187 Kruskal-Wallis test. To assess the plasmid-dynamics within each species, we compared plasmid



188 frequencies in the plasmid-recipient species population as the area under the curve. The integral  
189 estimates of the plasmid frequency in the recipient species were compared between the mercury  
190 conditions using the Kruskal-Wallis test. To assess the timing of chromosomal acquisition of the  
191 mercury resistance transposon Tn5042 in *P. putida* that differed between the plasmid host treatments,  
192 for each population we recorded the transfer number when we first observed plasmid-free transposon-  
193 containing genotypes of *P. putida*. We compared these values between the plasmid host treatments  
194 using the Kruskal-Wallis test. The species diversity of plasmid-carriers was calculated as the 1 - D  
195 Simpson's Index,  $1 - [\sum (\frac{n_i}{N})^2]$  where,  $n_i$  = the end-point population density of each plasmid-bearer  
196 species in community, and,  $N$  = the end-point population density of all plasmid-bearer species. We  
197 compared diversities between the plasmid host treatments and mercury conditions by using the Kruskal-  
198 Wallis test.

199

## 200 RESULTS

### 201 Original plasmid host species identity affects community-level plasmid abundance

202 The bacterial host species vary in their ability to stably maintain pQBR103 hierarchically as follows:  
203 *P. fluorescens* > *P. stutzeri* > *P. putida* (Kottara *et al.* 2018). We hypothesised therefore that the identity  
204 of the original plasmid host in a community is likely to affect the dynamics of the plasmid-encoded  
205 mercury resistance at the community-level. To test this, we quantified the total plasmid abundance in  
206 each community (Figure 1). Mercury selection increased total plasmid abundance (effect of mercury;  
207  $\chi^2(1, N=24)=17.28, p=3.226e-05$ ) and total plasmid abundance varied with original plasmid host  
208 identity, such that both with (effect of plasmid treatment;  $\chi^2(2, N=18)=11.556, p=0.003$ ) and without  
209 (effect of plasmid treatment;  $\chi^2(2, N=18)=11.474, p=0.003$ ) mercury selection, the total plasmid  
210 abundance was higher when the original plasmid host was *P. fluorescens*. Together these data suggest  
211 that community-level plasmid dynamics are affected by both the positive selection for plasmid-encoded  
212 traits and the identity of the original plasmid host species, being enhanced when plasmids are beneficial  
213 and carried by a proficient plasmid host.

214

### 215 **Species-level plasmid dynamics within communities**

216 To understand how the variation in community-level plasmid abundance was driven by original plasmid  
217 host identity, we next examined the species-level plasmid dynamics in each community. As predicted,  
218 when a proficient plasmid-host – *P. fluorescens* – was the original plasmid host it maintained the  
219 plasmid at high frequency within its population both with and without mercury (Figure 2). We detected  
220 plasmid dissemination from *P. fluorescens* to the other species at higher frequencies under mercury  
221 selection (effect of mercury;  $\chi^2(1, N=24)=4.653$ ,  $p=0.030$ ). This occurred to *P. putida* in all replicates  
222 and to *P. stutzeri* in 2/6 replicates with mercury selection and also to *P. stutzeri* at low levels in some  
223 of the communities without mercury selection. When *P. stutzeri* was the original plasmid host, it also  
224 maintained the plasmid within its own population both with and without mercury, and disseminated  
225 plasmids to the other species at a higher rate with mercury (effect of mercury;  $\chi^2(1, N=24)=11.644$ ,  
226  $p=0.0006$ ) (Figure 3). Variation in total plasmid abundance between replicate communities in this  
227 treatment appear to have been caused by whether or not *P. fluorescens* acquired the plasmid before it  
228 was outcompeted by mercury resistant competitors: where transmission to *P. fluorescens* occurred, total  
229 plasmid abundances were higher (Figure 3). Where *P. putida* was the original plasmid host, it did not  
230 maintain the plasmid within its own population: without mercury, the plasmid was simply lost, whereas,  
231 with mercury, plasmid-bearers were replaced by mutants that had inserted the Tn5042 encoding the *mer*  
232 operon into their chromosome (Figure 4). Chromosomal insertions of the Tn5042 in *P. putida* were  
233 observed in the other plasmid host treatments, but arose much later in these communities where *P.*  
234 *putida* had to acquire the plasmid horizontally from either *P. fluorescens* or *P. stutzeri* (effect of  
235 treatment;  $\chi^2(2, N=18)=10.947$ ,  $p= 0.004$ ). Although *P. putida* eventually lost the plasmid from its  
236 own population, prior to this loss it successfully disseminated the plasmid to *P. fluorescens* in 6/6  
237 replicates and to *P. stutzeri* in 3/6 replicates with mercury selection (Figure 4).

238

### 239 **Diversity of plasmid-carriers in communities**

240 Finally, we tested how the original plasmid host identity affected the diversity of plasmid-carriers at the  
241 end of the experiment. The diversity of plasmid-carriers was affected by both the original plasmid host

242 species identity (effect of plasmid treatment;  $\chi^2(2, N=36)=12.819, p=0.001$ ) and mercury selection  
243 (effect of mercury;  $\chi^2(1, N=36)=6.082, p=0.013$ ) (Figure 5). Without mercury selection the diversity  
244 of plasmid-carriers was highest when *P. stutzeri* was the original plasmid host. Whereas, with mercury  
245 selection, the diversity of plasmid-carriers was higher in communities where *P. fluorescens* or *P.*  
246 *stutzeri* were the original plasmid hosts compared to communities where *P. putida* was the original  
247 plasmid host. Consistent with our data on community-level plasmid abundance, these data show that  
248 the diversity of plasmid-carriers is likely to be higher when plasmids are beneficial and are introduced  
249 to the community by proficient plasmid hosts.

250

## 251 **DISCUSSION**

252 In natural microbial communities, broad host range plasmids are frequently transmitted to diverse host  
253 species thus highlighting the importance of plasmids in HGT and their role in the spread of resistance  
254 genes in the environment (Pukall, Tschäpe and Smalla 1996; Klümper *et al.* 2015). In this study, we  
255 aimed to understand the extent to which plasmid dynamics in a bacterial community are affected by the  
256 original plasmid host species identity within that community. Our findings suggest that plasmid  
257 abundance at the community-level was driven by the identity of the original plasmid host species. We  
258 observed that pQBR103 reached higher community-level abundance when hosted by a proficient  
259 plasmid-host, *P. fluorescens*. Dionisio *et al.* (2002) have previously shown the importance of species  
260 identity in shaping the plasmid dynamics in a community. This was further described by Hall *et al.*  
261 (2016) where a proficient plasmid-host could act as a source of the plasmid for a non-proficient host  
262 species in a two-species soil community. These plasmid dynamics were explained in terms of  
263 conjugative plasmids persisting in the community as infectious agents via interspecies transfer (Bahl,  
264 Hansen and Sørensen 2007). Here, we extend these results to a more complex three-species community,  
265 a different plasmid, and a wider range of plasmid host species and proficiencies.

266 The community-level plasmid abundance also varied according to mercury selection. In  
267 common with previous studies (Cairns *et al.* 2018), plasmids were observed at higher frequencies in  
268 recipient species in the presence versus absence of positive selection. Detecting HGT events is more

269 likely under positive selection, because, while individual conjugation events may be rare, positively  
270 selected horizontally acquired genes will rise to high frequency due to clonal expansion. This has led  
271 to a generally accepted, but probably incorrect view, that HGT is accelerated under positive selection  
272 (Aminov 2011; Fletcher 2015). By contrast, recent experimental data shows that horizontal  
273 transmission plays a more important role in plasmid stability in the absence of positive selection  
274 (Stevenson *et al.* 2017), leading to higher rates of gene mobilisation and transfer in these environments  
275 (Hall *et al.* 2017b). Mercury selection also drove the invasion of *P. putida* mutants that had lost the  
276 plasmid but captured the Tn5042 carrying the mercury resistance operon to the chromosome, an  
277 outcome rarely observed in the other host species. This confirms our previous data that the rate and/or  
278 propensity for transposition of traits from the plasmid to the chromosome is variable among  
279 *Pseudomonas* species (Kottara *et al.* 2018). We show here that the dynamics of this process are affected  
280 by the community context, specifically whether or not *P. putida* was the original plasmid host.  
281 Chromosomal capture of mercury resistance transposon in *P. putida* occurred earlier when it began the  
282 experiment with the plasmid, reflecting that transposition is random mutational event and thus more  
283 likely to occur in larger – plasmid-bearing – populations. Interestingly, however, our data also show  
284 that even low proficiency plasmid hosts, which rapidly capture useful traits and jettison the plasmid,  
285 can act as a source of plasmids for other species in community by transferring the plasmid to more  
286 proficient host species before it is lost.

287         In contrast to the study of Hall *et al.* (2016), which used a highly conjugative plasmid, pQBR57,  
288 the plasmid used here, pQBR103, has >1000-fold lower conjugation rate ( $\gamma$ , the rate parameter for  
289 conjugative plasmid transfer estimated by the end-point method as developed by Simonsen *et al.* (1990);  
290  $\text{Log}_{10}(\gamma) \text{ pQBR103} = \sim -13.8$ ,  $\text{Log}_{10}(\gamma) \text{ pQBR57} = \sim -10.8$ ; Hall *et al.* 2015). While previous studies of  
291 pQBR103 have focused on the importance compensatory evolution in its longer-term stability (Harrison  
292 *et al.* 2015), here we show an effect of between species conjugation increasing the community-level  
293 abundance of the plasmid. The role for interspecific conjugation in pQBR103 stability was most notable  
294 in communities where it was initially carried by a non-proficient original plasmid host, *P. putida*. Here,  
295 while the plasmid started in  $\sim 33\%$  of the population and went extinct in the *P. putida* population, it  
296 survived in the community by horizontal transmission, most commonly into *P. fluorescens*. Through

297 interspecific conjugation, pQBR103 increased the diversity of plasmid-carriers in communities,  
298 especially under mercury selection. However, this effect depended upon the original plasmid host  
299 species identity. Conjugation also depends on the population density, and in this case the higher  
300 population density of *P. fluorescens* could have enabled the plasmid transfer from *P. fluorescens*.  
301 Surprisingly, although with mercury selection more proficient plasmid host species (*i.e.* *P. fluorescens*  
302 and *P. stutzeri*) allowed higher diversities of plasmid-carriers, without mercury it was in communities  
303 where the moderately proficient plasmid host, *P. stutzeri*, was original plasmid host where the highest  
304 plasmid-carrier diversity was observed. This effect is likely to have been caused by the more equitable  
305 distribution of plasmid carriage in these communities, and specifically by higher rates of plasmid  
306 carriage in *P. stutzeri* itself compared to communities where this species had to obtain the plasmid via  
307 conjugation.

308           Soil microbial communities are highly diverse, which is thought to play a key role in their  
309 function (Torsvik and Øvreås 2002) and species diversity has been suggested to play a role in the  
310 dissemination of conjugative plasmids (Dionisio *et al.* 2002). Soil habitats are often characterised as  
311 hot spots for HGT (van Elsas and Bailey 2002; Sørensen *et al.* 2005) due to the spatially structured  
312 nature of such environments (Bahl, Hansen and Sørensen 2007; Fox *et al.* 2008; Røder *et al.* 2013).  
313 Here, we show that the identity of original plasmid host species determines the community-level  
314 abundance of conjugative plasmids in soil bacterial communities. Proficient plasmid hosts better  
315 maintain plasmids within their own population and transmit these plasmids to other species in the  
316 community. This implies that proficient plasmid host species could promote the robustness of  
317 communities by spreading potentially adaptive genes to more diverse species, allowing their survival  
318 upon environmental deterioration in the future.

319

## 320 **ACKNOWLEDGEMENTS**

321 The authors thank T. Daniell and D. Rozen for their comments on a previous version of this work.

322

## 323 **FUNDING**

324 This work was supported by funding from the European Research Council under the European Union's  
325 Seventh Framework Programme awarded to MB [grant number FP7/2007-2013/ERC grant StG-2012-  
326 311490–COEVOCON] and grants to MB and JPJH from the Natural Environment Research Council  
327 NE/R008825/1 and Biotechnology and Biological Sciences Research Council BB/R018154/1.

328

### 329 **AUTHOR CONTRIBUTIONS**

330 AK, JH and MB designed the study; AK performed the experiments and analysed the data; AK and MB  
331 drafted the manuscript.

332

333 **Conflict of interest.** The authors declare that there are no conflicts of interest.

334

### 335 **REFERENCES**

- 336 Aminov RI. Horizontal gene exchange in environmental microbiota. *Front Microbiol* 2011;**2**:158.
- 337 Bagdasarian M, Lurz R, Rückert B *et al.* Specific-purpose plasmid cloning vectors II. Broad host range,  
338 high copy number, RSF 1010-derived vectors, and a host-vector system for gene cloning in  
339 *Pseudomonas*. *Gene* 1981;**16**:237-247.
- 340 Bahl MI, Hansen LH, Sørensen SJ. Impact of conjugal transfer on the stability of IncP-1 plasmid pKJK5  
341 in bacterial populations. *FEMS Microbiol Lett* 2007;**266**:250-256.
- 342 Baltrus DA. Exploring the costs of horizontal gene transfer. *Trends Ecol Evol* 2013;**28**:489-495.
- 343 Bergstrom CT, Lipsitch M, Levin BR. Natural selection, infectious transfer and the existence conditions  
344 for bacterial plasmids. *Genetics* 2000;**155**:1505-1519.
- 345 Brockhurst MA, Harrison E, Hall JP *et al.* The ecology and evolution of pangenomes. *Curr*  
346 *Biol* 2019;**29**:R1094-R1103. DOI: 10.1016/j.cub.2019.08.012
- 347 Busquets A, Peña A, Gomila M *et al.* Genome sequence of *Pseudomonas stutzeri* strain JM300 (DSM  
348 10701), a soil isolate and model organism for natural transformation. *J Bacteriol* 2012;**194**:5477-  
349 5478.

350 Cairns J, Ruokolainen L, Hultman J *et al.* Ecology determines how low antibiotic concentration impacts  
351 community composition and horizontal transfer of resistance genes. *Commun Biol* 2018;**1**:1-8.

352 De Gelder L, Ponciano JM, Joyce P *et al.* Stability of a promiscuous plasmid in different hosts: no  
353 guarantee for a long-term relationship. *Microbiology* 2007;**153**:452-463.

354 Dionisio F, Matic I, Radman M *et al.* Plasmids spread very fast in heterogeneous bacterial communities.  
355 *Genetics* 2002;**162**:1525-1532.

356 Fletcher S. Understanding the contribution of environmental factors in the spread of antimicrobial  
357 resistance. *Environ Health Prev Med* 2015;**20**:243.

358 Fox RE, Zhong X, Krone SM *et al.* Spatial structure and nutrients promote invasion of IncP-1 plasmids  
359 in bacterial populations. *ISME J.* 2008;**2**:1024-1039.

360 Frost LS, Leplae R, Summers AO *et al.* Mobile genetic elements: the agents of open source evolution.  
361 *Nature Rev Microbiol.* 2005;**3**:722-732.

362 Gómez P, Buckling A. Bacteria-phage antagonistic coevolution in soil. *Science* 2011;**332**:106-109.

363 Hall JP, Harrison E, Lilley AK *et al.* Environmentally co-occurring mercury resistance plasmids are  
364 genetically and phenotypically diverse and confer variable context-dependent fitness effects.  
365 *Environ Microbiol* 2015;**17**:5008-5022.

366 Hall JP, Wood AJ, Harrison E *et al.* Source–sink plasmid transfer dynamics maintain gene mobility in  
367 soil bacterial communities. *P Natl Acad Sci USA* 2016;**113**:8260-8265.

368 Hall JP, Brockhurst MA, Harrison E. Sampling the mobile gene pool: innovation via horizontal gene  
369 transfer in bacteria. *Philos Trans R Soc B* 2017a;**372**:20160424. DOI: 10.1098/rstb.2016.0424

370 Hall JP, Williams D, Paterson S *et al.* Positive selection inhibits gene mobilization and transfer in soil  
371 bacterial communities. *Nat Ecol Evol* 2017b;**1**:1348-1353.

372 Harrison E, Brockhurst MA. Plasmid-mediated horizontal gene transfer is a coevolutionary process.  
373 *Trends Microbiol* 2012;**20**:262-267.

374 Harrison E, Guymer D, Spiers AJ *et al.* Parallel compensatory evolution stabilizes plasmids across the  
375 parasitism-mutualism continuum. *Curr Biol* 2015;**25**:2034-2039.

376 Heuer H, Smalla K. Plasmids foster diversification and adaptation of bacterial populations in soil.  
377 *FEMS Microbiol Rev* 2012;**36**:1083-1104.

378 Jurasinski G, Koebisch F, Hagemann U. Flux rate calculation from dynamic closed chamber  
379 measurements. *R package* 2012;version 0.2-1.

380 Klümper U, Riber L, Dechesne A *et al.* Broad host range plasmids can invade an unexpectedly diverse  
381 fraction of a soil bacterial community. *ISME J* 2015;**9**:934-945.

382 Kottara A, Hall JP, Harrison E *et al.* Variable plasmid fitness effects and mobile genetic element  
383 dynamics across *Pseudomonas* species. *FEMS Microbiol Ecol* 2018;**94**:fix172.

384 Lambertsen L, Sternberg C, Molin S. Mini-Tn7 transposons for site-specific tagging of bacteria with  
385 fluorescent proteins. *Environ Microbiol* 2004;**6**:726-732.

386 Levin BR, Stewart FM, Rice VA. The kinetics of conjugative plasmid transmission: fit of a simple mass  
387 action model. *Plasmid* 1979;**2**:247-260.

388 Lilley AK, Bailey MJ, Day MJ *et al.* Diversity of mercury resistance plasmids obtained by exogenous  
389 isolation from the bacteria of sugar beet in three successive years. *FEMS Microbiol Ecol*  
390 1996;**20**:211-227.

391 Lopatkin AJ, Meredith HR, Srimani JK *et al.* Persistence and reversal of plasmid-mediated antibiotic  
392 resistance. *Nat Commun* 2017;**8**:1-10.

393 Norman A, Hansen LH, Sørensen SJ. Conjugative plasmids: vessels of the communal gene pool. *Phil*  
394 *Trans R Soc B* 2009;**364**:2275-2289.

395 Pukall R, Tschäpe H, Smalla K. Monitoring the spread of broad host and narrow host range plasmids  
396 in soil microcosms. *FEMS Microbiol Ecol* 1996;**20**:53-66.

397 R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical  
398 Computing, Vienna, Austria. 2013;Online: <http://www.R-project.org>.



399 Rainey PB, Bailey MJ, Thompson IP. Phenotypic and genotypic diversity of fluorescent pseudomonads  
400 isolated from field-grown sugar beet. *Microbiology* 1994;**140**:2315-2331.

401 Røder HL, Hansen LH, Sørensen SJ *et al.* The impact of the conjugative IncP-1 plasmid pKJK5 on  
402 multispecies biofilm formation is dependent on the plasmid host. *FEMS Microbiol Lett*  
403 2013;**344**:186-192.

404 Sakuda A, Suzuki-Minakuchi C, Okada K *et al.* Conjugative selectivity of plasmids is affected by  
405 coexisting recipient candidates. *mSphere* 2018;**3**:e00490-18. DOI: 10.1128/mSphere.00490-18

406 Sambrook J, Fritsch EF, Maniatis T. Molecular cloning: a laboratory manual. *Cold Spring Harbor*  
407 *Laboratory Press* 1989; (No. Ed. 2).

408 San Millan A, Peña-Miller R, Toll-Riera M *et al.* Positive selection and compensatory adaptation  
409 interact to stabilize non-transmissible plasmids. *Nat Commun* 2014;**5**:1-11.

410 San Millan A, MacLean RC. Fitness costs of plasmids: a limit to plasmid transmission. *Microbiol Spectr*  
411 2017;**5**. DOI: 10.1128/microbiolspec.MTBP-0016-2017

412 Simonsen L, Gordon DM, Stewart FM *et al.* Estimating the rate of plasmid transfer: an end-point  
413 method. *Microbiology* 1990;**136**:2319-2325.


414 Sørensen SJ, Bailey M, Hansen LH *et al.* Studying plasmid horizontal transfer in situ: a critical review.  
415 *Nat Rev Microbiol* 2005;**3**:700-710.

416 Stevenson C, Hall JP, Harrison E *et al.* Gene mobility promotes the spread of resistance in bacterial  
417 populations. *ISME J* 2017;**11**:1930-1932.

418 Stewart FM, Levin BR. The population biology of bacterial plasmids: a priori conditions for the  
419 existence of conjugationally transmitted factors. *Genetics*. 1977;**87**:209-228.

420 Tett A, Spiers AJ, Crossman LC *et al.* Sequence-based analysis of pQBR103; a representative of a  
421 unique, transfer-proficient mega plasmid resident in the microbial community of sugar beet. *ISME*  
422 *J.* 2007;**1**:331-340.


- 423 Torsvik V, Øvreås L. Microbial diversity and function in soil: from genes to ecosystems. *Curr Opin*  
424 *Microbiol* 2002;**5**:240-245.
- 425 van Elsas JD, Bailey MJ. The ecology of transfer of mobile genetic elements. *FEMS Microbiol Ecol*  
426 2002;**42**:187-197.
- 427



428

429 **Figure 1.** Total plasmid density in the community throughout the selection experiment. Panels  
 430 data for communities that varied in mercury selection (without mercury, left-hand set; with  
 431 mercury, right-hand set) their initial original plasmid host (from left to right in each set: *P.*  
 432 *fluorescens*, *P. stutzeri* or *P. putida*). Brown shaded area shows the mean plasmid abundance  
 433 in the community  $\pm$  standard error (dotted line) from six replicates. Solid lines show the mean  
 434 total community bacterial density from six replicates.


435



436

437 **Figure 2.** Population density and mobile genetic element dynamics in communities where *P.*  
 438 *fluorescens* was the original plasmid host. A-F clonal populations evolving with or without  
 439 mercury. Lines show the population densities of *P. fluorescens* (blue); *P. stutzeri* (red); *P.*  
 440 *putida* (green). Brown areas show the density of plasmid carriers; Grey areas show the density  
 441 of cells that have retained the Tn5042 but lost the plasmid.


442



443

444 **Figure 3.** Population density and mobile genetic element dynamics in communities where *P.*  
 445 *stutzeri* was the original plasmid host. A-F clonal populations evolving with or without  
 446 mercury. Lines show the population densities of *P. fluorescens* (blue); *P. stutzeri* (red); *P.*  
 447 *putida* (green). Brown areas show the density of plasmid carriers; Grey areas show the density  
 448 of cells that have retained the Tn5042 but lost the plasmid.


449



450

451 **Figure 4.** Population density and mobile genetic element dynamics in communities where *P.*  
 452 *putida* was the original plasmid host. A-F clonal populations evolving with or without mercury.  
 453 Lines show the population densities of *P. fluorescens* (blue); *P. stutzeri* (red); *P. putida* (green).  
 454 Brown areas show the density of plasmid carriers; Grey areas show the density of cells that  
 455 have retained the Tn5042 but lost the plasmid.

456



457

458 **Figure 5.** Diversity of plasmid-carriers at the end of the experiment. Species diversity was  
 459 calculated as the 1-D Simpson's Index by using the end-point population densities of the  
 460 plasmid-carriers in each species in each original plasmid host community.

461