

OXFORD
UNIVERSITY PRESS

Journal of Travel Medicine

**Risk factors for acquisition of multidrug-resistant
Enterobacteriales among international travellers: A
synthesis of cumulative evidence**

Journal:	<i>Journal of Travel Medicine</i>
Manuscript ID	JTM-19-159.R1
Manuscript type:	Systematic reviews
Date Submitted by the Author:	n/a
Complete List of Authors:	Furuya-Kanamori, Luis; Australian National University, Research School of Population Health Stone, Jennifer; Australian National University Yakob, Laith; London School of Hygiene and Tropical Medicine Kirk, Martyn; Australian National University Collignon, Peter; Canberra Hospital Mills, Deborah; Queensland Children's Medical Research Institute, Lau, Colleen; Australian National University, Department of Global Health, Research School of Population Health; Travel Medicine Alliance Clinics,
Keywords:	antibiotic, antimicrobial, resistance, transmission, carriage, meta-analysis

SCHOLARONE™
Manuscripts

1
2
3 Systematic review and meta-analysis
4
5
6
7

8 Risk factors for acquisition of multidrug-resistant
9
10 *Enterobacterales* among international travellers: A synthesis of
11
12 cumulative evidence
13
14
15
16
17
18

19 **Running title:** Multidrug-resistant *Enterobacterales* and travellers
20
21
22

23
24 Luis Furuya-Kanamori,¹ Jennifer Stone,¹ Laith Yakob,² Martyn Kirk,¹ Peter Collignon,^{3,4}
25 Deborah J Mills,^{5*} Colleen L Lau^{1,5*}
26
27
28

29 ¹Research School of Population Health, College of Health and Medicine, Australian National
30 University, Canberra, Australia
31

32 ²Department of Disease Control, London School of Hygiene and Tropical Medicine, London,
33 UK
34

35 ³ACT Pathology, Canberra Hospital, Canberra, Australia
36

37 ⁴Medical School, College of Health and Medicine, Australian National University, Canberra,
38 Australia
39

40 ⁵Travel Medicine Alliance Clinics, Brisbane, Australia
41
42
43

44 * Joint senior authors
45
46
47
48
49
50

51 Corresponding author:

52 Dr Luis Furuya-Kanamori

53 A: 62 Mills Road, Research School of Population Health, Australian National University,
54 Acton, ACT, Australia

55 E: luis.furuya-kanamori@anu.edu.au
56
57
58
59
60

Abstract

Background: Recent studies have shown that over 50% of people travelling to South East Asia return colonised with multidrug-resistant *Enterobacterales* (MRE) including carbapenemase-producing *Enterobacterales*. Importation of MRE by travellers and subsequent spread to family members, communities, and healthcare facilities poses real risks that have not yet been adequately assessed. This systematic review and meta-analysis aims to quantify the risk factors and interventions for reducing the risk of MRE acquisition among international travellers.

Methods: A systematic search was conducted in PubMed, Web of Science, and Scopus for analytical epidemiological studies containing data post-2000 that assessed the risk factors to acquire and/or interventions to reduce the risk of MRE acquisition in travellers. Two researchers independently screened all the studies and extracted the information, and disagreements were resolved through consensus. The proportions of MRE acquisition by the region of destination and the odds ratio (OR) for the different risk factors and/or interventions were pooled using the inverse variance heterogeneity model.

Results: Twenty studies (5253 travellers from high-income countries) were included in the meta-analysis. South Asia (58.7%, 95%CI:44.5-72.5%) and Northern Africa (43.9%; 95%CI:37.6-50.3%) were the travel destinations with the highest proportion of MRE acquisition. Inflammatory bowel disease (OR 2.1; 95%CI:1.2-3.8), use of antibiotics (OR 2.4; 95%CI:1.9-3.0), traveller's diarrhoea (OR 1.7; 95%CI:1.3-2.3), and contact with the healthcare system overseas (OR 1.5; 95%CI:1.1-2.2) were associated with MRE colonisation. Vegetarians (OR 1.4; 95%CI:1.0-2.0) and backpackers (OR 1.5; 95%CI:1.2-1.8) were also at increased odds of MRE colonisation. Few studies (n=6) investigated preventive measures and found that consuming only bottled water/beverages, meticulous hand hygiene, and probiotics had no protective effect on MRE colonisation.

1
2
3 **Conclusions:** International travel is an important driver for MRE spread worldwide. Future
4
5 research needs to identify effective interventions to reduce the risk of MRE acquisition as well
6
7 as design strategies to reduce local transmission on return.
8
9

10
11
12 **Registration:** PROSPERO CRD42018076853.
13
14

15
16
17 **Keywords:** antibiotic; antimicrobial; resistance; transmission; carriage; meta-analysis
18
19

20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

Introduction

There is a large clinical and public health burden associated with antimicrobial resistance (AMR) in bacteria causing human infections. This burden is increasing over time and urgent action is required.^{1,2} In a report commissioned by the UK Prime Minister, it was estimated that globally AMR could lead to 10 million excess deaths per year, and a cumulative cost of US\$100 trillion by 2050.³ The problem of AMR is much more pronounced in low-income countries with poor water quality and sanitation, uncontrolled use of antibiotics, inadequate infrastructure, and poor governance (i.e. corruption) all being important contributors.^{4,5}

Although carriage of multidrug-resistant *Enterobacteriales* (MRE) is much more prevalent in communities in low-income countries (mainly in South East Asia, Western Pacific, and Eastern Mediterranean regions),⁶ high-income countries are not immune to this problem⁷ (particularly in hospitals and long-term care facilities where there is high prevalence of MRE), and multi-resistant bacteria may also spread from country to country.⁸ Global trade of food (including live animals) and international travel play major roles in the spread and transmission of resistant organisms.⁹ The number of persons crossing international borders has continued to increase over many decades, which has provided an important opportunity for dissemination of MRE to other countries. Asymptomatic carriers can transport MRE between countries in their gut at alarming rates e.g., studies have found that 89% of travellers returning from India and Sri Lanka to the United Kingdom were MRE carriers.¹⁰ These travellers, including medical tourists,¹¹ can transmit organisms to household contacts,^{12,13} and may introduce the organism into the community,¹⁴ healthcare facilities,¹⁵ and the environment. Once established, these resistant organisms are extremely expensive and difficult to control.

In addition to the public health perspective of the risk of introduction of multidrug resistant pathogens, special clinical considerations should be taken with returning travellers

1
2
3 that are carriers of MRE. For example, studies have reported an association between
4 international travel and extended-spectrum β -lactamase-producing (ESBL) *E. coli* bacteraemia
5
6 post trans-rectal ultrasound guided prostate biopsy^{16,17} raising the question of the role of
7
8 screening and antibiotic prophylaxis in this group of patients.
9
10

11
12 In 2015, Hassing et al. conducted a systematic review that compiled the evidence of
13 MRE detection among international travellers.¹⁸ However, they did not quantitatively analyse
14 the data. Additionally, since the publication of the systematic review, findings from the
15 COMBAT study,¹³ the largest study on MRE in travellers with over 2000 participants have
16 been published. In this systematic review and meta-analysis, we update the evidence by
17 including the most recent studies and quantifying the risk factors for MRE colonisation among
18 international travellers.
19
20
21
22
23
24
25
26
27
28
29
30

31 **Methods**

32
33 The protocol was prospectively registered in PROSPERO (CRD42018076853) and the
34 findings of this systematic review and meta-analysis are presented according to PRISMA
35 reporting guidelines.¹⁹
36
37
38
39
40
41

42 ***Search strategy and selection criteria***

43
44 The original search strategy was designed in PubMed and converted for use in Web of
45 Science and Scopus. The search included all articles published from 2000 until February 2019,
46 and the search was updated in July 2019. Search terms related to “travel”,
47
48 “*Enterobacteriaceae*”, and “drug resistance” were included. To achieve a comprehensive
49 evaluation of the published evidence, the systematic search was supplemented with a forwards
50 and backwards citation search as well as retrieving the first 20 similar articles from PubMed
51
52
53
54
55
56
57
58
59
60

1
2
3 for each of the studies included from the initial search. In addition, all references from relevant
4
5 previous systematic reviews were hand searched to identify possible missed studies.
6
7

8 Eligible studies were analytical epidemiological studies among international travellers
9
10 that reported the rate or risk (and protective) factors (i.e. compared MRE carriers versus non-
11
12 carriers) for MRE acquisition (e.g. due to contaminated food), colonisation (e.g. due to an
13
14 underlying condition such as inflammatory bowel disease), or selection (due to use of
15
16 antibiotics) on return, where data were available in an extractable format. Among the risk
17
18 factors of interest were demographic characteristics, travel characteristics, past medical history,
19
20 medication use and medical problems while overseas, food exposure, and reason for travel.
21
22 Potential protective factors included vaccines, probiotics, and hand hygiene. MRE included
23
24 ESBL, AmpC β -lactamase, or carbapenemase-producing *Enterobacterales*. For a study to be
25
26 included, it must have reported the country/region of travel and screened all participants on
27
28 return for MRE and not just targeting symptomatic participants (e.g. with diarrhoea or fever on
29
30 return). Given that the aim of the review was to describe the current situation of MRE among
31
32 international travellers, only studies containing data post-2000 were included.
33
34
35
36

37 Studies reporting MRE acquisition among military personnel, refugees and asylum
38
39 seekers, adoptees, travellers seeking medical attention, international inpatient transfers, or
40
41 during mass gatherings (e.g. Hajj, Olympic games) were not included as these travellers were
42
43 considered at higher risk and not deemed representative of the average traveller. Exclusion
44
45 criteria also included studies conducted on animals, in vitro studies, conference abstracts or
46
47 proceedings, descriptive studies (e.g. case series), and ecological studies. No language
48
49 restriction was imposed.
50
51
52
53
54
55

56 ***Study selection and data extraction***

57
58
59
60

1
2
3 Titles and abstracts of all papers were extracted and uploaded to the Rayyan platform
4
5 (<http://rayyan.qcri.org/>)²⁰ for screening. Two authors (LFK and JS) independently screened the
6
7 titles and abstracts on the Rayyan platform. The same authors examined the full-text papers for
8
9 eligibility in accordance with the review protocol. Any disagreements were resolved through
10
11 consensus.
12
13

14
15 Data from the included studies were extracted and summarised in a spreadsheet, the
16
17 recorded fields included 1) first author and year of publication; 2) country and study setting; 3)
18
19 study population characteristics (e.g. mean/median age, proportion of females, number of
20
21 travellers); 4) travel characteristics (e.g. travel destination, duration of travel); 5) rate of MRE
22
23 detection (i.e. number of travellers that tested positive for MRE on return among the number
24
25 of travellers that tested negative for MRE before travel); 6) risk (and protective) factors for
26
27 MRE acquisition, colonisation, or selection; and 7) MRE isolate characteristics (e.g. type of
28
29 bacteria and resistance genes). Adjusted effect estimates and 95% confidence intervals of the
30
31 risk factors were extracted. If adjusted effect estimates were not available, unadjusted estimates
32
33 were extracted or computed based on the sample size and number of events (i.e. MRE post-
34
35 travel) in the exposed and unexposed groups.
36
37
38
39
40
41

42 ***Statistical analysis***

44
45 Two effect measures of interest were examined, the proportion of MRE detected and
46
47 the odds ratio (OR) for risk factors. Previous reviews^{18,21,22} have reported large heterogeneity
48
49 in the proportion of detection across studies, suggesting the absence of a common worldwide
50
51 effect estimate for the proportion of MRE detection. Countries were grouped into geographical
52
53 regions (i.e. South Asia, Asia [excluding countries in South Asia], Northern Africa, Sub-
54
55 Saharan Africa, South and Central America, North America, Europe, and Oceania) as in a
56
57 previous systematic review by Hassing et al.¹⁸ The detailed list of countries per region can be
58
59
60

1
2
3 found in the supplementary material (S1). A meta-regression was fitted to examine if region of
4 destination was a predictor for the observed heterogeneity. It was found that region of
5 destination was the best predictor (accounted for over 45% of the observed variance) of MRE
6 detection. Therefore, the proportions of MRE detection were pooled by the region of
7 destination using the inverse variance heterogeneity (IVhet) model.²³ Adjusted and unadjusted
8 (when adjusted were not available) estimates along with their 95% confidence intervals were
9 pooled for the different risk factors using the IVhet model. The secondary objective of the
10 systematic review was to describe the most common bacteria isolated from the travellers as
11 well as report the distribution of resistance genes. Proportion meta-analysis using the IVhet
12 model was applied to pool the data from the types of bacteria and resistance genes in travellers.
13 Statistical heterogeneity was assessed using the I^2 . Publication bias was assessed using the Doi
14 plot and LFK index.²⁴ All the analyses were conducted in MetaXL version 5.3 (EpiGear Int
15 Pty Ltd; Sunrise Beach; Australia; <http://www.epigear.com>).

35 Results

37 *Studies identified*

39
40 The search identified 725 publications, 177 articles remained after the title and abstract
41 screening. Full-text review of 177 publications was conducted, and 28 met the eligibility
42 criteria. It was noted that there was overlap in participants in five sets of publications. Four^{13,25-}
43 ²⁷ and two^{28,29} publications used data from the COMBAT and VOYAG-R studies, respectively.
44 Three publications³⁰⁻³² reported results from a Finnish cohort of travellers, two publications^{33,34}
45 from a cohort of Australian healthcare volunteers, and two publications^{35,36} from a cohort of
46 Dutch travellers. Therefore, 20 datasets (from 28 publications) with 5253 travellers were
47 included in this meta-analysis (S2).
48
49
50
51
52
53
54
55
56
57
58
59
60

Characteristics of the included studies

All the studies were conducted in Europe (The Netherlands,^{12,13,35,37,38} Sweden,³⁹⁻⁴² Switzerland,^{43,44} France,^{28,45} Germany,^{38,46} Finland,³⁰ United Kingdom,¹⁰ and Denmark⁴⁷) except for three that were conducted in Australia,³³ Japan,^{48,49} and the USA.⁴⁹ The median age of the participants ranged from 25 to 66 years. Overall more females (n=3028, 57.6%) were included in the studies. The sample size ranged from 18 to 1847 participants. The median duration of travel ranged from 7 to 45 days, with 75% of the studies having a median duration of travel between 14 and 21 days. Thirteen studies collected stool samples,^{10,13,28,30,35,38-40,42,43,46,48,49} five studies used rectal swabs,^{12,33,41,44,45} and two studies use a combination of both types of samples.^{37,47} Collection of the post-travel sample was done within one month after return in 70% of the studies. The majority of studies (n=19, 95%) used selective media for MRE detection that were mainly followed by genotyping confirmation, while one study only used meta-genomic approach for MRE detection³⁵ (Table 1).

Quantitative analysis

Proportion of MRE detection by region of destination

Overall, there was a wide variability in the proportion of MRE detected, ranging from 13% to 88% and considerable heterogeneity across studies was observed. Heterogeneity substantially decreased when estimates were pooled by regions. South Asia (58.7%, 95%CI 44.5-72.5%), Northern Africa (43.9%; 95%CI 37.6-50.3%), and Asia (37.5%; 95%CI 23.6-51.9%) were the travel destinations with the highest proportion of MRE detection. These regions were followed by Sub-Saharan Africa (21.8%; 95%CI 12.0-32.4%), South and Central America (18.3%; 95%CI 8.8-28.9%), and North America (16.9%; 95%CI 2.6-35%); while the lowest proportion of detection were observed in Europe (10.3%; 95%CI 5.5-16.1%) and Oceania (6.9%; 95%CI 0-21.1%) (Figure 1).

Risk factors

MRE detection was not associated with sex, age, or duration of travel, but it was dependent on the travel destination. In terms of past medical history, it was found that inflammatory bowel disease (OR 2.1; 95%CI 1.2-3.8) was a significant risk factor for MRE colonisation. While overseas, use of antibiotics (OR 2.4; 95%CI 1.9-3.0) was the strongest risk factor; antimalarial prophylaxis (OR 1.0; 95%CI 0.8-1.4) was not associated with MRE colonisation, but it should be noted that commonly used antimalarial medications include doxycycline (a broad spectrum antibiotic) as well as other medications (e.g. mefloquine, atovaquone/proguanil, chloroquine) that would not be expected to have an effect on selection of AMR *Enterobacteriales*. Other factors associated with MRE colonisation were experiencing traveller's diarrhoea (OR 1.7; 95%CI 1.3-2.3) and having contact with the healthcare system either as inpatients or outpatients (OR 1.5; 95%CI 1.1-2.2). With regards to food exposure, having a vegetarian diet (OR 1.4; 95%CI 1.0-2.0) was found to increase the odds of acquiring MRE compared to other diets, and consuming only bottled water/beverages did not have an impact on MRE colonisation. Among the interventions that were examined for their potential impact in decreasing the risk of MRE colonisation, meticulous (as defined by primary study authors) hand hygiene and probiotics had no effect, while oral cholera vaccine (OR 1.6; 95%CI 1.0-2.5) was found to increase the odds of MRE colonisation, possible due to confounding by indication (see discussion). Backpacker travellers had a 50% (OR 1.5; 95%CI 1.2-1.8) increased odds of acquiring MRE compared to other types of travellers (Table 2).

Aetiology and resistance genes

Of the 2276 MRE isolates, the vast majority were *E. coli* (92.0%; 95%CI 84.9-97.3%) followed by *K. pneumoniae* (5.9%; 95%CI 1.2-11.8%). The remaining 2% of isolates included

Citrobacter spp. (*freundii* and *amalonaticus*), *Proteus* spp. (*vulgaris* and *mirabilis*), *Enterobacter cloacae*, *Enterococcus faecium*), *Acinetobacter baumannii*, *Morganella morganii*, and a case of non-Typhi *Salmonella enterica*.

*bla*_{CTX-M} (92.3%; 95%CI 44.6-100%) was the most frequent resistance gene found in the MRE isolates. CTX-M group 1 was the most common followed by CTX-M groups 9 and 4. *bla*_{SHV} and *bla*_{TEM} were found in 3.0% (95%CI 0-2.0%) and 2.0% (95%CI 0-1.8%) of the isolates, respectively. *Enterobacteriales* with carbapenem- (*bla*_{OXA}^{28,37,45} and *bla*_{NDM}^{28,44,45}) and colistin- (*mcr*⁴⁸) resistance genes were also isolated. Fifty isolates with carbapenem-resistant genes were identified mainly from travellers returning from Southeast Asia and the Indian subcontinent,^{28,37,44,45} while the three isolates with colistin-resistant genes were from travellers returning from Vietnam.⁴⁸

Discussion

In this systematic review and meta-analysis, we compiled data from 20 different studies with over 5000 travellers to update the evidence on the proportion and risk factors for MRE detection in international travellers. The highest proportions of MRE detection were observed from travellers returning from South Asia, Northern Africa, Asia, Sub-Saharan Africa, and South and Central America; our findings correspond with the global epidemiology of AMR⁴ and the ESBL producing *Enterobacteriales* colonisation prevalence among healthy individuals.⁵⁰ In a meta-analysis by Karanika et al., international travellers were found to be four times more likely to be colonised by ESBL producing *Enterobacteriales* than non-travellers.⁵⁰ These findings highlight the importance of international travel to low- and middle-income countries in the global spread of MRE.^{51,52} In recent years, dissemination of emerging pathogens such as NDM-1 carbapenem-resistant *K. pneumoniae* from India⁵³ and *mcr*-1 colistin-resistant *E. coli* from China⁵⁴ were facilitated by international travel.^{55,56} Of concern is

1
2
3 that our review revealed that carbapenem^{28,37,44,45} and colistin⁴⁸-resistant *Enterobacterales* are
4 being detected in international travellers. Investment in infrastructure to provide access to clean
5 water, and adequate sanitation and hygiene may reduce transmission in low- and middle-
6 income countries,⁴ and possibly reduce the global spread of AMR.
7
8
9
10
11

12 Contact with the healthcare system, traveller's diarrhoea, and antibiotic use while
13 travelling were strongly associated with MRE detection. The possibility that these factors are
14 associated through reverse causality with MRE cannot be excluded (i.e. MRE infection leading
15 to contact with the healthcare system, traveller's diarrhoea, and/or use of antibiotics). Even in
16 the event the observed associations are due to reverse causality, these factors are markers for
17 MRE detection and can be used to identify travellers at higher risk of being colonised by MRE.
18
19 Among the risk factors identified, one that travellers have control over is antibiotic
20 consumption, thus during pre-travel consultations, healthcare professionals need to emphasise
21 the use of antibiotics only when is required.
22
23
24
25
26
27
28
29
30
31
32

33 Acute diarrhoea is the most common condition affecting travellers to South and Central
34 Asia and the second most common among travellers worldwide.⁵⁷ Therefore, priority areas of
35 research have been identified to generate evidence-based recommendations for preventing
36 traveller's diarrhoea.⁵⁸ In line with the research priority areas, inflammatory bowel disease, a
37 vegetarian diet, and backpacking were identified as host risk factors for MRE detection.
38
39 Although it was not possible to examine the exposure to different classes and doses of
40 antibiotics in the current review, findings by Ruppe et al.²⁸ indicate that use of different classes
41 of antibiotics have distinct effects on the risk of MRE selection. They found that antibiotics
42 used for malaria prophylaxis (i.e. doxycycline) did not to increase the risk of MRE selection
43 (OR 0.9; 95%CI 0.6-1.6), while beta-lactams were associated with an increase in risk (OR 4.2;
44 95%CI 1.5-12.1).²⁸
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Non-antibiotic approaches to prevent MRE colonisation were also examined. Findings
4
5 from the current review align with a recent randomised controlled trial that found that
6
7 *Lactobacillus rhamnosus* GG did not reduce MRE colonisation in travellers.⁵⁹ The effect of
8
9 oral cholera vaccine was also assessed, and paradoxically it was found that cholera vaccine was
10
11 associated with an increase in the risk of MRE colonisation. Considering that the oral cholera
12
13 vaccine provides some cross-protection against *enterotoxigenic E. coli*, one of the most
14
15 common causes of traveller's diarrhoea, this association may seem contradictory. The most
16
17 likely explanation for the observation is that unadjusted estimates were analysed and the result
18
19 is affected by confounding by indication, i.e. travellers going to regions with higher risk of
20
21 MRE are more likely to receive oral cholera vaccine. In our previous study of MRE detection
22
23 in Australian travellers, those travelling to the Indian subcontinent were more likely to receive
24
25 oral cholera vaccine than travellers going to other destinations; when the analysis was adjusted
26
27 for travel destination and traveller's comorbidities, the point estimate indicated a protective
28
29 effect of oral cholera vaccine against MRE colonisation.⁶⁰ To prevent confounding by
30
31 indication, randomised controlled trials are needed to provide robust evidence on the effect of
32
33 potentially protective interventions.
34
35
36
37
38
39

40 It is clear that international travellers are a high risk group; over 50% of travellers
41
42 returning from South Asia are asymptomatic MRE carriers and there is increasing evidence of
43
44 subsequent household transmission of MRE.^{12,13} Cases of ESBL-producing *E. coli* bacteraemia
45
46 have been reported after prostate biopsies¹⁷ and international travel has been identified as the
47
48 risk factor for fluoroquinolone-resistant and ESBL-producing *E. coli* infections.^{16,61} Therefore,
49
50 the role of screening and contact precaution of returned travellers from clinical and public
51
52 health perspectives need to be further examined.⁶² It is unfeasible to screen all returning
53
54 travellers, but in a healthcare setting, it may be feasible to screen those who are planning to
55
56 undergo higher risk medical procedures; although current evidence suggest that ESBL-
57
58
59
60

1
2
3 producing *E. coli* is less likely to spread between patients in hospitals than other
4
5 *Enterobacterales* species.⁶³ Current studies on international patient transfers have shown that
6
7 targeted screening programs of high-risk patients may be a more cost-effective strategy than
8
9 mass screening⁶⁴ and contact precaution and environmental cleaning with chlorine-based
10
11 disinfectant can prevent transmission.⁶⁵ Given that prolonged MRE carriage is uncommon,
12
13 most travellers (~90%) are culture-negative within 6 months,^{28,33,46} strategies should aim to
14
15 reduce transmission and avoid elective surgical interventions within this period.
16
17
18

19 The findings of our systematic review and meta-analysis should be understood in light
20
21 of some of the limitations. Estimates extracted and synthesized from observational studies
22
23 included adjusted as well as unadjusted estimates, thus pooled estimates may be affected by
24
25 confounding variables. This may be the case of potential protective interventions, where
26
27 paradoxical findings were observed. To examine regional and within-country differences in
28
29 MRE detection, as well as the risk of different antibiotic classes and doses, more granular data
30
31 are required. Individual patient data meta-analysis may provide valuable information to answer
32
33 these questions. The current review examined the risk factors relevant to average travellers
34
35 from high-income countries, thus findings may not be applicable to special groups of travellers
36
37 such as military personnel, refugees and asylum seekers, or hospital transfers as they may have
38
39 different risk profiles. The median age of travellers in the included studies ranged from 25 to
40
41 66 years, so findings may not be generalizable to children. Some studies collected post-travel
42
43 samples between 3 and 12 months post travel, thus we cannot rule out the possibility that the
44
45 participants became colonised after their return.
46
47
48
49

50
51 In conclusion, international travel is an important driver for MRE spread worldwide.
52
53 South Asia as a travel destination is a major risk factor for MRE acquisition. Additional risk
54
55 factors for individuals to become colonised, include inflammatory bowel disease, use of
56
57 antibiotics, traveller's diarrhoea, contact with the healthcare system, having a vegetarian diet,
58
59
60

1
2
3 and backpacking. Future research needs to identify effective interventions to reduce the risk of
4
5 MRE colonisation and design strategies to reduce local transmission on return, including
6
7 transmission to household and community members, as well as wider dissemination into the
8
9 environment and animal populations
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

Acknowledgement

The authors acknowledge the assistance provided by Mr Tsheten Tsheten for creating the choropleth map.

Funding

LFK and CLL were supported by Australian National Health and Medical Research Council Early Career Fellowships (APP1158469 and APP1109035). MK was supported by an Australian National Health and Medical Research Council Career Development Fellowship (APP1145997). JS was supported by the Australian National University, Australia Degree by Research Scholarship.

Conflicts of interest

The authors do not have any conflicts of interest to declare.

Authors' contribution

Conception and design of the study: LFK, DJM, CLL

Collection and assembly of the data: LFK, JS

Analysis and interpretation: LFK, LY, DJM, CLL

Manuscript writing: All authors

Final approval of manuscript: All authors

References

1. de Kraker ME, Davey PG, Grundmann H. Mortality and hospital stay associated with resistant *Staphylococcus aureus* and *Escherichia coli* bacteremia: estimating the burden of antibiotic resistance in Europe. *PLoS Med* 2011; 8(10): e1001104.
2. Stewardson AJ, Allignol A, Beyersmann J, Graves N, Schumacher M, Meyer R, et al. The health and economic burden of bloodstream infections caused by antimicrobial-susceptible and non-susceptible Enterobacteriaceae and *Staphylococcus aureus* in European hospitals, 2010 and 2011: a multicentre retrospective cohort study. *Eurosurveillance* 2016; 21(33).
3. O'Neil J. Antimicrobial Resistance: Tackling a Crisis for the Health and Wealth of Nations. 2014. https://amr-review.org/sites/default/files/AMR%20Review%20Paper%20-%20Tackling%20a%20crisis%20for%20the%20health%20and%20wealth%20of%20nations_1.pdf (accessed Aug 2019).
4. Collignon P, Beggs JJ, Walsh TR, Gandra S, Laxminarayan R. Anthropological and socioeconomic factors contributing to global antimicrobial resistance: a univariate and multivariable analysis. *Lancet Planet Health* 2018; 2(9): e398-e405.
5. Ventola CL. The antibiotic resistance crisis: part 1: causes and threats. *P T* 2015; 40(4): 277-83.
6. Woerther PL, Burdet C, Chachaty E, Andremont A. Trends in human fecal carriage of extended-spectrum β -lactamases in the community: Toward the globalization of CTX-M. *Clin Microbiol Rev* 2013; 26(4): 744-58.
7. Klein EY, Tseng KK, Pant S, Laxminarayan R. Tracking global trends in the effectiveness of antibiotic therapy using the Drug Resistance Index. *BMJ Global Health* 2019; 4(2): e001315.
8. Kumarasamy KK, Toleman MA, Walsh TR, Bagaria J, Butt F, Balakrishnan R, et al. Emergence of a new antibiotic resistance mechanism in India, Pakistan, and the UK: a molecular, biological, and epidemiological study. *Lancet Infect Dis* 2010; 10(9): 597-602.
9. Collignon P. Antibiotic resistance: are we all doomed? *Intern Med J* 2015; 45(11): 1109-15.

10. Bevan ER, McNally A, Thomas CM, Piddock LJV, Hawkey PM. Acquisition and loss of CTX-M-producing and non-producing escherichia coli in the fecal microbiome of travelers to South Asia. *mBio* 2018; 9(6).
11. Chen LH, Wilson ME. The Globalization of Healthcare: Implications of Medical Tourism for the Infectious Disease Clinician. *Clin Infect Dis* 2013; 57(12): 1752-9.
12. Paltansing S, Vlot JA, Kraakman ME, Mesman R, Bruijning ML, Bernards AT, et al. Extended-spectrum β -lactamase-producing enterobacteriaceae among travelers from the Netherlands. *Emerg Infect Dis* 2013; 19(8): 1206-13.
13. Arcilla MS, van Hattem JM, Haverkate MR, Bootsma MCJ, van Genderen PJJ, Goorhuis A, et al. Import and spread of extended-spectrum beta-lactamase-producing Enterobacteriaceae by international travellers (COMBAT study): a prospective, multicentre cohort study. *Lancet Infect Dis* 2017; 17(1): 78-85.
14. Wilder-Smith A, Goh KT, Barkham T, Paton NI. Hajj-Associated Outbreak Strain of Neisseria meningitidis Serogroup W135: Estimates of the Attack Rate in a Defined Population and the Risk of Invasive Disease Developing in Carriers. *Clin Infect Dis* 2003; 36(6): 679-83.
15. Fournier S, Lepointeur M, Kassis-Chikhani N, Huang M, Brun-Buisson C, Jarlier V. Link between carbapenemase-producing Enterobacteria carriage and cross-border exchanges: eight-year surveillance in a large French multihospitals institution. *J Trav Med* 2012; 19(5): 320-3.
16. Patel U, Dasgupta P, Amoroso P, Challacombe B, Pilcher J, Kirby R. Infection after transrectal ultrasonography-guided prostate biopsy: increased relative risks after recent international travel or antibiotic use. *BJU Int* 2012; 109(12): 1781-5.
17. Williamson DA, Masters J, Freeman J, Roberts S. Travel-associated extended-spectrum beta-lactamase-producing Escherichia coli bloodstream infection following transrectal ultrasound-guided prostate biopsy. *BJU Int* 2012; 109(7): E21-2.
18. Hassing RJ, Alsmas J, Arcilla MS, van Genderen PJ, Stricker BH, Verbon A. International travel and acquisition of multidrug-resistant Enterobacteriaceae: a systematic review. *Eurosurveillance* 2015; 20(47).

19. Moher D, Liberati A, Tetzlaff J, Altman DG, The PG. Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. *PLoS Med* 2009; 6(7): e1000097.
20. Ouzzani M, Hammady H, Fedorowicz Z, Elmagarmid A. Rayyan-a web and mobile app for systematic reviews. *Syst Rev* 2016; 5(1): 210.
21. Ruppé E, Andremont A, Armand-Lefèvre L. Digestive tract colonization by multidrug-resistant Enterobacteriaceae in travellers: An update. *Travel Med Infect Dis* 2018; 21: 28-35.
22. Armand-Lefevre L, Andremont A, Ruppe E. Travel and acquisition of multidrug-resistant Enterobacteriaceae. *Med Mal Infect* 2018; 48(7): 431-41.
23. Doi SA, Barendregt JJ, Khan S, Thalib L, Williams GM. Advances in the meta-analysis of heterogeneous clinical trials I: The inverse variance heterogeneity model. *Contemp Clin Trials* 2015; 45(Pt A): 130-8.
24. Furuya-Kanamori L, Barendregt JJ, Doi SAR. A new improved graphical and quantitative method for detecting bias in meta-analysis. *Int J Evid Based Healthc* 2018; 16(4): 195-203.
25. Van Hattem JM, Arcilla MS, Bootsma MCJ, Van Genderen PJ, Goorhuis A, Grobusch MP, et al. Prolonged carriage and potential onward transmission of carbapenemase-producing Enterobacteriaceae in Dutch travelers. *Future Microbiol* 2016; 11(7): 857-64.
26. van Hattem JM, Arcilla MS, Grobusch MP, Bart A, Bootsma MC, van Genderen PJ, et al. Travel-related acquisition of diarrhoeagenic bacteria, enteral viruses and parasites in a prospective cohort of 98 Dutch travellers. *Travel Med Infect Dis* 2017; 19: 33-6.
27. van Hattem JM, Cabal A, Arcilla MS, Alvarez J, de Jong MD, Melles DC, et al. Risk of acquisition of human diarrhoeagenic Escherichia coli virulence genes in intercontinental travellers: A prospective, multi-centre study. *Travel Med Infect Dis* 2019.
28. Ruppe E, Armand-Lefevre L, Estellat C, Consigny PH, El Mniai A, Boussadia Y, et al. High Rate of Acquisition but Short Duration of Carriage of Multidrug-Resistant Enterobacteriaceae After Travel to the Tropics. *Clin Infect Dis* 2015; 61(4): 593-600.
29. Lorme F, Maataoui N, Rondinaud E, Esposito-Farese M, Clermont O, Ruppe E, et al. Acquisition of plasmid-mediated cephalosporinase producing Enterobacteriaceae after a travel to the tropics. *PLoS One* 2018; 13(12): e0206909.

- 1
2
3 30. Kantele A, Laaveri T, Mero S, Vilkmann K, Pakkanen SH, Ollgren J, et al.
4 Antimicrobials increase travelers' risk of colonization by extended-spectrum betalactamase-
5 producing Enterobacteriaceae. *Clin Infect Dis* 2015; 60(6): 837-46.
6
7
8
9 31. Kantele A, Mero S, Kirveskari J, Laaveri T. Fluoroquinolone antibiotic users select
10 fluoroquinolone-resistant ESBL-producing Enterobacteriaceae (ESBL-PE) - Data of a
11 prospective traveller study. *Travel Med Infect Dis* 2017; 16: 23-30.
12
13
14
15 32. Lääveri T, Vilkmann K, Pakkanen S, Kirveskari J, Kantele A. Despite antibiotic
16 treatment of travellers' diarrhoea, pathogens are found in stools from half of travellers at return.
17 *Travel Med Infect Dis* 2018; 23: 49-55.
18
19
20
21 33. Kennedy K, Collignon P. Colonisation with Escherichia coli resistant to "critically
22 important" antibiotics: A high risk for international travellers. *Eur J Clin Microbiol Infect Dis*
23 2010; 29(12): 1501-6.
24
25
26
27 34. Rogers BA, Kennedy KJ, Sidjabat HE, Jones M, Collignon P, Paterson DL. Prolonged
28 carriage of resistant E. coli by returned travellers: clonality, risk factors and bacterial
29 characteristics. *Eur J Clin Microbiol Infect Dis* 2012; 31(9): 2413-20.
30
31
32
33 35. von Wintersdorff CJ, Penders J, Stobberingh EE, Oude Lashof AM, Hoebe CJ,
34 Savelkoul PH, et al. High rates of antimicrobial drug resistance gene acquisition after
35 international travel, The Netherlands. *Emerg Infect Dis* 2014; 20(4): 649-57.
36
37
38
39 36. von Wintersdorff CJH, Wolffs PFG, van Niekerk JM, Beuken E, van Alphen LB,
40 Stobberingh EE, et al. Detection of the plasmid-mediated colistin-resistance gene mcr-1 in
41 faecal metagenomes of Dutch travellers. *J Antimicrob Chemother* 2016; 71(12): 3416-9.
42
43
44
45 37. Reuland EA, Sonder GJ, Stolte I, Al Naiemi N, Koek A, Linde GB, et al. Travel to Asia
46 and traveller's diarrhoea with antibiotic treatment are independent risk factors for acquiring
47 ciprofloxacin-resistant and extended spectrum beta-lactamase-producing Enterobacteriaceae-a
48 prospective cohort study. *Clin Microbiol Infect* 2016; 22(8): 731.e1-7.
49
50
51
52 38. Schaumburg F, Sertic SM, Correa-Martinez C, Mellmann A, Köck R, Becker K.
53 Acquisition and colonization dynamics of antimicrobial-resistant bacteria during international
54 travel: a prospective cohort study. *Clin Microbiol Infect* 2019 [Epub ahead of print].
55
56
57
58
59
60

- 1
2
3 39. Östholm-Balkhed A, Tärnberg M, Nilsson M, Nilsson LE, Hanberger H, Hällgren A.
4 Travel-associated faecal colonization with esbl-producing enterobacteriaceae: Incidence and
5 risk factors. *J Antimicrob Chemother* 2013; 68(9): 2144-53.
6
7
8
9 40. Tängdén T, Cars O, Melhus Å, Löwdin E. Foreign travel is a major risk factor for
10 colonization with Escherichia coli producing CTX-M-type extended-spectrum β -lactamases:
11 A prospective study with Swedish volunteers. *Antimicrob Agents Chemother* 2010; 54(9):
12 3564-8.
13
14
15
16 41. Vading M, Kabir MH, Kalin M, Iversen A, Wiklund S, Naucler P, et al. Frequent
17 acquisition of low-virulence strains of ESBL-producing Escherichia coli in travellers. *J*
18 *Antimicrob Chemother* 2016; 71(12): 3548-55.
19
20
21
22 42. Angelin M, Forsell J, Granlund M, Evengard B, Palmgren H, Johansson A. Risk factors
23 for colonization with extended-spectrum beta-lactamase producing Enterobacteriaceae in
24 healthcare students on clinical assignment abroad: A prospective study. *Travel Med Infect Dis*
25 2015; 13(3): 223-9.
26
27
28
29 43. Bernasconi OJ, Kuenzli E, Pires J, Tinguely R, Carattoli A, Hatz C, et al. Travelers can
30 import colistin-resistant Enterobacteriaceae, including those possessing the plasmid-mediated
31 mcr-1 gene. *Antimicrob Agents Chemother* 2016; 60(8): 5080-4.
32
33
34
35 44. Kuenzli E, Jaeger VK, Frei R, Neumayr A, DeCrom S, Haller S, et al. High colonization
36 rates of extended-spectrum beta-lactamase (ESBL)-producing Escherichia coli in Swiss
37 travellers to South Asia- a prospective observational multicentre cohort study looking at
38 epidemiology, microbiology and risk factors. *BMC Infect Dis* 2014; 14: 528.
39
40
41
42 45. Macaux L, Ndoye O, Cordel H, Pomares TB, Seytre D, Bouchaud O, et al. Extensively-
43 drug-resistant bacteria carriers among overseas travellers: one-third had not been hospitalized
44 previously. *Int J Antimicrob Agents* 2018; 52(3): 385-9.
45
46
47
48 46. Lübbert C, Straube L, Stein C, Makarewicz O, Schubert S, Mössner J, et al.
49 Colonization with extended-spectrum beta-lactamase-producing and carbapenemase-
50 producing Enterobacteriaceae in international travelers returning to Germany. *Int J Med*
51 *Microbiol* 2015; 305(1): 148-56.
52
53
54
55
56
57
58
59
60

- 1
2
3 47. Lausch KR, Fuursted K, Larsen CS, Storgaard M. Colonisation with multi-resistant
4 Enterobacteriaceae in hospitalised Danish patients with a history of recent travel: A cross-
5 sectional study. *Travel Med Infect Dis* 2013; 11(5): 320-3.
6
7
8
9 48. Nakayama T, Kumeda Y, Kawahara R, Yamaguchi T, Yamamoto Y. Carriage of
10 colistin-resistant, extended-spectrum β -lactamase-producing escherichia coli harboring the
11 mcr-1 resistance gene after short-term international travel to Vietnam. *Infect Drug Resist* 2018;
12 11: 391-5.
13
14
15
16 49. Weisenberg SA, Mediavilla JR, Chen L, Alexander EL, Rhee KY, Kreiswirth BN, et
17 al. Extended Spectrum Beta-Lactamase-Producing Enterobacteriaceae in International
18 Travelers and Non-Travelers in New York City. *PLoS One* 2012; 7(9).
19
20
21
22 50. Karanika S, Karantanos T, Arvanitis M, Grigoras C, Mylonakis E. Fecal Colonization
23 with Extended-spectrum Beta-lactamase-Producing Enterobacteriaceae and Risk Factors
24 among Healthy Individuals: A Systematic Review and Metaanalysis. *Clin Infect Dis* 2016;
25 63(3): 310-8.
26
27
28
29
30 51. Frost I, Van Boeckel TP, Pires J, Craig J, Laxminarayan R. Global Geographic Trends
31 in Antimicrobial Resistance: The Role of International Travel. *J Travel Med* 2019 [Epub ahead
32 of print].
33
34
35
36 52. Vila J. Multidrug-Resistant Bacteria Without Borders: Role of International Trips in
37 the Spread of Multidrug-Resistant Bacteria. *J Travel Med* 2015; 22(5): 289-91.
38
39
40 53. Yong D, Toleman MA, Giske CG, Cho HS, Sundman K, Lee K, et al. Characterization
41 of a new metallo-beta-lactamase gene, bla(NDM-1), and a novel erythromycin esterase gene
42 carried on a unique genetic structure in *Klebsiella pneumoniae* sequence type 14 from India.
43 *Antimicrob Agents Chemother* 2009; 53(12): 5046-54.
44
45
46
47 54. Liu YY, Wang Y, Walsh TR, Yi LX, Zhang R, Spencer J, et al. Emergence of plasmid-
48 mediated colistin resistance mechanism MCR-1 in animals and human beings in China: a
49 microbiological and molecular biological study. *Lancet Infect Dis* 2016; 16(2): 161-8.
50
51
52
53 55. Berrazeg M, Diene S, Medjahed L, Parola P, Drissi M, Raoult D, et al. New Delhi
54 Metallo-beta-lactamase around the world: an eReview using Google Maps. *Eurosurveillance*
55 2014; 19(20).
56
57
58
59
60

- 1
2
3 56. Schwarz S, Johnson AP. Transferable resistance to colistin: a new but old threat. *J*
4 *Antimicrob Chemother* 2016; 71(8): 2066-70.
5
6
7 57. Freedman DO, Weld LH, Kozarsky PE, Fisk T, Robins R, Von Sonnenburg F, et al.
8 Spectrum of disease and relation to place of exposure among ill returned travelers. *N Engl J*
9 *Med* 2006; 354(2): 119-30.
10
11
12
13 58. DuPont HL, Steffen R. Use of antimicrobial agents for treatment and prevention of
14 travellers' diarrhoea in the face of enhanced risk of transient fecal carriage of multi-drug
15 resistant enterobacteriaceae: setting the stage for consensus recommendations. *J Trav Med*
16 2017; 24(suppl_1): S57-s62.
17
18
19
20
21 59. Dall LB, Lausch KR, Gedebjerg A, Fuursted K, Storgaard M, Larsen CS. Do probiotics
22 prevent colonization with multi-resistant Enterobacteriaceae during travel? A randomized
23 controlled trial. *Trav Med Infect Dis* 2019; 27: 81-6.
24
25
26
27 60. Furuya-Kanamori L, Mills DJ, Robson J, Lau CL. Acquisition Rate and Risk Factors
28 for Multidrug Resistance Enterobacteriaceae among Australian Travellers. Poster presentation
29 at the 16th Conference of the ISTM. Washington DC, USA; 2019.
30
31
32
33 61. Knaapila J, Kallio H, Hakanen AJ, Syvänen K, Ettala O, Kähkönen E, et al. Antibiotic
34 susceptibility of intestinal Escherichia coli in men undergoing transrectal prostate biopsies: a
35 prospective, registered, multicentre study. *BJU Int* 2018; 122(2): 203-10.
36
37
38
39 62. Petersen E, Mohsin J. Should travelers be screened for multi-drug resistant (MDR)
40 bacteria after visiting high risk areas such as India? *Travel Med Infect Dis* 2016; 14(6): 591-4.
41
42
43 63. Duval A, Obadia T, Boëlle P-Y, Fleury E, Herrmann J-L, Guillemot D, et al. Close
44 proximity interactions support transmission of ESBL-K. pneumoniae but not ESBL-E. coli in
45 healthcare settings. *PLoS Comput Biol* 2019; 15(5): e1006496-e.
46
47
48
49 64. Kaspar T, Schweiger A, Droz S, Marschall J. Colonization with resistant
50 microorganisms in patients transferred from abroad: Who needs to be screened? *Antimicrob*
51 *Resist Infect Control* 2015; 4(1).
52
53
54 65. Chua KY, Grayson ML, Burgess AN, Lee JY, Howden BP. The growing burden of
55 multidrug-resistant infections among returned Australian travellers. *Med J Aust* 2014; 200(2):
56 116-8.
57
58
59
60

Table 1. Characteristics of the included studies

First author, year of publication	Country	Study period	Study population	Mean/median age (in years)	Proportion of females (%)	Sample size	Type of sample	Collection of sample post-travel	Laboratory detection method	Most common travel destinations	Mean/median duration of travel (in days)	Proportion of MRE detection (%)
Angelin 2015 ⁴²	Sweden	2010-2014	Healthcare students travelling for clinical courses	25	78	99	Stool sample	1-2 weeks after return	Selective media	89% South East Asia and Africa	45	35
Arcilla 2017 ¹³	The Netherlands	2012-2013	Patients attending travel clinics	51	54	1847	Stool sample	Immediately after return	Selective media + microarray genotyping	82% Asia and Africa	20	34
Bernasconi 2016 ⁴³	Switzerland	2015	People living in Switzerland	44	68	38	Stool sample	Within 1 week after return	Selective media + microarray genotyping	India	18	76
Bevan 2018 ¹⁰	UK	NR	Healthy volunteers	30	NR	18	Stool sample	NR	Selective media + PCR genotyping + whole genome sequencing	India and Sri Lanka	21	89
Kantele 2015 ³⁰	Finland	2009-2010	Patients attending a travel clinic	40	61	430	Stool sample	First or second stool after return	Selective media + PCR genotyping	83% Asian and Africa	19	21
Kennedy 2010 ³³	Australia	2008-2009	Hospital staff and contacts	45	62	102	Rectal swab	Within 2 weeks after return	Selective media + PCR genotyping	77% South East Asia and Europe	21	49
Kuenzli 2014 ⁴⁴	Switzerland	2012-2013	Patients attending travel clinics	41	56	170	Rectal swab	Directly after return	Selective media + microarray and PCR/DNA sequence genotyping	India, Bhutan, Nepal, and Sri Lanka	18	69
Lausch 2013 ⁴⁷	Denmark	2011	Inpatients with travel history in the last 3 months	37	57	88	Rectal swab and stool sample (if patient had diarrhoea)	Up to 3 months after return	Selective media	52% Asia and Africa	NR	13
Lubbert 2015 ⁴⁶	Germany	2013-2014	Patients attending a travel clinic	34	57	191	Stool sample	Within 1 week after return	Selective media + PCR genotyping	88% Asia and Africa	21	30
Macaux 2018 ⁴⁵	France	2014-2016	Inpatients with travel history in the last 12 months	62	34	138	Rectal swab	Up to 12 months after return	Selective media	83% Asia, Sub-Saharan Africa, and the Mediterranean	NR	NA
Nakayama 2018 ⁴⁸	Japan	2015-2016	Japanese travellers to Vietnam	37	42	19	Stool sample	Within 3 weeks after return	Selective media + PCR genotyping	Vietnam	7	88
Ostholm-Balkhed 2013 ³⁹	Sweden	2008-2009	Patients attending vacation clinics	54	59	226	Stool sample	Up to 4 months after return	Selective media + PCR genotyping	57% Sub-Saharan Africa and Asia	16	30

1	Paltansing 2013 ¹²	The Netherlands	2011	Patients attending a travel clinic	33	63	338	Rectal swabs	Immediately after return	Selective media + microarray genotyping	49% South East Asia and Central Africa	21	33
2													
3	Reuland 2016 ³⁷	The Netherlands	2012-2013	Patients attending a vacation clinic	33	58	418	Rectal swab or stool sample	Within 2 weeks after return	Selective media + microarray genotyping	Africa, Asia, and Latin America	14	23
4													
5													
6	Ruppe 2015 ²⁸	France	2012-2013	Patients attending international vaccination centres	36	61	574	Stool sample	Within 1 week after return	Selective media + PCR genotyping	Sub-Saharan Africa, Asia, and Latin America	20	50
7													
8													
9	Schaumburg 2019 ³⁸	Germany and The Netherlands	2016-2018	Patients attending vaccination centres	32	59	132	Stool sample	Within 1 week after return	Selective media + molecular assay	68% Asia and Africa	18	46
10													
11	Tangden 2010 ⁴⁰	Sweden	2007-2009	Patients attending travel clinics	43	55	100	Stool sample	NR	Selective media + PCR genotyping	65% Africa and Asia	14	24
12													
13	Vading 2016 ⁴¹	Sweden	2013-2015	Patients attending a travel clinic	49	68	175	Rectal swab	NR	Selective media + PCR genotyping	South East Asia, India, Northern Africa and Turkey	14	38
14													
15													
16													
17	von Wintersdorff 2014 ³⁵	The Netherlands	2010-2012	Patients attending travel clinics	43	58	122	Stool sample	Immediately after return	Meta-genomic	48% South East Asia and India	21	32
18													
19	Weisenberg 2012 ⁴⁹	USA	2009-2010	Patients attending a travel clinic	66	68	28	Stool sample	Within 1 week after return	Selective media + PCR genotyping	54% Sub- Saharan Africa and South Asia	16	25
20													
21													
22													

NR not reported; NA not applicable, the study was a matched case-control

Table 2. Risk factors for acquisition of multidrug-resistant *Enterobacterales*

	OR (95% CI)	I ² (%)	Number of studies with adjusted estimates/ Total number of studies
Demographic			
Sex (female)	0.91 (0.80-1.03)	0	2 / 14
Age (per 10 year increase)	1.01 (0.97-1.06)	0	2 / 5
Travel associated data			
Duration of travel (per week increase)	1.02 (0.99-1.06)	0	1 / 5
Travel destination (reference Europe)*	1	NA	0 / 10
South Asia	1.52 (1.31-1.76)		0 / 19
Northern Africa	1.33 (1.25-1.43)		0 / 11
Asia	1.26 (1.17-1.35)		0 / 16
Sub-Saharan Africa	1.12 (0.99-1.28)		0 / 15
Americas	1.11 (1.02-1.20)		0 / 16
Past medical history			
Inflammatory bowel disease	2.09 (1.16-3.77)	0	1 / 3
Chronic disease	1.01 (0.65-1.57)	57	2 / 5
Admitted to a hospital in the previous 3-6 months	1.79 (0.09-36.63)	68	1 / 2
Antibiotic use in the previous 3-12 months	1.00 (0.76-1.32)	12	3 / 4
Medical history while overseas			
Antibiotic use	2.38 (1.88-3.00)	0	4 / 12
Anti-acid use	1.08 (0.59-1.97)	25	0 / 5
Antimalarial prophylaxis	1.04 (0.75-1.44)	10	0 / 3
Travellers' diarrhoea	1.69 (1.25-2.30)	46	5 / 13
Contact with healthcare system	1.53 (1.09-2.15)	20	1 / 10
Food exposure			
Vegetarian diet	1.41 (1.01-1.96)	0	0 / 5
Only consume bottled water/beverages	1.29 (0.50-3.34)	77	1 / 6
Protective factors			
Oral cholera vaccine	1.61 (1.04-2.50)	0	0 / 2
Meticulous hand hygiene	1.10 (0.81-1.49)	0	1 / 4
Probiotics	1.06 (0.78-1.45)	0	0 / 2
Type of traveller			
Backpackers	1.46 (1.20-1.78)	0	2 / 7
Holiday makers	0.99 (0.70-1.38)	0	0 / 4
Visiting friends and relatives	0.95 (0.55-1.63)	29	0 / 5
Business travellers	0.78 (0.46-1.30)	0	0 / 4

OR odds ratio; CI confidence interval; NA not applicable

* OR obtained through meta-regression. Americas include North, Central, and South America. Oceania was excluded from the analysis because too few studies (3) reported data for this region.




Figure 1. Proportion of acquisition of multidrug-resistant *Enterobacterales* by region of destination

Supplementary material

S1. Countries by region

South Asia: Afghanistan, Bangladesh, Bhutan, India, Iran, Maldives, Nepal, Pakistan, Sri Lanka.

Asia (without South Asia): Armenia, Azerbaijan, Bahrain, Brunei, Cambodia, China, Cyprus, Georgia, Hong Kong, Indonesia, Iraq, Israel, Jordan, Japan, Kazakhstan, Kuwait, Kyrgyzstan, Laos, Lebanon, Mongolia, Malaysia, Myanmar, North Korea, Oman, Philippines, Qatar, Saudi Arabia, South Korea, Singapore, Palestine, Syria, Tajikistan, Thailand, Timor-Leste, Turkey, Turkmenistan, United Arab Emirates, Uzbekistan, Vietnam, Yemen.

Northern Africa: Algeria, Egypt, Libya, Morocco, Sudan, Tunisia, Western Sahara.


Sub-Saharan Africa: Angola, Benin, Botswana, Burkina Faso, Burundi, Cameroon, Cape Verde, Central African Republic, Chad, Comoros, Congo (Brazzaville), Côte d'Ivoire, Democratic Republic of the Congo, Djibouti, Equatorial Guinea, Eritrea, Ethiopia, Gabon, Ghana, Guinea, Guinea-Bissau, Kenya, Lesotho, Liberia, Madagascar, Malawi, Mali, Mauritania, Mauritius, Mozambique, Namibia, Niger, Nigeria, Reunion, Rwanda, São Tomé and Príncipe, Senegal, Seychelles, Sierra Leone, Somalia, South Africa, Sudan, Swaziland, Tanzania, The Gambia, Togo, Uganda, Zambia, Zimbabwe.

South and Central America: Anguilla, Antigua and Barbuda, Argentina, Aruba, Bahamas, Barbados, Belize, Bolivia, Bonaire, Sint Eustatius and Saba, Brazil, British Virgin Islands, Cayman Islands, Chile, Colombia, Costa Rica, Cuba, Curacao, Dominica, Dominican Republic, Ecuador, El Salvador, Falkland Islands, French Guiana, Grenada, Guadeloupe, Guatemala, Guyana, Haiti, Honduras, Jamaica, Martinique, Mexico, Montserrat, Nicaragua, Panama, Paraguay, Peru, Puerto Rico, Saint Kitts and Nevis, Saint Lucia, Saint Martin, Saint Vincent and the Grenadines, Saint-Barthelemy, Sint Maarten, Suriname, Trinidad and Tobago, Turks and Caicos Islands, US Virgin Islands, Uruguay, Venezuela.

North America: Bermuda, Canada, Greenland, Saint Pierre and Miquelon, United States.

Europe: Aland Islands, Albania, Andorra, Austria, Belarus, Belgium, Bosnia and Herzegovina, Bulgaria, Channel Islands, Croatia, Czech Republic, Denmark, Estonia, Faeroe Islands, Finland, the former Yugoslav Republic of Macedonia, France, Germany, Gibraltar, Greece, the Holy See, Hungary, Iceland, Ireland, Isle of Man, Italy, Latvia, Liechtenstein, Lithuania, Luxembourg, Malta, Monaco, Montenegro, Netherlands, Norway, Poland, Portugal, Moldova, Romania, Russia, San Marino, Serbia, Slovakia, Slovenia, Spain, Svalbard and Jan Mayen, Sweden, Switzerland, Ukraine, United Kingdom.

Oceania: American Samoa, Australia, Cook Islands, Fiji, French Polynesia, Guam, Kiribati, Marshall Islands, Micronesia, Nauru, New Caledonia, New Zealand, Niue, Norfolk Island, Northern Mariana Islands, Palau, Papua New Guinea, Pitcairn Islands, Samoa, Solomon Islands, Tokelau, Tonga, Tuvalu, Vanuatu, Wallis and Futuna.



22 **S2. PRISMA flow diagram for the selection of studies**

23
24 *28 studies comprising 20 datasets