

Data Papers

Ecology, 98(11), 2017, pp. 2979
© 2017 by the Ecological Society of America

ATLANTIC-CAMTRAPS: a dataset of medium and large terrestrial mammal communities in the Atlantic Forest of South America

FERNANDO LIMA,^{1,2,36} GABRIELLE BECA,¹ RENATA L. MUylaert,¹ CLINTON N. JENKINS,² MIRIAM L. L. PERILLI,³ ANA MARIA O. PASCHOAL,⁴ RODRIGO L. MASSARA,⁴ ADRIANO P. PAGLIA,⁴ ADRIANO G. CHIARELLO,⁵ MAURÍCIO E. GRAIPEL,⁶ JORGE J. CHEREM,⁷ ANDRÉ L. REGOLIN,¹ LUIZ GUSTAVO R. OLIVEIRA SANTOS,⁸ CARLOS R. BROCARDO,^{1,9} AGUSTÍN PAVIOLI,^{10,11} MARIO S. DI BITETTI,^{10,11,12} LEANDRO M. SCROSS,¹³ FABIANA L. ROCHA,¹⁴ ROBERTO FUSCO-COSTA,^{15,16} CLARISSA A. ROSA,¹⁷ MARINA X. DA SILVA,¹⁸ LUDMILA HUFNAGELL,⁴ PALOMA M. SANTOS,⁴ GABRIELA T. DUARTE,⁴ LUIZA N. GUIMARAES,⁴ LARISSA L. BAILEY,¹⁹ FLÁVIO HENRIQUE G. RODRIGUES,⁴ HEITOR M. CUNHA,²⁰ FELIPE M. FANTACINI,²¹ GRAZIELE O. BATISTA,²² JULIANO A. BOGONI,²³ MARCO A. TORTATO,²⁴ MICHELI R. LUIZ,²⁵ NIVALDO PERONI,⁶ PEDRO V. DE CASTILHO,²⁶ THIAGO B. MACCARINI,²⁷ VILMAR PICINATTO FILHO,²⁸ CARLOS DE ANGELO,^{10,11} PAULA CRUZ,^{10,11} VERÓNICA QUIROGA,^{10,11} MARÍA E. IEZZI,^{10,11} DIEGO VARELA,^{10,11} SANDRA M. C. CAVALCANTI,³ ALEXANDRE C. MARTENSEN,²⁹ ERICA V. MAGGIORINI,³⁰ FABIOLA F. KEESEN,³¹ ANDRÉ V. NUNES,²⁴ GISELE M. LESSA,³² PEDRO CORDEIRO-ESTRELA,³³ MAYARA G. BELTRÃO,^{33,34} ANNA CAROLINA F. DE ALBUQUERQUE,^{33,34} BIANCA INGBERMAN,¹⁵ CAMILA R. CASSANO,³⁵ LAURY CULLEN JUNIOR,² MILTON C. RIBEIRO,^{1,36} AND MAURO GALETTI^{1,36}

Abstract. Our understanding of mammal ecology has always been hindered by the difficulties of observing species in closed tropical forests. Camera trapping has become a major advance for monitoring terrestrial mammals in biodiversity rich ecosystems. Here we compiled one of the largest datasets of inventories of terrestrial mammal communities for the Neotropical region based on camera trapping studies. The dataset comprises 170 surveys of medium to large terrestrial mammals using camera traps conducted in 144 areas by 74 studies, covering six vegetation types of tropical and subtropical Atlantic Forest of South America (Brazil and Argentina), and present data on species composition and richness. The complete dataset comprises 53,438 independent records of 83 species of mammals, includes 10 species of marsupials, 15 rodents, 20 carnivores, eight ungulates and six armadillos. Species richness averaged 13 species (± 6.07 SD) per site. Only six species occurred in more than 50% of the sites: the domestic dog *Canis familiaris*, crab-eating fox *Cerdocyon thous*, tayra *Eira barbara*, south American coati *Nasua nasua*, crab-eating raccoon *Procyon cancrivorus* and the nine-banded armadillo *Dasypus novemcinctus*. The information contained in this dataset can be used to understand macroecological patterns of biodiversity, community, and population structure, but also to evaluate the ecological consequences of fragmentation, defaunation, and trophic interactions.

Key words: *Atlantic Forest; biodiversity hotspot; camera traps; forest fragmentation; invasive species; mammal communities; neotropical mammals.*

The complete data sets corresponding to abstracts published in the Data Papers section in the journal are published electronically as Supporting Information in the online version of this article at <http://onlinelibrary.wiley.com/doi/10.1002/ecy.1998/supinfo>

Manuscript received 5 May 2017; revised 31 July 2017; accepted 16 August 2017. Corresponding Editor: W. K. Michener.
³⁶E-mail: pardalismitis@gmail.com