

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

A novel strigolactone-miR156 module controls stomatal behaviour during drought recovery

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1764369> since 2020-12-16T10:54:06Z

Published version:

DOI:10.1111/pce.13758

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

1 **TITLE PAGE**

2 **i) Full title:**

3 **A novel strigolactone-miR156 module controls stomatal behaviour during drought recovery**

4

5 **ii) List of authors:**

6 Ivan Visentin¹ [ORCID: 0000 0002 5803 9034] ivan.visentin@unito.it

7 Chiara Pagliarani^{1, 2} [ORCID: 0000-0003-4656-6192] chiara.pagliarani@ipsp.cnr.it

8 Eleonora Deva^{1, 3} [ORCID: none] eleonora.deva@edu.unito.it

9 Alessio Caracci¹ [ORCID: none] alessio.caracci@edu.unito.it

10 Veronika Turečková⁴ [ORCID: none] hgradeckav@post.cz

11 Ondrej Novák⁴ [ORCID: 0000-0003-3452-0154] ondrej.novak@upol.cz

12 Claudio Lovisolo¹ [ORCID: 0000-0001-8825-2904] claudio.lovisolo@unito.it

13 Andrea Schubert¹ [ORCID: 0000-0001-7355-5865] andrea.schubert@unito.it

14 Francesca Cardinale^{1,*} [ORCID: 0000-0001-9892-9325] francesca.cardinale@unito.it

15 * **Corresponding author's further contact information:** phone +39 011 6708875

16

17 **iii) Authors' affiliations:**

18 ¹ Plant Stress Lab, DISAFA - Turin University, 10095 Grugliasco (TO) Italy

19 ² Current address: Institute for Sustainable Plant Protection, National Research Council, 10135
20 Torino, Italy

21 ³ StrigoLab Srl – 10125 Turin, Italy

22 ⁴ Laboratory of Growth Regulators, Centre of the Region Haná for Biotechnological and Agricultural
23 Research, Palacký University and Institute of Experimental Botany, Czech Academy of Sciences,
24 78371 Olomouc, Czech Republic

25

26 **iv) Funding**

27 The Ministry of Education, Youth and Sports, Czech Republic (Grant LO1204 from the National
28 Programme of Sustainability I) supported V.T. and O.N. The PlantStressLab team and E.D. have
29 received funding from the European Union's Horizon 2020 research and innovation programme
30 under the Grant Agreement No. [727929] (TOMRES). None of these funding sources were involved
31 in the design of the study and collection, analysis and interpretation of data, and in writing the
32 manuscript.

33

34 **v) Abstract**

35 miR156 is a conserved microRNA whose role and induction mechanisms under stress are poorly
36 known. Strigolactones are phytohormones needed in shoots for drought acclimation. They
37 promote stomatal closure ABA-dependently and independently; however, downstream effectors
38 for the former have not been identified. Linkage between miR156 and strigolactones under stress
39 has not been reported.

40 We compared ABA accumulation and sensitivity as well as performances of wt and miR156-
41 overexpressing (miR156-oe) tomato plants during drought. We also quantified miR156 levels in wt,
42 strigolactone-depleted and strigolactone-treated plants, exposed to drought stress.

43 Under irrigated conditions, miR156 overexpression and strigolactone treatment led to lower
44 stomatal conductance and higher ABA sensitivity. Exogenous strigolactones were sufficient for
45 miR156 accumulation in leaves, while endogenous strigolactones were required for miR156
46 induction by drought. The “after-effect” of drought, by which stomata do not completely re-open
47 after rewatering, was enhanced by both strigolactones and miR156. The transcript profiles of
48 several miR156 targets were altered in strigolactone-depleted plants.

49 Our results show that strigolactones act as a molecular link between drought and miR156 in
50 tomato, and identify miR156 as a mediator of ABA-dependent effect of strigolactones on the after-
51 effect of drought on stomata. Thus, we provide insights into both strigolactone and miR156 action
52 on stomata.

53

54 **vi) Key words**

55 Abscisic acid (ABA), After-effect of drought, Hormone signalling, Osmotic stress, *Solanum*
56 *lycopersicum*, Stomata, Stress-responsive microRNA

57

58 **vii) Running head: Novel strigolactone-miR156 module controls stomatal memory**

59

60 **viii) Acknowledgements:**

61 F.C. conceived of the work and designed research supported by I.V., C.L. and A.S.; I.V. performed
62 most of the experiments helped by C.P., A.C. and V.T.; I.V., C.P., C.L. and O.N. analysed data; I.V.
63 and F.C. wrote the article with contributions of all the authors, who read, commented and
64 approved the final manuscript. A.S. and F.C provided for financial support to the research work.

65 F.C. agrees to serve as the author responsible for contact and ensures communication. The
66 Ministry of Education, Youth and Sports, Czech Republic (Grant LO1204 from the National
67 Programme of Sustainability I) supported V.T. and O.N. The PlantStressLab team and E.D. have
68 received funding from the European Union's Horizon 2020 research and innovation programme
69 under the Grant Agreement No. [727929] (TOMRES). None of these funding sources were involved
70 in the design of the study and collection, analysis and interpretation of data, and in writing the
71 manuscript. The authors wish to thank Dr. Silva e Ferreira (Sao Paulo, Brazil) for sharing the
72 miR156-oe MicroTom line, Dr. Klee (University of Florida, USA) for the strigolactone-depleted line,
73 and Viviana Cisse and Cecilia Roche for technical support.

74

75

76 Word count, main body (Introduction to Acknowledgements): 5976

77 Number of figures: 6, of which 0 in colour

78 Number of supplementary figures: 1 of which 0 in colour

79 Number of supplementary tables: 1

80

81

82 **Main text**

83 **A novel strigolactone-miR156 module controls stomatal behaviour during drought recovery**

84

85 **ABSTRACT**

86 miR156 is a conserved microRNA whose role and induction mechanisms under stress are poorly
87 known. Strigolactones are phytohormones needed in shoots for drought acclimation. They
88 promote stomatal closure ABA-dependently and independently; however, downstream effectors
89 for the former have not been identified. Linkage between miR156 and strigolactones under stress
90 has not been reported.

91 We compared ABA accumulation and sensitivity as well as performances of wt and miR156-
92 overexpressing (miR156-oe) tomato plants during drought. We also quantified miR156 levels in wt,
93 strigolactone-depleted and strigolactone-treated plants, exposed to drought stress.

94 Under irrigated conditions, miR156 overexpression and strigolactone treatment led to lower
95 stomatal conductance and higher ABA sensitivity. Exogenous strigolactones were sufficient for
96 miR156 accumulation in leaves, while endogenous strigolactones were required for miR156
97 induction by drought. The “after-effect” of drought, by which stomata do not completely re-open
98 after rewatering, was enhanced by both strigolactones and miR156. The transcript profiles of
99 several miR156 targets were altered in strigolactone-depleted plants.

100 Our results show that strigolactones act as a molecular link between drought and miR156 in
101 tomato, and identify miR156 as a mediator of ABA-dependent effect of strigolactones on the after-
102 effect of drought on stomata. Thus, we provide insights into both strigolactone and miR156 action
103 on stomata.

104

105 **KEY WORDS**

106 Abscisic acid (ABA), After-effect of drought, Hormone signalling, Osmotic stress, *Solanum*
107 *lycopersicum*, Stomata, Stress-responsive microRNA

108

109 **INTRODUCTION**

110 It has been estimated that more than 90% of water uptake in plants is lost through transpiration.
111 Stomata represent a fundamental checkpoint balancing the entry of carbon dioxide and the exit of
112 water, and their regulation is at the core of the main acclimation strategy to water scarcity
113 (Matthews, Violet-Chabrand, & Lawson, 2017). Upon drought, tolerance mechanisms operate at

114 different spatial and temporal scales with rapid stomatal closure as the basis for preventing shoot
115 water loss (Tardieu, Simonneau, & Muller, 2018). MicroRNAs (miRNAs) and phytohormones have
116 been associated to the regulation of guard cell development and movement (Curaba, Singh, &
117 Bhalla, 2014; Ding, Tao, & Zhu, 2013).

118 miRNAs are a widespread class of endogenous, small RNA molecules (19-24 nt in length) that
119 negatively regulate gene expression at the transcriptional, post-transcriptional, and translational
120 levels (Nozawa, Miura, & Nei, 2012). In animals, under certain conditions, miRNAs are being
121 looked at as hormones, because of their cell-to-cell and also long-distance movement coupled to
122 signalling activity (Bayraktar, Van Roosbroeck, & Calin, 2017). While some plant miRNAs are
123 species-specific in terms of expression patterns and targets, others, including miR156, are very
124 conserved. The modulation of miR156 is crucial throughout development for correct leaf
125 formation, tillering/branching, plastochron, panicle/tassel architecture, and timing of age-
126 dependent reproductive transition along with fruit ripening and fertility (Wang & Wang, 2015). In
127 spite of a few differences among species, the pattern of mature miR156 accumulation in response
128 to environmental stimuli is also rather conserved (Khraiwesh, Zhu, & Zhu, 2012). Indeed, miR156 is
129 consistently induced by a variety of abiotic stresses such as drought, osmotic stress, heat, cold,
130 salinity, and macro-nutrient deficiency (Cui, Shan, Shi, Gao, & Lin, 2014; Ding, Fromm, &
131 Avramova, 2012; Hsieh et al., 2009; H. H. Liu, Tian, Li, Wu, & Zheng, 2008; Stief et al., 2014). As for
132 its role under stress, it was initially postulated that miR156 increase is merely needed to stall
133 flowering until stress is over. This effect would be exerted via miR156-mediated inhibition of the
134 age-dependent pathway to flowering. In support of this view, miR156-overexpressing plants
135 (miR156-oe hereafter) are late-flowering. Under long-day, inductive conditions the flowering time
136 in *Arabidopsis* (*Arabidopsis thaliana*) tends to inversely correlate with miR156 levels in a range of
137 environmental conditions (Cui et al., 2014; May et al., 2013). However, it is also becoming clear
138 that this miRNA may have more direct functions in stress acclimation. For instance, it was shown
139 that miR156-oe plants outperform wt controls both in *Medicago sativa* under drought (Arshad,
140 Feyissa, Amyot, Aung, & Hannoufa, 2017) and in *Arabidopsis* under osmotic/high salinity stress
141 (Cui et al., 2014) or recurring heat (Stief et al., 2014). These functions may be exerted by miR156
142 through post-transcriptional repression of one or more members of the wide family of SPL
143 (SQUAMOSA PROMOTER BINDING PROTEIN-LIKE) transcription factors (Arshad et al., 2017; Cui et
144 al., 2014; Stief et al., 2014), the best characterised among miR156 targets. However, these studies

145 do not clarify which cascade of molecular events induces miR156 under stress, and whether
146 miR156 can affect stomatal regulation.

147 Abscisic acid (ABA) is the best-characterised phytohormone among several affecting stomatal
148 functioning. Its direct application, as well as endogenous synthesis in phloem companion,
149 mesophyll and guard cells, trigger stomatal closure (Bauer et al., 2013; Kuromori, Sugimoto, &
150 Shinozaki, 2014; S. A. M. McAdam & Brodribb, 2018; Merilo et al., 2018). ABA signalling is known
151 to affect miRNA production (Lian et al., 2018; Speth, Willing, Rausch, Schneeberger, & Laubinger,
152 2013; Yan et al., 2017), but to our knowledge, no miRNAs have been demonstrated to affect ABA
153 sensitivity.

154 Strigolactones, the most recently discovered class of phytohormones, also play a role in this
155 scenario. ABA and strigolactones share a carotenoid precursor, from which strigolactone synthesis
156 proceeds through a partially known series of enzymes including DWARF27 (D27), CAROTENOID
157 CLEAVAGE DIOXYGENASE7 (CCD7) and 8 (CCD8), and MORE AXILLARY GROWTH1 (MAX1) to
158 produce bioactive strigolactones (Waters, Gutjahr, Bennett, & Nelson, 2017). Strigolactones
159 modulate several aspects of plant development and interactions with rhizosphere organisms
160 (Andreo-Jimenez, Ruyter-Spira, Bouwmeester, & Lopez-Raez, 2015; Cardinale, Korwin Krukowski,
161 Schubert, & Visentin, 2018; Lanfranco, Fiorilli, Venice, & Bonfante, 2018; Lopez-Raez, 2016).
162 Furthermore, plants with defective strigolactone synthesis or signalling in several dicot plants are
163 hypersensitive to drought, salt and osmotic stress (Cardinale et al., 2018). In *Arabidopsis*, *Lotus*
164 *japonicus* and tomato, strigolactones positively control stomatal movements as components of a
165 systemic stress signal (Tardieu, 2016). Indeed, mutants in strigolactone biosynthesis exhibited
166 reduced stomatal closure (Ha et al., 2014; J. Liu et al., 2015; Visentin et al., 2016). Conversely,
167 enhanced stomatal closure and drought tolerance was observed in plants treated with exogenous
168 strigolactones (Lv et al., 2018; Visentin et al., 2016; Y. Zhang, Lv, & Wang, 2018) or in which shoot
169 strigolactone biosynthesis is increased by grafting onto a low-strigolactone rootstock (Visentin et
170 al., 2016). The effect of strigolactones on stomatal closure depends, at least in part, on ABA
171 synthesis, transport and/or sensitivity. Accordingly, strigolactone depletion decreases sensitivity to
172 exogenous ABA in several species (Bu et al., 2014; Ha et al., 2014; J. Liu et al., 2015; Lv et al., 2018;
173 Visentin et al., 2016) and sensitivity to endogenous ABA in stressed tomato (Visentin et al., 2016).
174 On the other hand, treatment with the synthetic strigolactone analogue racemic GR24 (*rac*-GR24)
175 increases sensitivity to ABA in tomato (Visentin et al., 2016). It deserves attention that the above-
176 described model may be confined to dicot plants, as it does not apply to rice – most strigolactone-

177 biosynthetic mutants of which produce more ABA than the wild-type (wt) and thus are more
178 resistant to drought (Haider et al., 2018).

179 In this varied landscape, the possibility of a functional connection between miR156 and
180 strigolactones in the regulation of stomatal movements and drought avoidance has never been
181 investigated. miR156g was *in silico* predicted to directly target the transcripts of the strigolactone
182 biosynthetic gene *MAX1* in Arabidopsis (Marzec & Muszynska, 2015), but experimental proof of
183 this is currently lacking. Additionally, a functional link in the context of shoot development, via
184 stabilisation of specific SPL proteins by strigolactones, is already known in gramineous plants (J.
185 Liu, Cheng, Liu, & Sun, 2017; Song et al., 2017). The fact that the transcripts of several SPL factors
186 are targeted by miR156 offers a potential integration point between hormone- and miRNA-
187 mediated signalling (Kerr & Beveridge, 2017; M. Liu et al., 2017; Song et al., 2017). Nevertheless, it
188 is still unknown whether this or a similar mechanism is operational under stress.

189 In this work, we used different approaches to clarify the link between miR156 and strigolactones
190 with regards to stomatal regulation in tomato. We investigated the effect of miR156
191 overexpression on stomatal function and ABA metabolism/sensitivity as well as the transcript
192 stability of strigolactone biosynthetic genes. Moreover, the effect of strigolactones on miR156
193 levels were assessed by i) treatment with exogenous strigolactones coupled to the use of a
194 strigolactone-depleted transgenic line, and ii) the application of drought stress to increase both
195 strigolactone and miR156 levels in shoots. The results identified strigolactones as a molecular
196 component linking drought to miR156 accumulation. Furthermore, they allowed us to integrate
197 miR156 in a model that describes the connections between strigolactones and ABA in tomato
198 (Cardinale et al., 2018), thus offering insights into both strigolactone and miR156 action on
199 stomata.

200

201 **MATERIALS AND METHODS**

202 **Plant materials and growth conditions**

203 The tomato *CCD7*-silenced line 6936 (kind gift by Dr H. J. Klee, University of Florida) (Vogel et al.,
204 2010) and its wt M82 were grown in a growth room set at the following conditions: 16/8 h
205 day/night cycle, 25°C, 65% humidity, and 200 $\mu\text{mol s}^{-1} \text{m}^{-2}$ of photosynthetic photon flux density
206 (PPFD). Seeds were sterilized in 4% (v/v) sodium hypochlorite containing 0.02% (v/v) Tween 20,
207 rinsed thoroughly with sterile water, and plated on MS medium with 0.8% w:v agar, pH 5.9. Ten
208 days after, seedlings were transferred to an inert substrate and pots were watered with Hoagland

209 solution twice per week. Drought-stress kinetics were performed by uprooting plants of both wt
210 and *CCD7*-silenced genotypes and transferring them in vermiculite hydrated with only 20 ml of
211 water (stressed group), or transferred in wet vermiculite (well-watered controls). A sub-group of
212 wt plants were leaf-sprayed with GR24^{5DS} (see dedicated paragraph below) 24 h before the
213 beginning of stress, while all other plants were mock-treated with a solution of 0.01% v/v acetone
214 in water. Throughout the experiment, each condition and genotype was represented by 5 plants.
215 The tomato miR156-oe line overexpressing the *AtMIR156b* primary transcript and its wt
216 (MicroTom) were kindly provided by Dr. G. Silva (University of Sao Paulo, Brazil) (Ferreira e Silva et
217 al., 2014). Plants were grown in 1-liter pots filled with a commercial substrate (Terra Nature, NPK
218 12:14:24 imported by Raw Materials Europe, NL) composed of sandy-loam soil/expanded
219 clay/peat mixture (2:1:1 by weight) and maintained under greenhouse conditions. Two drought
220 time-courses were performed on 4-week-old plants by withholding water for 15 days - starting
221 from day zero - on 5 plants per line and treatment; an irrigated control group was kept for each
222 genotype. After 10 days, once severe water stress levels were reached, the plants were watered to
223 allow for recovery. Stomatal conductance was measured throughout the experiment, while leaf
224 samples were collected for water potential measurements at day zero, ten and fifteen (see below
225 for physiological analytical methods).

226

227 **Treatment with exogenous strigolactones**

228 A 5 μ M solution of GR24^{5DS} (StrigoLab Srl, Turin, Italy) in 0.01% v/v acetone in water was sprayed
229 on leaves of unstressed plants until runoff, while control plants were sprayed with a
230 corresponding solution of acetone only. This pure enantiomeric form of the synthetic strigolactone
231 analogue GR24 was preferred over commercial *rac*-GR24 due to the possibly confounding
232 bioactivity by the other enantiomer (GR24^{ent-5DS}) contained in the racemic mixture (Scaffidi et al.,
233 2014). Stomatal conductance (see below) was measured 2 and 24 h after treatment, while
234 samples for the quantification of mature miR156 were collected 2, 6 and 24 h after treatment (in
235 the absence of stress), deep frozen and stored at -80°C until analysis. When GR24^{5DS} treatment
236 was imposed on wt plants to be subsequently stressed, harvesting times were i) 24 h after
237 treatment (well-watered samples; with stress beginning immediately after harvest); ii) after 3 h of
238 stress (water-stressed samples; with re-watering and the beginning of recovery happening
239 immediately after harvest); iii) 24 h after the beginning of stress, i.e. 21 h into recovery (recovered
240 samples).

241

242 **Gene-transcript quantification and miR156 target-site detection**

243 For transcript quantification from axillary buds, at least 30 stem sections were excised at the level
244 of leaf insertion from 10-week-old MicroTom plants (eight plants each, for wt and miR156-oe
245 plants) to obtain two lots of 50 mg (fw) per genotype, and freeze-dried until analysis. For
246 quantification in shoot tissues, the same procedure was applied on 200 mg (fw) of tomato leaves.
247 Total RNA was extracted by using Spectrum™ Plant Total RNA Kit (SIGMA), and treated with DNase
248 I (ThermoScientific) at 37°C for 30 min to remove residual genomic DNA. First-strand cDNA was
249 synthesized from 1 µg of purified total RNA using the High-Capacity cDNA Reverse Transcription
250 Kit (Applied Biosystems, Monza, Italy) according to the manufacturer's instructions.

251 For targeted miR156 cDNA synthesis, a modified protocol with a stem-loop primer (Pagliarani et
252 al., 2017) was followed in samples of wt M82 and *CCD7*-silenced plants obtained from the
253 drought-stress experiments described above. For transcript quantification of candidate genes by
254 quantitative reverse-transcription PCR (qRT-PCR), random primers were used to reverse transcribe
255 total RNA. qRT-PCR analysis was carried out in a StepOnePlus system (Applied Biosystems) using
256 the SYBR Green (Applied Biosystems) method on 10 ng of cDNA (50 ng for *S/CCD7* transcripts). For
257 loci and primers (which were used at 400 nM), see Table S1. Transcript concentrations were
258 normalised on the geometric mean of *SlsnRU6* and *SIEF-1α* transcript concentrations used as
259 endogenous controls. Three independent biological replicates were analysed as a minimum, and
260 each qRT-PCR reaction was run in technical triplicates. Transcript amounts were quantified by the
261 $2^{-\Delta\Delta Ct}$ method. Putative target genes of miR156 were predicted *in silico* using the psRNATarget
262 algorithm with default setting parameters (<http://plantgrn.noble.org/psRNATarget/>) (Dai & Zhao,
263 2011).

264

265 **Stomatal conductance and aperture measurements**

266 During the drought-stress time-course on miR156-oe plants and their MicroTom wt, stomatal
267 conductance was measured daily between 10:00 and 12:00 am on two randomly selected, fully
268 developed apical leaves for each plant with a portable system (SC-1 Leaf Porometer for Stomatal
269 Conductance Measurements, Decagon Device, WA, USA). Leaf water potential was measured
270 using a Scholander-type pressure chamber (Soil Moisture Equipment Corp., Santa Barbara, CA,
271 USA) (Scholander, Bradstreet, Hemmingsen, & Hammel, 1965) on one leaf per plant, immediately
272 after gas exchange quantification. During the drought stress time-course on *CCD7*-silenced plants

273 and their M82 wt, stomatal conductance was measured on two randomly selected, fully
274 developed apical leaves for each plant with the same portable system as above. The selected time-
275 points for full measurements were: i) immediately before stress start, between 8:00 and 10:00 am;
276 ii) 3 h later, when stomatal conductance values were about 20% of the irrigated controls (after
277 which, the plants were watered to start recovering); iii) 6 and 24 h after the beginning of stress,
278 i.e. 3 and 21 h into the recovery period. In order to quantify guard-cell reactivity to exogenous ABA
279 treatments, leaves of 4-week-old wt (MicroTom) and miR156-oe plants were sprayed with varying
280 concentrations of ABA in water or with water alone until drip-off, then let dry for 1 h before
281 quantifying stomatal conductance as above. For the quantification of stomatal pore areas in the
282 two genotypes, mature excised leaves were pre-incubated in a MES-KCl buffer (10 mM MES-
283 KOH/50 mM KCl, pH 6.15) under light for 2 h to promote stomatal opening before treatments.
284 Leaves of three plants per line (wt and miR156-oe) were then incubated for 3 h in MES-KCl buffer
285 containing 10 μ M ABA. Subsequently, the abaxial epidermis of ABA-treated and untreated leaves
286 of both genotypes was peeled off following a published procedure (Hopper, Ghan, & Cramer,
287 2014), and stomatal apertures were measured with the ImageJ software on images captured using
288 a calibrated light microscope. The aperture areas of a minimum of 30 stomata from three
289 independent leaves per treatment were quantified.

290

291 **Extraction and quantification of ABA**

292 Plant tissues (each sample approximately 20 mg fw) were ground using 3-mm tungsten carbide
293 beads (Retsch GmbH & Co. KG, Haan, Germany) with a MM 301 vibration mill at a frequency of
294 27.0 Hz for 3 min (Retsch GmbH & Co. KG, Haan, Germany). One ml ice-cold
295 methanol/water/acetic acid (10/89/1, v/v) and internal standard mixtures [containing 20 pmol of
296 each of (-)-7',7',7'-²H₃-phaseic acid; (-)-7',7',7'-²H₃-dihydrophaseic acid; (-)-8',8',8'-²H₃-
297 neophaseic acid; (+)-4,5,8',8',8'-²H₅-ABA-GE; (-)-5,8',8',8'-²H₄-7'-OH-ABA; (+)-3',5',5',7',7',7'-²H₆-
298 ABA] were added to each sample. After 1 h shaking in the dark at 4°C, the homogenates were
299 centrifuged (20 000 *g*, 10 min, 4°C), and the pellets were then re-extracted in 0.5 ml extraction
300 solvent for 30 min. The combined extracts were purified by solid-phase extraction on Oasis® HLB
301 cartridges (60 mg, 3 ml, Waters, Milford, MA, USA), then evaporated to dryness in a Speed-Vac
302 (UniEquip) and finally analysed by UPLC-ESI(-/+)-MS/MS (Turečková, Novák, & Strnad, 2009).

303

304 **Statistical analysis**

305 Significant differences among treatments were statistically analysed by applying a one-way
306 ANOVA test, and Tukey's *HSD* post-hoc test was used for mean separation when ANOVA results
307 were significant ($P < 0.05$). Significant differences of pairwise comparisons were assessed by the
308 Student's *t*-test. The SPSS statistical software package (SPSS Inc., Cary, NC, USA, v.22) was used.

309

310 **RESULTS**

311 **miR156-oe plants have lower stomatal conductance and acclimate better to drought**

312 To test the hypothesis that miR156 induction under drought is relevant to the plant water balance,
313 the physiological responses of wt (MicroTom) and miR156-oe plants were analysed during a
314 drought and recovery time-course. Well-watered miR156-oe plants displayed a significantly lower
315 stomatal conductance than wt (day zero in Fig. 1A), while water potential in the same leaves did
316 not significantly differ between the two genotypes (well-watered samples in Fig. 1B). This pattern
317 of higher stomatal conductance in the wt was maintained throughout the experiment for the
318 irrigated controls. In stressed plants, miR156 overexpression correlated with lower stomatal
319 conductance in the initial phases of the dehydration kinetics (day two, Fig 1A). At the time of the
320 most severe drought stress level the situation reversed, as stomatal conductance of miR156-oe
321 plants was slightly but significantly higher than wt (day seven and ten, Fig. 1A). At this stage, leaf
322 water potential was significantly lower in wt than miR156-oe plants (water-stressed samples in Fig
323 1B, corresponding to day ten in Fig. 1A). At day fifteen, i.e. five days after re-irrigation, leaf water
324 potential of both wt and miR156-oe plants had recovered to levels comparable to those preceding
325 the drought spell (recovered samples in Fig. 1B). However, unlike wt plants, stomatal conductance
326 of miR156-oe tomato plants had completely failed to recover at this time (day fifteen, Fig. 1A).
327 These data suggest that: i) miR156 overexpression can lead to better acclimation to drought; ii)
328 the guard cells of miR156-oe plants respond more slowly to changes in water availability than
329 those of wt; and iii) under irrigated conditions, stomatal conductance is most likely controlled by a
330 non-hydraulic, miR156-related signal in tomato.

331

332 **The low stomatal conductance of miR156-oe plants is not due to increased free ABA**

333 Considering the pivotal role of ABA in stomatal closure, we reasoned that the physiological effects
334 of miR156 overexpression on stomatal opening observed in the above experiments may be due to
335 increased ABA levels. Hence, we set to quantify ABA and its main catabolites in the leaves of wt
336 and miR156-oe plants under well-watered conditions (day zero in Figure 1). Unexpectedly given

337 their low-transpiration phenotype, free ABA was significantly less concentrated in the leaves of
338 miR156-oe plants than in wt (Fig. 2A). Accordingly, transcripts of the ABA biosynthetic gene
339 *SINCE1* were somewhat lower in miR156-oe plants, albeit not significantly (Fig. 2B). While
340 glycosylated ABA levels were similar in wt and miR156-oe plants, the ABA catabolite phaseic acid
341 showed a non-significant trend towards higher concentrations in miR156-oe plants than in the wt
342 (Fig. 2A). Consistently, transcript amounts of the ABA-hydroxylating genes *SICYP707A1* and
343 *SICYP707A2*, which catalyse a key step in phaseic acid production from ABA, were significantly
344 higher in miR156-oe than wt leaves (Fig. 2B). Other related metabolites, such as dihydro-phaseic
345 acid, neo-phaseic acid and 7'-hydroxy-ABA, were below the limit of detection in both genotypes
346 (n.d. in Fig. 2A). Thus, miR156 overexpression seems to decrease endogenous ABA levels, at least
347 partly by inducing degradation. These data therefore exclude that the low stomatal conductance
348 rates observed in miR156-oe plants under well-watered conditions are due to increased ABA
349 content in total leaf tissues.

350

351 **miR156 promotes stomatal closure in response to ABA**

352 Since leaves of the miR156-oe genotype contained less free ABA than those of the wt and yet they
353 had more closed stomata, we hypothesized that miR156 may increase guard cell sensitivity to
354 ABA. To further address this point, we quantified stomatal conductance in wt and miR156-oe
355 plants 1 h after treatment with increasing concentrations of ABA, in comparison with the
356 corresponding mock treatment with water. The results clearly showed that unlike the wt, guard
357 cells of miR156-oe plants are oversensitive to ABA. In fact, in miR156-oe plants the decrease in
358 average conductance values was already significant at 1 μ M ABA concentration. Also at higher ABA
359 concentrations the decrease of stomatal conductance relative to the mock-treated controls was
360 more marked in miR156-oe plants than in the wt (Fig. 3). To confirm physiological data at the
361 morphological level, we as well assessed stomatal closure by determining stomatal pore areas
362 after 1 h exposure to 10 μ M exogenous ABA. In the absence of ABA in the floating medium,
363 stomatal pore areas were significantly wider in wt than miR156-oe leaves (Fig. S1A), confirming
364 the stomatal conductance differences occurring between the two genotypes under unstressed and
365 untreated conditions (Fig. 1 and 3). Upon ABA treatment, average stomatal pore areas in miR156-
366 oe plants decreased by about 60% with respect to non-treated stomata in the same genotype. By
367 contrast, mean stomatal aperture in wt plants decreased by about 30% only, compared to the
368 untreated control (Fig. S1B and C). These results indicate that guard cells have higher sensitivity to

369 exogenous ABA in miR156-oe than wt plants. This in turn supports the notion that the lower
370 stomatal conductance in well-watered miR156-oe vs wt plants might be due, at least partly, to
371 higher sensitivity to endogenous ABA, which would more than compensate for the lower ABA
372 content.

373

374 **Strigolactones are needed and sufficient for miR156 induction in tomato, both under normal** 375 **conditions and under drought**

376 As outlined in the introduction, both the miR156 and strigolactone biosynthetic pathways are
377 induced by drought in leaves; both are needed for full resistance to abiotic stress; and both
378 enhance guard cell sensitivity to ABA in tomato. This was shown above for miR156 and in a
379 previous work for strigolactones (Visentin et al., 2016). Thus, we set to explore if not only a
380 correlation but also a causal link exists between strigolactones and miR156 that may be relevant
381 to stomatal functioning.

382 As a first approach, we compared the levels of mature miR156 in leaf tissues 2, 6 and 24 h after
383 treatment with the strigolactone analogue GR24^{5DS} with mock-treated controls. Results from this
384 experiment indicate that exogenous strigolactones are sufficient to increase the concentration of
385 the mature miRNA starting a few hours after treatment and up to 24 h later (Fig. 4A).

386 We next tested whether endogenous strigolactones are required for miR156 induction under
387 stress, by quantifying mature miR156 in leaves of wt (M82) and *CCD7*-silenced tomato plants
388 under irrigated and stress conditions. Severe stress levels are achieved quickly (within a few hours)
389 in this experimental set-up, by transferring the plants to dry or wet vermiculite substrate for the
390 stressed and control group, respectively. The two genotypes showed a very similar, low
391 concentration of miR156 under irrigated conditions, while wt leaves treated 24 h earlier with
392 GR24^{5DS} had higher concentrations than mock-treated leaves (Fig. 4B), consistently with the
393 results in Fig. 4A. At the most severe stress point, the concentration of mature miR156 increased
394 in wt leaves, as expected. Conversely, in plants impaired in strigolactone biosynthesis, no
395 induction could be observed (water-stressed samples in Fig. 4B). Additionally, the results in Fig. 4B
396 clearly show a powerful synergic effect of GR24^{5DS} pre-treatment and drought on mature miR156
397 levels. This is obvious both in stressed samples, and especially after re-watering (for 24 h in the
398 recovered samples in Fig. 4B), when miR156 amounts decrease towards pre-stress levels in the
399 leaves of mock-treated wt plants, , while they steadily increase up to 450 folds in the GR24^{5DS}-pre-
400 treated leaves.

401

402 **Strigolactones promote sustained stomatal closure (“after effect”) during recovery from drought**

403 Based on the above data, miR156 appears to promote the after-effect of drought, and its
404 inducibility by stress seems to completely depend on strigolactones in tomato. Therefore, we
405 reasoned that if the strigolactone-miR156 module is operational in guard cells, then strigolactones
406 should positively affect the extent of the after-effect of drought as well. To elucidate this point,
407 stomatal conductance data (Fig. 4C) were collected during the same experiment reported in Fig.
408 4B. Data showed clearly that GR24^{5DS}-treated plants displayed an enhanced after-effect of
409 drought, thus mimicking the physiological response of miR156-oe plants. Consistently, *CCD7*-
410 silenced plants did the opposite and recovered their stomatal conductance faster than wt plants.
411 Overall, these data attested that strigolactones promote sustained stomatal closure during
412 recovery from drought (the so-called “after-effect”) in tomato, as miR156 does.

413 Finally, we tested whether exogenous strigolactones could directly induce stomatal closure. In
414 *Arabidopsis*, treatment with *rac*-GR24 induces a fast stomatal closure that has been demonstrated
415 to be ABA-independent (Lv et al., 2018). Thus, we measured stomatal conductance in unstressed,
416 wt tomato leaves treated with GR24^{5DS} compared with the mock-treated controls. Our results
417 clearly showed that stomatal conductance levels of wt tomato decreased significantly within 2 h of
418 GR24^{5DS} treatment, while they they started recovering at 24 h (Fig. 4D).

419

420 **Strigolactones affect the transcript accumulation profiles of putative SPL factors**

421 The above results, together with previous observations on the activation of the strigolactone
422 biosynthetic pathway in leaves under drought (Visentin et al., 2016), suggested that a
423 strigolactone-miR156 module does exist in tomato. This module may be fully operational under
424 and after drought, thanks to an initial stress-induced increase of leaf strigolactones. If so, then we
425 should be able to identify transcripts of putative miR156 targets that are dysregulated, in a stress-
426 and strigolactone-dependent fashion. Among all loci encoding putative SPL factors in tomato
427 (Salinas, Xing, Hohmann, Berndtgen, & Huijser, 2012), four (*Solyc03g114850*, *Solyc07g062980*,
428 *Solyc10g009080* and *Solyc04g045560*) have been reported as candidate targets of miR156 under
429 drought (M. Liu et al., 2017). Accordingly, in our experiments, the transcripts of all of them
430 followed a decreasing trend in droughted wt tomatoes. However, there was no difference
431 between wt and *CCD7*-silenced leaves at this time-point, suggesting that neither strigolactones nor
432 miR156-mediated post-transcriptional gene silencing are needed for this decrease. Conversely

433 during recovery, accumulation patterns of *Solyc07g062980*, *Solyc10g009080* and *Solyc04g045560*
434 in wt and *CCD7*-silenced plants diverged from those of miR156. This finding is consistent with the
435 hypothesis that after drought - i.e. when the stomatal after-effect becomes apparent - these
436 transcripts are destabilised by miR156, which in turn is induced by strigolactones (Fig. 5). On the
437 contrary, the transcript profiles of *COLORLESS NON-RIPENING (CNR, Solyc02g077920)* and of the
438 orthologue of *SPL9/15* in Arabidopsis (*Solyc10g078700*) (Silva et al., 2019) showed no significant
439 changes during drought and they both increased during recovery in the two genotypes, mirroring
440 the related miRNA expression trend (Fig. 5). A significant increase of steady-state transcript
441 concentrations of both *CNR* and *SPL9/15* was instead observed in *CCD7*-silenced plants under pre-
442 stress conditions. It should be noted here that transcripts from all of these genes are validated
443 targets of miR156 in unstressed wt plants, and they are almost undetectable in miR156-oe plant
444 tissues (Silva et al., 2019; X. Zhang et al., 2011).

445

446 **miR156 does not target the transcripts of strigolactone-biosynthetic genes for degradation in** 447 **tomato**

448 The transcripts of the strigolactone-biosynthetic gene *MAX1* have been predicted *in silico* to be
449 direct targets of miR156 in Arabidopsis (Marzec & Muszynska, 2015). This would in principle lead
450 to transcript degradation and thus lower strigolactone production in miR156-oe plants. Indeed, in
451 axillary buds of miR156-oe potatoes, the strigolactone content was lower than in wt (Bhogale et
452 al., 2014), but the molecular underpinnings were not investigated at that time. To search for all
453 possible links between miR156 and strigolactones, and given that strigolactones are known to
454 feedback regulate their own synthesis at the transcriptional level, we also checked whether
455 miR156 might affect the stability of strigolactone biosynthetic genes in tomato. We addressed this
456 issue *in silico* first, by searching for miR156 target sequences on transcripts of *D27*, *CCD7*, *CCD8*
457 and *MAX1* in Arabidopsis and tomato. However, no acceptable predictions satisfying the
458 requirements for identification of miRNA targets in plants were obtained (Axtell & Meyers, 2018;
459 Dai & Zhao, 2011; Jones-Rhoades & Bartel, 2004). This contrasts with available information for
460 *MAX1* and miR156g in Arabidopsis (Marzec & Muszynska, 2015).

461 Additionally, we quantified transcripts of strigolactone-biosynthetic genes in roots, where they are
462 mostly expressed, and axillary buds of miR156-oe and wt plants. However, no evidence of miR156-
463 driven transcript destabilisation was found, and even higher transcript concentrations in the
464 miR156-oe plants was observed for some genes in either tissue (Fig. 6A and B). Therefore,

465 negative regulation by miR156 on strigolactone biosynthetic genes at the transcript stability level
466 was not observed in tomato. Of course, indirect effects at the protein or metabolite level cannot
467 be excluded at this stage. Nonetheless, the latter information would not change the answer to the
468 question being asked here, i.e. if miR156 directly destabilizes the transcripts of strigolactone-
469 biosynthetic genes, as suggested by Marzec and Muszynska (2015).

470

471 **DISCUSSION**

472 **miR156 induction by drought requires strigolactones in tomato**

473 Almost nothing is known about the molecular cues modulating miR156 levels under any
474 conditions. Even though specific changes in the epigenetic landscape at the *MIR156a* and *c* loci are
475 reported to be important for correct miR156 expression during Arabidopsis development (Xu,
476 Zhang, & Wu, 2018), no information are available about the signalling path connecting it to stress.
477 Partly filling this gap, we found that miR156 induction by drought was completely dependent on
478 the efficient synthesis of endogenous strigolactones. We also observed that exogenous
479 strigolactones increased mature miR156 levels in tomato leaves, both in the absence and
480 especially in combination with drought stress. These findings make strigolactones the first
481 identified molecular component in the drought-triggered pathway leading to miR156 induction,
482 and to its stress-related effects (Cui et al., 2014; Stief et al., 2014).

483

484 **The role of the strigolactone-miR156 module in stomata**

485 Given the demonstrated effects of strigolactones on stomata in several plant species (Bu et al.,
486 2014; Ha et al., 2014; J. Liu et al., 2015; Lv et al., 2018; Visentin et al., 2016), and the above
487 described functional connection between strigolactones and miR156, we sought to investigate
488 whether miR156 may act at the stomatal level specifically. When we compared the physiological
489 performances of miR156-oe with those of wt plants, stomatal conductance of the former was
490 indeed lower under irrigated conditions, suggesting a positive control of miR156 on stomatal
491 closure. This trait of miR156-oe plants is opposite to strigolactone-related mutants, and
492 reminiscent of tomato plants whose shoots experience high strigolactones. This latter situation
493 can be achieved either by treating with exogenous strigolactones (which induce miR156, as stress
494 does), or grafting a wt scion onto a strigolactone-depleted rootstock (which induces higher
495 strigolactone biosynthesis in the shoot, higher ABA sensitivity in guard cells and lower stomatal
496 conductance) (Ha et al., 2014; J. Liu et al., 2015; Visentin et al., 2016). It is noteworthy that in

497 miRNA156-oe plants subjected to severe stress, stomatal conductivity was higher than in the wt
498 (as observed in miR156-oe *M. sativa* by Arshad et al., 2017), reversing the pattern observed upon
499 irrigation. This could be explained by non-ABA-dependent signal(s) acting downstream of hydraulic
500 signals generated by severe drought. Consequently, miR156-oe plants lost less water under severe
501 drought. This was attested by their water potential dropping less than wt, and by the fact that
502 their better water status overweighed higher ABA sensitivity in terms of stomatal closure.
503 The effects of the strigolactone-miR156 module under recovery might be executed via the post-
504 transcriptional repression of SPL genes such as *Solyc10g009080*, which most convincingly showed
505 a divergent transcript profile with miR156 in this phase. Their ultimate function under stress is
506 worthy of further experimental investigation. It should be noted here that, in drought stress
507 conditions, the transcripts of all four SPL genes suggested to be miR156 targets decreased in
508 *CCD7*-silenced leaves as much as in the wt. As in these transgenic tissues miR156 levels did not rise
509 as a consequence of drought, we must conclude that their regulation during stress is more likely to
510 occur at the transcriptional rather than post-transcriptional level, contrarily to previous
511 suggestions (M. Liu et al., 2017).

512

513 **miR156 is a possible mediator of ABA-dependent effects of strigolactones on stomata**

514 Strigolactone effects on stomata are known to be both ABA-dependent and independent. Indeed,
515 treatment with exogenous strigolactones induces fast, ABA-independent closure of stomata but
516 also higher sensitivity to ABA (Brun et al., 2019; Li et al., 2017; J. Liu et al., 2015; Lv et al., 2018;
517 Visentin et al., 2016; Y. Zhang et al., 2018), while strigolactone-related mutants have impaired
518 ABA-dependent responses to osmotic stress (reviewed in Cardinale et al., 2018).

519 We found several indications rather pointing to miR156 affecting the ABA-dependent subset of
520 strigolactone effects on stomata, under irrigated conditions. Firstly, stomatal closure in response
521 to exogenous strigolactone treatment (ABA-independent in *Arabidopsis*) (Lv et al., 2018) was
522 achieved within 2 h in tomato, when miR156 induction was not significant yet; and started
523 decreasing by 24 h, when miR156 levels were highest (Fig. 4A vs Fig. 4D). Secondly, under irrigated
524 conditions, miR156-oe plants had lower stomatal conductance but also significantly less free ABA
525 in leaves, possibly due to weaker biosynthesis and especially accelerated ABA conversion to
526 phaseic acid. Accordingly, the activation of ABA catabolism is a reported effect of increased
527 strigolactone synthesis in the shoot, and of exogenous strigolactone treatment in different tissues
528 (Ferrero et al., 2018; Lechat et al., 2012; Toh et al., 2012; Visentin et al., 2016). Thirdly, stomatal

529 closure by exogenous ABA was more complete in miR156-oe plants than in wt, which may be
530 explained by higher sensitivity to, or more effective transport of ABA. Higher ABA sensitivity would
531 more than compensate lower ABA levels and justify reduced stomatal conductance in miR156-oe
532 plants. As in other priming phenomena, the strigolactone-dependent increase in ABA sensitivity
533 may be exerted at or downstream the perception and signalling level. Consistently, several ABA-
534 responsive genes are less induced by drought in the *max2* signalling mutant of Arabidopsis than in
535 the wt (Ha et al., 2014). Finally, but not less importantly, stomatal conductance of miR156-oe
536 plants recovered much less than wt following drought stress. This response was opposite to *CCD7*-
537 silenced plants, while resembling that of plants treated with exogenous strigolactones prior to
538 stress imposition. This set of features converges on the idea that the strigolactone-miR156
539 pathway promotes the “after-effect” of drought, which is thought to be part of a wider stress-
540 memory mechanism (Lämke & Bäurle, 2017). After rewatering in fact, stomatal conductance never
541 quite reaches the levels of unstressed plants even though water potential has, and this response
542 could depend on non-hydraulic signals such as ABA (Ding et al., 2012; Galmes, Medrano, & Flexas,
543 2007; Lovisolo, Perrone, Hartung, & Schubert, 2008). Thus, genetic and pharmacological evidence
544 indicates that both strigolactones and miR156 increase sensitivity to ABA in a number of species,
545 including tomato. This reinforces the possibility of a strigolactone-miR156 module setting basal
546 stomata sensitivity to ABA in accordance with the water-balance history of the plant. Of course,
547 our data do not exclude that part of miR156 effects may also be exerted ABA-independently.

548
549 In conclusion, the results of the present work provided insight into the induction of miR156 under
550 drought. They also demonstrated a cause-effect link between miR156 accumulation and the
551 regulation of water relations and stomatal functioning. Particularly, we suggest that miR156 might
552 not directly mediate the short-term, transient and likely ABA-independent stomatal closure
553 triggered upon GR24^{5DS} treatment. Rather, the strigolactone-miR156 module seems to set basal
554 sensitivity of guard cells to ABA. This is most obvious after a drought period has occurred, when
555 the module has been activated but hydraulic signals are no longer active. We propose that the
556 module is also relevant for the establishment of specific drought responses, since its activation
557 might promote acclimation to (recurring) water-limiting conditions by increasing sensitivity to ABA
558 and drought avoidance. Tomato plants conditionally depleted of miR156, or overexpressing it, will
559 allow discriminating between the stress-related, strigolactone-dependent effects of miR156 on
560 stomata, and the long-term developmental consequences of its constitutive dysregulation.

561

562

563 **ACKNOWLEDGEMENTS**

564 F.C. conceived of the work and designed research supported by I.V., C.L. and A.S.; I.V. performed
565 most of the experiments helped by C.P., A.C. and V.T.; I.V., C.P., C.L. and O.N. analysed data; I.V.
566 and F.C. wrote the article with contributions of all the authors, who read, commented and
567 approved the final manuscript. A.S. and F.C provided for financial support to the research work.
568 F.C. agrees to serve as the author responsible for contact and ensures communication. The
569 Ministry of Education, Youth and Sports, Czech Republic (Grant LO1204 from the National
570 Programme of Sustainability I) supported V.T. and O.N. The PlantStressLab team and E.D. have
571 received funding from the European Union's Horizon 2020 research and innovation programme
572 under the Grant Agreement No. [727929] (TOMRES). None of these funding sources were involved
573 in the design of the study and collection, analysis and interpretation of data, and in writing the
574 manuscript. The authors wish to thank Dr. Silva e Ferreira (Sao Paulo, Brazil) for sharing the
575 miR156-oe MicroTom line, Dr. Klee (University of Florida, USA) for the *CCD7*-silenced line, and
576 Viviana Cisse and Cecilia Roche for technical support. The authors declare no conflicts of interest.

REFERENCES

- Andreo-Jimenez, B., Ruyter-Spira, C., Bouwmeester, H. J., & Lopez-Raez, J. A. (2015). Ecological relevance of strigolactones in nutrient uptake and other abiotic stresses, and in plant-microbe interactions below-ground. *Plant and Soil*, 394, 1-19.
- Arshad, M., Feyissa, B., Amyot, L., Aung, B., & Hannoufa, A. (2017). MicroRNA156 improves drought stress tolerance in alfalfa (*Medicago sativa*) by silencing *SPL13*. *Plant Science*, 258, 122-136.
- Axtell, M. J., & Meyers, B. C. (2018). Revisiting criteria for plant microRNA annotation in the era of big data. *Plant Cell*, 30(2), 272-284.
- Bauer, H., Ache, P., Lautner, S., Fromm, J., Hartung, W., Al-Rasheid, K. A., . . . Hedrich, R. (2013). The stomatal response to reduced relative humidity requires guard cell-autonomous ABA synthesis. *Current Biology*, 23(1), 53-57.
- Bayraktar, R., Van Roosbroeck, K., & Calin, G. A. (2017). Cell-to-cell communication: microRNAs as hormones. *Molecular Oncology*, 11(12), 1673-1686.
- Bhogale, S., Mahajan, A. S., Natarajan, B., Rajabhoj, M., Thulasiram, H. V., & Banerjee, A. K. (2014). MicroRNA156: a potential graft-transmissible microRNA that modulates plant architecture and tuberization in *Solanum tuberosum* ssp. *andigena*. *Plant Physiology*, 164(2), 1011-1027.
- Brun, G., Thoiron, S., Braem, L., Pouvreau, J. B., Montiel, G., Lechat, M. M., . . . Delavault, P. (2019). CYP707As are effectors of karrikin and strigolactone signalling pathways in *Arabidopsis thaliana* and parasitic plants. *Plant, Cell & Environment*, 42(9), 2612-2626.
- Bu, Q., Lv, T., Shen, H., Luong, P., Wang, J., Wang, Z., . . . Huq, E. (2014). Regulation of drought tolerance by the F-box protein MAX2 in *Arabidopsis*. *Plant Physiology*, 164(1), 424-439.
- Cardinale, F., Korwin Krukowski, P., Schubert, A., & Visentin, I. (2018). Strigolactones: mediators of osmotic stress responses with a potential for agrochemical manipulation of crop resilience. *Journal of Experimental Botany*, 69(9), 2291-2303.
- Christmann, A., Weiler, E. W., Steudle, E., & Grill, E. (2007). A hydraulic signal in root-to-shoot signalling of water shortage. *Plant Journal*, 52(1), 167-174.
- Cui, L. G., Shan, J. X., Shi, M., Gao, J. P., & Lin, H. X. (2014). The miR156-SPL9-DFR pathway coordinates the relationship between development and abiotic stress tolerance in plants. *Plant Journal*, 80(6), 1108-1117.
- Curaba, J., Singh, M. B., & Bhalla, P. L. (2014). miRNAs in the crosstalk between phytohormone signalling pathways. *Journal of Experimental Botany*, 65(6), 1425-1438.
- Dai, X., & Zhao, P. (2011). psRNATarget: a plant small RNA target analysis server. *Nucleic Acids Research*, 39, W155-W159.
- Ding, Y., Fromm, M., & Avramova, Z. (2012). Multiple exposures to drought 'train' transcriptional responses in *Arabidopsis*. *Nature Communications*, 3, 740.
- Ding, Y., Tao, Y., & Zhu, C. (2013). Emerging roles of microRNAs in the mediation of drought stress response in plants *Journal of Experimental Botany*, 64(1), 3077-3086.
- Ferreira e Silva, G. F., Silva, E. M., Azevedo Mda, S., Guivin, M. A., Ramiro, D. A., Figueiredo, C. R., . . . Nogueira, F. T. (2014). MicroRNA156-targeted SPL/SBP box transcription factors regulate tomato ovary and fruit development. *Plant Journal*, 78(4), 604-618.
- Ferrero, M., Pagliarani, C., Novak, O., Ferrandino, A., Cardinale, F., Visentin, I., & Schubert, A. (2018). Exogenous strigolactone interacts with abscisic acid-mediated accumulation of anthocyanins in grapevine berries. *Journal of Experimental Botany*, 69(9), 2391-2401.
- Galmes, J., Medrano, H., & Flexas, J. (2007). Photosynthetic limitations in response to water stress and recovery in Mediterranean plants with different growth forms. *New Phytologist*, 175(1), 81-93.
- Ha, C. V., Leyva-Gonzalez, M. A., Osakabe, Y., Tran, U. T., Nishiyama, R., Watanabe, Y., . . . Tran, L. S. (2014). Positive regulatory role of strigolactone in plant responses to drought and salt stress. *Proceedings of the National Academy of Sciences of the United States of America*, 111(2), 851-856.
- Haider, I., Andreo-Jimenez, B., Bruno, M., Bimbo, A., Flokova, K., Abuauf, H., . . . Ruyter-Spira, C. (2018). The interaction of strigolactones with abscisic acid during the drought response in rice. *J Exp Bot*, 69(9), 2403-2414.

- Holbrook, N. M., Shashidhar, V. R., James, R. A., & Munns, R. (2002). Stomatal control in tomato with ABA-deficient roots: response of grafted plants to soil drying. *Journal of Experimental Botany*, 53(373), 1503-1514.
- Hopper, D. W., Ghan, R., & Cramer, G. R. (2014). A rapid dehydration leaf assay reveals stomatal response differences in grapevine genotypes. *Horticulture Research*, 1, 2.
- Hsieh, L. C., Lin, S. I., Shih, A. C., Chen, J. W., Lin, W. Y., Tseng, C. Y., . . . Chiou, T. J. (2009). Uncovering small RNA-mediated responses to phosphate deficiency in *Arabidopsis* by deep sequencing. *Plant Physiology*, 151(4), 2120-2132.
- Jones-Rhoades, M., & Bartel, D. (2004). Computational identification of plant microRNAs and their targets, including a stress-induced miRNA. *Molecular Cell*, 14, 787-799.
- Kerr, S. C., & Beveridge, C. A. (2017). IPA1: a direct target of SL signaling. *Cell Research*, 27, 1191–1192.
- Khraiweh, B., Zhu, J. K., & Zhu, J. (2012). Role of miRNAs and siRNAs in biotic and abiotic stress responses of plants. *Biochimica et Biophysica Acta*, 1819(2), 137-148.
- Kuromori, T., Sugimoto, E., & Shinozaki, K. (2014). Inter-tissue signal transfer of abscisic acid from vascular cells to guard cells. *Plant Physiology*, 164, 1587-1592.
- Lämke, J., & Bäurle, I. (2017). Epigenetic and chromatin-based mechanisms in environmental stress adaptation and stress memory in plants. *Genome Biology*, 18(1), 124.
- Lanfranco, L., Fiorilli, V., Venice, F., & Bonfante, P. (2018). Strigolactones cross the kingdoms: plants, fungi, and bacteria in the arbuscular mycorrhizal symbiosis. *Journal of Experimental Botany*, 69(9), 2175-2188.
- Lechat, M. M., Pouvreau, J. B., Peron, T., Gauthier, M., Montiel, G., Veronesi, C., . . . Delavault, P. (2012). *PrCYP707A1*, an ABA catabolic gene, is a key component of *Phelipanche ramosa* seed germination in response to the strigolactone analogue GR24. *Journal of Experimental Botany*, 63(14), 5311-5322.
- Li, W., Nguyen, K. H., Chu, H. D., Ha, C. V., Watanabe, Y., Osakabe, Y., . . . Tran, L. P. (2017). The karrikin receptor KAI2 promotes drought resistance in *Arabidopsis thaliana*. *PLoS Genetics*, 13(11), e1007076.
- Lian, C., Yao, K., Duan, H., Li, Q., Liu, C., Yin, W., & Xia, X. (2018). Exploration of ABA-responsive miRNAs reveals a new hormone signaling crosstalk pathway regulating root growth of *Populus euphratica*. *International Journal of Molecular Sciences*, 19(5), 1481.
- Liu, H. H., Tian, X., Li, Y. J., Wu, C. A., & Zheng, C. C. (2008). Microarray-based analysis of stress-regulated microRNAs in *Arabidopsis thaliana*. *RNA*, 14(5), 836-843.
- Liu, J., Cheng, X., Liu, P., & Sun, J. (2017). MiR156-targeted SBP-box transcription factors interact with DWARF53 to regulate *TEOSINTE BRANCHED1* and *BARREN STALK1* expression in bread wheat. *Plant Physiology*, 174(3), 1931-1948.
- Liu, J., He, H., Vitali, M., Visentin, I., Charnikhova, T., Haider, I., . . . Cardinale, F. (2015). Osmotic stress represses strigolactone biosynthesis in *Lotus japonicus* roots: exploring the interaction between strigolactones and ABA under abiotic stress. *Planta*, 241(6), 1435-1451.
- Liu, M., Yu, H., Zhao, G., Huang, Q., Lu, Y., & Ouyang, B. (2017). Profiling of drought-responsive microRNA and mRNA in tomato using high-throughput sequencing. *BMC Genomics*, 18(1), 481.
- Lopez-Raez, J. A. (2016). How drought and salinity affect arbuscular mycorrhizal symbiosis and strigolactone biosynthesis? *Planta*, 243(6), 1375-1385.
- Lovisolo, C., Perrone, I., Hartung, W., & Schubert, A. (2008). An abscisic acid-related reduced transpiration promotes gradual embolism repair when grapevines are rehydrated after drought. *New Phytologist*, 180(3), 642-651.
- Lv, S., Zhang, Y., Li, C., Liu, Z., Yang, N., Pan, L., . . . Wang, G. (2018). Strigolactone-triggered stomatal closure requires hydrogen peroxide synthesis and nitric oxide production in an abscisic acid-independent manner. *New Phytologist*, 217(1), 290–304.
- Marzec, M., & Muszynska, A. (2015). *In silico* analysis of the genes encoding proteins that are involved in the biosynthesis of the RMS/MAX/D pathway revealed new roles of strigolactones in plants. *International Journal of Molecular Sciences*, 16(4), 6757-6782.
- Matthews, J. S. A., Violet-Chabrand, S. R. M., & Lawson, T. (2017). Diurnal variation in gas exchange: the balance between carbon fixation and water loss. *Plant Physiology*, 174(2), 614-623.

- May, P., Liao, W., Wu, Y., Shuai, B., McCombie, W. R., Zhang, M. Q., & Liu, Q. A. (2013). The effects of carbon dioxide and temperature on microRNA expression in Arabidopsis development. *Nature Communications*, 4, 2145.
- McAdam, S. A., Manzi, M., Ross, J. J., Brodribb, T. J., & Gomez-Cadenas, A. (2016). Uprooting an abscisic acid paradigm: Shoots are the primary source. *Plant Signaling & Behavior*, 11(6), e1169359.
- McAdam, S. A. M., & Brodribb, T. J. (2018). Mesophyll cells are the main site of abscisic acid biosynthesis in water-stressed leaves. *Plant Physiology*, 177(3), 911-917.
- Merilo, E., Yarmolinsky, D., Jalakas, P., Parik, H., Tulva, I., Rasulov, B., . . . Kollist, H. (2018). Stomatal VPD response: there is more to the story than ABA. *Plant Physiology*, 176(1), 851-864.
- Nozawa, M., Miura, S., & Nei, M. (2012). Origins and evolution of microRNA genes in plant species. *Genome Biology and Evolution*, 4(3), 230-239.
- Pagliarani, C., Vitali, M., Ferrero, M., Vitulo, N., Incarbone, M., Lovisolo, C., . . . Schubert, A. (2017). The accumulation of miRNAs differentially modulated by drought stress is affected by grafting in grapevine. *Plant Physiology*, 173(4), 2180-2195.
- Salinas, M., Xing, S., Hohmann, S., Berndtgen, R., & Huijser, P. (2012). Genomic organization, phylogenetic comparison and differential expression of the SBP-box family of transcription factors in tomato. *Planta*, 235(6), 1171-1184.
- Scaffidi, A., Waters, M. T., Sun, Y. K., Skelton, B. W., Dixon, K. W., Ghisalberti, E. L., . . . Smith, S. M. (2014). Strigolactone hormones and their stereoisomers signal through two related receptor proteins to induce different physiological responses in Arabidopsis. *Plant Physiology*, 165(3), 1221-1232.
- Scholander, P. F., Bradstreet, E. D., Hemmingsen, E. A., & Hammel, H. T. (1965). Sap pressure in vascular plants. Negative hydrostatic pressure can be measured in plants. *Science*, 148, 339-346.
- Silva, G. F. F., Silva, E. M., Correa, J. P. O., Vicente, M. H., Jiang, N., Notini, M. M., . . . Nogueira, F. T. S. (2019). Tomato floral induction and flower development are orchestrated by the interplay between gibberellin and two unrelated microRNA-controlled modules. *New Phytologist*, 221(3), 1328-1344.
- Song, X., Lu, Z., Yu, H., Shao, G., Xiong, J., Meng, X., . . . Li, J. (2017). IPA1 functions as a downstream transcription factor repressed by D53 in strigolactone signaling in rice. *Cell Research*, 27(9), 1128-1141.
- Speth, C., Willing, E. M., Rausch, S., Schneeberger, K., & Laubinger, S. (2013). RACK1 scaffold proteins influence miRNA abundance in Arabidopsis. *Plant Journal*, 76(3), 433-445.
- Stief, A., Altmann, S., Hoffmann, K., Pant, B. D., Scheible, W. R., & Baurle, I. (2014). Arabidopsis miR156 regulates tolerance to recurring environmental stress through SPL transcription factors. *Plant Cell*, 26(4), 1792-1807.
- Tardieu, F. (2016). Too many partners in root-shoot signals. Does hydraulics qualify as the only signal that feeds back over time for reliable stomatal control? *New Phytologist*, 212(4), 802-804.
- Tardieu, F., Simonneau, T., & Muller, B. (2018). The physiological basis of drought tolerance in crop plants: a scenario-dependent probabilistic approach. *Annual Review of Plant Biology*, 69, 733-759.
- Toh, S., Kamiya, Y., Kawakami, N., Nambara, E., McCourt, P., & Tsuchiya, Y. (2012). Thermoinhibition uncovers a role for strigolactones in Arabidopsis seed germination. *Plant and Cell Physiology*, 53(1), 107-117.
- Turečková, V., Novák, O., & Strnad, M. (2009). Profiling ABA metabolites in *Nicotiana tabacum* L. leaves by ultra-performance liquid chromatography-electrospray tandem mass spectrometry. *Talanta*, 80(1), 390-399.
- Visentin, I., Vitali, M., Ferrero, M., Zhang, Y., Ruyter-Spira, C., Novak, O., . . . Cardinale, F. (2016). Low levels of strigolactones in roots as a component of the systemic signal of drought stress in tomato. *New Phytologist*, 212(4), 954-963.
- Vogel, J. T., Walter, M. H., Giavalisco, P., Lytovchenko, A., Kohlen, W., Charnikhova, T., . . . Klee, H. J. (2010). SICCD7 controls strigolactone biosynthesis, shoot branching and mycorrhiza-induced apocarotenoid formation in tomato. *Plant Journal*, 61(2), 300-311.
- Wang, H., & Wang, H. (2015). The miR156/SPL module, a regulatory hub and versatile toolbox, gears up crops for enhanced agronomic traits. *Molecular Plant*, 8(5), 677-688.
- Waters, M. T., Gutjahr, C., Bennett, T., & Nelson, D. C. (2017). Strigolactone signaling and evolution. *Annual Review of Plant Biology*, 68, 291-322.

- Xu, Y., Zhang, L., & Wu, G. (2018). Epigenetic regulation of juvenile-to-adult transition in plants. *Frontiers in Plant Science*, 9, 1048.
- Yan, J., Wang, P., Wang, B., Hsu, C. C., Tang, K., Zhang, H., . . . Zhu, J. K. (2017). The SnRK2 kinases modulate miRNA accumulation in Arabidopsis. *PLoS Genetics*, 13(4), e1006753.
- Zhang, X., Zou, Z., Zhang, J., Zhang, Y., Han, Q., Hu, T., . . . Ye, Z. (2011). Over-expression of *Sly-miR156a* in tomato results in multiple vegetative and reproductive trait alterations and partial phenocopy of the *sft* mutant. *FEBS Letters*, 585(2), 435-439.
- Zhang, Y., Lv, S., & Wang, G. (2018). Strigolactones are common regulators in induction of stomatal closure in planta. *Plant Signaling & Behavior*, 13(3), e1444322.

FIGURE LEGENDS

Figure 1. Effects of miR156 overexpression on tomato physiological performances during a drought and recovery time-course. **(A)** Stomatal conductance (g_s) of wt (MicroTom, empty symbols) and miR156-oe plants (black symbols) under well-watered conditions (ww, squares) and water stress followed by recovery (ws/rec, circles). Water withdrawal started at day zero, while recovery started by irrigation ten days later. Differences in g_s for well-watered plants were statistically significantly different at all points except day seven, but were not reported on the graph for better clarity. **(B)** Leaf water potential (Ψ_{leaf}) of wt and miR156-oe plants under different water conditions (well-watered = day zero, water-stressed = day ten and recovered plants = day fifteen of time-course in A). Data represent the mean \pm SE of $n = 5$ biological replicates from two independent experiments. In (A) and (B), * indicates significant differences between genotypes for the same time-point and condition as determined by the Student's t -test, while in (B), lower case letters mark significant differences between different conditions in wt or miR156-oe plants separately, as determined by a one-way ANOVA test ($P < 0.05$).

Figure 2. Effect of miR156 overexpression on ABA metabolism in unstressed wt (MicroTom) and miR156-oe plants (day zero in Fig. 1). The concentrations of **(A)** free ABA, abscisic- β -D-glucosyl ester (ABA-GE), phaseic acid (PA), dihydro-phaseic acid (DPA), neo-phaseic acid (neo-PA), 7'-hydroxy-ABA (7'OH-ABA) and of **(B)** transcripts of the ABA catabolic genes *SICYP707A1* and *SICYP707A2* and of the ABA biosynthetic gene *SINCED1* were quantified. Gene transcript abundance was normalised to endogenous *SIEF-1 α* and *SlsnRU6*, and presented as fold-change value over mean values of wt plants, which were set to 1. Data represent the mean \pm SE of $n = 5$ biological replicates. * indicates significant differences between genotypes, as determined by a Student's t -test ($P < 0.05$). n.d. = not detectable.

Figure 3. Effects of miR156 overexpression on stomatal responses to ABA treatment. Induced stomatal closure was quantified by measuring stomatal conductance 1 h after treatment with increasing concentrations of ABA in wt (MicroTom) and miR156-oe leaves, and expressed as a percentage of mean values for mock-treated leaves in the same genotype, which were set to 100. Data are the means \pm SE of $n = 5$ biological replicates. Different letters indicate significant differences between different treatments in the same genotype as determined by a one-way ANOVA test, while * indicates significant differences between genotypes for the same condition, as attested by Student's t -test ($P < 0.05$).

Figure 4. Strigolactones affect miR156 production and promote the after-effect of drought. **(A)** Effect of GR24^{5DS} (5 μ M) compared to mock treatment on the concentration of mature miR156 in leaf tissues during a short-term time-course in unstressed wt plants (0, 2, 6 and 24 h after treatment, $n = 3$ each sample being a pool of 3 leaflets). **(B)** Mature miR156 levels in leaves of wt (treated with exogenous strigolactones or mock-treated), and mock-treated *CCD7*-silenced plants during a quick drought time-course. Stress was imposed by uprooting plants at time zero (well-watered samples) and transferring them into dry vermiculite, with irrigated controls being transferred into wet vermiculite. Water-stressed and recovered samples were harvested 3 or 24 h (respectively) after the beginning of stress, having been re-watered right after stress peaking. For strigolactone and mock treatment, leaves were sprayed as in (A) 24 h before time zero (the beginning of stress). Data represent the mean \pm SE of $n = 3$ biological replicates from three independent experiments. Target RNA abundance was normalised to endogenous *SIEF-1 α* and *SlsnRU6* transcripts and presented as fold-change value over mock-treated wt tissues, which were set to 1. Different letters indicate significant differences as determined by a one-way ANOVA test ($P < 0.05$). **(C)** Stomatal conductance values for the experiment in (B). One-way ANOVA test ($P < 0.05$) was applied to detect differences among genotypes within a given time point. **(D)** Normalised stomatal conductance of wt (M82) plants upon treatment with GR24^{5DS}. The aerial parts of plants were sprayed with a 5 μ M GR24^{5DS} solution, and stomatal conductance (g_s , $\text{mmol H}_2\text{O m}^{-2} \text{s}^{-1}$) was measured 2 and 24 h after the treatment. Data are presented as percentage of stomatal conductance over average g_s values of mock-treated plants, which were set to 100%. Data represent the mean \pm SE of $n = 4$ biological replicates from three independent experiments. Different letters indicate statistically significant differences among treatments as determined by a one-way ANOVA test ($P < 0.05$).

Figure 5. Effects of drought on the transcript concentration of six miR156 targets in leaves of wt (M82) and *CCD7*-silenced (SL-) tomato plants. *SICCD7* transcripts are also displayed as a control for stress and transgene effect on this key strigolactone-biosynthetic gene. Gene transcripts were normalised to endogenous *SIEF-1 α* and *SlsnRU6* in stressed and recovered plants from the drought time-course of Fig. 4B. Stress was imposed by uprooting plants at time zero (well-watered samples) and transferring them into dry vermiculite. Water-stressed and recovered samples were harvested 3 or 24 h (respectively) after the beginning of stress, having been re-watered right after stress peaking. Data are presented as fold change over the mean values for well-watered plants (time zero) of the same genotype, which were set to 1. Data represent the means \pm SE of $n = 5$

(Soly07g062980, Soly10g009080 and Soly10g078700) or $n = 3$ (Soly03g114850, Soly04g045560, Soly01g090660 and Soly02g077920) biological replicates. Different letters indicate significant differences between conditions and genotypes within a given bar cluster as determined by a one-way ANOVA test ($P < 0.05$).

Figure 6. Effects of miR156 overexpression on transcripts of strigolactone-biosynthetic genes. Transcript amounts of the putative (*SID27*) or confirmed (*SIMAX1*, *SICCD7* and *SICCD8*) strigolactone-biosynthetic genes were quantified in **(A)** roots or **(B)** axillary buds of wt (MicroTom) and miR156-oe plants. Gene transcripts were normalised to endogenous *SIEF-1 α* and *SlsnRU6* and presented as fold change over mean wt values, which were set to 1. In (A) data represent the means \pm SE of $n = 4$ biological replicates from two independent experiments; in (B), each of two replicates was the pool of at least 30 axillary buds from 8 plants. *SICCD7* transcript was not detectable in samples represented in (B). * indicates significant differences between genotypes as determined by a Student's *t*-test ($P < 0.05$).

SUPPLEMENTARY DATA

The following supplementary data are available:

Supplementary Figure S1. Effects of miR156 overexpression on stomatal reactivity to exogenous ABA expressed as pore areas.

Supplementary Table S1. List of primer pairs used in this work, with target gene names and/or *Solyc* codes.





Figure 1





Figure 2




Figure 3




Figure 4





Figure 5





Figure 6




Figure S1. Effects of miR156 overexpression on stomatal reactivity to exogenous ABA. **(A)** Representative images of stomata from wt (MicroTom, left-hand panels) and miR156-oe plants (right-hand panels), before (upper panels) or after treatment (lower panels) with ABA 10 μM for 1 h. **(B)** Raw and **(C)** normalised stomatal aperture areas measured on abaxial epidermal peels of the two genotypes, before and after ABA treatment. Data represent the mean and \pm SE of $n > 30$ biological replicates from two independent experiments. * indicates significant differences between genotypes under the same conditions, while different letters mark significant differences between treated and untreated leaves of the same genotype as determined by a Student's t -test ($P < 0.05$).

Table S1. List of primers used in this work, with target gene names and/or *Solyc* codes.

primer/target name and/or <i>Solyc</i> code	sequence	reference
stem-loop miR156	5'-GTCGTATCCAGTGCAGGGTCCGAGGTATTCGCACTGGATACGACGTGCTC-3'	Chen <i>et al.</i> (2005)
Mature miR156	5'-GTCGTATCCAGTGCAGGGT-3'	this work
MIMAT0009138 (from miRBase)	5'-TTGACAGAAGATAGAGAGCACG-3'	
<i>SINCED1</i> Solyc08g016720.1	5'-AGGCAACAGTGAAACTTCCATCAAG-3' 5'-TCCATTAAAGAGGATATTACCGGGGAC-3'	Sun <i>et al.</i> (2012)
<i>SICYP707A1</i> Solyc04g078900.3	5'-CCCAGAGTTCTTTCCTGATCCACAA-3' 5'-GAATGCCACTACCAGATCTACCAC-3'	Sun <i>et al.</i> (2012)
<i>SICYP707A2</i> Solyc08g075320.3	5'-TCGAAAAAGGATACAATTCGATGCC-3' 5'-CTGCAATTTGTTCGTGAGTCC-3'	Sun <i>et al.</i> (2012)
<i>SICNR</i> Solyc02g077920.2	5'-GGCAGCAAATAACCTACTTTC-3' 5'-CAATTTGCTTAGAAATCCGGG-3'	this work
<i>SISPL15</i> Solyc10g078700.1	5'-TCAGCTACCAGGACCAGTTATCAT-3' 5'-CGTCGATTCTTGATCCCC-3'	this work
putative miR156 target 1 Solyc03g114850.2	5'-CATTCAAGGGCTATCAGGGA-3' 5'-TGGACATTCTACCTGCCGAA-3'	this work
putative miR156 target 2 Solyc07g062980.2	5'-CAAGACACCACTGCCATTG-3' 5'-AAAGGTCCAACAACGATGCC-3'	this work
putative miR156 target 3 Solyc10g009080.2	5'-TCCAGGAGAAGGGTCAAGTT-3' 5'-TCGAAGTACCAACAGACAAAAGT-3'	this work
putative miR156 target 4 Solyc04g045560.2	5'- TAACTCCACAAGCACGAGGT-3' 5'- AGTGGTCCCTGAAAGCTTGA-3'	this work
putative <i>SID27</i> Solyc09g065750.3	5'-TGTTCTTCTCATGCAGGCAAAT-3' 5'-GCTGTTGCAATCTGCTTGGT-3'	this work
<i>SICCD7</i> Solyc01g090660.3	5'-GTTGCTTTACCAATGGTTCAATTT-3' 5'-TACATTCATCATGGAAGGATCAAAGTT-3'	Kohlen <i>et al.</i> (2012)
<i>SICCD8</i> Solyc08g066650.3	5'-CCAATTGCCTGTAATAGTTC-3' 5'-GCCTTCAACGACGAGTTCTC-3'	Kohlen <i>et al.</i> (2012)
<i>SIMAX1</i> Solyc08g062950.3	5'-CGCCCTTAGTTCAGAGAA-3' 5'-GCCAACCAACCCATGTTCC-3'	this work
<i>SlsnRU6</i>	5'-GGGAACGATACAGAGAAGATTAGC-3' 5'-ACCATTCTCGATTTGTGCGT-3'	this work
<i>SIEF-1α</i> Solyc06g005060.3	5'-CTCCATTGGGTCG TTTTGCT-3' 5'-GGTCACCTTGGC ACCAGTTG-3'	Digilio <i>et al.</i> (2010)

References not listed in the main bibliographic section are:

Chen C, Ridzon DA, Broomer AJ, Zhou Z, Lee DH, Nguyen JT, Barbisin M, Xu NL, Mahuvakar VR, Andersen MR, Lao KQ, Livak KJ, Guegler KJ. 2005. Real-time quantification of microRNAs by stem-loop RT-PCR. *Nucleic Acids Research* **33**(20), e179.

Digilio MC, Corrado G, Sasso R, Coppola V, Iodice L, Pasquariello M, Bossi S, Maffei ME, Coppola M, Pennacchio F, Rao R, Guerrieri E. 2010. Molecular and chemical mechanisms involved in aphid resistance in cultivated tomato. *New Phytologist* **187**(4), 1089-1101.

Sun L, Sun Y, Zhang M, Wang L, Ren J, Cui M, Wang Y, Ji K, Li P, Li Q, Chen P, Dai S, Duan C, Wu Y, Leng P. 2012. Suppression of *9-cis-Epoxycarotenoid dioxygenase*, which encodes a key enzyme in abscisic acid biosynthesis, alters fruit texture in transgenic tomato. *Plant Physiology* **158**(1), 283–298.