

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**Development of a sensitive TaqMan qPCR assay for detection and quantification of venturia inaequalis in apple leaves and fruit and in air samples**

**This is a pre print version of the following article:**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1770316> since 2021-01-31T22:39:01Z

*Published version:*

DOI:10.1094/PDIS-10-19-2160-RE

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

# Development and validation of a highly sensitive real-time PCR TaqMan® assay for specific detection and quantification of *Venturia inaequalis* in apple leaves and fruit and in air samples

Simona Prencipe, Fabiano Sillo, Angelo Garibaldi, Maria Lodovica Gullino, Davide Spadaro

The first, second, fourth and fifth authors: Department of Agricultural, Forestry and Food Sciences (DiSAFA), University of Torino, via Paolo Braccini 2, 10095, Grugliasco, TO, Italy; First, third, fourth and fifth authors: Centre of Competence for the Innovation in the Agro-environmental Sector - AGROINNOVA, University of Turin, via Paolo Braccini 2, 10095, Grugliasco, TO, Italy.

## Abstract

A TaqMan real-time PCR assay, based on the translation elongation factor 1- $\alpha$  gene, was developed for the quantification of *Venturia inaequalis* in leaves and fruits of *Malus x domestica* and in spore trap samples. The designed primers and probe amplified a specific 86 bp fragment for *V. inaequalis*. The specificity of the assay was tested using 35 strains of *V. inaequalis* and 20 different fungal species, including common pathogens of apple and other species of *Venturia*. The limit of detection was 20 fg, which is lower than a single genome of *V. inaequalis*. The selectivity of the assay was tested using DNA from three cultivars of *Malus x domestica* and no influence on pathogen amplification was found. The assay was also validated for repeatability and reproducibility. With this assay, it was possible to detect and quantify *V. inaequalis* in four cultivars ('Ambrosia', 'Fiorina', 'Golden Delicious' and 'Mondial Gala') in both symptomatic and asymptomatic leaves, and in symptomatic 'Golden Delicious' apple fruit, stored for 2 months. Furthermore, the assay was successfully tested on air samples coming from apple orchards. The quantification of the molecular assay, when compared with the estimated number of *V. inaequalis* cells using an optical microscope, showed a correlation coefficient of 0.8186. The developed technique could be used to detect *V. inaequalis* in asymptomatic samples and could be a promising tool for timely application of fungicides in orchards and to improve the efficacy of disease management.

**Keywords:** apple scab, *Malus x domestica*, real-time PCR, TaqMan, *Venturia inaequalis*.

## 1. Introduction

Apple scab is a worldwide disease affecting apple cultivation (*Malus x domestica*, Bork H.) and it is caused by the ascomycete *Venturia inaequalis* (Cooke) G. Winter (Sutton et al., 2014). The disease is particularly severe in temperate climate regions characterized by humid and cool springs (Bowen

et al., 2011) and, if not appropriately managed, can cause huge economic losses (MacHardy, 1996). Disease control requires an integrated strategy, based on prophylaxis practice, application of fungicides, and increasing use of resistant cultivars (MacHardy et al., 2001). *V. inaequalis* has a hemibiotrophic life cycle. Ascospores, which are released from pseudothecia and form during winter in the fallen leaves, are the sexual reproductive structures of the pathogen, and cause primary infections during the growing season (spring - early summer). If the weather conditions are favorable, asexual conidia are released and cause secondary infections during the growing season (Carrisse et al., 2000; Bowen et al., 2011). The entire life cycle of *V. inaequalis* is strongly influenced by the humidity and temperature conditions (MacHardy, 1996). In order to reduce scab infections, fungicide application programs are scheduled based on the local weather conditions, on disease prediction models and on the level of infection in spy plants (Mills and Laplante 1951; Gadoury and MacHardy 1986; Meitz-Hopkins et al., 2014; Carrisse et al., 2000). The evaluation of the level of infection is based on a visual scoring, which is an operator-dependent method used to attribute a disease severity index (Bock et al., 2010; Gusberti et al., 2012). The detection and quantification of pathogens, even in asymptomatic tissues, currently rely on molecular techniques. Different studies have been reported for specific amplification of *V. inaequalis* using conventional PCR (Schnabel et al. 1999; Stehmann et al. 2001, Koh et al., 2013). Daniëls et al. (2012) developed a qPCR assay using the housekeeping genes ATP-binding cassette transporter 2 (ABC2) and the elongation factor (EF1), while Gusberti et al. (2012) developed a qPCR assay, based on primers and probe designed on the internal transcribed spacer (ITS) to detect *V. inaequalis*. Spore traps were used in combination with qPCR in order to quantify the air-borne inoculum of different plant pathogens (Carrisse et al., 2009; Klosterman et al., 2014; Huang et al., 2016), including *V. inaequalis* (Meitz-Hopkins et al., 2014). To date, no studies have been carried out using the TaqMan technology combined with spore traps. The aim of the current work was to develop and validate a specific TaqMan quantitative PCR (qPCR) assay that could be used to detect and quantify *V. inaequalis* on different *Malus x domestica* cultivars, both in the leaves and in the fruits. *V. inaequalis* isolates from different fields and growing seasons were used to evaluate the assay specificity. Moreover, different plant materials (symptomatic and asymptomatic) and artificially inoculated leaves were tested. The assay was validated for specificity on 20 different species, including *Venturia asperata*, which has recently been reported as a pathogen on scab-resistant varieties of apple having the *Rvi6* gene (Caffier et al., 2012; Turan et al., 2019). Furthermore, the assay was tested on spore trap samples in order to detect and quantify the airborne inoculum of *V. inaequalis*.

## 2. Materials and methods

### Fungal strains

Strains of *V. inaequalis* were isolated from leaves of apple tree (*Malus x domestica*) ‘Ambrosia’, ‘Golden Delicious’ and ‘Mondial Gala’, cultivated in northern Italy (Piedmont) during 2015 and 2016. Thirty-five isolates were selected for this study. Other apple pathogenic or commonly present in orchard fungal species were isolated from apple leaves. All the isolates were identified through the amplification of the ribosomal DNA internal transcribed spacer (ITS) region, following the protocol of White et al. (1990). Reference strains (*V. inaequalis* CBS 815.69, *V. asperata* IRHS 2345, *V. pirina* CBS 120.825, *V. nashicola* CBS 794.84, *V. cerasi* CBS 444.54 and *Fusicladium carpophilum* CBS 497.62) and strains from the Agroinnova collection were used to verify the assay specificity. The strains used in this study are summarized in Table 1. The strains were maintained as monoconidial cultures in tubes of malt extract agar (MEA) (Sigma Aldrich, Germany) and stored at 4° C.

### Fungal DNA extraction and quantification

Strains of *Venturia* spp. were grown on MEA for 30 days at 20±1 °C in the dark, while the other fungal species were grown for 10 days at 25±1 °C. The fungal DNA was extracted from approximately 200 mg of fresh-weight mycelium using an Omega E.Z.N.A. Fungal DNA Mini Kit (VWR, USA), according to the manufacturer’s instructions. The DNA quality and the concentration were measured using a Nanodrop 2000 Spectrophotometer (Thermo Scientific, Wilmington, DE, USA).

### Sequence analysis of the translation elongation factor gene and design of primers and TaqMan probe

The translation elongation factor 1 alpha (*EF1-α*) sequences were amplified, for the 35 *V. inaequalis* strains, using EF1 (CGAGAAGTTCGAGAAGGT) and EF2 (CCAATGACGGTGACATAG) primers. PCR was carried out in a total volume of 25 µL containing 2.5 µL of Buffer 10 X, 0.5 µL of MgCl<sub>2</sub>, 0.75 µL of dNTPs (10 mM), 1 µL of each primer (10 mM), 0.2 µL of Taq DNA polymerase (Qiagen, Germany) and 20 ng of template DNA. The thermal cycling program was performed according to Gladioux et al. (2010). After agarose gel electrophoresis, the PCR products were purified using a QIAquick® PCR purification Kit (Qiagen), and sequenced in both directions by Macrogen, Inc. (The Netherlands). The consensus sequences were obtained by assembling forward and reverse sequences, using DNA Baser (Heracle Biosoft, Romania). The consensus sequences obtained for the *V. inaequalis* strains were compared with those deposited in GenBank and a multi alignment was

103 performed using the CLUSTALW algorithm, through Molecular Evolutionary Genetics Analysis  
104 (MEGA6) software, version 6.0. The alignment was used to design the primers and probe used in this  
105 study.

106 Six sets of primers and two probes were designed using Primer Express™ software 3.0 (Applied  
107 Biosystem, Foster City, USA) and are listed in **Table 2**. The TaqMan probes were labelled at the 5'-  
108 end with the 6-carboxyfluorescein dye (6-FAM) reporter and Black Hole Quencher (BHQ1) or the  
109 non-fluorescent quencher minor groove binder (NF-MGB) at the 3'-end. OligoCalc tool  
110 (<http://biotools.nubic.northwestern.edu/OligoCalc.html>) was used to verify the presence of hairpins  
111 and potential secondary structures, while *in silico* specificity was verified using the BLASTN tool of  
112 the National Centre of Biotechnology Information (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>).  
113 Invitrogen (Carlsbad, USA) and Metabion (Steinkirchen, Germany) synthesized primers and probes,  
114 respectively.

115

#### 116 **Conventional end-point PCR and qPCR optimization and amplification**

117 The primer pair sets were first assessed by means of conventional end-point PCR using DNA from  
118 *V. inaequalis* strains and other selected species. The PCR endpoints were performed in a 25 µL  
119 reaction, composed of 2 µL of Buffer 10x, 0.8 µL of MgCl<sub>2</sub>, 1 µL of dNTPs (10 mM), 1 µL of each  
120 primer (10 mM), 0.2 µL of Taq Platinum Pfx DNA polymerase (Invitrogen, USA) and 20 ng of DNA.  
121 The PCR thermal cycler conditions were 3 min at 95°C followed by 30 cycles of 95°C for 45 s, 54°C  
122 for 45 s, 72°C for 1 min and a final extension of 5 min. The PCR products were run on 1% agarose  
123 gel in a TBE buffer, and visualized under UV transilluminator using the Quantity One software  
124 (BioRad Labs, Hercules, USA). After performing end-point PCR, the primer pairs that gave the best  
125 results were selected and used in qPCR with SYBR Green in order to compare them with the  
126 specificity and sensitivity of the TaqMan assay. Real-time reactions were performed using a  
127 StepOnePlus qPCR system (Applied Biosystems) with 96 well-plates (Optical reaction plate, Applied  
128 Biosystems) sealed with MicroAmp optical adhesive film (Applied Biosystems).

129 SYBR Green reactions were carried out using 10 µL of Power SYBR Green Mastermix 10x (Applied  
130 Biosystems), 1 µL of each primer (3 µM) and 1 µL of template DNA. Amplification conditions were  
131 95 °C for 10 min, followed by 40 cycles of 57 °C for 1 min and 95 °C for 15 s. The melting curves  
132 were acquired after each run at the following conditions: 95 °C for 15 s, 60 °C for 15 s and 95 °C for  
133 15 s. Sterile water was used as a negative control. In order to determine the sensitivity of the SYBR  
134 Green assay, a standard curve was obtained with *V. inaequalis* DNA using 1b14 strain 8-fold serially  
135 diluted (from 20 ng to 2 fg).

136 The TaqMan assay was performed using TaqMan Universal Mastermix 2X (Applied Biosystems).  
137 Each 96-well plate was loaded with a negative control, standard DNA and a positive control in  
138 triplicate. Different primer concentrations (from 3  $\mu$ M to 0.3  $\mu$ M) and different temperatures were  
139 initially tested at the annealing stage (57 to 60 °C). Reactions were carried out in a final volume of  
140 20  $\mu$ l, with 1  $\mu$ l of DNA, 0.4  $\mu$ l of each primer (3  $\mu$ M) and 0.2  $\mu$ l of probe (5  $\mu$ M). Sterile water was  
141 used as a negative control. Amplification conditions were 95 °C for 10 min, 40 cycles of 57 °C for 1  
142 min and 95 °C for 15 s. The newly developed TaqMan assay was compared, in terms of sensitivity  
143 and specificity, with the TaqMan assay previously published by [Gusberti et al., \(2012\)](#).  
144 The Ct values generated by qPCR were compared with the standard curve to quantify the DNA of *V.*  
145 *inaequalis* in target samples. The approximate number of cells was calculated by dividing the DNA  
146 quantity by the weight of the genome of *V. inaequalis* (0.0000597 ng; [Deng et al., 2017](#)), thereby  
147 obtaining the number of target cells (*V. inaequalis*) per  $\mu$ L of reaction.  
148

#### 149 **Specificity, sensitivity, selectivity, repeatability and reproducibility of the TaqMan qPCR**

150 The TaqMan assay was validated according to the international EPPO standard PM 7/98 ([EPPO](#)  
151 [2014](#)). In order to evaluate the specificity of the assay for *V. inaequalis*, 20 different species, including  
152 other pathogenic *Venturia* species that affect different hosts and *V. asperata*, which has recently been  
153 reported in Italy on apples, were used ([Table 1](#)). The presence of aspecific amplification signals was  
154 evaluated after 40 cycles in different experiments.

155 In order to determine the sensitivity of the TaqMan assay, a standard calibration curve was obtained  
156 with *V. inaequalis* DNA using the 1b14 strain 8-fold serially diluted (from 20 ng to 2 fg) in sterile  
157 deionized water. Furthermore, in order to verify the influence of the host DNA on *V. inaequalis*  
158 amplification, the pathogen DNA was 8-fold diluted in *Malus x domestica* ‘Ambrosia’, ‘Mondial  
159 Gala’ and ‘Golden Delicious’ DNA. The standard curve reaction was carried out in triplicate and used  
160 as an internal control in order to quantify the target DNA in different samples. Repeatability was  
161 checked by running three independent assays of the test. Two different operators tested the  
162 reproducibility of the assay in two different laboratories and on different days.  
163

#### 164 **Detection of *V. inaequalis* in naturally infected *Malus x domestica* leaves and in apple fruit**

165 During the 2015-2017 seasons, leaves from *Malus x domestica* plants showing symptoms of apple  
166 scab were harvested and stored for DNA extraction. From ‘Ambrosia’, ‘Golden Delicious’ and  
167 ‘Mondial Gala’ leaves, respectively 11, 11 and 5 samples were collected. Furthermore, four apples  
168 that had been stored for 2 months and which showed apple scab symptoms were selected for DNA  
169 extraction. DNA from the infected plants/fruit was extracted from approximately 100 mg of fresh-



weight leaves or fruit. The samples were previously freeze-dried, ground with liquid nitrogen and then extracted with E.Z.N.A. Plant DNA kit (VWR, USA), following the manufacturer's instructions.

#### **Detection of *V. inaequalis* in asymptomatic *Malus x domestica* leaves**

A total of 30 samples of asymptomatic *Malus x domestica* 'Ambrosia', 'Fiorina' and 'Golden Delicious' leaves were analyzed in May 2018 to assess the presence of *V. inaequalis* using the TaqMan assay. The leaves were divided into two equal parts using sterile blades and two different extractions were performed (100 mg each). The DNA extraction was performed as previously described.

#### **Detection of *V. inaequalis* in artificially inoculated *Malus x domestica* leaves**

*V. inaequalis* conidia were obtained from a single spore strain (1b14), according to [Parker et al. \(2005\)](#). Leaves were obtained from *Malus x domestica* 'Fiorina' plants. The absence of *V. inaequalis* in the plant material was confirmed using a binocular microscope (Nikon Eclipse 55i, Tokyo, Japan). Leaf disks (1 cm diameter) were collected for the assay, disinfected with a 10% solution of sodium hypochlorite, washed by immersion in sterile deionized water and air-dried. The inoculation of the leaf disks was performed on the adaxial leaf surface with an initial concentration of  $10^2$  conidia/mL of *V. inaequalis*, which was subsequently serially diluted (1:2, 1:3, 1:5, 1:10, 1:15, 1:20, 1:50 and 1:100). Control leaves were prepared in a similar way with sterile deionized water. Two replicates were tested for each conidial concentration. DNA was extracted from artificially inoculated leaves, following previously described procedures.

#### **Detection of *V. inaequalis* from spore trap samples**

A volumetric spore sampler (Burkard Manufacturing Co. Ltd., Rickmansworth, Hertfordshire, UK), placed in an apple orchard in Manta (Cuneo, Italy; 44.609217; 7.502627), was used to detect the airborne conidia of *V. inaequalis*. The tapes were collected at 24 h intervals for 14 days. Each daily tape was first visualized under an optical microscope to count the *V. inaequalis* cells at 40× magnification. The same segments were subsequently placed in 50 mL tubes and stored at 4 °C until processing. An aliquot of 25 mL of a polyethylene glycol (PEG) alkaline buffer (50 g/L PEG average Mn 4600; 20 mM KOH; pH 13.5) was added to each tube and vortexed for 20 minutes for the DNA extraction. After incubation of 1 h at 65 °C, the samples were vortexed for 20 min and centrifuged for 30 min at 8,000 g. The supernatant was collected and the DNA was extracted using an E.Z.N.A. Fungal DNA Mini kit, according to the manufacturer's protocol, by adjusting the buffer volumes.

203 Repeatability of the assay on air samples was evaluated by performing 6 independent reactions using  
204 the same set of samples.

205

206 **Data analysis.** StepOne™ software was used to automatically generate the baseline range and the  
207 qPCR standard curves, as well as to determine the Ct values. Student's t-test was used to analyze the  
208 reproducibility of the assay.

209

### 210 **3. Results**

211

#### 212 **TaqMan assay optimization**

213 The *EF1-α* gene was selected because of the presence of a conserved region and the deletion of 31  
214 nucleotides in the *V. inaequalis* species, able to differentiate from the other *Venturia* species,  
215 including *V. asperata*. The alignment of partial *EF1-α* gene sequences from different *V. inaequalis*  
216 strains and 5 other *Venturia* species was used to design the primers and probes reported in **Table 2**.  
217 The F1/R11 primer pair, which gave the best results with conventional end-point PCR, was selected  
218 for the real-time assays. The optimization of the TaqMan qPCR assay was tested using different  
219 primer and probe concentrations and considering different cycling conditions. The best conditions  
220 were found for a primer concentration of 3 μM at 57 °C for 1:00 for the annealing stage. The Ven1  
221 probe and F1/R11 primer pair were selected for the TaqMan assay as they showed the most suitable  
222 amplification.

223

#### 224 **Specificity, sensitivity, selectivity, repeatability and reproducibility of the TaqMan qPCR**

225 The TaqMan qPCR, with the designed primer pair and probe, was able to amplify the 86 bp fragment  
226 of the *EF1-α* gene in different experiments from 35 *V. inaequalis* strains (**ST1**). No amplification  
227 signal was detected after 40 cycles for *V. asperata*, *V. pirina*, *V. carpophila*, *V. cerasi*, *V. naschicola*  
228 or for the other tested species.

229 The DNA of *V. inaequalis*, serially diluted from 20 ng to 0.2 fg in sterile distilled water, was used to  
230 build a standard curve in order to evaluate the limit of detection (LOD). The pathogen was  
231 quantifiable from 20 ng to 20 fg (**Fig. 1**), and a LOD threshold cycle (Ct), ranging between 36 and  
232 37, was obtained. The LOD of 20 fg is lower than a single genome of *V. inaequalis*, (0.0000597 ng,  
233 **Deng et al., 2017**). The mean value of the regression slope was -3.28, and the mean relative efficiency  
234 was between 99% and 110%. No influence was observed on the selectivity of the TaqMan assay when  
235 *V. inaequalis* DNA, serially diluted in *Malus x domestica* 'Ambrosia', 'Golden Delicious' and  
236 'Mondial Gala' DNA, was used (**Fig. 2**). The amplifications showed similar PCR efficiencies and a



237 reliable correlation between the Ct values and the amount of measured *V. inaequalis* DNA (Fig. 2).  
238 No statistical differences ( $p>0.05$ ) were found for the results of the student's t-test, which was used  
239 to analyze the repeatability and reproducibility of the assay.

240

#### 241 **Specificity and sensitivity comparison of the SYBR Green and TaqMan assays**

242 The SYBR Green assay by using the selected F1/R11 primer pair provided positive results for the  
243 target DNA from different cultivars, and no amplification was observed for the other *Venturia* species  
244 after 40 cycles. However, aspecific signals were detected for some species used as negative controls,  
245 i.e. *Cladosporium cladosporioides*, *Colletotrichum fiorinae*, *Fusarium equiseti*, *Penicillium*  
246 *expansum* and *Phoma* sp., after 34 to 36 cycles. The sensitivity of the assay with SYBR Green  
247 revealed a 10-fold higher LOD (200 fg) than the TaqMan assay designed on the *EF1-α* gene.

248 The TaqMan assay previously developed by Gusberti et al. (2012) gave a cross-reaction for the strain  
249 MALT1 *Alternaria* sp. isolated from apple leaves, while no other amplification was obtained for the  
250 other tested species. Sensitivity, assessed using the method of Gusberti et al. (2012) on the *V.*  
251 *inaequalis* 1b14 strain, showed a LOD of 100 fg (ST2).

252

#### 253 **Detection of *V. inaequalis* in naturally infected *Malus x domestica* leaves and fruit**

254 The TaqMan assay was used to quantify *V. inaequalis* in naturally infected leaves of three *Malus x*  
255 *domestica* cultivars and apple fruit. No influence of the host DNA was detected, and all the analyzed  
256 samples resulted positive for the target amplification (Fig. 3A and ST3). The average number of  
257 cells/ $\mu$ L was  $5.26 \times 10^4$  for the 'Ambrosia' samples,  $4.28 \times 10^4$  for the 'Mondial Gala' samples and  
258  $3.76 \times 10^4$  for the 'Golden Delicious' samples. The assay was able to detect 2.83 *V. inaequalis*  
259 cells/ $\mu$ L in the naturally infected leaves (mean Ct 34.35) and 182 cells/ $\mu$ L in the naturally infected  
260 fruit (mean Ct 28.38) (Fig. 3A and ST3).

261

#### 262 **Detection of *V. inaequalis* in asymptomatic *Malus x domestica* leaves**

263 The TaqMan assay was able to detect *V. inaequalis* in both the asymptomatic resistant and susceptible  
264 cultivars (Fig. 3B and ST4). The lowest concentration was 3.21 cells/ $\mu$ L and it was found in the As7  
265 sample (resistant 'Fiorina'), while the highest concentration was found in the susceptible 'Ambrosia'  
266 samples, with  $4.50 \times 10^3$  cells/ $\mu$ L. The mean *V. inaequalis* concentration was  $1.90 \times 10^2$  cells/ $\mu$ L for  
267 'Fiorina', while  $3.46 \times 10^2$  cells/ $\mu$ L were found for 'Golden Delicious' and  $1.91 \times 10^3$  cells/ $\mu$ L for  
268 'Ambrosia'.

269

#### 270 **Detection of *V. inaequalis* in artificially inoculated *Malus x domestica* leaves**

271 In order to test the developed TaqMan assay on environmental-like samples, *Malus x domestica*  
272 leaves were artificially inoculated with a *V. inaequalis* conidial suspension. The results obtained for  
273 the quantification of *V. inaequalis* are reported in Figure 4. All the samples amplified with a clear  
274 amplification signal, with Ct values ranging from 28, for the samples inoculated with highest  
275 concentration, to 36 for the lowest. No amplification was obtained for the negative controls. The  
276 amplification showed linearity in the serial dilutions, and the assay allowed us to quantify from  $1.65 \times 10^2$  cells/ $\mu$ L, for the initial sample, to 1.21 cells/ $\mu$ L, for the 100-fold diluted sample.

278

#### 279 **Detection of *V. inaequalis* from spore trap samples**

280 The TaqMan assay was used to detect and quantify the presence of airborne inoculum of *V. inaequalis*  
281 in the spore trap samples. The estimated conidial concentrations, based on microscope counts, were  
282 higher than the estimated concentrations calculated from the amount of *V. inaequalis* DNA detected  
283 by the TaqMan assay. Only one of the 14 analyzed samples showed no amplification (Table 3). The  
284 estimated mean number of cells/ $\mu$ L obtained from the microscope count was  $1.01 \times 10^4$ , while it was  
285  $5.26 \times 10^3$  for the TaqMan assay. The correlation coefficient between the two assays was positive,  
286 with an  $R^2$  value of 0.8186. The repeatability of the assay was variable, with at least one positive  
287 amplification per sample over six reactions (Table 3).

288

#### 289 **4. Discussion**

290 The molecular techniques currently applied for the detection of plant pathogens are often used to  
291 specifically identify and quantify fungal species in crops and food commodities (Postollec et al.,  
292 2001; Capote et al., 2012; Aslam et al., 2017). In this study, a highly sensitive TaqMan real-time  
293 assay has been developed for the specific detection and quantification of *V. inaequalis*, and it has  
294 successfully been used with symptomatic and asymptomatic leaves, fruit and air samples.

295 Conventional molecular methods, based on PCR, showed specificity when used for the detection of  
296 *Venturia* species (Schnabel et al., 1999; Koh et al., 2013), but there is a lack of effective quantitative  
297 results (Suarez et al., 2005). Real-time PCR gives more specific and sensitive results than  
298 conventional PCR and allows pathogens to be quantified (Selma et al., 2008; Mirmajlessi et al., 2015;  
299 Baskarathevan et al., 2016; Kuzdraliński et al., 2017). Real-time PCR was used in previous studies  
300 to detect and quantify *V. inaequalis* in different types of samples. Both SYBR Green I and the  
301 TaqMan assay were utilized. The first technique exploited a fluorescent measurement for DNA  
302 amplification through double strand DNA binding dyes (SYBR Green I®), although there were some  
303 limits pertaining to aspecific amplifications and difficulties in the interpretation of the results after  
304 the melting curve (Martinez et al., 2011). These issues could be solved by using the second technique,

305 i.e. TaqMan chemistry, which includes specific fluorescent probes into the PCR (Amaral Carneiro et  
 306 al., 2017).

307 In preliminary studies, different target sequences were explored for the development of the TaqMan  
 308 real-time assay, following multiple sequence alignments (data not shown). The used sequences were  
 309 obtained from strains isolated in Italy and from sequences available in the GenBank public database.  
 310 Due to the high intraspecific variability, the ITS region and the beta-tubulin gene were discarded (data  
 311 not shown), whereas the *EFL-α* gene proved a highly conserved species-specific region for *V.*  
 312 *inaequalis* and it was therefore selected to design primers and probes.

313 The developed assay was compared using SYBR Green I and with the TaqMan assay, previously  
 314 published (Gusberti et al., 2012). The primers designed on the *EFL-α* gene cross-reacted with various  
 315 tested species when used with SYBR Green I. Cross-reaction was also observed in the work of Meitz-  
 316 Hopkins et al. (2014), where two qPCR assays, based on the CYP51A1 gene and the ITS region, were  
 317 developed. The study of Daniëls et al. (2012), using a qPCR assay based on the ITS region, also  
 318 showed cross-reaction. The main fungal species that showed cross-amplification with the  
 319 aforementioned molecular tools were *Alternaria* sp. and *V. asperata*, *V. nascicola*, *V. pirina*, *V. cerasi*  
 320 and *V. carpophila*. Our data demonstrated that the use of SYBR Green allowed to reach a sensitivity  
 321 of 200 fg, which in turn results in a 5 times lower than the detection limit obtained by Meitz-Hopkins  
 322 et al. (2014), but 2 times higher than that reported by Daniëls et al. (2012).

323 When the specificity was tested using the TaqMan probe technology, no cross-reaction was detected  
 324 for the other *Venturia* species, including *V. asperata*, or for other pathogens. On the contrary, when  
 325 the specificity was tested with the TaqMan probe developed by Gusberti et al. (2012), a positive  
 326 amplification was obtained for a strain of *Alternaria* sp. from apple leaves. A blast search of the  
 327 primers and probes used in the study of Gusberti et al. (2012) gave 100% homology and 100%  
 328 coverage with *A. solani* (accession number CPO22033.1) and 100% homology and 72% coverage  
 329 with *A. alternata* (accession number XM\_01852723.1).

330 Because of the high intraspecific variability that exists within the species *V. inaequalis* (Tenzer and  
 331 Gessler, 1999 ; Tenzer et al., 1999; Ebrahimi et al., 2016), the specificity of the assay was confirmed  
 332 on DNA extracted from a large number of pure cultures of *V. inaequalis* strains, isolated from  
 333 different cultivars over different years, and positive amplifications were observed for all the strains.  
 334 The assay showed a low variation in the Ct values obtained in independent experiments and resulted  
 335 to be highly reproducible.

336 When the sensitivity of the assays was compared, the TaqMan assay resulted in a lower detection  
 337 limit than the SYBR Green assay. The advantage of using the TaqMan assay has been reported in a  
 338 qPCR comparative study by Soltany-Rezaee-Rad et al. (2015). A qPCR with SYBR Green could

339 inhibit the Taq DNA polymerase, thereby reducing the sensitivity of the assay (Kermekchiev et al.,  
 340 2009). The LOD of our TaqMan assay (20 fg) proved to be more sensitive than that reported by  
 341 Gusberti et al. (2012) for which the LOD was 100 fg. The LOD of the TaqMan assay (20 fg) is similar  
 342 to those reported for other species i.e. *Botrytis cinerea* (Suarez et al., 2005), *Fusarium solani* (Bernal-  
 343 Martínez et al., 2012), *Aspergillus fumigatus* (Fernandez-Molina et al., 2014), *Lichtheimia*  
 344 *corymbifera* (Springer et al., 2016), *Fusarium fujikuroi* (Amaral Carneiro et al., 2017) and *Fusarium*  
 345 *culmorum* (Bilska et al., 2018).

346 The TaqMan assay was also tested to quantify the target DNA in the presence of the plant material.  
 347 The sensitivity was not affected by the presence of the DNA of *Malus x domestica* from different  
 348 cultivars, and the assay allowed to detect and quantify *V. inaequalis* in inoculated samples with  
 349 significant linearity. Positive amplifications were also obtained from leaf samples, collected both  
 350 from susceptible and resistant cultivars, that were, respectively, symptomatic or asymptomatic. In  
 351 addition, the assay proved to be useful for the detection of the pathogen in fruit samples.

352 When the assay was performed on DNA obtained from the spore trap, 13 samples out of 14 resulted  
 353 positive to *V. inaequalis*, even with a low conidial concentration. In our study, the overall  
 354 concentration estimated with a microscope count resulted higher than that estimated with the TaqMan  
 355 assay, but with a linear relationship ( $R^2$ : 0.8186), as previously reported by Carisse et al. (2009),  
 356 pertaining to the quantification of *Botrytis squamosa*. However, the repeatability of the assay was  
 357 variable for air samples, as inhibition of PCR may occur when DNA is extracted from air samples,  
 358 thus false-negative plate readings and a reduced amplification efficiency may be obtained. The  
 359 reasons for these false negatives could be linked to competition in the amplification due to: high  
 360 amounts of non-target DNA, inoculum density, co-extraction of contaminants (PCR inhibitors) or  
 361 unequal distribution of conidia on tapes (McDevitt et al., 2007; Rogers et al., 2009; Williams et al.  
 362 2001; Bilodeau et al 2012; Klosterman et al., 2014; Dung et al., 2015).

363 In conclusion, the development of a highly sensitive species-specific assay is important to detect  
 364 pathogen at low concentrations, even during latent infection or in asymptomatic samples. In our study,  
 365 the use of a TaqMan real-time assay increased the sensitivity of the molecular tool and led to the  
 366 advantage of being able to detect less than a single target cell. The assay proved to be specific and  
 367 highly sensitive for the detection of *V. inaequalis*, both in symptomatic and asymptomatic apple  
 368 leaves. The developed qPCR could be used in apple scab risk management systems to quantify the  
 369 inoculum in the field and to plan phytosanitary treatments. Furthermore, the technique proved to be  
 370 sensitive with air samples, though assessed on a limited number of samples.

371 This newly developed TaqMan assay could be a useful tool, combined with a weather-based data  
 372 model, to plan phytosanitary treatments in order to control apple scab. The here developed TaqMan

373 assay could also be used, together with nanobiosensor technology, to quantify airborne inoculum in  
374 the field, without time consuming DNA extraction and processing, and could therefore be used to  
375 implement a decision support system for apple scab management.

376

### 377 **Acknowledgments**

378 The Authors wish to thank Fondazione Cassa di Risparmio di Cuneo for its financial support through  
379 the projects “FRUITSENSOR – Converging technologies for sustainable precision fruit-growing”  
380 and “SMART APPLE - Innovative and SMART technologies for sustainable APPLE production”.

381 The Authors gratefully acknowledge Dr. Luca Nari and Dr. Graziano Vittone from AGRION for  
382 providing the samples and Dr. Sara Franco Ortega for her valuable help in the development of the  
383 assay.

## Literature cited

1. Amaral Carneiro, G., Matic, S., Ortu, G., Garibaldi, A., Spadaro, D., and Gullino, M.L. 2017. Development and validation of a TaqMan real-time PCR assay for the specific detection and quantification of *Fusarium fujikuroi* in rice plants and seeds. *Phytopathology*. 107:885-892.
2. Aslam, S., Tahir, A., Aslam, M. F., Alam, M. W., Shedayi, A. A., and Sadia, S. 2017. Recent advances in molecular techniques for the identification of phytopathogenic fungi - a mini review. *J. Plant Interact.* 12:493-504.
3. Baskarathevan, J., Taylor, R. K., Ho, W., McDougal, R. L., Shivas, R. G., and Alexander, B. J. R. 2016. Real-time PCR assays for the detection of *Puccinia psidii*. *Plant Dis.* 100:617-624.
4. Bernal-Martínez, L., Buitrago, M. J., Castelli, M. V., Rodríguez-Tudela, J. L., and Cuenca-Estrella, M. 2012. Detection of invasive infection caused by *Fusarium solani* and non-*Fusarium solani* species using a duplex quantitative PCR-based assay in a murine model of fusariosis. *Med. Mycol.* 50:270-275.
5. Bilodeau, G. J., Koike, S. T., Uribe, P., and Martin, F. N. 2012. Development of an assay for rapid detection and quantification of *Verticillium dahliae* in soil. *Phytopathology* 102:331-343.
6. Bilska, K., Kulik, T., Ostrowska-Kołodziejczak, A., Buśko, M., Pasquali, M., Beyer, M., Baturo-Cieśniewska, A., Juda, M., Załuski, D., Treder, K., Denekas, J., and Perkowski, J. 2018. Development of a highly sensitive FcMito qPCR assay for the quantification of the toxigenic fungal plant pathogen *Fusarium culmorum*. *Toxins*. 10:211.
7. Bock, C. H., Poole, G. H., Parker, P. E., and Gottwald, T. R. 2010. Plant disease severity estimated visually, by digital photography and image analysis, and by hyperspectral imaging. *Crit. Rev. Plant Sci.* 29:59-107.
8. Bowen, J. K., Mesarich, C. H., Bus, V. G. M., Beresford, R. M., Plummer, K. M., and Templeton, M. D. 2011. *Venturia inaequalis*: the causal agent of apple scab. *Mol. Plant Pathol.* 12:105-122.
9. Caffier, V., Le Cam, B., Expert, P., Tellier, M., Devaux, M., Giraud, M., and Chevalier, M. 2012. A new scab-like disease on apple caused by the formerly saprotrophic fungus *Venturia asperata*. *Plant Pathol.* 61:915-924.
10. Capote, N., Pastrana, A. M., Aguado, A., and Sánchez-Torres, P. 2012. Molecular tools for detection of plant pathogenic fungi and fungicide resistance. Pages 151-202 in C. J. Cumagun (eds.), *Plant pathology*. InTech, Rijeka, Croatia.



- 416 11. Carisse, O., McCartney, H. A., Gagnon, J. A., and Brodeur, L. 2009. Quantification of  
417 airborne inoculum as an aid in the management of leaf blight of onion caused by *Botrytis*  
418 *squamosa*. Plant Dis. 89:726-733.
- 419 12. Carisse, O. Pillion, V., Rolland, D., and Bernier, J. 2000. Effect of fall application of fungal  
420 antagonists on spring ascospore production of apple scab pathogen, *Venturia inaequalis*.  
421 Phytopathology 90:31-37.
- 422 13. Daniëls, B., De Landtsheer, A., Dreesen, R., Davey, M. W., and Keulemans, J. 2012. Real-  
423 time PCR as a promising tool to monitor growth of *Venturia* spp. in scab-susceptible and -  
424 resistant apple leaves. Eur. J. Plant Pathol. 134:821-833.
- 425 14. Deng, C. H., Plummer, K. M., Jones, D. A. B., Mesarich, C. H., Shiller, J., Taranto, A. P., et  
426 al. 2017. Comparative analysis of the predicted secretomes of *Rosaceae* scab pathogens  
427 *Venturia inaequalis* and *V. pirina* reveals expanded effector families and putative  
428 determinants of host range. BMC Genomics. 18:339.
- 429 15. Dung, J. K. S., Scott, J. C., Alderman, S.C., Kaur, N., Walenta, D. L., Frost, K. E. and Hamm,  
430 P. B. 2015. Development of a DNA-based protocol to detect airborne ergot spores in cool-  
431 season grass seed fields. Pages 31-34 in: Seed Production Research Report at Oregon State  
432 University USDA-ARS Cooperating, Ext/CrS 152. Anderson, N. P., Hulting, A. G., Walenta,  
433 D. L., Flowers M. D, and C. S. Sullivan (eds.). Oregon State Univ., Corvallis.
- 434 16. Ebrahimi, L., Fotuhifar, K. B., Nikkhah, M.J., Naghavi, M. R., and Baisakh, N. 2016.  
435 Population genetic structure of apple scab (*Venturia inaequalis* (Cooke) G. Winter) in Iran.  
436 PLoS One 11:e0160737.
- 437 17. EPPO. 2014. PM 7/98 (2) Specific requirements for laboratories preparing accreditation for a  
438 plant pest diagnostic activity. EPPO Bulletin 44:117-114.
- 439 18. Fernandez-Molina, J. V., Abad-Diaz-de-Cerio, A., Sueiro-Olivares, M., Pellon, A., Ramirez-  
440 Garcia, A., Garaizar, J., Pemán J., Hernando, F. L., Rementeria A. 2014. Rapid and specific  
441 detection of section *Fumigati* and *Aspergillus fumigatus* in human samples using a new  
442 multiplex real-time PCR. Diagn. Microbiol. Infect. Dis. 80:111-118.
- 443 19. Gadoury, D. M., and Machardy, W. E. 1986. Forecasting ascospore dose of *Venturia*  
444 *inaequalis* in commercial apple orchards. Phytopathology. 76:112-118.
- 445 20. Gladieux, P., Caffier, V., Devaux, M., and Le Cam, B. 2010. Host-specific differentiation  
446 among populations of *Venturia inaequalis* causing scab on apple, pyracantha and loquat.  
447 Fung. Genet. Biol. 47:511-521.

21. Gusberti, M., Patocchi, A., Gessler, C., and Brogini, G. A. L. 2012. Quantification of *Venturia inaequalis* growth in *Malus × domestica* with quantitative real-time polymerase chain reaction. *Plant Dis.* 96:1791-1797.
22. Huang, C. M., Liao, D. J, Wu, H. S., Shen, W. C., and Chung, C. L. 2016. Cyclone-based spore trapping, quantitative real-time polymerase chain reaction and high resolution melting analysis for monitoring airborne inoculum of *Magnaporthe oryzae*. *Ann. Appl. Biol.* 169:75-90.
23. Kermekchiev, M. B., Kirilova, L., Vail, E. E., and Barnes, W. M. 2009. Mutants of Taq DNA polymerase resistant to PCR inhibitors allow DNA amplification from whole blood and crude soil samples. *Nucleic Acids Res.* 37:e40.
24. Klosterman, S. J., Anchieta, A. G., McRoberts, N., Koike, S. T., Subbarao, K. V., VoglMayr, H., Choi, Y. J., Thines, M., and Martin, F. N. 2014. Coupling spore traps and quantitative PCR assays for detection of the downy mildew pathogens of spinach (*Peronospora effusa*) and beet (*Peronospora schachtii*) *Phytopathology*, 104:1349-1359.
25. Koh, H. S., Sohn, S. H., Lee, Y. S., Koh, Y. J., Song, J. H., and Jung, J.S. 2013. Specific and sensitive detection of *Venturia nashicola*, the scab fungus of Asian pears, by nested PCR. *Plant Pathol. J.* 29:357-363.
26. Kuzdraliński, A., Kot, A., Szczerba, H., Nowak, M., and Muszyńska, M. 2017. A review of conventional PCR assays for the detection of selected phytopathogens of wheat. *J. Mol. Microbiol. Biotech.* 27:175-189.
27. MacHardy, W.E. 1996. Models to predict ascospore maturity. In *Apple scab: biology, epidemiology, and management*. Pages 251 in MacHardy W.E. (eds.), American Phytopathological Society (APS Press), St. Paul, Minnesota USA.
28. MacHardy, W. E., Gadoury, D. M., and Gessler, C. 2001. Parasitic and biological fitness of *Venturia inaequalis*: relationship to disease management strategies. *Plant Dis.*85: 1036-1051.
29. Martínez, N., Martín, M. C., Herrero, A., Fernández, M., Alvarez, M. A., and Ladero, V. 2011. qPCR as a powerful tool for microbial food spoilage quantification: significance for food quality. *Trends Food Sci. Technol.* 22:367-376.
30. McDevitt, J. J., Lees, P. S. J., Merz, W. G. and Schwab, K. J. 2007. Inhibition of quantitative PCR analysis of fungal conidia associated with indoor air particulate matter. *Aerobiologia*, 23:35-45.
31. Meitz-Hopkins, J. C., von Diest, S. G., Koopman, T. A., Bahramisharif, A., and Lennox, C. L. 2014. A method to monitor airborne *Venturia inaequalis* ascospores using volumetric spore traps and quantitative PCR *Eur. J. Plant Pathol.* 140:527-541.

32. Mills, W. D., and Laplante, A. A. 1951. Diseases and insects in the orchard. Cornell Extension Bulletin, 711, 20-27.
33. Mirmajlessi, S., Loit, E., Mand, M., and Mansouripour, S. 2015. Real-time PCR applied to study on plant pathogens: potential applications in diagnosis - a review. Plant Protect. Sci. 51:177-90.
34. Parker, D. M., Hilber, U. W., Bodmer, M., Smith, F. D., Yao, C., and Köller, W. 1995. Production and transformation of conidia of *Venturia inaequalis*. Phytopathology 85:87-91.
35. Postollec, F., Falentin, H., Pavan, D., Combrisson, J., and Sohier, D. 2011. Recent advances in quantitative PCR (qPCR) applications in food microbiology. Food Microbiol. 28:848-61.
36. Rogers, S. L., Atkins, S. D. and West, J. S. 2009. Detection and quantification of airborne inoculum of *Sclerotinia sclerotiorum* using quantitative PCR Plant Pathol. 58:324-331.
37. Schnabel, G., Schnabel, E. L., and Jones, A. L. 1999. Characterization of ribosomal DNA from *Venturia inaequalis* and its phylogenetic relationship to rDNA from other tree-fruit *Venturia* species. Phytopathology. 89:100-108.
38. Selma, M. V., Martínez-Culebras, P. V., and Aznar, R., 2008. Real-time PCR based procedures for detection and quantification of *Aspergillus carbonarius* in wine grapes. Int. J. Food Microbiol. 122:126-134.
39. Soltany-Rezaee-Rad, M., Sepehrizadeh, Z., Mottaghi-Dastjerdi, N., Yazdi, M.T., and Seyatesh N. 2015. Comparison of SYBR Green and TaqMan real-time PCR methods for quantitative detection of residual CHO host-cell DNA in biopharmaceuticals Biologicals 43:130-135.
40. Springer, J., Goldenberger, D., Schmidt, F., Weisser, M., Wehrle-Wieland, E., Einsele, H., Frei, R., and Löffler, J. 2016. Development and application of two independent real-time PCR assays to detect clinically relevant Mucorales species. J. Med. Microbiol. 65:227-234.
41. Stehmann, C., Pennycook, S., and Plummer, K. M. 2001. Molecular identification of a sexual interloper: The pear pathogen, *Venturia pirina*, has sex on apple. Phytopathology 91:633-641.
42. Suarez, M.B., Walsh, K., Boonham, N., O'Neill, T., Pearson, S., and Barker, I. 2005. Development of real-time PCR (TaqMan (R)) assays for the detection and quantification of *Botrytis cinerea* in planta. Plant Physiol. Biochem. 43:890-899.
43. Sutton, T.B., Aldwinckle, H.S., Agnello, A.M., and Walgenbach, J.F. 2014. Compendium of apple and pear diseases and pests. 2<sup>nd</sup> Ed. American Phytopathological Society, St Paul, MN, USA, 218 pp.
44. Tenzer, I. and Gessler, C. 1999. Genetic diversity of *Venturia inaequalis* across Europe. Eur. J. Plant Pathol. 105:545-552.

45. Tenzer, I., degli Ivanissevich, S., Morgante, M., and Gessler, C. 1999. Identification of microsatellite markers and their application to population genetics of *Venturia inaequalis*. *Phytopathology* 89:748-753.
46. Turan, C., Menghini, M., Gazzetti, K., Ceredi, G., Mari, M., and Collina M. 2019. First identification of *Venturia asperata* from atypical scab-like symptoms in Italian apple orchards. *E. J. Plant Pathol.* 153:299-305.
47. White, T. J., Bruns, T., Lee, S., and Taylor, J. 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. Pages 315-322, in: *PCR Protocols: a Guide to Methods and Applications*. Innis, M. A., Gelfand, D. H., Sninsky, J. J., White, T. J. (eds.), Academic Press, San Diego, USA.
48. Williams, R. H., Ward, E., and McCartney, H. A., 2001. Methods for integrated air sampling and DNA analysis for detection of airborne fungal spores. *Appl. Environ. Microbiol.* 67:2453-2459.

530  
531  
532

## Tables

Table 1 – Strain ID, host, source of isolation and origin of the fungal species used in this study to develop the TaqMan real-time PCR assay.

Species	Strain ID	Host	Source	Origin
<i>Venturia inaequalis</i>	1b1	<i>Malus x domestica</i> ‘Ambrosia’	Leaf	Italy
<i>Venturia inaequalis</i>	1b5	<i>Malus x domestica</i> ‘Ambrosia’	Leaf	Italy
<i>Venturia inaequalis</i>	1b10	<i>Malus x domestica</i> ‘Ambrosia’	Leaf	Italy
<i>Venturia inaequalis</i>	1b7	<i>Malus x domestica</i> ‘Ambrosia’	Leaf	Italy
<i>Venturia inaequalis</i>	1b6	<i>Malus x domestica</i> ‘Ambrosia’	Leaf	Italy
<i>Venturia inaequalis</i>	116_c2	<i>Malus x domestica</i> ‘Ambrosia’	Leaf	Italy
<i>Venturia inaequalis</i>	216_b4	<i>Malus x domestica</i> ‘Golden Delicious’	Leaf	Italy
<i>Venturia inaequalis</i>	3b6	<i>Malus x domestica</i> ‘Ambrosia’	Leaf	Italy
<i>Venturia inaequalis</i>	3b2	<i>Malus x domestica</i> ‘Ambrosia’	Leaf	Italy
<i>Venturia inaequalis</i>	3b5	<i>Malus x domestica</i> ‘Ambrosia’	Leaf	Italy
<i>Venturia inaequalis</i>	516_6	<i>Malus x domestica</i> ‘Ambrosia’	Leaf	Italy
<i>Venturia inaequalis</i>	516_2	<i>Malus x domestica</i> ‘Ambrosia’	Leaf	Italy
<i>Venturia inaequalis</i>	516_3	<i>Malus x domestica</i> ‘Ambrosia’	Leaf	Italy
<i>Venturia inaequalis</i>	3aa	<i>Malus x domestica</i> ‘Golden Delicious’	Leaf	Italy
<i>Venturia inaequalis</i>	3a1	<i>Malus x domestica</i> ‘Golden Delicious’	Leaf	Italy
<i>Venturia inaequalis</i>	3ac	<i>Malus x domestica</i> ‘Golden Delicious’	Leaf	Italy
<i>Venturia inaequalis</i>	416_a12	<i>Malus x domestica</i> ‘Mondial Gala’	Leaf	Italy
<i>Venturia inaequalis</i>	416_a5	<i>Malus x domestica</i> ‘Mondial Gala’	Leaf	Italy
<i>Venturia inaequalis</i>	416_a9	<i>Malus x domestica</i> ‘Mondial Gala’	Leaf	Italy
<i>Venturia inaequalis</i>	516_h	<i>Malus x domestica</i> ‘Ambrosia’	Leaf	Italy
<i>Venturia inaequalis</i>	516_4	<i>Malus x domestica</i> ‘Ambrosia’	Leaf	Italy
<i>Venturia inaequalis</i>	516_5	<i>Malus x domestica</i> ‘Ambrosia’	Leaf	Italy
<i>Venturia inaequalis</i>	3af	<i>Malus x domestica</i> ‘Golden Delicious’	Leaf	Italy
<i>Venturia inaequalis</i>	416_b2	<i>Malus x domestica</i> ‘Golden Delicious’	Leaf	Italy
<i>Venturia inaequalis</i>	416_a1	<i>Malus x domestica</i> ‘Mondial Gala’	Leaf	Italy
<i>Venturia inaequalis</i>	516_10	<i>Malus x domestica</i> ‘Ambrosia’	Leaf	Italy
<i>Venturia inaequalis</i>	1b2	<i>Malus x domestica</i> ‘Ambrosia’	Leaf	Italy
<i>Venturia inaequalis</i>	416_a13	<i>Malus x domestica</i> ‘Mondial Gala’	Leaf	Italy
<i>Venturia inaequalis</i>	1b9	<i>Malus x domestica</i> ‘Ambrosia’	Leaf	Italy
<i>Venturia inaequalis</i>	3b3	<i>Malus x domestica</i> ‘Ambrosia’	Leaf	Italy
<i>Venturia inaequalis</i>	3b4	<i>Malus x domestica</i> ‘Ambrosia’	Leaf	Italy
<i>Venturia inaequalis</i>	3b10	<i>Malus x domestica</i> ‘Ambrosia’	Leaf	Italy
<i>Venturia inaequalis</i>	1b14	<i>Malus x domestica</i> ‘Ambrosia’	Leaf	Italy
<i>Venturia inaequalis</i>	416_a3	<i>Malus x domestica</i> ‘Mondial Gala’	Leaf	Italy
<i>Venturia inaequalis</i>	1b13	<i>Malus x domestica</i> ‘Ambrosia’	Leaf	Italy
<i>Venturia inaequalis</i>	CBS 815.69	<i>Malus sylvestris</i>	Fruit	The Netherlands
<i>Venturia cerasi</i>	CBS 444.54	<i>Prunus cerasus</i>	Unknown	Germany
<i>Venturia pirina</i>	CBS120.825	<i>Pyrus communis</i>	Unknown	Brazil

Species	Strain ID	Host	Source	Origin
<i>Venturia nashicola</i>	CBS 794.84	<i>Pyrus serotina</i> var <i>Culta</i>	Unknown	Japan
<i>Venturia asperata</i>	IRHS 2345	<i>Malus x domestica</i>	Fruit	France
<i>Venturia carpophila</i>	CBS 497.62	<i>Prunus mirabelle</i>	Unknown	Switzerland
<i>Monilia fructicola</i>	MON1	<i>Malus x domestica</i>	Leaf	Italy
<i>Monilia laxa</i>	LAXA3	<i>Prunus persica</i>	Fruit	Italy
<i>Botryosphaeria dothidea</i>	BOTRYO1	<i>Malus x domestica</i>	Leaf	Italy
<i>Sclerotinia sclerotiorum</i>	SCLA2	Unknown	Unknown	Italy
<i>Colletotrichum fioriniae</i>	COLLRIV	<i>Malus x domestica</i>	Fruit	Italy
<i>Epicoccum nigrum</i>	EPI2	<i>Malus x domestica</i>	Leaf	Italy
<i>Stemphylium</i> sp.	STEM	<i>Pyrus communis</i>	Fruit	Italy
<i>Cladosporium cladosporioides</i>	CLAD1	<i>Malus x domestica</i>	Leaf	Italy
<i>Alternaria mali</i>	CBS 106.24	<i>Malus sylvestris</i>	Unknown	The USA
<i>Alternaria</i> sp.	MALT1	<i>Malus x domestica</i>	Leaf	Italy
<i>Botrytis cinerea</i>	BOT1	<i>Malus x domestica</i>	Leaf	Italy
<i>Phoma</i> sp.	PH5	<i>Malus x domestica</i>	Leaf	Italy
<i>Fusarium equiseti</i>	3FEQS	Unknown	Unknown	Italy
<i>Penicillium expansum</i>	PEX16	<i>Malus x domestica</i>	Leaf	Italy
<i>Ramularia mali</i>	RM2	<i>Malus x domestica</i>	Fruit	Italy

533

534



535 Table 2. Primer pairs and probes, designed on *EF1-α* gene used for the detection of *V. inaequalis* with  
 536 the TaqMan real-time assay.

Primers and Probes	Amplicon (bp)	Sequence (5'-3')
F 3.2/R11	191 bp	ACCCGGATTTCATTTCGAAACT GCAATCGTTAGCATCGTCATAGTG
F 4.4/R11	154 bp	TTTTGCACTGTGGCAGCCC GCAATCGTTAGCATCGTCATAGTG
F 1/R11	86 bp	CACTTCCCCGCTATTACAGT GCAATCGTTAGCATCGTCATAGTG
F2.3/R6	127 bp	TTGCCCCTCCAAAATTACAGTG GGCGGCTTCCTATTGCAATC
F1/R6	100 bp	CACTTCCCCGCTATTACAGT GGCGGCTTCCTATTGCAATC
F4.3/R6	171 bp	AAATTTTGCACTGTGGCAGC GGCGGCTTCCTATTGCAATC
F 2.2/R6	127 bp	TTGCCCCTCCAAAATTACAGT GGCGGCTTCCTATTGCAATC
Vina	-	[FAM] - AGCCCAACTTTCTCCGGTCC - [NF-MGB]
Ven1	-	[FAM] - CTCAAGGCAGCCCAACTTTCTCCGGT - [BHQ1]

537

538 **Table 3** – Number of *V. inaequalis* cells estimated using the optical microscope compared with the  
539 number of cells estimated with the TaqMan real-time assay for the analyses of spore trap samples,  
540 and repeatability of the assay, expressed as the number of positive amplifications in six different  
541 reactions.

Sample	Estimated number of cells (optical microscope)	Estimated number of cells (TaqMan assay)	Number of positive results
ST1	$2.73 \times 10^4$	$5.49 \times 10^3$	6/6
ST2	$5.21 \times 10^4$	$6.88 \times 10^3$	6/6
ST3	$1.31 \times 10^4$	$1.12 \times 10^4$	5/6
ST4	$7.94 \times 10^3$	$4.42 \times 10^3$	6/6
ST9	$1.30 \times 10^3$	$2.72 \times 10^3$	3/6
ST10	$2.05 \times 10^3$	$4.80 \times 10^3$	3/6
ST10/2	$4.67 \times 10^2$	*	*
ST11	$8.92 \times 10^2$	$1.66 \times 10^3$	2/6
ST12	$8.51 \times 10^3$	$9.34 \times 10^3$	1/6
ST13	$1.37 \times 10^3$	$2.58 \times 10^3$	4/6
ST17	$2.83 \times 10^3$	$5.14 \times 10^3$	2/6
ST21	$1.09 \times 10^4$	$7.93 \times 10^3$	2/6
ST27	$1.36 \times 10^3$	$3.70 \times 10^3$	2/6
ST28	$2.21 \times 10^3$	$2.52 \times 10^3$	2/6

542  
543 \*the extraction failed for the ST10/2 sample.

## Captions

Figure 1. Standard curve obtained with genomic DNA of the *V. inaequalis* 1b14 strain showing the correlation between the *V. inaequalis* DNA amount and the Ct values (the standard deviation values range from 0.04 to 0.22, thus the standard deviation bars are too small to display on the graph).

Figure 2. Standard curve obtained with genomic DNA of the *V. inaequalis* 1b14 strain diluted in *Malus x domestica* DNA ‘Ambrosia’, ‘Golden Delicious’ and Mondial Gala. The *V. inaequalis* DNA amount is plotted against the Ct values (the standard deviation values range from 0.03 to 0.53, thus the standard deviation bars are too small to display on the graph).

Figure 3. Detection of *V. inaequalis* with the TaqMan real-time PCR on: **A.** naturally infected leaves of *Malus x domestica* ‘Ambrosia’, ‘Golden Delicious’ and ‘Mondial Gala’; **B.** asymptomatic leaves of *Malus x domestica* ‘Ambrosia’, ‘Golden Delicious’ and ‘Fiorina’. The values are expressed as the mean  $\pm$ SD (n=9).

Figure 4. Detection of *V. inaequalis* with the TaqMan real-time PCR on *Malus x domestica* ‘Ambrosia’ leaf disks artificially inoculated at different concentrations. The values are expressed as the mean  $\pm$ SD (n=9).

562 **Supplementary Tables**  
563 **Supplementary Table 1** - Results of the q-PCR amplification (Ct values  $\pm$  standard deviation) with  
564 the target DNA ( $\cong$  20 ng) of different *V. inaequalis* strains used to evaluate the specificity of the  
565 TaqMan real-time assay.

Sample ID	Species	Cultivar	Ct Mean $\pm$ SD*
1b1	<i>V. inaequalis</i>	Ambrosia	18.57 $\pm$ 0.04
1b5	<i>V. inaequalis</i>	Ambrosia	18.23 $\pm$ 0.24
1b10	<i>V. inaequalis</i>	Ambrosia	17.42 $\pm$ 0.04
1b7	<i>V. inaequalis</i>	Ambrosia	17.13 $\pm$ 0.23
1b6	<i>V. inaequalis</i>	Ambrosia	17.58 $\pm$ 0.19
116_c2	<i>V. inaequalis</i>	Ambrosia	18.12 $\pm$ 0.30
216_b4	<i>V. inaequalis</i>	Ambrosia	17.32 $\pm$ 0.20
3b6	<i>V. inaequalis</i>	Ambrosia	17.14 $\pm$ 0.20
3b2	<i>V. inaequalis</i>	Ambrosia	18.22 $\pm$ 0.11
3b5	<i>V. inaequalis</i>	Ambrosia	17.20 $\pm$ 0.13
516_6	<i>V. inaequalis</i>	Ambrosia	18.35 $\pm$ 0.30
516_2	<i>V. inaequalis</i>	Ambrosia	17.41 $\pm$ 0.13
516_3	<i>V. inaequalis</i>	Ambrosia	17.81 $\pm$ 0.17
3aa	<i>V. inaequalis</i>	Golden Delicious	17.49 $\pm$ 0.22
3a1	<i>V. inaequalis</i>	Golden Delicious	17.79 $\pm$ 0.03
3ac	<i>V. inaequalis</i>	Golden Delicious	17.52 $\pm$ 0.12
416_a12	<i>V. inaequalis</i>	Mondial Gala	18.49 $\pm$ 0.10
416_a5	<i>V. inaequalis</i>	Mondial Gala	17.92 $\pm$ 0.31
416_a9	<i>V. inaequalis</i>	Mondial Gala	18.29 $\pm$ 0.30
516_h	<i>V. inaequalis</i>	Ambrosia	17.44 $\pm$ 0.37
516_4	<i>V. inaequalis</i>	Ambrosia	18.14 $\pm$ 0.37
516_5	<i>V. inaequalis</i>	Ambrosia	18.09 $\pm$ 0.22
3af	<i>V. inaequalis</i>	Golden Delicious	17.75 $\pm$ 0.31
416_b2	<i>V. inaequalis</i>	Golden Delicious	17.76 $\pm$ 0.36
416_a1	<i>V. inaequalis</i>	Mondial Gala	17.76 $\pm$ 0.39
516_10	<i>V. inaequalis</i>	Ambrosia	18.45 $\pm$ 0.32
1b2	<i>V. inaequalis</i>	Ambrosia	17.66 $\pm$ 0.23
416_a13	<i>V. inaequalis</i>	Mondial Gala	17.47 $\pm$ 0.24
1b9	<i>V. inaequalis</i>	Ambrosia	17.94 $\pm$ 0.11
3b3	<i>V. inaequalis</i>	Ambrosia	17.11 $\pm$ 0.24
3b4	<i>V. inaequalis</i>	Ambrosia	17.68 $\pm$ 0.11
3b10	<i>V. inaequalis</i>	Ambrosia	17.13 $\pm$ 0.25
1b14	<i>V. inaequalis</i>	Ambrosia	18.00 $\pm$ 0.39
416_a3	<i>V. inaequalis</i>	Mondial Gala	17.85 $\pm$ 0.28
1b13	<i>V. inaequalis</i>	Ambrosia	17.34 $\pm$ 0.07

566  
567 \* The values are expressed as the mean  $\pm$ SD (n=9)

568 **Supplementary Table 2** – Standard curve obtained for the quantification of the *V. inaequalis* 1b14  
569 strain using the TaqMan real-time PCR assay developed by Gusberti et al. (2012). Ct mean values ±  
570 standard deviation.

<i>V. inaequalis</i> DNA concentration (ng)	Ct values ± SD
10	20.13 ± 0.11
1	23.24 ± 0.12
0.1	26.74 ± 0.10
0.01	30.10 ± 0.04
0.001	33.69 ± 0.71
0.0001	36.21 ± 0.86

571

572 **Supplementary Table 3** – Ct mean values  $\pm$  standard deviation and the estimated number of *V.*  
573 *inaequalis* cells/ $\mu$ L  $\pm$  standard deviation for the symptomatic *Malus x domestica* ‘Ambrosia’,  
574 ‘Mondial Gala’ and ‘Golden Delicious’ leaves obtained using the TaqMan real-time PCR assay.

Sample ID	Cultivar	Ct mean $\pm$ SD*	<i>V. inaequalis</i> cells/ $\mu$ L $\pm$ SD*
1b	Ambrosia	21.93 $\pm$ 0.10	1.67 x 10 <sup>4</sup> $\pm$ 1137.31
3b	Ambrosia	25.15 $\pm$ 0.20	1.59 x 10 <sup>3</sup> $\pm$ 209.73
Amba17	Ambrosia	19.89 $\pm$ 0.27	8.63 x 10 <sup>4</sup> $\pm$ 15945.86
316s2	Ambrosia	18.86 $\pm$ 0.18	1.25 x 10 <sup>5</sup> $\pm$ 15918.36
216	Ambrosia	25.38 $\pm$ 0.54	1.91 x 10 <sup>3</sup> $\pm$ 777.22
516	Ambrosia	22.35 $\pm$ 0.64	2.19 x 10 <sup>4</sup> $\pm$ 8523.70
Amb1	Ambrosia	23.89 $\pm$ 0.34	5.59 x 10 <sup>3</sup> $\pm$ 1262.95
Amb4	Ambrosia	25.29 $\pm$ 0.90	1.53 x 10 <sup>3</sup> $\pm$ 829.27
Col1	Ambrosia	18.71 $\pm$ 0.29	1.36 x 10 <sup>5</sup> $\pm$ 26719.38
Col2	Ambrosia	20.37 $\pm$ 0.38	5.90 x 10 <sup>4</sup> $\pm$ 1673.62
Col3	Ambrosia	19.12 $\pm$ 0.48	1.23 x 10 <sup>5</sup> $\pm$ 40605.49
416a1	Gala	27.96 $\pm$ 0.12	3.20 x 10 <sup>3</sup> $\pm$ 476.83
416a2	Gala	18.12 $\pm$ 0.05	2.48 x 10 <sup>5</sup> $\pm$ 8191.52
GalaA	Gala	20.76 $\pm$ 0.79	4.31 x 10 <sup>4</sup> $\pm$ 17722.88
Gala1	Gala	25.81 $\pm$ 0.03	9.33 x 10 <sup>2</sup> $\pm$ 261.42
Gala2	Gala	24.85 $\pm$ 0.24	1.82 x 10 <sup>3</sup> $\pm$ 311.25
Gala3	Gala	24.98 $\pm$ 0.18	2.01 x 10 <sup>3</sup> $\pm$ 399.74
Gala4	Gala	27.79 $\pm$ 0.49	2.38 x 10 <sup>2</sup> $\pm$ 73.01
Gold11	Golden Delicious	21.04 $\pm$ 0.39	3.16 x 10 <sup>4</sup> $\pm$ 8540.33
116	Golden Delicious	18.32 $\pm$ 0.24	2.07 x 10 <sup>5</sup> $\pm$ 34735.35
1a	Golden Delicious	34.35 $\pm$ 0.37	0.28 x 10 <sup>1</sup> $\pm$ 0.67
416b2	Golden Delicious	22.58 $\pm$ 0.17	1.13 x 10 <sup>4</sup> $\pm$ 1400.69
416b3	Golden Delicious	23.53 $\pm$ 0.23	5.85 x 10 <sup>3</sup> $\pm$ 977.49
Gold22	Golden Delicious	21.26 $\pm$ 0.35	3.35 x 10 <sup>4</sup> $\pm$ 8582.51
3a	Golden Delicious	24.25 $\pm$ 0.23	3.46 x 10 <sup>3</sup> $\pm$ 582.48
Gold3a	Golden Delicious	20.17 $\pm$ 0.31	6.19 x 10 <sup>4</sup> $\pm$ 13818.83
GoldA	Golden Delicious	20.29 $\pm$ 0.14	5.79 x 10 <sup>4</sup> $\pm$ 5549.25
Gold4	Golden Delicious	31.65 $\pm$ 0.65	3.30 x 10 <sup>1</sup> $\pm$ 337.26
Gold1	Golden Delicious	26.12 $\pm$ 0.43	1.05 x 10 <sup>3</sup> $\pm$ 337.26
Mti2	Golden Delicious	36.36 $\pm$ 0.84	7.58 x 10 <sup>2</sup> $\pm$ 359.66
MtiA	Golden Delicious	26.11 $\pm$ 0.44	3.21 x 10 <sup>4</sup> $\pm$ 9293.85
MtiB	Golden Delicious	21.51 $\pm$ 0.06	7.04 x 10 <sup>3</sup> $\pm$ 294.32
MtiC	Golden Delicious	28.38 $\pm$ 0.07	1.82 x 10 <sup>2</sup> $\pm$ 0.39

575 \* Values are expressed as the mean  $\pm$ SD (n=9).



576 **Supplementary Table 4** – Ct mean values  $\pm$  standard deviation and the estimated number of *V.*  
577 *inaequalis* cells/ $\mu$ L  $\pm$  standard deviation for the asymptomatic *Malus x domestica* ‘Fiorina’,  
578 ‘Ambrosia’ and ‘Golden Delicious’ leaves obtained using the TaqMan real-time PCR assay.

Sample ID	Cultivar	Ct mean $\pm$ SD*	<i>V. inaequalis</i> cells/ $\mu$ L $\pm$ SD*
As1	Fiorina	27.21 $\pm$ 0.42	4.23 x 10 <sup>2</sup> $\pm$ 56.27
As2	Fiorina	28.86 $\pm$ 0.11	1.30 x 10 <sup>2</sup> $\pm$ 10.25
As3	Fiorina	27.68 $\pm$ 0.12	2.98 x 10 <sup>2</sup> $\pm$ 26.45
As4	Ambrosia	25.56 $\pm$ 0.07	1.31 x 10 <sup>3</sup> $\pm$ 6.18
As5	Ambrosia	23.81 $\pm$ 0.21	4.50 x 10 <sup>3</sup> $\pm$ 175.73
As6	Fiorina	26.20 $\pm$ 0.18	8.40 x 10 <sup>2</sup> $\pm$ 10.92
As7	Fiorina	34.48 $\pm$ 0.07	0.03 x 10 <sup>2</sup> $\pm$ 0.22
As8	Fiorina	33.18 $\pm$ 0.12	0.06 x 10 <sup>2</sup> $\pm$ 0.55
As9	Fiorina	34.94 $\pm$ 0.13	0.02 x 10 <sup>2</sup> $\pm$ 0.29
As10	Ambrosia	24.32 $\pm$ 0.16	3.14 x 10 <sup>3</sup> $\pm$ 157.47
As11	Ambrosia	24.15 $\pm$ 0.16	3.53 x 10 <sup>3</sup> $\pm$ 7.29
As12	Ambrosia	24.16 $\pm$ 0.15	3.51 x 10 <sup>3</sup> $\pm$ 15.75
As13	Fiorina	33.54 $\pm$ 0.95	0.06 x 10 <sup>2</sup> $\pm$ 1.12
As14	Fiorina	28.89 $\pm$ 0.18	1.28 x 10 <sup>2</sup> $\pm$ 15.92
As15	Golden Delicious	28.11 $\pm$ 0.17	2.21 x 10 <sup>2</sup> $\pm$ 25.84
As16	Golden Delicious	29.87 $\pm$ 0.51	6.65 x 10 <sup>1</sup> $\pm$ 22.08
As17	Fiorina	29.91 $\pm$ 0.33	6.31 x 10 <sup>1</sup> $\pm$ 13.57
As18	Ambrosia	28.91 $\pm$ 0.05	1.26 x 10 <sup>2</sup> $\pm$ 33.85
As19	Ambrosia	27.57 $\pm$ 0.14	3.21 x 10 <sup>2</sup> $\pm$ 30.04
As20	Ambrosia	25.90 $\pm$ 0.20	1.03 x 10 <sup>3</sup> $\pm$ 43.97
As21	Golden Delicious	26.72 $\pm$ 0.13	5.82 x 10 <sup>2</sup> $\pm$ 39.72
As22	Golden Delicious	27.85 $\pm$ 0.07	2.64 x 10 <sup>2</sup> $\pm$ 13.54
As23	Golden Delicious	26.17 $\pm$ 0.25	8.61 x 10 <sup>2</sup> $\pm$ 15.32
As24	Golden Delicious	27.64 $\pm$ 0.07	3.04 x 10 <sup>2</sup> $\pm$ 15.71
As25	Golden Delicious	26.60 $\pm$ 0.19	6.34 x 10 <sup>2</sup> $\pm$ 89.10
As26	Ambrosia	25.55 $\pm$ 0.67	1.43 x 10 <sup>3</sup> $\pm$ 6.01
As27	Golden Delicious	27.31 $\pm$ 0.19	3.86 x 10 <sup>2</sup> $\pm$ 51.03
As28	Golden Delicious	28.00 $\pm$ 0.18	2.39 x 10 <sup>2</sup> $\pm$ 29.63
As29	Golden Delicious	30.81 $\pm$ 0.30	3.37 x 10 <sup>1</sup> $\pm$ 6.58
As30	Golden Delicious	29.47 $\pm$ 0.21	8.50 x 10 <sup>1</sup> $\pm$ 12.54

579  
580 \* Values are expressed as the mean  $\pm$ SD (n=9).