

Washington University in St. Louis

## Washington University Open Scholarship

---

Biology Faculty Publications & Presentations

Biology

---

1-2009

### Small RNA-directed heterochromatin formation in the context of development: what flies might learn from fission yeast

Kathryn Huisinga

*Washington University in St. Louis*

Sarah C.R. Elgin

*Washington University in St. Louis, [selgin@wustl.edu](mailto:selgin@wustl.edu)*

Follow this and additional works at: [https://openscholarship.wustl.edu/bio\\_facpubs](https://openscholarship.wustl.edu/bio_facpubs)



Part of the [Biology Commons](#)

---

#### Recommended Citation

Huisinga, Kathryn and Elgin, Sarah C.R., "Small RNA-directed heterochromatin formation in the context of development: what flies might learn from fission yeast" (2009). *Biology Faculty Publications & Presentations*. 195.

[https://openscholarship.wustl.edu/bio\\_facpubs/195](https://openscholarship.wustl.edu/bio_facpubs/195)

This Article is brought to you for free and open access by the Biology at Washington University Open Scholarship. It has been accepted for inclusion in Biology Faculty Publications & Presentations by an authorized administrator of Washington University Open Scholarship. For more information, please contact [digital@wumail.wustl.edu](mailto:digital@wumail.wustl.edu).

Published in final edited form as:

*Biochim Biophys Acta*. 2009 January ; 1789(1): 3–16. doi:10.1016/j.bbagr.2008.08.002.

## Small RNA directed heterochromatin formation in the context of development: what flies might learn from fission yeast

Kathryn L. Huisinga<sup>1</sup> and Sarah C.R. Elgin

Department of Biology, Washington University, St Louis, MO 63130, USA

### Summary

A link between the RNAi system and heterochromatin formation has been established in several model organisms including *S. pombe* and *A. thaliana*. However, the data to support a role for small RNAs and the associated machinery in transcriptional gene silencing in animal systems is more tenuous. Using the *S. pombe* system as a model, we analyze the role of small RNA pathway components and associated small RNAs in regulating transposable elements and potentially directing heterochromatin formation at these elements in *Drosophila melanogaster*.

### Keywords

heterochromatin; *Drosophila melanogaster*; Small RNAs; *S. pombe*; transposable elements; cell-cycle

### Small non-coding RNA pathways

Pathways to generate small non-coding RNAs exist in a wide-range of eukaryotic organisms, from the yeast *Schizosaccharomyces pombe* to plants to humans. The only known exception among the established eukaryotic model organisms is *Saccharomyces cerevisiae*, which lacks the RNAi processing system, but does use antisense and non-coding RNAs to help regulate cellular processes[1,2]. Small RNAs and the pathways that generate them are implicated in a variety of epigenetic processes, including control of transposable elements, genome rearrangement, RNA-directed DNA methylation, quelling, heterochromatin formation, and nucleolar dominance and are studied in a wide variety of organisms, including *S. pombe*, *Tetrahymena*, *Drosophila melanogaster*, *Caenorhabditis elegans*, *Arabidopsis thaliana*, and mammalian systems. Three types of proteins are primarily involved in generating small RNAs. The first category is the RNaseIII family of nucleases, which include the Dicers and Drosha [3–5]. These enzymes are responsible for cutting longer double-stranded RNA (dsRNA) into smaller fragments of 20–30nt. Second are the dsRNA binding (DRB) proteins, which act as partners to the RNaseIII enzymes, and include loquacious (loqs), R2D2, and Pasha in *Drosophila*[6–11]. The third category is the Argonaute family of PAZ/Piwi domain containing proteins, which bind to the small dsRNAs and utilize that information to direct a variety of gene silencing effects. The PAZ domain is important for binding the small RNA, while the Piwi domain mediates the endonucleolytic cleavage, aka “slicer” activity, responsible for cutting the target RNAs identified by the small bound RNA. The Argonaute family proteins

<sup>1</sup>Corresponding author e-mail: khuising@wustl.edu, fax: (314) 935-4432.

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

are divided into two clades, the Piwi clade and the Ago clade, each having distinct functions as discussed below. Metazoans contain members from both clades, while the sole Argonaute in fission yeast and those in plants are all from the Ago clade [reviewed in 12].

To date, there have been three small RNA pathways described in animals (see Figure 1) [reviewed in 13]. The siRNA pathway is most familiar, as this is the pathway employed in RNAi knockdown experiments. This pathway processes long dsRNA using a Dicer enzyme (Dicer-2 in *Drosophila*) to generate short 21–23nt dsRNAs that are incorporated into an Argonaute family member-containing siRNA-induced silencing complex (siRISC, containing AGO2 in *Drosophila*). This complex utilizes the slicer activity of its Argonaute to destroy mRNA transcripts having sequence complimentary to the small RNAs. In flies, as in plants and worms, the endogenous function of this pathway includes acting as a viral defense system, as mutations in this pathway result in increased susceptibility to viral infection [reviewed in 14]. Very recent publications have implicated components of this pathway (Ago2 and Dcr2) in the generation of endogenous siRNAs (esiRNAs) derived from transposable elements, convergent transcripts, and long hairpin RNA genes in *Drosophila* embryos and somatic cells [15–20]. These papers are discussed in a recent review [21]. The details of this pathway and how it interfaces with the other pathways in *Drosophila* remain to be elucidated. The miRNA pathway uses RNA polymerase II (RNAPII) transcribed RNAs that fold back on themselves to form imperfect hairpins that are the source of dsRNA. In flies, these hairpins are processed first by the Drosha/Pasha complex and then by the Dicer-1 endonuclease and its DRB partner, loqs, to create short 21–23nt RNAs. These RNAs are loaded into the AGO1 containing miRISC, which blocks translation of mRNAs with imperfect pairing to the miRNA present in the complex. Depending upon the amount of base-pairing between the miRNA and the target mRNA, slicing of the mRNA can occur as well. The primary function of this pathway appears to be fine-tuning the expression of genes in a developmental context [reviewed in 22]. The third small RNA pathway present in animals is the piRNA pathway in which members of the Piwi clade of Argonaute proteins produce somewhat longer small RNAs (24–30nt) generated (at least in part) from specific genomic regions, termed piRNA clusters. Repeat associated siRNAs (rasiRNAs) identified in *Drosophila* are one class generated by the piRNA pathway [23]. piRNA clusters in flies are composed primarily of imperfect copies of transposable elements, including both retrotransposons and DNA transposons [24]. In mice there are different piRNA clusters expressed at different developmental stages, some of which are enriched for transposable elements, some of which are not (see below). The piRNA pathway appears to be critical for maintenance of germline stability, as mutations in this pathway result in fertility defects [25–29].

In this review we focus on the proposed role of small RNAs and of the proteins they are associated with in directing and maintaining the formation of heterochromatin. This relationship is well established in *S. pombe* [30], which serves as our paradigm for discussing the possible mechanisms. Often this function of small RNAs and the RNAi machinery is referred to as *transcriptional gene silencing*, as the end result is to shut off transcription via modification of chromatin. However, in the best-understood system, *S. pombe*, there is a post-transcriptional aspect to this process in that the regions of the genome destined to be transcriptionally silenced (i.e. the *dg-dh* pericentromeric repeats and the MAT locus) are actually transcribed during S-phase of the cell-cycle (see below). Here, we will consider the mechanisms that ultimately result in changes in chromatin structure with subsequent decreases in gene expression at a transcriptional level, as *transcriptional gene silencing*, while acknowledging that transcription of the silenced region and subsequent processing of these transcripts, can be involved in this process. One can, however, differentiate this mechanism from straightforward post-transcriptional gene silencing mechanisms, which function solely by degrading transcripts and do not result in any changes in chromatin structure or in the transcription level of the homologous region. In addition to *S. pombe*, links between small

RNAs and heterochromatin formation have been detected in several animal models, including flies, zebrafish, and mammalian systems. We review what is known about the relationship between small RNAs and heterochromatin formation and examine some unanswered questions about this relationship by focusing on the *S. pombe* and *Drosophila* systems, but include a brief examination of the links in the mammalian systems. Other recent reviews have examined the roles for small RNAs in epigenetic regulation in other model organisms in more detail [12, 31–33].

## Formation of heterochromatin in *S. pombe* utilizes cell-cycle regulated small RNAs

Heterochromatic regions of the genome in the model organism *S. pombe* include the silent mating type (MAT) loci, subtelomeres, the ribosomal DNA (rDNA), and the regions surrounding the centromeres (pericentric domains)[34]. Reporter genes located in these regions exhibit a variegating phenotype, characteristic of heterochromatin[35]. The chromatin of these regions is enriched for Swi6 [the *S. pombe* Heterochromatin Protein 1 (HP1) homologue], histone 3 methylated at lysine 9 (H3K9me), and Clr4, the *S. pombe* homologue of the Su(var) 3–9 histone methyltransferase [34]. All of these biochemical marks are associated with heterochromatin. A long-standing question is how these marks are targeted to the specific regions of a genome, which are appropriately heterochromatic, but not to other regions. Much work in fission yeast over the past several years has elucidated a role for the RNAi machinery and small RNAs in directing the formation of heterochromatin in the pericentric regions, where this pathway is required for heterochromatin formation[36]. The RNAi pathway is also required for establishment of heterochromatin at the silent mating-type loci, but here it plays a redundant role with the ATF/CREB pathway in maintaining the heterochromatic state[37,38].

Small RNAs (20–22nt) with identity to an endogenous heterochromatic sequence, the *dh* repeats at the pericentromere, were the first identified in *S. pombe*[39]. Subsequent characterization of the *S. pombe* small RNA population has identified not only RNAs with sequence homology to the repetitive *dg* and *dh* elements present at the outer (*otr*) regions of *S. pombe* centromeres, but also RNAs matching the inner (*inr*) centromeric repeats, the CenH region of the MAT locus, the rDNA and the subtelomeres[34,39]. A self-reinforcing feedback loop model for the generation of these small RNAs, suggesting how the RNAi processing machinery works to recruit heterochromatin factors to the pericentromere, has been proposed (Figure2)[40,41]. Transcription of the *dg-dh* repeats in the *otr* region of the centromeres is dependent on RNAPII, as mutations in subunits of RNAPII result in a loss of small RNAs and a loss of silencing[42,43]. At the *dg-dh* repeats, both the forward and reverse strands can be transcribed[36]. The reverse strand is transcribed even in the presence of heterochromatin, but transcription of the forward strand is repressed transcriptionally by Swi6 recruitment[36,42]. Ago1, the sole Argonaute family protein in *S. pombe*, interacts with the polymerase and plays a role in processing these transcripts via its slicing activity[44,45]. Mutations which disrupt the catalytic activity of Ago1 result in disruption of heterochromatin formation and the loss of spreading of heterochromatin into reporter genes inserted in the *otr* region[46]. Ago1 is a component of the RNA-induced transcriptional silencing (RITS) complex, which also includes Chp1, a chromodomain-containing protein which is dependent upon Clr4 mediated H3K9 methylation for localization to heterochromatic regions[47], and Tas3, a protein that appears to play a bridging role between Ago1 and Chp1 in the complex[48–50]. In addition, the RITS complex contains small RNAs with homology to the *dg-dh* repeats[50]. RITS interacts with a second complex, the RNA-directed RNA polymerase complex (RDRC), which contains Rdp1, an RNA-dependent RNA polymerase (RdRp) that uses the RNAPII transcript as a template to generate dsRNA[40,51]. RDRC contains two additional components, Cid12, a member of the polyA polymerase family of enzymes, and Hrr1, a putative helicase[51]. The Dicer enzyme interacts with RDRC[52] and processes the dsRNAs produced, thereby generating additional

small RNAs, which can then feed back into the RITS complex to be bound by Ago1. A second Ago1 containing complex, Argonaute siRNA chaperone (ARC), plays a role in this process by regulating the conversion of the small double-stranded RNA generated by Dicer into the single-stranded guide RNAs present in RITS[53]. The two additional proteins in ARC, Arb1 and Arb2, appear to play a role in regulating this conversion[53]. This process is thought to occur in *cis* at the pericentromeric region being transcribed. Chromatin immunoprecipitation experiments have demonstrated that many of these factors (RNAPII, RITS, and RDRC) are present at the silenced heterochromatic regions in *S. pombe*[34,41,43,47]. Evidence for a *cis* mechanism comes from the observation that tethering the RITS complex (via a Tas3- $\Delta$ N peptide fusion protein which binds to its cognate RNA binding site, the BoxB) to the nascent transcript of a *ura4* reporter gene drives formation of heterochromatin and transcriptional gene silencing, but only in *cis*, not in *trans*[54]. This *cis* restriction is regulated by the ribonuclease Eri1 (which presumably degrades excess siRNA), as Eri1 deletion strains are able to silence a *ura4* reporter in *trans* as well as the *ura4* reporter where the Tas3- $\Delta$ N peptide is tethered[54].

Small RNA processing is required for the formation of heterochromatin at the centromeric repeats in *S. pombe*; deletion of components of the RNAi pathway results in the loss of silencing of reporters inserted into this region and the loss of heterochromatic marks (i.e. Swi6 & Clr4 binding, H3K9me)[36,50]. It is proposed that the targeting of RITS, via the small RNAs with homology to the repeats, promotes the binding of Chp1 to methylated H3K9[50] (see Figure 2). However, the recruitment of the RNAi machinery and the Clr4 histone methyltransferase (HMT) which deposits this mark at the pericentromere are co-dependent, in that loss of one results in the loss of the other[34,40,41,55]. Additional insight into Clr4 recruitment comes from the observation that Clr4 exists in a protein complex, designated CLRC, with several additional factors including Rik1, a WD repeat containing protein that has been proposed to help recruit Clr4 to heterochromatic regions[56–58]. Also present in the complex are the cullin protein Cul4, a component of the cullin-dependent ligase (CDL) family of E3 ubiquitin ligases, and two additional components of the CDL family, Ned8, a neddylation factor which modifies cullins, and Pip1, a Ring finger protein[56,57]. Two additional proteins, Dos1/Clr8 (Raf1/Cmc1) and Dos2/Clr7 (Raf2/Cmc2), which were identified in screens for mutants that disrupt Swi6 localization and heterochromatin formation, are also in the CLRC complex[56,57,59, 60]. Loss of components of the CLRC complex results in disruption of heterochromatin formation at the pericentromere. A recent study has shown that recruitment of the Rik1 component of CLRC is dependent upon the RNAi pathway[55]. Mutations that disrupt Ago1 or Rdp1 activity result in the loss of Rik1 recruitment. Mutations in Cul4 disrupt the recruitment of Clr4, but not of Rik1, indicating a potential role for Cul4 in bridging a Rik1-Clr4 interaction. The binding of Rik1 correlates with increased transcription of repeat regions and coincides with RNAPII occupancy, lending support to the model that Rik1 plays an upstream, targeting role in the recruitment of Clr4. Therefore, it appears that the Rik1 subunit of CLRC mediates recruitment of the complex in an RNAi dependent manner. In addition, CLRC interacts with the RITS complex in a manner dependent upon the Tas3 subunit[55], which may also facilitate recruitment of CLRC to heterochromatic regions. Taken together, this data supports a model where the recruitment of CLRC and RITS are co-dependent.

A puzzling aspect of this model is exactly how RNAPII dependent transcription of the pericentric repeats occurs in a heterochromatic region, as heterochromatin packaging is generally considered inhibitory to the RNAPII transcription machinery[reviewed in 61,62]. Recent publications showing that RNAi-mediated heterochromatin formation appears to be regulated in a cell-cycle dependent manner, with transcription of the heterochromatic regions occurring during S-phase when the DNA is not yet completely packaged as heterochromatin, help to explain some of the apparent contradictions[63,64]. As cells enter mitosis, H3K9me and Swi6 binding are reduced; this reduction persists until S phase. H3K9me and Swi6 binding then gradually increase again until G2 when binding is maximal (Figure 2). There is an inverse

relationship between Swi6 binding and the presence of histone 3 phosphorylated at serine 10 (H3S10ph), which is highest during mitosis. The increase in H3S10ph coincides with the recruitment of condensin, which is implicated in mitotic chromosome condensation. Binding of RNAPII to the heterochromatic regions and the generation of heterochromatic transcripts occurs during S phase[63,64]. However, in a *clr4Δ* strain, the binding of RNAPII increases throughout the cell cycle, indicating that heterochromatin assembly does normally limit RNAPII occupancy as one would predict[63]. Rik1, which is critical for loading CLRC, also shows peak binding during S phase, coinciding with the RNAPII peak, whereas RITS binding increases during S phase and persists into G2, consistent with the data showing that Ago1 interacts with RNAPII. Clr4 and the CLRC component Raf2 show profiles similar to H3K9me and Swi6, with the least binding during mitosis, gradually increasing during S phase, with the peak of binding during G2. Concurrent with transcription by RNAPII during S phase is an enrichment of H3K36me and the Alp13/Eaf3 protein[63]. Analysis of RNAPII transcription at euchromatic genes has demonstrated that RNAPII targets the H3K36 methyltransferase Set2 to the transcribed region[65], which in turn recruits histone deacetylase complexes (HDACs) via interaction of the Alp13/Eaf3 chromodomain with the H3K36 methylated histone[66,67]. This recruitment presumably facilitates reassembly of chromatin following transcription. The report of H3K36me and Alp13/Eaf3 at the heterochromatic repeats indicates that a similar process is used when RNAPII is transcribing heterochromatic regions. These findings indicate that S phase is permissive for transcription of heterochromatic regions, but that the transcripts are processed to small RNAs by recruitment of Ago1 and other components, which then facilitate the recruitment of the heterochromatin machinery that is present at these regions during G2. These findings have significant implications for understanding heterochromatin formation and stability in multicellular organisms, which contain different cell types that are undergoing division at different rates or are terminally differentiated and have exited the cell cycle.

A key question in understanding the function of heterochromatin formation in the centromere region is the relationship between pericentric heterochromatin and the functional centromere. Recent work using a minichromosome transformation assay has added to the understanding of this relationship by demonstrating that heterochromatin proteins Swi6, Chp1, and Clr4, and the RNAi machinery, in particular Dicer, are necessary to promote CENP-A deposition and kinetochore assembly over the central domain of a fission yeast centromere on a naïve template [68]. This appears to be an absolute requirement only for establishing the centromere, as loss of these factors does not affect CENP-A deposition on endogenous centromeres or on minichromosome centromeres which have already incorporated CENP-A in a previous generation. Consistent with the requirement for heterochromatin, the *otr* region (which contains the *dg-dh* repeats), in addition to the *cnt* region where CENP-A binds, must be present on the naked template[68]. These results are in agreement with the previous observation that cells deleted for Ago1, Dicer or Rdp1, while viable, have defects in chromosome segregation[69, 70]. Interestingly, mutations in some of the *Drosophila* RNAi components with a potential role in heterochromatin formation exhibit defects in chromosome segregation as well as in gene silencing (see below).

## Generation of rasiRNAs in *Drosophila melanogaster*

Cloning and sequencing *Drosophila* small RNA populations over a developmental time course first identified the rasiRNA class of small RNAs in flies. These small RNAs, which can be considered a subclass of the larger piRNA class of small RNAs, are ~24–30nt long and have homology to repetitive sequences including transposable elements[71]. Additional characterization of the rasiRNAs has shown that they have only a 5' phosphate group; based on their resistance to periodate oxidation/ $\beta$ -elimination, it was concluded that they lack either a 2' or 3' hydroxyl group at the 3' terminus[72]. Subsequent studies have shown that the 3' end

of rasiRNAs is 2'-O-methylated by the *Drosophila* homologue of *Arabidopsis thaliana* HEN1, DmHen1/Pimet[73,74]. This class of small RNAs is primarily present in the ovaries, testes, and early embryos of flies, although a few RNAs from this class can be isolated from later stages of development[71]. Analysis of two specific types of rasiRNAs, those homologous to the *Suppressor of Stellate* locus (which controls expression of the X-linked *Stellate* gene in an RNAi dependent manner[75,76]) and those homologous to the LTR-containing retrotransposon *roo* (found in the germline of adult flies), demonstrated that the generation of these rasiRNAs in ovaries or testes was independent of several key genes in the siRNA pathway, including *Dcr-2*, its DRB partner *R2D2*, and *Ago2*, as well as *loqs*, encoding the DRB protein required to produce miRNAs[72]. Dependency upon *Dcr-1* was more difficult to examine, as flies homozygous null for *Dcr-1* are not viable. Nevertheless, using mitotic mutant clones in ovaries, it was demonstrated that *Dcr-1* as well is not required for production of the *roo* rasiRNA. The lack of dependency upon *Dcr-1* and *Dcr-2* is consistent with the fact that these enzymes previously had been shown to produce small RNAs of ~22nt[3]. Subsequent studies showed that the presence of wild type levels of these rasiRNAs is dependent upon two different members of the Piwi clade in the Argonaute family, *piwi* and *aubergine* (*aub*), as well as two putative RNA helicases, *spn-E* (aka *hls*), a member of the DE-H family of RNA-dependent ATPases[77], and *armitage* (*armi*), a homologue of the *Arabidopsis* SDE3 protein that contains an ATP-dependent domain distinct from DEA(H/D) box helicases[72,78]. Previous studies have demonstrated a role for *armi* in assembly of the siRISC complex from ovaries[79] and a requirement for *spn-E* in *Stellate* silencing[80] and RNAi activation during oocyte maturation[81].

Further understanding of how rasiRNAs are generated and what their biological role might be has come from sequencing the small RNAs associated with the Argonaute proteins from the Piwi clade in flies, Piwi, Aub, and AGO3. Purification and subsequent sequencing of the small RNAs associated with these proteins in ovaries or testes has shown that the large majority share identity with transposable elements (including both class I retrotransposons and class II DNA transposons) and other repetitive sequences present at heterochromatic loci[24,82–85]. Using the rasiRNAs that map to only one genomic location (i.e. uniquely mapping rasiRNAs), several hundred discrete loci where rasiRNAs are produced have been identified[24,85]. These loci are termed “piRNA clusters” or “rasiRNA clusters” and vary in size from several Kb to hundreds of Kb. The majority of these clusters are located in the  $\beta$ -heterochromatin present at pericentric regions, the fourth chromosome, and at telomeres, where there are numerous partial, inactive transposable elements. These clusters account for more than 92% of all of the sequenced piRNAs[24]. The largest cluster, located at cytological region 42AB, has homology to ~30% of all of the sequenced piRNAs and is 240kb in length[24]. Other prominent clusters include telomeric clusters on most arms (X, 2R, 2L, 3R and 4) that have homology to both telomere associated sequence (TAS) repeats and the *Het-A* and *TART* transposons, and a cluster present at 20A (*flamenco*, discussed in more detail below) which previously had been identified as a region involved in transposable element (TE) control[86,87]. It appears that there is a high level of complexity in piRNAs, as the three sequencing projects which examined Piwi-associated RNAs from ovaries show little overlap among the actual sequences of the small RNA populations[85], although these studies all identified clusters in many of the same genomic regions.

When the sequences of the rasiRNAs associated with the three different Piwi clade Argonaute proteins are compared, a strong strand-specific asymmetry is observed. Piwi and Aub are associated with antisense strands while Ago3 is found with sense strands, although this asymmetry appears to be reversed for some families of TEs. Upon examining the relationship between these two classes of rasiRNAs, one observes that Aub and Piwi associated rasiRNAs exhibit a strong bias for U at the 5' end, while AGO3 associated rasiRNAs have a preference for A at nucleotide 10. Based upon this sequence bias and comparisons between the cloned

sequences of the different classes, a “ping-pong” model for the generation of these small RNAs has been proposed (Figure 3) [24,82]. In this model, a primary piRNA is generated by transcription of the piRNA cluster. This primary piRNA transcript is then processed by a mechanism that uses rasiRNAs from the transposable elements oriented in the antisense direction which are bound by Piwi or Aub. These Piwi/Aub/RNA complexes pair with sense transcripts, possibly generated from active transposons or potentially from TEs oriented in the sense direction in piRNA clusters, and guide the slicing of the 5' end of the sense transcript. (The mechanism for generating the 3' end is currently unknown.) The sense piRNAs are bound by AGO3 and can then pair back with antisense transcripts from the piRNA clusters and other sources and guide the slicing of their 5' end. These antisense piRNAs are bound again by Piwi/Aub and can be used to continue the cycle. In addition to the strand bias, the cellular localization of these three Argonaute proteins in ovaries also differs. While Piwi is localized in the nucleus of both nurse cells and follicle cells, Aub and AGO3 are located in the cytoplasm, and show enrichment in the nuage, an electron-dense perinuclear structure present in the germline of many animals. In flies, many factors involved in small RNA processing are localized in the nuage (Table 1). All three proteins are loaded into the developing oocyte, as shown by the fact that all are present in 0–2h embryos; Aub shows an accumulation in the posterior pole, where the primordial germline forms, of stage 10 developing oocytes and persists there, being found exclusively in germ line cells in the embryo[24]. Interestingly, the greatest amount of complementarity between rasiRNAs is seen between those bound to AGO3 and Aub, both of which localize to the nuage.

While the ping-pong model explains much of the data, in particular how small RNAs are generated independent of Dicer proteins, many unanswered questions remain. It is unclear how the 3' ends of rasiRNAs are generated. The model requires that the sliced rasiRNA is cleaved by a yet-to-be-identified nuclease. Potential candidates for this nuclease include the products of *zucchini* (*zuc*) and *squash* (*sqw*), which have been identified to play a role in rasiRNA production and localize to the nuage[88]. However, whether or not these or other nucleases are used remains to be determined. A more complicated question is how the strand bias is achieved among the different Argonaute proteins. Most piRNA clusters are bidirectionally transcribed, generating a mix of sense and antisense transcripts[23,24]. A mechanism by which an active TE, which would presumably be transcribed in the sense direction, could be biased for regulation by one member of the Piwi clade can be envisioned by potential co-recruitment of an Argonaute family member with the transcription machinery. As mentioned previously, in fission yeast Ago1 interacts with RNAPII [44,45], and studies of the plant RNA Pol IV polymerase found interactions between RNAP IV and Argonaute 4 through the functionally conserved WG/GW motif[89,90]. A more complicated mechanism would need to be invoked for establishing the strand bias for transcripts generated from piRNA clusters, where the model proposes the generation of one long transcript encompassing the entire cluster[24]. In the case of the *flammenco/COM* locus at cytological band 20A, most of the TEs within the cluster are oriented in the antisense direction, but this bias does not occur in the majority of piRNA clusters, where partial and degenerate TEs are oriented in both the sense and antisense direction [23,24]. In this circumstance, the co-recruitment of an Argonaute family member with the polymerase, while quite possible, is insufficient to completely explain the strand bias. Finally, the source of the sense transcripts used to initiate the process has not been fully elucidated. Full-length active TEs, either located in euchromatin or buried in heterochromatin, or partial TEs, located in piRNA clusters and oriented in the sense direction, could potentially give rise to such RNAs.



## Role of rasiRNA machinery in silencing transposable elements in the germline

Investigations into rasiRNAs and the proteins involved in generating rasiRNAs have identified a role for these factors in regulating the expression and transposition of transposable elements in the male and female germlines. In terms of this discussion, it is helpful to clarify the definition of “germline”. In the strict sense, this term refers solely to the cell lineage producing the gametes. However, experimentalists in *Drosophila* have used the term to include all types of cells present in ovaries or testes. Thus in the case of the ovary, this definition encompasses the oocyte, the nurse cells, and the surrounding follicle cells; the latter are actually a somatic cell type, although they play a critical role in germline function. In-depth analysis of the piRNA cluster located at cytological band 20A has demonstrated that mutations disrupting the locus cause up-regulation and mobilization of the TEs sharing sequence identity with the locus. Previous studies isolated mutations in region 20A as *flamenco(flam)/COM*, a controlling region for mobilization of the LTR-containing retrotransposons *gypsy*, *Zam*, and *Idefix*. *gypsy* is a retrovirus that is expressed in follicle epithelium, where it is packaged into a virus-like particle and can invade the ovary, while *Zam* is expressed in posterior follicular cells, and *Idefix* is expressed in the germarium.

Regulation of mobilization of these transposons is defined by two different genetic states; one allows mobilization [referred to as Permissive (P), in the case of *gypsy* regulation, or Unstable (U) in the case of *Zam* or *Idefix* regulation], and one that does not allow mobilization [classified as Restrictive (R, for *gypsy*) or Stable (S, for *Zam/Idefix*)]. Mutations have been identified which cause a switch from an R strain to a P strain, leading to increased accumulation of *gypsy* RNA in the follicle cells, loss of control of *gypsy* and *Zam* mobilization (*Idefix* was not tested in this assay), and subsequent female sterility[87,91]. These mutations were mapped to region 20A; upon identification of this locus as a piRNA cluster, it was demonstrated that these *flam* mutants had reduced levels of piRNAs that map to the 20A cluster and that the generation of the primary piRNA transcript had been disrupted[24,91,92]. Additional studies demonstrated that mutations in *piwi*, the only Argonaute family member present in the follicle cells, also caused a global decrease in *gypsy* rasiRNAs and a corresponding increase in expression of *gypsy* transcripts in the germline[92,93]. A second body of work, focused on the regulation of *Zam* and *Idefix*, demonstrated similar effects of up regulation and mobilization of these TEs in response to mutations in *flam/COM*, and a dependency upon *piwi* for control of these elements in the germline[86,94]. Together, these studies have linked three observed effects: change in the level of rasiRNAs with homology to a TE, increased expression of the TE in the germline, and mobilization of the TE in the germline. Either mutations in *flam/COM*, which disrupt the primary piRNA transcripts, or mutations in *piwi*, which appears to be required for processing these transcripts, can cause these three effects. The results suggest that the piRNA system limits the presence of TE transcripts, which in turn will limit transposition. Clearly there is a component of this mechanism which utilizes post-transcriptional processing of the TE transcripts. It remains to be determined whether this part of the shut-down mechanism then directs the changes in chromatin structure which occur at these regions, similar to what is observed in *S. pombe*, or whether another separate mechanism directs the changes in chromatin structure, while the ping-pong mechanism is only concerned with removing TE transcripts in the germline by post-transcriptional gene silencing (i.e. RNA degradation). If this post-transcriptional silencing of the TEs is linked to the chromatin modifications of these regions, this would imply that the *cis* requirement observed for processing transcripts in *S. pombe* does not apply in *Drosophila*, given that some of the ping-pong pathway components involved in processing the rasiRNAs are not even present in the nucleus (i.e. Aub & AGO3). Additional evidence that the *cis* requirement may not apply in flies can be found in the fact that loss of *Snipper*, the putative *Drosophila* homologue of the *EriI* exonuclease (which imparts

the *cis* restriction in *S. pombe*), does not affect gene silencing mediated by RNAi in flies[95]. It is possible that the ping-pong pathway generates rasiRNAs that are used in the early embryo to re-establish heterochromatic domains and silence TEs in somatic cells (see below).

Additional studies have demonstrated a role for other components of the RNAi machinery in regulating the expression of a variety of different TEs and in the generation of rasiRNAs in the germline. Table 1 summarizes the effects observed in the ovaries on the expression of a subset of TEs in the presence of mutations in different factors linked to regulation of the rasiRNA pathway. These include many germline-specific factors that localize to the nuage, such as the products of *aub*, *krimper*, *maelstrom*, and *cutoff*, as well as the putative nucleases, *zuc* and *squ*, mentioned previously[72,88,96–98]. Also included are *spn-E* and *armi*, two helicase family members that have been implicated in a variety of roles in RNAi-based regulation[79, 81]. As there are no mutations available in the *AGO3* gene, a requirement for AGO3 in regulation of TE expression has not been tested. The effect of mutations in the different components varies to some extent depending upon which TE is examined. This variability may indicate a division of labor within the system and/or may reflect cell specific differences. However, no up-regulation of TEs in the female germline is observed with mutations in the siRNA pathway components *AGO2*, *Dcr-2*, or *R2D2*, or in the miRNA pathway component *loqs*, in agreement with their lack of effect on rasiRNA production[72].

In addition to increased TE expression, many lines with mutations in rasiRNA components are female sterile and show defects in dorsal-ventral patterning of the embryo. Investigation using *armi* and *aub* mutants has found that these embryonic patterning defects can be suppressed by mutations in *mei-41* and/or *mnk*, which encode the fly ATR and Chk2 kinases activated by DNA double-strand breaks (DSBs)[99]. In agreement with this result is the observation that  $\gamma$ H2A.v foci are present in the germline of the *armi* and *aub* mutants. ( $\gamma$ H2A.v is a phosphorylated form of the H2A histone variant that is associated with DNA DSBs.) While the *ATR/Chk2* mutants suppress the patterning defects, they do not alter the defects in TE silencing associated with mutations in these rasiRNA pathway members. A similar suppression of patterning defects is observed between *mnk* mutations and *cuff* and *squ* mutations; however, there is no suppression of patterning defects by *mnk* mutations in *zuc*, *spn-E* or *piwi* mutants [88,97,99]. It has been proposed that this suppression of patterning defects occurs due to the bypass of the DSB checkpoint that would normally be activated when TE expression is up-regulated and uncontrolled mobilization of TEs occurs, resulting in DSBs throughout the genome. The lack of suppression with *zuc*, *spn-E*, and *piwi* mutations indicates an additional role for these factors in a pathway outside of that mediated by Chk2.

## Relationship between transposable element silencing and heterochromatin formation in the germline

There is significant overlap in the *Drosophila* genome between the location of TEs and the packaging of DNA into heterochromatin, as defined by the presence of HP1 and H3K9me2, and as assessed by the recovery of variegating reporter inserts. While there is abundant evidence that rasiRNAs and the associated processing machinery regulate the expression of transposable elements in the germline, this process might be based solely on a post-transcriptional gene silencing mechanism that is unrelated and independent from the chromatin modifications in these regions. However, control of TE expression and mobilization could also occur via a transcriptional gene silencing mechanism, potentially similar to that used in *S. pombe*, where transcription of a repeat region helps to drive the formation of heterochromatin. Based on the *S. pombe* model and the data available from flies, one can propose a model in which the rasiRNAs function to direct the heterochromatin machinery (HP1, HMTs, and other factors) to these regions in a manner that is dependent upon transcription through the rasiRNA cluster to establish heterochromatin in the germline. As discussed above, data from *S. pombe* indicate

that this transcription of the heterochromatin regions occurs predominantly during S phase of the cell cycle[64]. There are interesting implications for the role of transcriptional gene silencing in the Drosophila germline, since while the oocyte is going through meiosis germline nurse cells undergo endoreduplication. In addition to the piRNA clusters, partial (presumably inactive) TEs are scattered throughout heterochromatic domains in Drosophila, including regions in pericentric heterochromatin and on the fourth chromosome, which have not been implicated as piRNA loci. It is unclear whether the piRNA system can impact heterochromatin formation at these partial TEs and if so, whether there are transcripts through these regions, or if this impact occurs through a separate mechanism. The possibilities are intriguing.

The strongest evidence for a transcriptional gene silencing mechanism in the germline comes from studies with mutations in *spn-E* in the ovaries of adult flies. Chromatin immunoprecipitation assays have demonstrated that mutations in *spn-E* affect chromatin structure at three different endogenous transposable elements (*HeT-A*, *I* element, and *copia*) in ovarian tissue isolated from homozygous females. A decrease in the repressive marks H3K9me2, H3K9me3 and HP1 and an increase in the active mark H3K4me2 are observed at both the promoter and ORF regions of these elements[100]. These observations are the first to actually link a mutation in a component of the rasiRNA pathway, with its associated decrease in rasiRNAs, to a change in chromatin structure and an increase in TE expression, all in one (complex) tissue type, the ovary.

In spite of this observation, other data indicate that this mechanism may be more complicated than a simple extrapolation from the case of *S. pombe* would suggest. The chromatin structure changes reported above are not seen in adult somatic cells of *spn-E* mutations (as assayed in the “carcass” tissue, the remainder of the fly after dissecting out the ovaries), suggesting that once set, heterochromatin can be maintained in the absence of RNAi in differentiated somatic tissues. Further, mutations in *Su(var)2–5* (HP1a) or *Su(var)3–9* (H3-K9 HMT), two key components of heterochromatin formation in flies, have no effect on the expression of the *copia* or *mdg1* TEs in testes[101]. While this data could indicate a post-transcriptional gene silencing mechanism, differences in the chromatin system in the germline cells may also be critical. Possible explanations for this result are the presence of a testes specific HP1 variant, HP1e[102], which may be functionally redundant to HP1a. There is also an ovary specific HP1, HP1d/Rhino, whose possible role in heterochromatin formation in the ovary is largely unexplored[102,103]. Furthermore, there are at least three active HMTs in Drosophila[104]; while *Su(var)3–9* was the first identified, it does not perform all of the critical functions of histone H3 K9 methylation as shown by the fact that flies with null mutations in *Su(var)3–9* are viable and fertile. This survival is distinct from mutations in *Su(var)2–5*, which are larval lethal[105], and mutations in rasiRNA pathway components, which are generally female sterile and/or maternal effect lethal (see Table 1 and references therein). While *Su(var)3–9* mutations impact heterochromatin formation, as assayed by position effect variegation, in the pericentric regions, they do not cause loss of silencing at the primarily heterochromatic 4<sup>th</sup> chromosome [106]. In contrast, the *dSETDB1/Egless* HMT, which plays an important role in oogenesis, also has a major role in regulating heterochromatin formation on the 4<sup>th</sup> chromosome, as well as in other regions of the genome[107–110]. As high levels of dSETBD are present in testes [109], it is possible that this HMT plays the primary role in H3K9 methylation and heterochromatin formation in the male germline, which would explain the lack of effect on TE expression in *Su(var)3–9* mutants. However, a functional role for either dSETDB or the third Drosophila HMT, G9a, in the male germline has not yet been explored.

Studies on the mobilization of *HeT-A* and *TART* elements, which are two non-LTR retrotransposon-like elements arrayed at Drosophila telomeres, may also help to elucidate the relationship between components of the rasiRNA pathway and components of the heterochromatin pathway. Flies heterozygous for mutations in the *spn-E* or *aub* genes have an

increased frequency of transposition of *HeT-A* and *TART* elements to broken chromosome ends [111]. Upon further analysis of the newly repaired chromosomes, it was discovered that the majority of the attachments were a *TART* element and that some of the repairs from the same cross have identical breakpoint junctions, indicating that the repair may have taken place at a premeiotic stage of oogenesis. This increase in transposition corresponds with the greater increase in *TART* expression relative to *HeT-A* in the ovaries of heterozygous *spn-E* mutants, although both TEs show strong up-regulation in the ovaries of homozygous *aub* or *spn-E* mutant flies [111]. The same assay had been used previously to demonstrate a role for HP1a in regulating the elongation of telomeres [112]. Mutations in *Su(var)2-5* result in an increased frequency of *HeT-A* or *TART* attachment to the broken chromosome end, again corresponding to an increased level of *HeT-A* transcript in RNA isolated from female *Su(var)2-5* mutant flies indicating a role for HP1a in regulating expression and transposition of *HeT-A* and *TART* in the female germline [112]. However, these studies were not able to distinguish whether the repair of the broken end occurs through an increase in *HeT-A* or *TART* transposition to that end, or by recombination with other telomeres.

## A role for the rasiRNA pathway in heterochromatin formation in somatic cells?

The links between heterochromatin and “RNAi” become less clear when one examines whether or not the rasiRNA pathway plays a role in heterochromatin formation outside of the germline. Examination of the literature covering the role of rasiRNA pathway members in somatic tissues leaves one with some apparently contradictory observations. Unfortunately, maternal loading of both proteins and RNA into the oocyte blurs the line between germline and somatic cells, making genetic dissection between these two cell types difficult in *Drosophila*. In this section, we attempt to address some of these apparent contradictions and examine the role of rasiRNA pathway members in the context of the development of the organism.

In examining how rasiRNA pathway components may be involved in heterochromatin formation, one must consider the developmental regulation of the process. Establishment of heterochromatin in the somatic tissues of the fly apparently begins in the early embryo during syncytial blastoderm. HP1 is enriched in the intensely DAPI-staining regions corresponding to heterochromatic regions of the genome, as first visualized at ca. nuclear replication cycle 10, in nuclei located at the apical surface of the embryo [113]. This enrichment increases significantly during nuclear cycle 14 when cellularization of the blastoderm occurs [113]. Heterochromatic silencing of a *lacZ* reporter gene is initiated at the onset of gastrulation, around stage 6, which occurs within nuclear cycle 14 [114]. A high degree of silencing is maintained through the mitotic cycles of larval development, until a relaxation occurs at the end of the 3<sup>rd</sup> instar larval stage (associated with the onset of differentiation of adult tissues), leading to a variegating phenotype [114]. The pattern of silencing observed after this relaxation during 3<sup>rd</sup> instar correlates with the pattern of silencing in the adult organism [115]. The presence of heterochromatin in the organism must be considered in the context of these stages, in that heterochromatin formation must be initiated correctly in the early embryo and maintained in order to observe a heterochromatic-responsive read-out in observations of the late larval or adult fly tissues. Mutations that disrupt the establishment, as well as mutations which fail to maintain heterochromatin or which cause misregulation of the relaxation stage, would be expected to affect the read-out observed. Given that robust zygotic transcription occurs starting in the interphase of nuclear cycle 14 [116] (albeit some zygotic transcripts are observed as early as nuclear cycle 7–8), it is likely that the components involved in the early embryonic establishment of heterochromatin are deposited into the egg by the mother.

Initial evidence for a role of the rasiRNA machinery in heterochromatin formation in somatic cells came from the observation that mutations in *piwi*, *aub*, and *spn-E* function as weak *Su*

(var)s when four different variegating insertions of the *white* gene are examined[117]. Consistent with the fact that the immediate sequence surrounding these reporters is different, due to different genomic insertion sites and the presence of transgene arrays at two of the reporters, the different rasiRNA components suppress variegation of the inserts to different degrees. For example, the reporter on the arm of chromosome 4 shows a higher impact with *piwi* mutations whereas a reporter in the pericentric region of the 4<sup>th</sup> chromosome is more strongly impacted by *aub* and *spn-E* mutations. This variability of suppression is also true for a fifth *white* reporter insertion, which contains a copy of the transposable element *1360* in addition to the reporter. In this case, mutations in *aub* show the highest level of suppression [118]. The differences in suppression of PEV by mutations in the rasiRNA pathway become even more complex when a *white* reporter inserted into a piRNA cluster at the 3R TAS is examined. In this case, mutations in *piwi* cause enhancement of variegation[85]. Given these varying effects one must keep in mind what is known about the regulation of the surrounding heterochromatic sequence. What TEs are present in the immediate vicinity and are there piRNA clusters at or adjacent to the insertion site? Are there other factors that might also play a role in regulating heterochromatin at this site? One such possibility would be binding sites for the DNA-binding protein D1, which is involved in heterochromatin formation at AT-rich satellite DNA[119]. As more information is elucidated about heterochromatin it is apparent that it is not homogeneous, and therefore one should hesitate to draw conclusions about heterochromatin as a whole from the observations made with a handful of reporters. Nonetheless, extensive investigation of these systems can certainly lead to important discoveries.

As noted above, examination of adult female carcass tissues showed no increase in *HeT-A*, *copia*, or *I* element expression in *spn-E*, *armi*, or *piwi* mutant lines and no change in the chromatin structure of these TEs in *spn-E* mutants[100]. These observations are consistent with a previous report of no effect on *HeT-A* transcript levels in *spn-E*, *piwi*, or *armi* mutant male larvae. In the same study, no effect on *HeT-A* expression with *Su(var)3-9* mutations was observed, but an increase in expression with *Su(var)2-5* mutations was seen[120]. The increase of *HeT-A* expression in *Su(var)2-5* mutant male larvae is consistent with the effect observed in *Su(var)2-5* mutant female flies[112] and demonstrates the importance of HP1 in repressing the expression of TEs in germline and somatic cells. The lack of any effect on *HeT-A* expression in somatic cells in a *Su(var)3-9* mutant may simply be due to the redundancy of HMTs as discussed above. The silencing of a *copia* LTR-*LacZ* reporter in ovarian tissue has been reported to be dependent upon *spn-E*, *armi*, and *piwi*, but silencing of this same transgene was not affected in somatic tissues of the larvae (larval brain, haltere imaginal disc, and salivary gland)[100]. A similar situation is observed with a GFP reporter whose expression is regulated by a fragment of the *Zam* or *Idefix* TE[94]. These reporters are silenced in the ovaries in a manner dependent upon *piwi* and the *flam/COM* locus. However, the silencing of these GFP transgenes outside of the germline, in larvae, pupal and adult stages, is independent of *piwi*. Interestingly, this *piwi* independent silencing outside the germline is still dependent upon *flam/COM*, consistent with rasiRNAs playing a role in the early embryo[94].

There are several issues to consider when analyzing these seemingly contradictory results. The regulation of a reporter inserted into a euchromatic region whose expression is controlled by some component of a TE, such as the *copia* LTR-*LacZ* reporter or the *Zam/Idefix* GFP reporter, may be different from that seen using the more traditional PEV assays. In the latter case, one is looking at a normally euchromatic gene inserted into a heterochromatic region, which may show different sensitivities to disruption of components in the rasiRNA system. Single copies of *P* element based reporters are not silenced when inserted into the euchromatic arms, even if the construct includes an additional TE fragment; either proximity to heterochromatic masses or a high local TE repeat density appears to be required[118]. In addition, the requirement for “establishment” of heterochromatin domains at the early embryo stage in the somatic tissue

may differ from regulation of TEs in the germline tissues. As most null mutations of the rasiRNA pathway members result in female sterility, parents heterozygous for the rasiRNA component are generally crossed to generate the homozygous individuals that are used in many of these assays. Therefore, a potential “half dose” of any hypothetical rasiRNA targeting complex will generally have been loaded into the embryo of the homozygous mutant individual. These complexes may be sufficient to initiate the silencing of reporters under the regulation of TE sequences, but the low dose may result in defects in heterochromatin stability, resulting in less silencing of “euchromatic gene” based reporters embedded into heterochromatin. It is also possible there are different requirements for assembly of heterochromatin in the somatic cells and germline tissues of an organism. Not all of the effects of mutations in the rasiRNA pathway on heterochromatin formation can be explained by loss of components loaded into the embryo from the female germline, as a mutation passed through the male germline to heterozygous offspring can exhibit defects in PEV assays as well [117]. This observation argues that either there is a role for the rasiRNA components after the “establishment” of heterochromatin in the early embryo (possibly at the relaxation phase), that the onset of expression of these factors is such that they are expressed in the developing zygote before or during the stage of heterochromatin formation that they are required for, or that there is some sort of “mark” transmitted through the mutant male that causes a disruption in forming heterochromatin.

Support for a rasiRNA-based targeting complex in the early embryo comes from the observation that Piwi, the only Argonaute protein present in the nucleus, can interact with HP1a [121]. Evidence for this interaction was first obtained with a yeast-two-hybrid (Y2H) screen of an ovary cDNA library using Piwi as bait. This interaction involves contact between a PxVxL motif at the N-terminus of Piwi and the chromo shadow domain of HP1a. Piwi and HP1a can be co-immunoprecipitated from embryo nuclear extract, and co-localize in the nuclei of early embryos at the time of heterochromatin formation [121]. The functional relevance of this interaction for heterochromatin formation was demonstrated by a PEV assay where a V30A mutation in *piwi* (shown to impact the interaction with HP1 in the Y2H assay) resulted in less silencing at a variegating *white* reporter previously shown to be sensitive to *piwi* mutations [121]. However, there have been no reports to date of the isolation of an HP1a-Piwi complex as such, or investigations of any HP1a-coprecipitating small RNAs.

### A role for siRNA pathway components in *Drosophila* chromatin structure?

Much of the data suggesting that small RNAs may be involved in heterochromatin formation in flies comes from the evidence that the rasiRNA pathway components contribute to silencing of TEs and of reporters that are located in heterochromatic regions of the fly genome. However, there are several observations that indicate cross-talk between the rasiRNA and the siRNA pathways which suggest that components of the siRNA pathway could also play a role in heterochromatin formation in flies. Factors involved in the siRNA pathway have been isolated from S2 cells in complexes capable of degrading a transcript *in vitro* via an RNAi based mechanism [122,123]. Complex components include siRNAs, the Argonaute family member AGO2, the *Drosophila* homolog of the Fragile X mental retardation protein (dFMR1), VIG (vasa intronic gene), the *Drosophila* homolog of p68 RNA helicase (also known as Dmp68/Rm62/Lip), TudorSN, and two ribosomal proteins, L5 and L11, along with 5S RNA [123, 124]. This pathway has been shown to play a role in viral defense, as mutations in components of this pathway, as well as some components of the rasiRNA pathway (*piwi*, *aub*, *armi*), cause an increased sensitivity relative to wild-type flies upon viral exposure (as measured by percent survival several days post infection) [125–128]. Interestingly, components of the rasiRNA pathway, including *armi*, *aub*, and *spn-E*, have also been implicated in a role in classic post-transcriptional “RNAi” silencing mediated primarily via components of the siRNA pathway [78,79,81,129], suggesting some “cross-talk” between these pathways (Figure 1). Furthermore,

the recent identification of endogenous siRNAs present in both somatic and germline tissue which interact with Ago2, require Dcr2 for processing, and have homology to transposable elements supports potential “cross-talk” between these pathways in heterochromatin formation and maintenance [15–18].

Three components of the siRISC complex, Dmp68, AGO2, and dFMR1 have been linked to the formation of heterochromatin. The *Drosophila* p68 homologue (*Lip*) was first identified genetically as a suppressor of position effect variegation and of several *white* mutations caused by retrotransposon insertions[130]. *Lip* mutations also showed changes in the level of *cop* transcript present in the fly and alterations in the level of transcript at the *white* allele used in the initial screen[130]. These changes in transcript levels can potentially be explained by a role for Dmp68 in transcriptional deactivation[131]. The Dmp68 protein is an RNA helicase required for transcript clearance at sites of active transcription; mutations in *Dmp68* result in RNA export defects. These functions seem to be important for deactivation of highly transcribed genes, such as the *hsp70* gene following heat shock[131]. AGO2 and dFMR1 also play a role in chromosome function during early embryogenesis[132,133]. Mutations in either *AGO2* or *dFMR1* cause asynchronous nuclear division cycles in the early embryo. These embryos show a mislocalization of HP1 relative to wild-type embryos and have defects in assembling centromeric heterochromatin, demonstrated by the fact that mutations in either factor show a suppression of PEV for *white* reporters inserted into pericentric heterochromatin. In addition, *AGO2* mutants exhibit mislocalization of the centromere-specific H3 variant CID. Furthermore, both factors are required for normal development of the pole cells in the early embryo. However, none of these defects are fully penetrant, as *AGO2* homozygous mutants produce viable, fertile adults, and *dFMR1* null mutants are viable, although viability is reduced and there are associated neuronal defects. Possible explanations for this lack of penetrance are that the mutants used for these studies are not actually complete nulls or that there are multiple redundant pathways involved[132,133]. A model involving components of both the siRNA pathway and the rasiRNA pathway in the establishment of heterochromatin in the early embryo may be needed (Figure 3), although at present there is no direct evidence of complexes between siRNA components and heterochromatin components. Given the evidence that components of the rasiRNA pathway are also involved in viral defense in the fly and that components of the rasiRNA pathway are involved in siRNA silencing via a double-stranded RNA hairpin knock-down, it seems likely that cross-talk between these two pathways can occur when the organism is presented with different types of RNA, either viral or that expressed from TEs, that may challenge its survival (Figure 1). The detailed mechanisms behind this cross-talk remain to be worked out.

Further evidence for a link between the siRNA pathway and heterochromatin stability comes from studies showing that mutations in *dcr-2*, the primary Dicer of the siRNA pathway, show disorganized and multiple nucleoli, and extrachromosomal circles of ribosomal DNA (rDNA) and satellite DNA in salivary gland and diploid nuclei[134]. Interestingly, a second siRNA component, *ago2*, and the rasiRNA pathway components *piwi*, *aub*, and *spn-E* also show an increased number of nucleoli per nucleus. Similar defects are observed in lines with mutations in the heterochromatin components *Su(var)3-9* or *Su(var)2-5*. These changes in nuclear organization correspond to a decrease in H3K9me at these loci in a *dcr-2* mutant and appear to result from the generation of extra-chromosomal circular DNA. While all of the loci examined reside in heterochromatin, it is not entirely clear whether the formation of heterochromatin at the rDNA or at satellite sequences is similar to or somewhat different from that at transposable elements. However, the results do indicate a role for small RNA pathways in forming and maintaining heterochromatin at the rDNA and satellite sequences. A similar requirement for the RNAi system in maintaining nucleolar integrity has also been found in *Arabidopsis*[reviewed in 135]. Thus, the RNAi system is required to maintain the integrity of the genome in somatic cells in domains with a high density of repetitive sequences, most

likely through the establishment and maintenance of heterochromatin to minimize internal recombination events.

## Observations supporting a role for small RNAs in heterochromatin formation in vertebrates

Do small RNAs play a role in directing heterochromatin formation in mammals? Several groups have identified piRNA clusters in the male germline of mice and rats that are important for spermatogenesis [23,136–140]. These clusters appear to be developmentally regulated, as some are expressed in the pachytene stage and some in pre-pachytene stages. The clusters expressed in pre-pachytene are more transposon-rich and have been implicated in transposon control in the germline, presumably using a mechanism similar to the rasiRNA pathway in flies [23]. Additional support for a role of the RNAi machinery in transposon control in mammals comes from disruption of the genes for the mouse *piwi* homologues *Mili* or *Miwi2*, which results in derepression and loss of methylation of transposons in the male germline [23,25]. Analysis of mouse oocytes found small RNA of the siRNA class (~20–24nt) derived from retroelements [138], implicating this second class of small RNAs in transposon control (at least in the oocyte) as well. Initial studies on mouse embryonic stem cells with a deletion of the *Dicer* gene reported an increased accumulation of centromeric transcripts from the major and minor satellite repeats; however no changes in DNA methylation or H3K9 methylation were observed at these loci [141]. More recent studies found changes in DNA methylation in *Dicer* mutants, but these effects appear to be mediated through the microRNA pathway via regulation of DNA methyltransferases by *Rbl2* and the *mir-290* cluster, indicating an indirect role for these small RNAs in epigenetic regulation [142,143]. Thus, there is accumulating evidence of a role for small RNAs in controlling repetitive sequences in the germline, but as yet little or no evidence for a direct role in somatic cells.

## A model for heterochromatin formation in flies

Given the overall body of evidence it seems highly likely that small RNAs are playing a role in directing heterochromatin formation in *Drosophila*, but clearly there are many details that remain to be elucidated, and additional evidence is required to confirm this role. Two pathways could be operating in parallel in flies, one in the germline and a second in the early embryo, the latter required to set up heterochromatin structure in somatic nuclei and maintain it throughout development of the organism. However, it seems likely that components of a germline system are present, and could help direct the embryonic/somatic pathway, potentially by depositing small RNAs and associated proteins into the embryo. It also appears likely that a transcription-based mechanism could be used, similar to that in *S. pombe*, but potentially without the *cis* restriction (Figure 3). Exactly what “RNAi” machinery is used may depend upon the type of transposable element/repeat to be silenced and what tissue/cell type (germline or somatic) is examined, as these factors may influence which components are recruited. In the germline, it appears that the rasiRNA pathway components play the primary, if not only, role in regulating TE expression and mobilization, likely through a post-transcriptional mechanism but also by controlling the formation/maintenance of heterochromatin, whereas in the early embryo both the rasiRNA pathway and siRNA pathway components may play a role in regulating TEs and directing heterochromatin formation. As preserving germline integrity is crucial for the success of future generations, TE expression/mobilization in this tissue may be under more stringent control than in somatic tissue. However, regulation of TEs in somatic tissues is presumably required to prevent mutations caused by mobilization as well as destabilization of heterochromatin that could decrease survival and hence the ability of the organism to propagate. As formation of heterochromatin is critical for basic cellular processes in the organism, it is quite possible that there are additional pathways outside of a small RNA based mechanism, such as the DNA-protein interactions observed at the MAT loci in *S.*



*pombe*, which direct heterochromatin formation. It is likely that some of the primary players, such as HP1a and H3K9me, will function as the “end-point” for multiple pathways in setting up heterochromatin in the organism. Once heterochromatin is formed, it appears to be sufficient to maintain TEs in a silent state, as mutations in the rasiRNA pathway no longer have a major impact. The requirement for the rasiRNA pathway in the germline may indicate that heterochromatin formation is less efficacious there; the dynamic state of the germline could be designed to inoculate the embryo against viral challenges later in life. The siRNA/rasiRNA system is complex, as required to fill its various roles: defense against invading viruses, protection against endogenous transposable elements, maintenance of genome stability in regions with high levels of repetitious sequences, accomplished in both germ line and somatic cells. Sorting out the various pathways involved, including cross-talk between components will take many more years of effort by numerous investigators.

## Acknowledgements

We thank Brent Brower-Toland, Nicole Riddle, and Elena Gracheva for critical comments on this manuscript. K. Huisinga is supported by a Kirschstein NRSA Postdoctoral Fellowship from NIH; work from the Elgin lab cited above is supported by NIH grants R01 GM68388 and GM073190 to SCRE.

## References

1. Camblong J, Iglesias N, Fickentscher C, Dieppois G, Stutz F. Antisense RNA stabilization induces transcriptional gene silencing via histone deacetylation in *S. cerevisiae*. *Cell* 2007;131:706–717. [PubMed: 18022365]
2. Martens JA, Laprade L, Winston F. Intergenic transcription is required to repress the *Saccharomyces cerevisiae* SER3 gene. *Nature* 2004;429:571–574. [PubMed: 15175754]
3. Bernstein E, Caudy AA, Hammond SM, Hannon GJ. Role for a bidentate ribonuclease in the initiation step of RNA interference. *Nature* 2001;409:363–366. [PubMed: 11201747]
4. Lee YS, Nakahara K, Pham JW, Kim K, He Z, Sontheimer EJ, Carthew RW. Distinct roles for *Drosophila* Dicer-1 and Dicer-2 in the siRNA/miRNA silencing pathways. *Cell* 2004;117:69–81. [PubMed: 15066283]
5. Pham JW, Pellino JL, Lee YS, Carthew RW, Sontheimer EJ. A Dicer-2-dependent 80s complex cleaves targeted mRNAs during RNAi in *Drosophila*. *Cell* 2004;117:83–94. [PubMed: 15066284]
6. Denli AM, Tops BB, Plasterk RH, Ketting RF, Hannon GJ. Processing of primary microRNAs by the Microprocessor complex. *Nature* 2004;432:231–235. [PubMed: 15531879]
7. Forstemann K, Tomari Y, Du T, Vagin VV, Denli AM, Bratu DP, Klattenhoff C, Theurkauf WE, Zamore PD. Normal microRNA maturation and germ-line stem cell maintenance requires Loquacious, a double-stranded RNA-binding domain protein. *PLoS Biol* 2005;3:e236. [PubMed: 15918770]
8. Jiang F, Ye X, Liu X, Fincher L, McKearin D, Liu Q. Dicer-1 and R3D1-L catalyze microRNA maturation in *Drosophila*. *Genes Dev* 2005;19:1674–1679. [PubMed: 15985611]
9. Landthaler M, Yalcin A, Tuschl T. The human DiGeorge syndrome critical region gene 8 and Its D. melanogaster homolog are required for miRNA biogenesis. *Curr Biol* 2004;14:2162–2167. [PubMed: 15589161]
10. Liu Q, Rand TA, Kalidas S, Du F, Kim HE, Smith DP, Wang X. R2D2, a bridge between the initiation and effector steps of the *Drosophila* RNAi pathway. *Science* 2003;301:1921–1925. [PubMed: 14512631]
11. Saito K, Ishizuka A, Siomi H, Siomi MC. Processing of pre-microRNAs by the Dicer-1-Loquacious complex in *Drosophila* cells. *PLoS Biol* 2005;3:e235. [PubMed: 15918769]
12. Girard A, Hannon GJ. Conserved themes in small-RNA-mediated transposon control. *Trends Cell Biol* 2008;18:136–148. [PubMed: 18282709]
13. Matranga C, Zamore PD. Small silencing RNAs. *Curr Biol* 2007;17:R789–793. [PubMed: 17878043]
14. Ding SW, Voinnet O. Antiviral immunity directed by small RNAs. *Cell* 2007;130:413–426. [PubMed: 17693253]

15. Chung WJ, Okamura K, Martin R, Lai EC. Endogenous RNA interference provides a somatic defense against *Drosophila* transposons. *Curr Biol* 2008;18:795–802. [PubMed: 18501606]
16. Czech B, Malone CD, Zhou R, Stark A, Schlingehayde C, Dus M, Perrimon N, Kellis M, Wohlschlegel JA, Sachidanandam R, Hannon GJ, Brennecke J. An endogenous small interfering RNA pathway in *Drosophila*. *Nature* 2008;453:798–802. [PubMed: 18463631]
17. Ghildiyal M, Seitz H, Horwich MD, Li C, Du T, Lee S, Xu J, Kittler EL, Zapp ML, Weng Z, Zamore PD. Endogenous siRNAs derived from transposons and mRNAs in *Drosophila* somatic cells. *Science* 2008;320:1077–1081. [PubMed: 18403677]
18. Kawamura Y, Saito K, Kin T, Ono Y, Asai K, Sunohara T, Okada TN, Siomi MC, Siomi H. *Drosophila* endogenous small RNAs bind to Argonaute 2 in somatic cells. *Nature* 2008;453:793–797. [PubMed: 18463636]
19. Okamura K, Balla S, Martin R, Liu N, Lai EC. Two distinct mechanisms generate endogenous siRNAs from bidirectional transcription in *Drosophila melanogaster*. *Nat Struct Mol Biol* 2008;15:581–590. [PubMed: 18500351]
20. Okamura K, Chung WJ, Ruby JG, Guo H, Bartel DP, Lai EC. The *Drosophila* hairpin RNA pathway generates endogenous short interfering RNAs. *Nature* 2008;453:803–806. [PubMed: 18463630]
21. Obbard DJ, Finnegan DJ. RNA Interference: Endogenous siRNAs Derived from Transposable Elements. *Curr Biol* 2008;18:R561–563. [PubMed: 18606126]
22. Stefani G, Slack FJ. Small non-coding RNAs in animal development. *Nat Rev Mol Cell Biol* 2008;9:219–230. [PubMed: 18270516]
23. Aravin AA, Hannon GJ, Brennecke J. The Piwi-piRNA pathway provides an adaptive defense in the transposon arms race. *Science* 2007;318:761–764. [PubMed: 17975059]
24. Brennecke J, Aravin AA, Stark A, Dus M, Kellis M, Sachidanandam R, Hannon GJ. Discrete small RNA-generating loci as master regulators of transposon activity in *Drosophila*. *Cell* 2007;128:1089–1103. [PubMed: 17346786]
25. Carmell MA, Girard A, van de Kant HJ, Bourc'his D, Bestor TH, de Rooij DG, Hannon GJ. MIWI2 is essential for spermatogenesis and repression of transposons in the mouse male germline. *Dev Cell* 2007;12:503–514. [PubMed: 17395546]
26. Deng W, Lin H. miwi, a murine homolog of piwi, encodes a cytoplasmic protein essential for spermatogenesis. *Dev Cell* 2002;2:819–830. [PubMed: 12062093]
27. Harris AN, Macdonald PM. Aubergine encodes a *Drosophila* polar granule component required for pole cell formation and related to eIF2C. *Development* 2001;128:2823–2832. [PubMed: 11526087]
28. Kuramochi-Miyagawa S, Kimura T, Ijiri TW, Isobe T, Asada N, Fujita Y, Ikawa M, Iwai N, Okabe M, Deng W, Lin H, Matsuda Y, Nakano T. Mili, a mammalian member of piwi family gene, is essential for spermatogenesis. *Development* 2004;131:839–849. [PubMed: 14736746]
29. Cox DN, Chao A, Baker J, Chang L, Qiao D, Lin H. A novel class of evolutionarily conserved genes defined by piwi are essential for stem cell self-renewal. *Genes Dev* 1998;12:3715–3727. [PubMed: 9851978]
30. Buhler M, Moazed D. Transcription and RNAi in heterochromatic gene silencing. *Nat Struct Mol Biol* 2007;14:1041–1048. [PubMed: 17984966]
31. Hutvagner G, Simard MJ. Argonaute proteins: key players in RNA silencing. *Nat Rev Mol Cell Biol* 2008;9:22–32. [PubMed: 18073770]
32. Matzke M, Kanno T, Huettel B, Daxinger L, Matzke AJ. Targets of RNA-directed DNA methylation. *Curr Opin Plant Biol* 2007;10:512–519. [PubMed: 17702644]
33. Weil C, Martienssen R. Epigenetic interactions between transposons and genes: lessons from plants. *Curr Opin Genet Dev*. 2008
34. Cam HP, Sugiyama T, Chen ES, Chen X, FitzGerald PC, Grewal SI. Comprehensive analysis of heterochromatin- and RNAi-mediated epigenetic control of the fission yeast genome. *Nat Genet* 2005;37:809–819. [PubMed: 15976807]
35. Grewal SI. Transcriptional silencing in fission yeast. *J Cell Physiol* 2000;184:311–318. [PubMed: 10911361]
36. Volpe TA, Kidner C, Hall IM, Teng G, Grewal SI, Martienssen RA. Regulation of heterochromatic silencing and histone H3 lysine-9 methylation by RNAi. *Science* 2002;297:1833–1837. [PubMed: 12193640]

37. Hall IM, Shankaranarayana GD, Noma K, Ayoub N, Cohen A, Grewal SI. Establishment and maintenance of a heterochromatin domain. *Science* 2002;297:2232–2237. [PubMed: 12215653]
38. Jia S, Noma K, Grewal SI. RNAi-independent heterochromatin nucleation by the stress-activated ATF/CREB family proteins. *Science* 2004;304:1971–1976. [PubMed: 15218150]
39. Reinhart BJ, Bartel DP. Small RNAs correspond to centromere heterochromatic repeats. *Science* 2002;297:1831. [PubMed: 12193644]
40. Sugiyama T, Cam H, Verdel A, Moazed D, Grewal SI. RNA-dependent RNA polymerase is an essential component of a self-enforcing loop coupling heterochromatin assembly to siRNA production. *Proc Natl Acad Sci U S A* 2005;102:152–157. [PubMed: 15615848]
41. Noma K, Sugiyama T, Cam H, Verdel A, Zofall M, Jia S, Moazed D, Grewal SI. RITS acts in cis to promote RNA interference-mediated transcriptional and post-transcriptional silencing. *Nat Genet* 2004;36:1174–1180. [PubMed: 15475954]
42. Djupedal I, Portoso M, Spahr H, Bonilla C, Gustafsson CM, Allshire RC, Ekwall K. RNA Pol II subunit Rpb7 promotes centromeric transcription and RNAi-directed chromatin silencing. *Genes Dev* 2005;19:2301–2306. [PubMed: 16204182]
43. Kato H, Goto DB, Martienssen RA, Urano T, Furukawa K, Murakami Y. RNA polymerase II is required for RNAi-dependent heterochromatin assembly. *Science* 2005;309:467–469. [PubMed: 15947136]
44. Schramke V, Sheedy DM, Denli AM, Bonila C, Ekwall K, Hannon GJ, Allshire RC. Retraction: RNA-interference-directed chromatin modification coupled to RNA polymerase II transcription. *Nature* 2005;437:1057. [PubMed: 16222304]
45. Schramke V, Sheedy DM, Denli AM, Bonila C, Ekwall K, Hannon GJ, Allshire RC. RNA-interference-directed chromatin modification coupled to RNA polymerase II transcription. *Nature* 2005;435:1275–1279. [PubMed: 15965464]
46. Irvine DV, Zaratiegui M, Tolia NH, Goto DB, Chitwood DH, Vaughn MW, Joshua-Tor L, Martienssen RA. Argonaute slicing is required for heterochromatic silencing and spreading. *Science* 2006;313:1134–1137. [PubMed: 16931764]
47. Sadaie M, Iida T, Urano T, Nakayama J. A chromodomain protein, Chp1, is required for the establishment of heterochromatin in fission yeast. *EMBO J* 2004;23:3825–3835. [PubMed: 15372076]
48. DeBeauchamp JL, Moses A, Noffsinger VJ, Ulrich DL, Job G, Kosinski AM, Partridge JF. Chp1-Tas3 interaction is required to recruit RITS to fission yeast centromeres and for maintenance of centromeric heterochromatin. *Mol Cell Biol* 2008;28:2154–2166. [PubMed: 18212052]
49. Partridge JF, DeBeauchamp JL, Kosinski AM, Ulrich DL, Hadler MJ, Noffsinger VJ. Functional separation of the requirements for establishment and maintenance of centromeric heterochromatin. *Mol Cell* 2007;26:593–602. [PubMed: 17531816]
50. Verdel A, Jia S, Gerber S, Sugiyama T, Gygi S, Grewal SI, Moazed D. RNAi-mediated targeting of heterochromatin by the RITS complex. *Science* 2004;303:672–676. [PubMed: 14704433]
51. Motamedi MR, Verdel A, Colmenares SU, Gerber SA, Gygi SP, Moazed D. Two RNAi complexes, RITS and RDRC, physically interact and localize to noncoding centromeric RNAs. *Cell* 2004;119:789–802. [PubMed: 15607976]
52. Colmenares SU, Buker SM, Buhler M, Dlakic M, Moazed D. Coupling of double-stranded RNA synthesis and siRNA generation in fission yeast RNAi. *Mol Cell* 2007;27:449–461. [PubMed: 17658285]
53. Buker SM, Iida T, Buhler M, Villen J, Gygi SP, Nakayama J, Moazed D. Two different Argonaute complexes are required for siRNA generation and heterochromatin assembly in fission yeast. *Nat Struct Mol Biol* 2007;14:200–207. [PubMed: 17310250]
54. Buhler M, Verdel A, Moazed D. Tethering RITS to a nascent transcript initiates RNAi- and heterochromatin-dependent gene silencing. *Cell* 2006;125:873–886. [PubMed: 16751098]
55. Zhang K, Mosch K, Fischle W, Grewal SI. Roles of the Clr4 methyltransferase complex in nucleation, spreading and maintenance of heterochromatin. *Nat Struct Mol Biol*. 2008
56. Hong EJ, Villen J, Gerace EL, Gygi SP, Moazed D. A cullin E3 ubiquitin ligase complex associates with Rik1 and the Clr4 histone H3-K9 methyltransferase and is required for RNAi-mediated heterochromatin formation. *RNA Biol* 2005;2:106–111. [PubMed: 17114925]

57. Horn PJ, Bastie JN, Peterson CL. A Rik1-associated, cullin-dependent E3 ubiquitin ligase is essential for heterochromatin formation. *Genes Dev* 2005;19:1705–1714. [PubMed: 16024659]
58. Jia S, Kobayashi R, Grewal SI. Ubiquitin ligase component Cul4 associates with Clr4 histone methyltransferase to assemble heterochromatin. *Nat Cell Biol* 2005;7:1007–1013. [PubMed: 16127433]
59. Li F, Goto DB, Zaratiegui M, Tang X, Martienssen R, Cande WZ. Two novel proteins, dos1 and dos2, interact with rik1 to regulate heterochromatic RNA interference and histone modification. *Curr Biol* 2005;15:1448–1457. [PubMed: 16040243]
60. Thon G, Hansen KR, Altes SP, Sidhu D, Singh G, Verhein-Hansen J, Bonaduce MJ, Klar AJ. The Clr7 and Clr8 directionality factors and the Pcu4 cullin mediate heterochromatin formation in the fission yeast *Schizosaccharomyces pombe*. *Genetics* 2005;171:1583–1595. [PubMed: 16157682]
61. Grewal SI, Elgin SC. Transcription and RNA interference in the formation of heterochromatin. *Nature* 2007;447:399–406. [PubMed: 17522672]
62. Huisinga KL, Brower-Toland B, Elgin SC. The contradictory definitions of heterochromatin: transcription and silencing. *Chromosoma* 2006;115:110–122. [PubMed: 16506022]
63. Chen ES, Zhang K, Nicolas E, Cam HP, Zofall M, Grewal SI. Cell cycle control of centromeric repeat transcription and heterochromatin assembly. *Nature* 2008;451:734–737. [PubMed: 18216783]
64. Kloc A, Zaratiegui M, Nora E, Martienssen R. RNA Interference Guides Histone Modification during the S Phase of Chromosomal Replication. *Curr Biol*. 2008
65. Morris SA, Shibata Y, Noma K, Tsukamoto Y, Warren E, Temple B, Grewal SI, Strahl BD. Histone H3 K36 methylation is associated with transcription elongation in *Schizosaccharomyces pombe*. *Eukaryot Cell* 2005;4:1446–1454. [PubMed: 16087749]
66. Li B, Carey M, Workman JL. The role of chromatin during transcription. *Cell* 2007;128:707–719. [PubMed: 17320508]
67. Nicolas E, Yamada T, Cam HP, Fitzgerald PC, Kobayashi R, Grewal SI. Distinct roles of HDAC complexes in promoter silencing, antisense suppression and DNA damage protection. *Nat Struct Mol Biol* 2007;14:372–380. [PubMed: 17450151]
68. Folco HD, Pidoux AL, Urano T, Allshire RC. Heterochromatin and RNAi are required to establish CENP-A chromatin at centromeres. *Science* 2008;319:94–97. [PubMed: 18174443]
69. Hall IM, Noma K, Grewal SI. RNA interference machinery regulates chromosome dynamics during mitosis and meiosis in fission yeast. *Proc Natl Acad Sci U S A* 2003;100:193–198. [PubMed: 12509501]
70. Volpe T, Schramke V, Hamilton GL, White SA, Teng G, Martienssen RA, Allshire RC. RNA interference is required for normal centromere function in fission yeast. *Chromosome Res* 2003;11:137–146. [PubMed: 12733640]
71. Aravin AA, Lagos-Quintana M, Yalcin A, Zavolan M, Marks D, Snyder B, Gaasterland T, Meyer J, Tuschl T. The small RNA profile during *Drosophila melanogaster* development. *Dev Cell* 2003;5:337–350. [PubMed: 12919683]
72. Vagin VV, Sigova A, Li C, Seitz H, Gvozdev V, Zamore PD. A distinct small RNA pathway silences selfish genetic elements in the germline. *Science* 2006;313:320–324. [PubMed: 16809489]
73. Horwich MD, Li C, Matranga C, Vagin V, Farley G, Wang P, Zamore PD. The *Drosophila* RNA methyltransferase, DmHen1, modifies germline piRNAs and single-stranded siRNAs in RISC. *Curr Biol* 2007;17:1265–1272. [PubMed: 17604629]
74. Saito K, Sakaguchi Y, Suzuki T, Siomi H, Siomi MC. Pimet, the *Drosophila* homolog of HEN1, mediates 2'-O-methylation of Piwi-interacting RNAs at their 3' ends. *Genes Dev* 2007;21:1603–1608. [PubMed: 17606638]
75. Aravin AA, Klenov MS, Vagin VV, Bantignies F, Cavalli G, Gvozdev VA. Dissection of a natural RNA silencing process in the *Drosophila melanogaster* germ line. *Mol Cell Biol* 2004;24:6742–6750. [PubMed: 15254241]
76. Aravin AA, Naumova NM, Tulin AV, Vagin VV, Rozovsky YM, Gvozdev VA. Double-stranded RNA-mediated silencing of genomic tandem repeats and transposable elements in the *D. melanogaster* germline. *Curr Biol* 2001;11:1017–1027. [PubMed: 11470406]

77. Gillespie DE, Berg CA. Homeless is required for RNA localization in *Drosophila* oogenesis and encodes a new member of the DE-H family of RNA-dependent ATPases. *Genes Dev* 1995;9:2495–2508. [PubMed: 7590230]
78. Cook HA, Koppetsch BS, Wu J, Theurkauf WE. The *Drosophila* SDE3 homolog armitage is required for oskar mRNA silencing and embryonic axis specification. *Cell* 2004;116:817–829. [PubMed: 15035984]
79. Tomari Y, Du T, Haley B, Schwarz DS, Bennett R, Cook HA, Koppetsch BS, Theurkauf WE, Zamore PD. RISC assembly defects in the *Drosophila* RNAi mutant armitage. *Cell* 2004;116:831–841. [PubMed: 15035985]
80. Stapleton W, Das S, McKee BD. A role of the *Drosophila* homeless gene in repression of Stellate in male meiosis. *Chromosoma* 2001;110:228–240. [PubMed: 11513298]
81. Kennerdell JR, Yamaguchi S, Carthew RW. RNAi is activated during *Drosophila* oocyte maturation in a manner dependent on aubergine and spindle-E. *Genes Dev* 2002;16:1884–1889. [PubMed: 12154120]
82. Gunawardane LS, Saito K, Nishida KM, Miyoshi K, Kawamura Y, Nagami T, Siomi H, Siomi MC. A slicer-mediated mechanism for repeat-associated siRNA 5' end formation in *Drosophila*. *Science* 2007;315:1587–1590. [PubMed: 17322028]
83. Nishida KM, Saito K, Mori T, Kawamura Y, Nagami-Okada T, Inagaki S, Siomi H, Siomi MC. Gene silencing mechanisms mediated by Aubergine piRNA complexes in *Drosophila* male gonad. *RNA* 2007;13:1911–1922. [PubMed: 17872506]
84. Saito K, Nishida KM, Mori T, Kawamura Y, Miyoshi K, Nagami T, Siomi H, Siomi MC. Specific association of Piwi with rasiRNAs derived from retrotransposon and heterochromatic regions in the *Drosophila* genome. *Genes Dev* 2006;20:2214–2222. [PubMed: 16882972]
85. Yin H, Lin H. An epigenetic activation role of Piwi and a Piwi-associated piRNA in *Drosophila melanogaster*. *Nature* 2007;450:304–308. [PubMed: 17952056]
86. Desset S, Meignin C, Dastugue B, Vaury C. COM, a heterochromatic locus governing the control of independent endogenous retroviruses from *Drosophila melanogaster*. *Genetics* 2003;164:501–509. [PubMed: 12807771]
87. Pelisson A, Song SU, Prud'homme N, Smith PA, Bucheton A, Corces VG. Gypsy transposition correlates with the production of a retroviral envelope-like protein under the tissue-specific control of the *Drosophila* flamenco gene. *EMBO J* 1994;13:4401–4411. [PubMed: 7925283]
88. Pane A, Wehr K, Schupbach T. zucchini and squash encode two putative nucleases required for rasiRNA production in the *Drosophila* germline. *Dev Cell* 2007;12:851–862. [PubMed: 17543859]
89. El-Shami M, Pontier D, Lahmy S, Braun L, Picart C, Vega D, Hakimi MA, Jacobsen SE, Cooke R, Lagrange T. Reiterated WG/GW motifs form functionally and evolutionarily conserved ARGONAUTE-binding platforms in RNAi-related components. *Genes Dev* 2007;21:2539–2544. [PubMed: 17938239]
90. Li CF, Pontes O, El-Shami M, Henderson IR, Bernatavichute YV, Chan SW, Lagrange T, Pikaard CS, Jacobsen SE. An ARGONAUTE4-containing nuclear processing center colocalized with Cajal bodies in *Arabidopsis thaliana*. *Cell* 2006;126:93–106. [PubMed: 16839879]
91. Mevel-Ninio M, Pelisson A, Kinder J, Campos AR, Bucheton A. The flamenco locus controls the gypsy and ZAM retroviruses and is required for *Drosophila* oogenesis. *Genetics* 2007;175:1615–1624. [PubMed: 17277359]
92. Pelisson A, Sarot E, Payen-Groschene G, Bucheton A. A novel repeat-associated small interfering RNA-mediated silencing pathway downregulates complementary sense gypsy transcripts in somatic cells of the *Drosophila* ovary. *J Virol* 2007;81:1951–1960. [PubMed: 17135323]
93. Sarot E, Payen-Groschene G, Bucheton A, Pelisson A. Evidence for a piwi-dependent RNA silencing of the gypsy endogenous retrovirus by the *Drosophila melanogaster* flamenco gene. *Genetics* 2004;166:1313–1321. [PubMed: 15082550]
94. Desset S, Buchon N, Meignin C, Coiffet M, Vaury C. In *Drosophila melanogaster* the COM Locus Directs the Somatic Silencing of Two Retrotransposons through both Piwi-Dependent and -Independent Pathways. *PLoS ONE* 2008;3:e1526. [PubMed: 18253480]

95. Kupsco JM, Wu MJ, Marzluff WF, Thapar R, Duronio RJ. Genetic and biochemical characterization of *Drosophila* Snipper: A promiscuous member of the metazoan 3'hExo/ERI-1 family of 3' to 5' exonucleases. *RNA* 2006;12:2103–2117. [PubMed: 17135487]
96. Lim AK, Kai T. Unique germ-line organelle, nuage, functions to repress selfish genetic elements in *Drosophila melanogaster*. *Proc Natl Acad Sci U S A* 2007;104:6714–6719. [PubMed: 17428915]
97. Chen Y, Pane A, Schupbach T. Cutoff and aubergine mutations result in retrotransposon upregulation and checkpoint activation in *Drosophila*. *Curr Biol* 2007;17:637–642. [PubMed: 17363252]
98. Findley SD, Tamanaha M, Clegg NJ, Ruohola-Baker H. Maelstrom, a *Drosophila* spindle-class gene, encodes a protein that colocalizes with Vasa and RDE1/AGO1 homolog, Aubergine, in nuage. *Development* 2003;130:859–871. [PubMed: 12538514]
99. Klattenhoff C, Bratu DP, McGinnis-Schultz N, Koppetsch BS, Cook HA, Theurkauf WE. *Drosophila* rasiRNA pathway mutations disrupt embryonic axis specification through activation of an ATR/Chk2 DNA damage response. *Dev Cell* 2007;12:45–55. [PubMed: 17199040]
100. Klenov MS, Lavrov SA, Stolyarenko AD, Ryazansky SS, Aravin AA, Tuschl T, Gvozdev VA. Repeat-associated siRNAs cause chromatin silencing of retrotransposons in the *Drosophila melanogaster* germline. *Nucleic Acids Res* 2007;35:5430–5438. [PubMed: 17702759]
101. Kalmykova AI, Klenov MS, Gvozdev VA. Argonaute protein PIWI controls mobilization of retrotransposons in the *Drosophila* male germline. *Nucleic Acids Res* 2005;33:2052–2059. [PubMed: 15817569]
102. Vermaak D, Henikoff S, Malik HS. Positive selection drives the evolution of rhino, a member of the heterochromatin protein 1 family in *Drosophila*. *PLoS Genet* 2005;1:96–108. [PubMed: 16103923]
103. Volpe AM, Horowitz H, Grafer CM, Jackson SM, Berg CA. *Drosophila* rhino encodes a female-specific chromo-domain protein that affects chromosome structure and egg polarity. *Genetics* 2001;159:1117–1134. [PubMed: 11729157]
104. Mis J, Ner SS, Grigliatti TA. Identification of three histone methyltransferases in *Drosophila*: dG9a is a suppressor of PEV and is required for gene silencing. *Mol Genet Genomics* 2006;275:513–526. [PubMed: 16622709]
105. Lu BY, Emtage PC, Duyf BJ, Hilliker AJ, Eissenberg JC. Heterochromatin protein 1 is required for the normal expression of two heterochromatin genes in *Drosophila*. *Genetics* 2000;155:699–708. [PubMed: 10835392]
106. Haynes KA, Gracheva E, Elgin SC. A Distinct type of heterochromatin within *Drosophila melanogaster* chromosome 4. *Genetics* 2007;175:1539–1542. [PubMed: 17194780]
107. Clough E, Moon W, Wang S, Smith K, Hazelrigg T. Histone methylation is required for oogenesis in *Drosophila*. *Development* 2007;134:157–165. [PubMed: 17164421]
108. Seum C, Reo E, Peng H, Rauscher FJ 3rd, Spierer P, Bontron S. *Drosophila* SETDB1 is required for chromosome 4 silencing. *PLoS Genet* 2007;3:e76. [PubMed: 17500594]
109. Stabell M, Bjorkmo M, Aalen RB, Lambertsson A. The *Drosophila* SET domain encoding gene dEset is essential for proper development. *Hereditas* 2006;143:177–188. [PubMed: 17362353]
110. Tzeng TY, Lee CH, Chan LW, Shen CK. Epigenetic regulation of the *Drosophila* chromosome 4 by the histone H3K9 methyltransferase dSETDB1. *Proc Natl Acad Sci U S A* 2007;104:12691–12696. [PubMed: 17652514]
111. Savitsky M, Kwon D, Georgiev P, Kalmykova A, Gvozdev V. Telomere elongation is under the control of the RNAi-based mechanism in the *Drosophila* germline. *Genes Dev* 2006;20:345–354. [PubMed: 16452506]
112. Savitsky M, Kravchuk O, Melnikova L, Georgiev P. Heterochromatin protein 1 is involved in control of telomere elongation in *Drosophila melanogaster*. *Mol Cell Biol* 2002;22:3204–3218. [PubMed: 11940677]
113. Kellum R, Raff JW, Alberts BM. Heterochromatin protein 1 distribution during development and during the cell cycle in *Drosophila* embryos. *J Cell Sci* 1995;108(Pt 4):1407–1418. [PubMed: 7615662]
114. Lu BY, Ma J, Eissenberg JC. Developmental regulation of heterochromatin-mediated gene silencing in *Drosophila*. *Development* 1998;125:2223–2234. [PubMed: 9584122]


115. Lu BY, Bishop CP, Eissenberg JC. Developmental timing and tissue specificity of heterochromatin-mediated silencing. *EMBO J* 1996;15:1323–1332. [PubMed: 8635465]
116. Foe, VE.; Odell, GM.; Edgar, BA. *The Development of Drosophila melanogaster*. Cold Spring Harbor Laboratory Press; 1993. Mitosis and morphogenesis in the Drosophila embryo: point and counterpoint.
117. Pal-Bhadra M, Leibovitch BA, Gandhi SG, Rao M, Bhadra U, Birchler JA, Elgin SC. Heterochromatic silencing and HP1 localization in Drosophila are dependent on the RNAi machinery. *Science* 2004;303:669–672. [PubMed: 14752161]
118. Haynes KA, Caudy AA, Collins L, Elgin SC. Element 1360 and RNAi components contribute to HP1-dependent silencing of a pericentric reporter. *Curr Biol* 2006;16:2222–2227. [PubMed: 17113386]
119. Aulner N, Monod C, Mandicourt G, Jullien D, Cuvier O, Sall A, Janssen S, Laemmler UK, Kas E. The AT-hook protein D1 is essential for Drosophila melanogaster development and is implicated in position-effect variegation. *Mol Cell Biol* 2002;22:1218–1232. [PubMed: 11809812]
120. Perrini B, Piacentini L, Fanti L, Altieri F, Chichiarelli S, Berloco M, Turano C, Ferraro A, Pimpinelli S. HP1 controls telomere capping, telomere elongation, and telomere silencing by two different mechanisms in Drosophila. *Mol Cell* 2004;15:467–476. [PubMed: 15304225]
121. Brower-Toland B, Findley SD, Jiang L, Liu L, Yin H, Dus M, Zhou P, Elgin SC, Lin H. Drosophila PIWI associates with chromatin and interacts directly with HP1a. *Genes Dev* 2007;21:2300–2311. [PubMed: 17875665]
122. Caudy AA, Myers M, Hannon GJ, Hammond SM. Fragile X-related protein and VIG associate with the RNA interference machinery. *Genes Dev* 2002;16:2491–2496. [PubMed: 12368260]
123. Ishizuka A, Siomi MC, Siomi H. A Drosophila fragile X protein interacts with components of RNAi and ribosomal proteins. *Genes Dev* 2002;16:2497–2508. [PubMed: 12368261]
124. Caudy AA, Ketting RF, Hammond SM, Denli AM, Bathoorn AM, Tops BB, Silva JM, Myers MM, Hannon GJ, Plasterk RH. A micrococcal nuclease homologue in RNAi effector complexes. *Nature* 2003;425:411–414. [PubMed: 14508492]
125. Galiana-Arnoux D, Dostert C, Schneemann A, Hoffmann JA, Imler JL. Essential function in vivo for Dicer-2 in host defense against RNA viruses in drosophila. *Nat Immunol* 2006;7:590–597. [PubMed: 16554838]
126. van Rij RP, Saleh MC, Berry B, Foo C, Houk A, Antoniewski C, Andino R. The RNA silencing endonuclease Argonaute 2 mediates specific antiviral immunity in Drosophila melanogaster. *Genes Dev* 2006;20:2985–2995. [PubMed: 17079687]
127. Wang XH, Aliyari R, Li WX, Li HW, Kim K, Carthew R, Atkinson P, Ding SW. RNA interference directs innate immunity against viruses in adult Drosophila. *Science* 2006;312:452–454. [PubMed: 16556799]
128. Zamboni RA, Vakharia VN, Wu LP. RNAi is an antiviral immune response against a dsRNA virus in Drosophila melanogaster. *Cell Microbiol* 2006;8:880–889. [PubMed: 16611236]
129. Specchia V, Benna C, Mazzotta GM, Piccin A, Zordan MA, Costa R, Bozzetti MP. aubergine Gene Overexpression in Somatic Tissues of auberginestng Mutants Interferes With the RNAi Pathway of a yellow Hairpin dsRNA in Drosophila melanogaster. *Genetics* 2008;178:1271–1282. [PubMed: 18385112]
130. Csink AK, Linsk R, Birchler JA. The Lighten up (Lip) gene of Drosophila melanogaster, a modifier of retroelement expression, position effect variegation and white locus insertion alleles. *Genetics* 1994;138:153–163. [PubMed: 8001783]
131. Buszczak M, Spradling AC. The Drosophila P68 RNA helicase regulates transcriptional deactivation by promoting RNA release from chromatin. *Genes Dev* 2006;20:977–989. [PubMed: 16598038]
132. Deshpande G, Calhoun G, Schedl P. Drosophila argonaute-2 is required early in embryogenesis for the assembly of centric/centromeric heterochromatin, nuclear division, nuclear migration, and germ-cell formation. *Genes Dev* 2005;19:1680–1685. [PubMed: 16024657]
133. Deshpande G, Calhoun G, Schedl P. The drosophila fragile X protein dFMR1 is required during early embryogenesis for pole cell formation and rapid nuclear division cycles. *Genetics* 2006;174:1287–1298. [PubMed: 16888325]

134. Peng JC, Karpen GH. H3K9 methylation and RNA interference regulate nucleolar organization and repeated DNA stability. *Nat Cell Biol* 2007;9:25–35. [PubMed: 17159999]
135. Preuss S, Pikaard CS. rRNA gene silencing and nucleolar dominance: insights into a chromosome-scale epigenetic on/off switch. *Biochim Biophys Acta* 2007;1769:383–392. [PubMed: 17439825]
136. Grivna ST, Beyret E, Wang Z, Lin H. A novel class of small RNAs in mouse spermatogenic cells. *Genes Dev* 2006;20:1709–1714. [PubMed: 16766680]
137. Lau NC, Seto AG, Kim J, Kuramochi-Miyagawa S, Nakano T, Bartel DP, Kingston RE. Characterization of the piRNA complex from rat testes. *Science* 2006;313:363–367. [PubMed: 16778019]
138. Watanabe T, Takeda A, Tsukiyama T, Mise K, Okuno T, Sasaki H, Minami N, Imai H. Identification and characterization of two novel classes of small RNAs in the mouse germline: retrotransposon-derived siRNAs in oocytes and germline small RNAs in testes. *Genes Dev* 2006;20:1732–1743. [PubMed: 16766679]
139. Aravin A, Gaidatzis D, Pfeffer S, Lagos-Quintana M, Landgraf P, Iovino N, Morris P, Brownstein MJ, Kuramochi-Miyagawa S, Nakano T, Chien M, Russo JJ, Ju J, Sheridan R, Sander C, Zavolan M, Tuschl T. A novel class of small RNAs bind to MILI protein in mouse testes. *Nature* 2006;442:203–207. [PubMed: 16751777]
140. Girard A, Sachidanandam R, Hannon GJ, Carmell MA. A germline-specific class of small RNAs binds mammalian Piwi proteins. *Nature* 2006;442:199–202. [PubMed: 16751776]
141. Murchison EP, Partridge JF, Tam OH, Cheloufi S, Hannon GJ. Characterization of Dicer-deficient murine embryonic stem cells. *Proc Natl Acad Sci U S A* 2005;102:12135–12140. [PubMed: 16099834]
142. Benetti R, Gonzalo S, Jaco I, Munoz P, Gonzalez S, Schoeftner S, Murchison E, Andl T, Chen T, Klatt P, Li E, Serrano M, Millar S, Hannon G, Blasco MA. A mammalian microRNA cluster controls DNA methylation and telomere recombination via Rbl2-dependent regulation of DNA methyltransferases. *Nat Struct Mol Biol* 2008;15:268–279. [PubMed: 18311151]
143. Sinkkonen L, Hugenschmidt T, Berninger P, Gaidatzis D, Mohn F, Artus-Revel CG, Zavolan M, Svoboda P, Filipowicz W. MicroRNAs control de novo DNA methylation through regulation of transcriptional repressors in mouse embryonic stem cells. *Nat Struct Mol Biol* 2008;15:259–267. [PubMed: 18311153]




**Post Transcriptional Gene Silencing**

**Transcriptional Gene Silencing?**




**Figure 1.**

Overview of small RNA pathways in *Drosophila*. Members of the RNase III family of nucleases are in light red, members of the DRB family are in light blue, and members of the Argonaute family of proteins are in green/yellow/orange. The cross-talk between Dcr-1 and Dcr-2[4] and potentially between the siRNA and rasiRNA pathways, discussed in the text, is indicated.



**Figure 2.** Schematic model for cell-cycle dependent regulation of heterochromatin formation at *S. pombe* pericentric regions. The self-reinforcing loop of heterochromatin formation is illustrated during S-phase, when transcription of the *dg-dh* region occurs. Adapted from figures in [55, 63,64]. The diagram at M phase has been simplified to emphasize the transition in Swi6 binding.



**Figure 3.** Model for small RNA mediated regulation of TE silencing in the *Drosophila* germline and potential links to heterochromatin formation in germline and early embryo. The proposed ping-pong mechanism of TE regulation in the germline is illustrated on the left. Links between rasiRNA machinery (much of which is present in the nuage of the female germline) and heterochromatin formation are diagrammed. Other RNAi components implicated in heterochromatin formation or maintenance are listed in Table 1.

Table 1

**Regulation of TE expression in Ovaries**

Expression of various transposable elements as determined by Northern, RT-PCR, or *in situ* hybridization in the ovaries of flies carrying mutations in various components of small RNA processing systems relative to that observed in the ovaries of wild-type flies. An asterisk indicates analysis in larval gonads. Discrepancies between different publications are highlighted; N/A = not analyzed.

Factor	Description	Cellular localization	Tissue type	Mutant viability/sterility	Non-LTR retrotransposon			LTR retrotransposon				Repeat locus		
					I element	HeT-A	TART	F-element	Mdgl	Roo	Gypsy		Copia	Gate
<i>piwi</i>	Argonaute	Nucleus	Somatic and germline	Female sterile, male sterile	Yes, small <sup>1,6</sup>	No <sup>1,6</sup>		Yes, large <sup>1</sup>	Yes, small <sup>1</sup>	Yes <sup>1,6*</sup>	Yes <sup>6</sup>			Yes, large <sup>1</sup>
<i>aub</i>	Argonaute	Nuage-cyto	Germline	Female sterile	Yes, large <sup>10,1</sup>	Yes <sup>1,8,10,12</sup>	Yes <sup>8</sup> , yes <sup>12</sup> , no (2, heterozygotes)	Yes-modest <sup>14</sup>	No <sup>1</sup> , yes-modest <sup>14</sup>	No <sup>1,6*</sup>		Yes <sup>14</sup>	Yes <sup>8</sup> , yes-modest <sup>14</sup>	
<i>AGO3</i>	Argonaute	Nuage-cyto	Germline											
<i>spn-E</i>	RNA helicase	N/A	Germline (female)	Female sterile	Yes, medium <sup>6,10,1</sup>	Yes <sup>6,10,1,12</sup>	Yes <sup>12</sup>	Yes <sup>14</sup>	Yes, small <sup>1</sup> ; Yes <sup>14</sup>	Yes, medium <sup>1</sup>	Yes <sup>6,10</sup>	Yes <sup>9</sup>	Yes <sup>14</sup>	Yes, small <sup>1</sup> ; yes <sup>14</sup>
<i>arni</i>	RNA helicase	Nuage-cyto	Germline	Female sterile	Yes, medium <sup>1,6</sup>	Yes <sup>1,8,6</sup>	Yes <sup>8</sup>	Yes, small <sup>1</sup>	Yes, small <sup>1</sup> ; no <sup>8</sup>	V. slight <sup>1</sup>	No <sup>6</sup>		Yes, medium <sup>1</sup>	
<i>AGO2</i>	Argonaute	Nuage-cyto	Somatic	Viable	No <sup>1</sup>	No <sup>1</sup>		No <sup>1</sup>	V. slight <sup>1</sup>	No <sup>1</sup>			No <sup>1</sup>	
<i>Dcr-2</i>	Dicer	Cyto	Somatic	Viable	No <sup>1</sup>	No <sup>1</sup>		No <sup>1</sup>	No <sup>1</sup>	No <sup>1</sup>			No <sup>1</sup>	
<i>R2D2</i>	DRB	Cyto	Somatic	Viable	No <sup>1</sup>	No <sup>1</sup>		No <sup>1</sup>	No <sup>1</sup>	No <sup>1</sup>			No <sup>1</sup>	
<i>Dcr-1 (mutant clone)</i>	Dicer	Cyto	Somatic	Lethal	No <sup>1</sup>	No <sup>1</sup>		No <sup>1</sup>	No <sup>1</sup>	No <sup>1</sup>			No <sup>1</sup>	
<i>Loqs</i>	DRB	Cyto	Somatic	Viable	No <sup>1</sup>	No <sup>1</sup>	Yes <sup>2</sup>	No <sup>1</sup>	No <sup>1</sup>	No <sup>1</sup>			No <sup>1</sup>	
<i>zuc</i>	Nuclease	Nuage-cyto	Germline	Female sterile	Yes <sup>2</sup>	Yes <sup>3</sup>								
<i>squ</i>	Nuclease	Nuage-cyto	Germline	Female sterile	Yes <sup>3</sup>									
<i>ciff</i>	Homologue of yeast Rai1	Partial nuage and cyto	Germline	Female sterile										
<i>mael</i>	Partial potential HMG-box	Shuttles- <i>nuage</i> /cyto and nucleus	Somatic and germline	Female sterile	Yes <sup>8</sup>	Yes <sup>8</sup>	Yes <sup>8</sup>		No <sup>8</sup>				Yes <sup>8</sup>	
<i>krimp</i>	Predicted Tudor domain	Nuage	Germline	Female sterile	Yes <sup>8</sup>	Yes <sup>8</sup>	No <sup>8</sup>		No <sup>8</sup>				Yes <sup>8</sup>	

<sup>1</sup> V. V. Vagin, A. Sigova, C. Li, H. Setz, V. Gvozdev, P.D. Zamore, A distinct small RNA pathway silences selfish genetic elements in the germline, *Science* 313 (2006) 320–324.

<sup>2</sup> A. Pane, K. Wehr, T. Schupbach, zucchini and squash encode two putative nucleases required for rasiRNA production in the *Drosophila* germline, *Dev Cell* 12 (2007) 851–862.

<sup>3</sup> Y. Chen, A. Pane, T. Schupbach, Cutoff and aubergine mutations result in retrotransposon upregulation and checkpoint activation in *Drosophila*, *Curr Biol* 17 (2007) 637–642.

<sup>4</sup> S.D. Findley, M. Tamanaha, N.J. Clegg, H. Ruohola-Baker, Maelstrom, a *Drosophila* spindle-class gene, encodes a protein that colocalizes with Vasa and RDE1/AGO1 homologs, in nuage, *Development* 130 (2003) 859–871.

- 5 C. Klattenhoff, D.P. Bratu, N. McGinnis-Schultz, B.S. Koppetsch, H.A. Cook, W.E. Theurkauf, Drosophila rasiRNA pathway mutations disrupt embryonic axis specification through activation of an ATR/Chk2 DNA damage response, *Dev Cell* 12 (2007) 45–55.
- 6 M.S. Klenov, S.A. Lavrov, A.D. Stolyarenko, S.S. Ryazansky, A.A. Aravin, T. Tuschl, V.A. Gvozdev, Repeat-associated siRNAs cause chromatin silencing of retrotransposons in the *Drosophila melanogaster* germline, *Nucleic Acids Res* 35 (2007) 5430–5438.
- 7 A.I. Kalmykova, M.S. Klenov, V.A. Gvozdev, Argonaute protein Piwi controls mobilization of retrotransposons in the *Drosophila* male germline, *Nucleic Acids Res* 33 (2005) 2052–2059.
- 8 A.K. Lim, T. Kai, Unique germ-line organelle, nuage, functions to repress selfish genetic elements in *Drosophila melanogaster*, *Proc Natl Acad Sci U S A* 104 (2007) 6714–6719.
- 9 G.L. Kogan, A.V. Tulin, A.A. Aravin, Y.A. Abramov, A.I. Kalmykova, C. Maisonhaute, V.A. Gvozdev, The GATE retrotransposon in *Drosophila melanogaster*: mobility in heterochromatin and aspects of its expression in germline tissues, *Mol Genet Genomics* 269 (2003) 234–242.
- 10 V.V. Vagin, M.S. Klenov, A.I. Kalmykova, A.D. Stolyarenko, R.N. Kotelnikov, V.A. Gvozdev, The RNA interference proteins and vasa locus are involved in the silencing of retrotransposons in the female germline of *Drosophila melanogaster*, *RNA Biol* 1 (2004) 54–58.
- 11 E. Sarot, G. Payen-Groschene, A. Bucheton, A. Pelisson, Evidence for a piwi-dependent RNA silencing of the gypsy endogenous retrovirus by the *Drosophila melanogaster* flamenco gene, *Genetics* 166 (2004) 1313–1321.
- 12 M. Savitsky, D. Kwon, P. Georgiev, A. Kalmykova, V. Gvozdev, Telomere elongation is under the control of the RNAi-based mechanism in the *Drosophila* germline, *Genes Dev* 20 (2006) 345–354.
- 13 A.A. Aravin, M.S. Klenov, V.V. Vagin, F. Bantignies, G. Cavalli, V.A. Gvozdev, Dissection of a natural RNA silencing process in the *Drosophila melanogaster* germ line, *Mol Cell Biol* 24 (2004) 6742–6750.
- 14 A.A. Aravin, N.M. Naumova, A.V. Tulin, V.V. Vagin, Y.M. Rozovsky, V.A. Gvozdev, Double-stranded RNA-mediated silencing of genomic tandem repeats and transposable elements in the *D. melanogaster* germline, *Curr Biol* 11 (2001) 1017–1027.
- 15 W. Stapleton, S. Das, B.D. McKee, A role of the *Drosophila* homeless gene in repression of Stellate in male meiosis, *Chromosoma* 110 (2001) 228–240.
- 16 A. Pelisson, E. Sarot, G. Payen-Groschene, A. Bucheton, A novel repeat-associated small interfering RNA-mediated silencing pathway downregulates complementary sense gypsy transcripts in somatic cells of the *Drosophila* ovary, *J Virol* 81 (2007) 1951–1960.