

RESEARCH ARTICLE

EARTH SCIENCES

Global blue carbon accumulation in tidal wetlands increases with climate change

Faming Wang^{1,2,3,4†*}, Christian J. Sanders^{4,5†*}, Isaac R. Santos^{5,6}, Jianwu Tang⁴, Mark Schurech⁷, Matthew L. Kirwan⁸, Robert E. Kopp⁹, Kai Zhu¹⁰, Xiuzhen Li⁴, Jiacan Yuan^{9,11}, Wenzhi Liu^{2,12}, Zhian Li^{1,2,3}

¹ Xiaoliang Research Station for Tropical Coastal Ecosystems, Key Laboratory of Vegetation Restoration and Management of Degraded Ecosystems, and the CAS engineering Laboratory for Ecological Restoration of Island and Coastal Ecosystems, South China Botanical Garden, Chinese Academy of Sciences, Guangzhou 510650, China

² Center of Plant Ecology, Core Botanical Gardens, Chinese Academy of Sciences, Guangzhou 510650, China

³ Southern Marine Science and Engineering Guangdong Laboratory (Guangzhou), Guangzhou 511458, China

⁴ State Key Laboratory of Estuarine and Coastal Research and Institute of Eco-Chongming, East China Normal University, Shanghai 201100, China

⁵ National Marine Science Centre, School of Environment, Science and Engineering, Southern Cross University, Coffs Harbour NSW 2450, Australia

⁶ Department of Marine Sciences, University of Gothenburg, Gothenburg 40530, Sweden

⁷ Lincoln Centre for Water and Planetary Health, School of Geography, University of Lincoln, Lincoln LN67TS, UK

⁸ Virginia Institute of Marine Science, College of William and Mary, Gloucester Point, VA 23185, USA

⁹ Department of Earth and Planetary Sciences and Rutgers Institute of Earth, Ocean, and Atmospheric Sciences, Rutgers University, New Brunswick, NJ 08854, USA,

¹⁰ Department of Environmental Studies, University of California, Santa Cruz, CA 95064, USA

¹¹ Department of Atmospheric and Oceanic Sciences, Fudan University, Shanghai 200433, China

¹² CAS Key Laboratory of Aquatic Botany and Watershed Ecology, Wuhan Botanical Garden, Chinese Academy of Sciences, Wuhan 430074, China

*Correspondence to: Dr. Christian J. Sanders (Christian.Sanders@scu.edu.au), and Dr. Faming Wang (wangfm@scbg.ac.cn).

†Equally contributed to this work.

Abstract:

Coastal tidal wetlands produce and accumulate significant amounts of organic carbon (C) that help to mitigate climate change. However, previous data limitations have prevented a robust evaluation of the global rates and mechanisms driving C accumulation. Here, we go beyond recent soil C stock estimates to reveal global tidal wetland C accumulation and predict changes under relative sea-level rise, temperature and precipitation. We use data from literature study sites and our new observations spanning wide latitudinal gradients and 20 countries. Globally, tidal wetlands accumulate 53.65 (95%CI: 48.52-59.01) Tg C yr⁻¹, which is ~30% of the organic C buried on the ocean floor. Modelling based on current climatic drivers and under projected emissions scenarios revealed a net increase in the global C accumulation by 2100. This rapid increase is driven by sea-level rise in tidal marshes, and higher temperature and precipitation in mangroves. Countries with large areas of coastal wetlands, like Indonesia and Mexico, are more susceptible to tidal wetland C losses under climate change, while regions such as Australia, Brazil, the USA and China will experience a significant C accumulation increase under all projected scenarios.

Keywords: Coastal wetlands, Blue C, C burial rate, Global change

Introduction

Mangroves and tidal marshes are highly productive wetlands that photosynthetically sequester atmospheric CO₂ as organic carbon (C) [1]. A varying fraction of this C is buried in tidally inundated suboxic and anoxic sediments and thereby largely prevented from returning to the atmosphere [2]. Sediments within these wetlands do not become saturated with C, because

sea level rise expands soil volume and creates accommodation space, accelerating burial of organic matter, and ultimately enhance the long term preservation of sedimentary C [3-5]. These coastal wetlands C has been regarded as Blue Carbon (BC) since a decade ago to describe their disproportionately large contribution to global C sequestration [6, 7]. The preservation and restoration of BC ecosystems such as mangroves and saltmarshes have been suggested as effective approaches to mitigate climate change [1]. Although the role of BC in climate change mitigation and adaptation has reached international prominence [7-9], several questions remained unanswered in BC studies [7]: What is the global extent and spatial distribution of coastal wetlands BC? What factors influence their BC burial rates? How does climate change impact C accumulation in these BC ecosystems?

Feedbacks between vegetation and soil accretion help mangroves and tidal marshes to remain within the intertidal zone in the face of sea-level rise [3, 4] through the accumulation of mineral and organic sediments [10] space (i.e. the inland space available for sediments to accumulate and be colonized by wetland vegetation) [3, 11]. However, human activities are imposing many pressures on tidal wetlands, including climate change-induced accelerated sea-level rise, subsidence through groundwater as well as the extraction of other substances, and the reduction of sediment supply from the construction of systems that reduce sediment flow such as dams. Furthermore, a lack of accommodation space, which is the space made available for organic and inorganic material to deposit, in relation to an increase in relative sea level rise rate (RSLR) may limit the sediment accumulation rates in specific tidal wetlands [11, 12]. As a result, there is growing concern that a lack of space for inland migration brought about through land use change or steep landward slopes may prevent mangrove and tidal marsh wetland adaptation to fast rates of sea-level rise [13, 14].

Blue Carbon accumulation in tidal wetlands is potentially driven by climatic factors (e.g., temperature, rainfall, and evapotranspiration) [15-17], coastal oceanographic processes (e.g., tidal amplitude, currents and geomorphology) [13, 14] and nutrient availability [18]. Identifying the controlling factors of C accumulation dynamics is critical for understanding the fate of the C buried by these wetlands under future climate change scenarios on global and regional scales. If climate change alters the fundamental drivers of organic C accumulation, the amount of sequestered C may also be modified and provide a different feedback to global warming.

Here, we estimate global tidal wetlands Blue C accumulation rates in different regions and countries. We rely on literature (n=564) data supplemented by new (n=49) soil core data from previously unaccounted areas in Indonesia, tropical South America and Africa (Fig.1). Soil cores were dated with the radiometric geochronologies (^{137}Cs , $^{239+240}\text{Pu}$ and ^{210}Pb) or Sediment Elevation Table (SET) methods covering time scales of sub-decade to decades. Both methods account for surface and subsurface processes, including organic and mineral sedimentation, sediment compaction and organic matter decomposition [19, 20], and are widely used for sediment and carbon accretion rate measurements [21]. We first evaluate the spatial patterns and drivers of global soil accretion and C accumulation building on recent work focusing on carbon stocks [22, 23]. We then predict C accumulation rates (CAR) in global tidal wetlands under differing emission scenarios based on CMIP5 projections. We further combine projected CAR with a future global tidal wetland area change dataset [11] to develop the first projections of how global tidal wetland carbon accumulation will change in response to anthropogenic CO_2 emissions. This approach focuses on a centennial time scale of anthropogenic climate change and prevents bias from other methods operating at different time scales [24]. Previous studies have

shown that C accumulation increases with RSLR at specific locations [12, 15], but there are no projections of how RSLR will drive global tidal wetland C accumulation.

Results and Discussion

Global coastal C accumulation rates

The arithmetic average C accumulation rates (CAR) were estimated to be $194 \pm 15 \text{ g C m}^{-2} \text{ yr}^{-1}$ for mangroves and $168 \pm 7 \text{ g C m}^{-2} \text{ yr}^{-1}$ for tidal marshes, based on all compiled coastal wetlands sites worldwide. The tidal marsh CAR in this study was much lower than the previous reports by Ouyang and Lee [25], who estimated the global coastal marsh CAR at $245 \text{ g C m}^{-2} \text{ yr}^{-1}$ based on 143 global sites data. The mangrove CAR in our study was similar to previous estimate of $163 \text{ g C m}^{-2} \text{ yr}^{-1}$ by Breithaupt, Smoak [2] based on 66 mangrove sites, but much lower than an earlier estimate ($210 \text{ g C m}^{-2} \text{ yr}^{-1}$) by Chmura, Anisfeld [26], likely due to the increased dataset in the more recent studies (Fig. 1).

Tidal wetland C accumulation rates in the Northern and Southern hemispheres were not significantly different ($p < 0.05$). However, pooling the data into broad latitudinal bands revealed that tropical mangroves between 0° and 10° had the highest C accumulation per unit area ($428 \pm 7 \text{ g C m}^{-2} \text{ yr}^{-1}$ for Northern hemisphere and $384 \pm 11 \text{ g C m}^{-2} \text{ yr}^{-1}$ for Southern hemisphere) and temperate tidal marshes between 30° and 40° had the lowest ($144 \pm 6 \text{ g C m}^{-2} \text{ yr}^{-1}$ for Northern hemisphere and $88.7 \pm 3.5 \text{ g C m}^{-2} \text{ yr}^{-1}$ for Southern hemisphere) (Fig. 1). Mangroves located in the 0° to 10° latitudinal band, where $\sim 50\%$ of mangrove living biomass occurs [27], were underrepresented in previous global datasets [2, 25, 28]. Our new observations in Brazil, Indonesia, and India help to fill this gap. For example, Indonesian mangroves had the highest soil accretion (SAR, $36 \pm 22 \text{ mm yr}^{-1}$) and C accumulation rates ($1722 \pm 183 \text{ g C m}^{-2} \text{ year}^{-1}$) [29].

Drivers of global coastal C accumulation

A variety of environmental factors can drive wetland soil carbon sequestration [15-17]. We collected climatic and environmental factors for each site, including mean annual temperature (MAT), annual precipitation (MAP), tidal range, elevation, RSLR, total suspended matters (TSM), and tropical cyclone frequency, to attempt in discovery the principal environmental factors that drive soil C accumulation in tidal marshes and mangrove forests (See details in M&M section). Our study contains a robust dataset to assess the main drivers of CAR on global and local scale. We developed linear mixed-effect models (see M&M section and Table S1) that provides results for the detected main environmental factors on CAR (Table S2) as standardized coefficients (Fig. S1). These coefficients indicate the proportional change in CAR in response to one SD change in an environmental factor. The model fit to observations is good (Fig. S2) and the residuals were normally distributed without obvious bias (Fig. S3 and S4).

The linear mixed model using RSLR and MAT as covariate effects totally explained 51% of the variability in CAR in tidal marshes (Table S1), which is consistent with model and regional field evidence linking CAR with these environmental factors [12, 15]. In mangroves, MAT and mean annual precipitation (MAP) explained 57 % of the variation in CAR (Table S1, and the detailed model section were listed in the M&M section). The positive relationship between mangrove CAR and temperature (Fig. S5) supports recent studies demonstrating how warming increases plant production, soil carbon stocks, and soil surface elevation [5, 16]. Precipitation is another important driver of soil C accumulation in warmer climate mangrove soils, but not in tidal marshes found in cooler regions (Fig. S5). This finding was also consistent with recent reports on how precipitation controls canopy height, aboveground biomass C, and function of mangrove forests [30, 31]. In mangrove soils, precipitation regulates organic C decomposition by

modifying the oxygen supply to the soil [9, 32, 33] and increases plant productivity and growth by providing freshwater and nutrients [34]. Soil inundation following rainfall also increases mangrove belowground biomass [35]. Therefore, increasing precipitation in tropical coastal regions due to climate change [36] should also increase the C accumulation capacity in some of the world's largest mangrove systems including in the Indo-Pacific and tropical South Americas regions.

Local scale environmental factors, like tidal range, marsh elevation and tropical cyclone frequency are also regarded as critical factors that affect the CAR in coastal wetlands. In this study, we extracted tidal range for each site from a recently developed global tidal range dataset [37, 38]. However, the tidal range variables did not significantly affect the CAR in either tidal marshes or mangroves globally (Table S1 and Fig S6).

Besides tidal ranges, elevation was also regarded as a local environmental factor directly related to the tidal wetland CAR. As a result of the apparent elevation distribution, compiled tidal marsh data can be further be divided into high and low marsh in some sites. However, when we compared the CAR differences between upper tidal (high) marsh and lower tidal (low) marsh no significant difference was found (Fig S7), which was similar as our previous observations in the United States [5]. Similarly, there were some mangroves sites that reported the relative location, we thus separated these sites into high and low tidal sites with apparent elevation difference. However, we also did not observe any significant difference between these tidal areas (Fig S7). As most of our compiled studies did not report site elevations, we further extracted the elevation data for each site from CoastalDEM database, which is a digital terrain model providing bare earth elevations for coastal areas with 90 m horizontal resolution [39]. Again, we found that the elevation did not affect the CAR in either tidal marsh or mangroves (Fig. S6b).

As tropical cyclones have been reported to greatly affect the mangrove aboveground biomass and plant height [31], we also evaluated the tropical cyclone effect on mangrove carbon accumulation rates (CAR) based on the Global Cyclone Hazard Frequency and Distribution, v1 (1980–2000) [40]. Our results (Table S1 and Fig. S6c) also suggested that the tropical cyclone frequency did not significantly affect mangrove CAR. Although higher tropical cyclone frequency was reported to damage the mangrove canopy height, and reduce the aboveground biomass [31], the soil C accumulation in mangroves was a balance between biomass C inputs and soil C decomposition. It was therefore reasonable that tropical cyclone frequency has a limited influence on the mangrove soil CAR.

Global extrapolation and projection

The global extrapolations were based on the 12148 coastline segments from the Dynamic Interactive Vulnerability Assessment (DIVA) modeling framework [41], in which the existing coastal wetlands, as reported by the United Nations Environment Programme World Conservation Monitoring Center (UNEP WCMC), were combined [42]. Our global extrapolations show that the tidal wetland accumulates 53.65 (95%CI: 48.52-59.01) Tg C yr⁻¹ (Figure 1, and Table 1), comparable to the carbon burial in global lakes [43], 30% of the carbon burial in the oceans [44], and 0.5% [45] of the current rate of anthropogenic carbon dioxide emissions. We further separated the data between vegetation types (Table S3), and found that tidal marsh contributed 12.63 Tg C yr⁻¹. The value was comparable to previous reports for the global tidal marsh. For example, Ouyang and Lee [25] estimated ~10.6 Tg C yr⁻¹ sequestered in global 41657 km² salt marsh, based on 143 sites data. However, our global mangrove estimation (41 Tg C yr⁻¹) was generally higher than previous estimates (18.4 to 34 Tg C yr⁻¹) [26, 46], likely because the new soil core data came from tropical mangrove contain some of the highest

mangrove C accumulation rates on the globe (Fig. 1). In terms of C sequestration per country, Indonesia was found to contain the greatest with 14.7 Tg C yr⁻¹, and Australia the second most at 6.86 Tg C yr⁻¹ (Table 1 and Table S3).

To estimate changes in global CAR under projected RCP4.5 and RCP8.5 scenarios by 2100, we applied the linear changes in CAR that is a function of MAP, MAT, and RSLR for tidal marshes and mangroves (Models 1 and 2, see details in M&M section). The coastal wetland area changes under corresponding scenarios were extracted from a recent integrated global model developed by Schuerch, Spencer [11]. The model considered both the ability of coastal wetlands to vertically build up through sediment accretion, and the accommodation space indicated by population density, namely, the vertical and lateral space available for fine sediments to accumulate and colonized by wetland vegetation under different climate change scenarios [11]. We estimate that global tidal wetland C accumulation (Figure 3) will increase by at least 35%, i.e. from 53.65 to 67.5 Tg yr⁻¹ (95%CI: 60-83.7 Tg yr⁻¹) in 2100 under the high-emissions RCP8.5 scenario (Table 1), even though the total area will decrease by 30% as a result of highly limited accommodation space[11]. Under the moderate-emissions RCP4.5 scenario, total C accumulation will increase from 10% to 34% relative to current values, i.e. from 58.8 to 71.6 Tg C yr⁻¹ in 2100, depending on the projected total wetland area regulated by population density [11] (Table 1 & Figure 3&4). We thus suggest that even with a decrease in wetland area, the total C accumulation in tidal wetlands will keep increasing until the end of the 21st century due to increased temperature and precipitation in mangroves, and higher RSLR in tidal marshes. Our estimated increases in soil C accumulation would have numerous implications for coastal C budgets and the assessment of their sink potential. Given increasing atmospheric C and global

temperatures, our modeling implies that the soil organic C accumulation and wetland extent will be more resilient to sea-level rise by 2100 than previously suggested [3, 11].

At the national scale, Australia will have the greatest C sequestration in 2100, ranging from 12 to 27 Tg C yr⁻¹ depending on scenario assumptions (Table 1 and Table S3). Our projections indicate that Indonesia, Malaysia and Mexico are vulnerable to climate change under the most limited lateral accommodation space [11], while countries such as Australia, USA, Brazil and China will have significant gains in tidal wetland C sequestration capacities under the RCP 4.5 and RCP 8.5 scenarios due to expanding wetland area [11, 47].

Most of the C that accumulated in the tidal wetland sediments is derived from autochthonous organic matter, particularly root material [48, 49]. However, the methods used in this study do not account for the carbon that accumulates along the active root zone. The changes in carbon attributed to root mass increases can be important to the ecosystem scale carbon stocks of coastal wetlands of which is outside the scope of the sediment dating techniques employed in this study [50, 51]. Furthermore, much of the organic material produced in tidal wetlands can be exported to the ocean or buried in adjacent areas such as tidal flats [2] and the coastal ocean [51]. For instance, mudflats near tidal wetlands may have comparable sediment C accumulation rates for which tidal vegetated wetlands are often a major source of organic matter [2, 52]. In addition, lateral exports of bicarbonate from tidal marsh [53] and mangrove [51] soils into the coastal ocean can exceed soil accumulation and represents an overlooked C sequestration mechanism on time scales of thousands of years. Under warming conditions, mangrove forests are migrating in a poleward direction, encroaching on tidal marshes and thereby increasing the C accumulation capacity of global tidal wetlands [17, 54, 55]. Therefore, our global C sequestration predictions are conservative since they do not include C that accumulates in active roots, exported to nearby

ecosystems and the increased C sequestration due to the ongoing poleward extension of mangroves.

Our global extrapolations and projections of literature and new data under RCP4.5 and RCP 8.5 contextualize the empirical relationships between CAR and environment factors [56]. The projection of global tidal wetland C sequestration considered both wetland area changes [11] and the CAR response to future climate change factors (mean annual temperature (MAT), mean annual precipitation (MAP) and relative sea-level rise rate (RSLR)). Although the model we used for future tidal wetland area is contested among researchers [11, 38, 57], the results of our global projection mainly rely on the increasing CAR under future climate change factors. For example, the global C sequestration is projected to increase by at least 35% towards the final decade of this century (Figure 3) even though the total area of tidal wetlands will decrease by 30% as predicted by Schuerch, Spencer [11]. Moreover, the patterns of CAR in response to climate change that we observed in this study are in line with a previous modeling stimulation of salt marsh C accumulation in response to climate change [15]. We thus believe that our study captures the actual CAR responses to altered RSLR, precipitation and temperature, which may benefit future process-based models.

Conclusion

Overall, our results highlight the feedbacks between climate change and C sequestration in tidal wetlands. Projected increases in precipitation, temperature and RSLR will drive a net increase in global C sequestration in wetlands during the 21st century. The projected 59 to 100 Tg C yr⁻¹ total global tidal wetland C accumulation in 2100 (Table 1) represents an additional sink of 5 to 46 Tg of atmospheric CO₂-C yr⁻¹ for a total of 53.65 Tg C yr⁻¹. Even though these

global tidal wetlands only occupy <0.1% of the global area, they could offset at least 0.5% of the current anthropogenic CO₂ emission rates, a spatial efficiency that is 15 times higher than terrestrial and 50 times higher than the open ocean per unit area [46]. Therefore, our results demonstrate that preserving and rehabilitating mangroves and salt marshes will remain an effective approach to tackle global climate change with significant regional benefits in tidal wetland-rich countries.

Materials and Methods

We compiled data on sediment accretion rates (SAR) measured in tidal marshes and mangroves worldwide, using ¹³⁷Cs, ²³⁹⁺²⁴⁰Pu, ²¹⁰Pb geochronologies or SET methods to generate the most robust dataset on tidal wetland carbon sequestration rates to date. We used the Web of Science (Thomson Reuters, New York, NY) and Google Scholar (Google Inc., Mountain View, CA) to search the literature using the terms: (accretion)+(tidal wetlands or tidal marsh or mangrove* or coastal wetland or coastal marsh)+(soil or sediments). To be included in our dataset, studies had to cover the temporal scale of sub-decade to decades, which are compatible with anthropogenic climate change and future CMIP5 projections. Data were extracted until September 2018 from published studies or obtained via personal communication. We compiled a total of 102 studies with 564 reported sites that matched our criteria (Table S2).

Besides the above-reported data, we also took 49 sites by our own survey, which contained some regions that had never been reported before, like Africa, and some poor data regions such as India, Brazil, and Indonesia. The sediment core intervals were sealed and cooled for transport. In the laboratory, cores were sectioned at 2 cm intervals; samples with known volumes were weighed and freeze-dried. The dry bulk density was calculated by dividing the dry weight of the

sediment by the initial volume. The total organic carbon was analyzed in a Flash Element Analyzer. For each of these study sites, we collected information on latitude, longitude, mean annual temperature (MAT), mean annual precipitation (MAP), tidal range, mean tidal height, and soil or sediment properties.

Soil C content data in some studies were derived from the measurement of loss on ignition (LOI). LOI measurement of mangrove soils was transformed into organic C content dividing by 1.724 [58]. For tidal marsh soils, we applied the quadratic relationship specific to tidal marshes reported by Craft, Seneca [59]: $TOC=0.04 \times LOI+0.0025 \times LOI^2$. Bulk density (BD) was also not reported in some sites. The missing BD was calculated based on a mixing model that describes the bulk density (BD) as a function of LOI in intertidal wetland sediments [60]. The model assumes that the bulk volume of sediment is equal to the sum of self-packing volumes of organic and mineral components or $BD = 1/[LOI/k_1 + (1-LOI)/k_2]$, where k_1 and k_2 are the self-packing densities of the pure organic and inorganic components, respectively. The values of k_1 and k_2 were estimated to be $0.085 \pm 0.0007 \text{ g cm}^{-3}$ and $1.99 \pm 0.028 \text{ g cm}^{-3}$, respectively [60].

Some studies have directly reported the carbon accumulation rates (CAR), which was calculated by SAR multiplying the soil C density. The vertical SAR represented average soil accretion rates from sub-decade to decades depending on the different dating methods. The ^{137}Cs , $^{239+240}\text{Pu}$ and ^{210}Pb dating methods, and the Sediment Elevation Table (SET) methods were employed in this study to determine average rates of vertical sediment accretion [61, 62]. Where reports made available both SAR and C density, the CAR was calculated in this study. Some studies have reported the SAR and C density or C content for multiple layers, reflecting their changes over time. For these studies, we averaged their C density and SAR over up to 30 cm soils which recorded the most recent C accumulation (less than 100 yrs for most sites).

Besides the soil C related parameters, other environmental variables, i.e., mean annual temperature (MAT), mean annual precipitation (MAP), relative sea-level rise rate (RSLR), total suspended matters (TSM), tidal range, elevation, and tropical cyclone frequency were also collected. MAT and MAP for each site were acquired from world climate data [63]. RSLR data was collected from the Permanent Service for Mean Sea Level (PSMSL) database [64]. Total suspended matter (TSM, mg/L) is derived from MEdium Resolution Imaging Spectrometer (MERIS) satellite data, processed in the framework of the GlobColour project (<http://globcolour.info>) [11]. We use the monthly averages from April 2002 to April 2012 that has a horizontal resolution of $1/24^\circ$. We extracted tidal range for each site from a newly developed global tidal range data set [37, 38], representing the tidal range (that is, the difference between mean low water and mean high water), mean high water neap (MHWN) and mean high water spring (MHWS) tidal levels. Most of the compiled studies did not report the site elevations, and we thus extracted the elevation data for each site from the CoastalDEM database, which is a digital terrain model providing bare earth elevations for coastal areas with 90 m horizontal resolution [39]. We assumed that all the tidal wetlands located between the lowest and highest tidal levels, and removed the elevation data which were much higher or lower than the low and high tidal levels.

As tropical cyclones have been reported to greatly affect the mangrove aboveground biomass and plant height [31], we also evaluated cyclone effects on mangrove carbon accumulation rates (CAR) based on the Global Cyclone Hazard Frequency and Distribution, v1 (1980–2000) [65]. The dataset is a 2.5-minute global grid based on more than 1,600 storm tracks for the period 1 January 1980 through 31 December 2000 for the Atlantic, Pacific, and Indian Oceans that were

assembled and modeled at UNEP/GRID-Geneva PreView [65]. The site-specific cyclone hazard risk was extracted from the dataset.

Statistical analysis

Linear mixed models (LMMs) were used to evaluate the factors that may drive the measured C accumulation rate (CAR) and sediments accretion rate (SAR). The study reference was included as a random factor because clustering replicates by study location could introduce spatial autocorrelation. The LMMs were fit assuming a Gaussian error distribution using the “lme4” package for the R statistical program [66]. We constructed the LMMs separately for tidal marsh and mangrove (Table S1). There were many missing elevation data in our dataset, and Fig S2 and Fig S3 have shown that elevation has no effect on tidal wetland CAR in either tidal marsh or mangrove globally. As a result, we thus did not include elevation in our LMMs analysis. Initial tidal marsh LMMs included all of the putative explanatory variables (including vegetation types, longitude, latitude, Mean Annual Temperature (MAT), Mean Annual Precipitation (MAP), relative sea-level rise rate (RSLR), tidal range and total suspended matter (TSM) to explain the variations in CAR (see Table S1), while the initial mangrove LMMs also have the tropical cyclone risk variable (See the details in Table S1). Given the positive skew in the distribution of CAR, it was log-transformed prior to use in the models. All the environmental variables (MAT, MAP, RSLR, Tidal range, TSM, Cyclone risk) were standardized (subtracting the mean and dividing by standard deviation). The standardization makes coefficients comparable among environmental factors, which we show them in Fig S3. Neither of these data transformations significantly altered the statistical outputs, so were retained in the final models.

Model selection was performed using the Akaike information criterion (AIC) of competing models. AIC measures the relative quality of models for a given set of data, by estimating the

information lost when the model represents the process that generates the data [67, 68]. Practically, AIC rewards goodness of fit (likelihood) and penalizes overfitting (number of parameters). A lower AIC indicates a preferred model. Table S1 shows the AIC values of all the models. In tidal marshes, RSLR and MAT were included in the most parsimonious final model (Model 1), while only MAT and MAP were included in the final mangrove model (Model 2). All reported P-values from LMMs are generated by “r.squaredGLMM” function from “MuMIn” package. Variance explained by the model was estimated by calculating R^2 values for the minimally-adequate LMM following Nakagawa and Schielzeth to retain the random effects structure [68].

Tidal Marsh: $\log(\text{CAR}) \sim \text{RSLR} + \text{MAT} + (1|\text{Studies})$ ----- (Model 1)

Mangrove: $\log(\text{CAR}) \sim \text{MAP} + \text{MAT} + (1|\text{Studies})$ ----- (Model 2)

To assess model performance, we compared modeled vs. observed CAR data, summarized by goodness-of-fit measures. The modeled values were calculated as the expected (or fitted) CAR given environmental variables in the tidal marsh or mangrove datasets. These modeled values were then compared with the observed values. Fig. S4 shows that the modeled and observed log-transformed CAR values are close to the 1:1 reference line. We then summarized their goodness of fit by Pearson correlation coefficient, which is a measure between -1 and $+1$, where a higher value indicates a better fit. For all vegetation groups, the modeled and observed CAR are highly correlated ($R^2 = 0.51$ for the tidal marsh model, and 0.47 for the mangrove model). This assessment validates that the model can predict the data well.

To check model assumptions, we performed residual diagnostics. Residual diagnostics in Fig. S5 show that the residuals are not correlated with CAR values. Fig. S6 validates the normality assumption of the model residuals, supporting log-transformed CAR variables.

We then used the final linear models without the random effects to evaluate the strength of the simple statistical relationship for global tidal marsh and mangrove CAR projections. In tidal marshes, CAR was a linear function of RSLR and MAT (Model 1). However, CAR generally reached a maximum rate if SAR can not increase anymore. In our dataset, SAR is a function of RSLR and total suspended matter (TSM, Linear model: $R^2=0.20$, $p<0.01$, Model 3). We assumed that the TSM for each segment would not change until 2100 [11]. There will be a critical RSLR point for each segment when SAR equals the RSLR in our future projection [11]. We thus assumed that SAR would reach its maximum in the scenario when RSLR is higher than the critical RSLR point, and after this point the SAR will stop increasing with higher RSLR.

Tidal Marsh: $SAR=0.571 RSLR+0.22 TSM+1.62$ ----- (Model 3)

Extrapolation and projection

The global extrapolation was based on the 12148 coastline segments from the Dynamic Interactive Vulnerability Assessment (DIVA) modeling framework [41]. Within each coastline segment, the existing coastal wetlands, as reported by the United Nations Environment Programme World Conservation Monitoring Center (UNEP WCMC), were combined [42]. The geostatistical principle assumes that vegetation distribution gradually changes with environmental factors, like latitude, longitude, temperature and precipitation [69]. We thus assumed that C accumulation at each segment has the highest similarity to that at the closest sampled site. Spatial extrapolation methods thus were used in this study to calculate the CAR of each of the tidal wetland segments: the CAR of each tidal wetland segment was estimated based on the nearest five reported CAR sites. However, due to the rare sampling points in Africa, Europe and high latitude regions, we stress that there will be less accurate for these regions in the extrapolation.

To estimate changes in global CAR under projected RCP4.5 and RCP8.5 scenarios by 2100, we applied the linear changes in CAR that is a function of MAP, MAT, and RSLR for tidal marshes and mangroves. We use projected temperature and precipitation from the NASA Earth Exchange Global Daily Downscaled Projections (NEX-GDDP) dataset [70], a high-resolution ($0.25^\circ \times 0.25^\circ$) dataset of global climate projections downscaled and bias-corrected from the output of 21 general circulation models (GCMs) in the Coupled Model Intercomparison Project Phase 5 (CMIP5) archive [71] under the Representative Concentration Pathway (RCP) 4.5 (represents a moderate-emission scenario in which global-averaged radiative forcing is around 4.5 W m^{-2} in 2100) and RCP8.5 (a comparatively high-emission scenario in which global-averaged radiative forcing is about 8.5 W m^{-2} in 2100) scenarios in the period between 2020 and 2100. However, the ensemble of the 21 realizations described above is an “ensemble of opportunity,” as these GCMs are not independent with each other [72, 73]. Thus, it fails to cover the full uncertainty of future projections and underestimates the extreme values in tails. To provide a probabilistic ensemble of climate projections and cover the tails of the probability distribution that are missing from the GCM ensemble, we use the surrogate/model mixed ensemble (SMME) method [74] to assign probabilistic weights to GCM output and generate surrogate models based on the probability distribution of global mean surface temperature (GMST) produced by a simple climate model. Generally speaking, we first assign the weights by comparing the GMST from the 21 GCM outputs with the probability distribution of GMST produced by a simple climate model (e.g. MAGGIC6, Meinshausen, Raper [75]). The number and weights of model surrogates are also determined in this step to cover the tails of distribution that are not captured by the 21 GCM outputs. We then generate model surrogates by scaling the GMSTs in the tail bins of the GMST probability distribution by the forced component from a

selected GCM output and adding the unforced component from the same GCM. Same with Carleton et al [76], we generate 12 model surrogates. Together with the 21 GCM outputs, this approach provides us 33 climate projections in total and 33 weights assigned to each projection.

We extracted the full distribution of future global RSLR data from Kopp, Horton [77] at decadal intervals for locations of each coastal wetland segment under RCP 4.5 and RCP 8.5 trajectories. Site-to-site differences in the RSLR projections [77] originate from varying non-climatic background uplift or subsidence, oceanographic effects, and spatially variable responses of the geoid and the lithosphere to shrinking land ice.

The projected CAR for tidal marshes and mangroves was thus calculated based on their (MAT, MAP and RSLR) differences between current values and future values by Model 1 and Model 2. The climate and RSLR projections captured physical uncertainty in the climate system. Besides the physical uncertainty, there is an important second source of uncertainty arising from the statistic estimates of the linear models (Model 1 and 2). To capture both sources of uncertainty, we firstly conduct a Monte Carlo procedure. For each of the coastal segments, we randomly draw a set of parameters that compose the CAR in the statistic models from empirical normal distributions defined by the means and confidence intervals. Second, we randomly sample a paired MAT and MAP data from the 33 ensemble members under RCP 4.5 and RCP8.5 scenarios, respectively. Each sampled projection is weighted by corresponding weight which is normalized by the sum of weights of all the samples. Third, we randomly sample an RSLR projection from the full distribution of RSLR projections⁴³. Fourth, a projection of CAR is estimated by combing the parameter sets with corresponding predictors for each equation (e.g. MAT and RSLR for Model 1 and MAT and MAP for Model 2). Finally, we repeat this process to obtain 1000 projection estimates for each equation. The uncertainty of CAR projections is

estimated based on the 1000 projections at each coastal segments and each type of wetland. In this study, we reported their median value and the 95% (2.5%-97.5%) quantiles of the probability distribution.

The future coastal wetland area changes under RCP4.5 and RCP 8.5 were extracted from an integrated global model that considers both the ability of coastal wetlands to build up vertically by sediment accretion, and the accommodation space which is driven by population density in each segment [11]. The projected global coastal wetland C accumulation amount was then calculated as the product of projected future CAR and coastal wetland area for each coastline segment.

The data analysis, global extrapolation and projection was performed by R version 3.6 [78] and Matlab R2016a (The MathWorks, Inc. Natick, MA, USA).

References

1. Duarte CM, Losada IJ and Hendriks IE *et al.* The role of coastal plant communities for climate change mitigation and adaptation. *Nat Clim Change* 2013;**3**:961-968.
2. Breithaupt JL, Smoak JM and Smith TJ *et al.* Organic carbon burial rates in mangrove sediments: Strengthening the global budget. *Global Biogeochem Cycles* 2012;**26**:2012GB004375.
3. Kirwan ML and Megonigal JP. Tidal wetland stability in the face of human impacts and sea-level rise. *Nature* 2013;**504**:53-60.
4. Mudd SM, D'Alpaos A and Morris JT. How does vegetation affect sedimentation on tidal marshes? Investigating particle capture and hydrodynamic controls on biologically mediated sedimentation. *J Geophys Res Earth Surf* 2010;**115**: 2009JF001566.
5. Wang F, Lu X and Sanders CJ *et al.* Tidal wetland resilience to sea level rise increases their carbon sequestration capacity in United States. *Nat Commun* 2019;**10**:5434.
6. McLeod E, Chmura GL and Bouillon S *et al.* A blueprint for blue carbon: toward an improved understanding of the role of vegetated coastal habitats in sequestering CO₂. *Front Ecol Environ* 2011;**9**:552-560.
7. Macreadie PI, Anton A and Raven JA *et al.* The future of Blue Carbon science. *Nat Commun* 2019;**10**:3998.
8. Zinke L. The colours of carbon. *Nat Rev Earth Environ* 2020;**1**:141-141.
9. Spivak AC, Sanderman J and Bowen JL *et al.* Global-change controls on soil-carbon accumulation and loss in coastal vegetated ecosystems. *Nat Geosci* 2019;**12**:685-692.

10. Schuerch M, Vafeidis A and Slawig T *et al.* Modeling the influence of changing storm patterns on the ability of a salt marsh to keep pace with sea level rise. *J Geophys Res Earth Surf* 2013;**118**:84-96.
11. Schuerch M, Spencer T and Temmerman S *et al.* Future response of global coastal wetlands to sea-level rise. *Nature* 2018;**561**:231-234.
12. Rogers K, Kelleway JJ and Saintilan N *et al.* Wetland carbon storage controlled by millennial-scale variation in relative sea-level rise. *Nature* 2019;**567**:91-95.
13. Lovelock CE, Cahoon DR and Friess DA *et al.* The vulnerability of Indo-Pacific mangrove forests to sea-level rise. *Nature* 2015;**526**:559-563.
14. Parkinson RW, Craft C and Delaune RD *et al.* Marsh vulnerability to sea-level rise. *Nat Clim Chang* 2017;**7**:756.
15. Kirwan ML and Mudd SM. Response of saltmarsh carbon accumulation to climate change. *Nature* 2012;**489**:550-554.
16. Osland MJ, Gabler CA and Grace JB *et al.* Climate and plant controls on soil organic matter in coastal wetlands. *Global Chang Bio* 2018;**24**:5361-5379.
17. Coldren GA, Langley JA and Feller IC *et al.* Warming accelerates mangrove expansion and surface elevation gain in a subtropical wetland. *J Ecol* 2019;**107**:79-90.
18. Lovelock CE, Feller IC and Reef R *et al.* Variable effects of nutrient enrichment on soil respiration in mangrove forests. *Plant Soil* 2014;**379**:135-148.
19. Webb EL, Friess DA and Krauss KW *et al.* A global standard for monitoring coastal wetland vulnerability to accelerated sea-level rise. *Nat Clim Change* 2013;**3**:458-465.
20. Breithaupt JL, Smoak JM and Byrne RH *et al.* Avoiding timescale bias in assessments of coastal wetland vertical change. *Limnol Oceanogr* 2018;**63**:S477-S495.
21. Howard J, Hoyt S and Isensee K *et al.* *Coastal Blue Carbon. Methods for assessing carbon stocks and emissions factors in mangroves, tidal salt marshes, and seagrasses.* Conservation International, Intergovernmental Oceanographic Commission of UNESCO, International Union for Conservation of Nature. Arlington, Virginia, US. 2014.
22. Donato DC, Kauffman JB and Murdiyarso D *et al.* Mangroves among the most carbon-rich forests in the tropics. *Nat Geosci* 2011;**4**:293-297.
23. Atwood TB, Connolly RM and Almahasheer H *et al.* Global patterns in mangrove soil carbon stocks and losses. *Nat Clim Chang* 2017;**7**:523-528.
24. Breithaupt JL, Smoak JM and Byrne RH *et al.* Avoiding timescale bias in assessments of coastal wetland vertical change. *Limnol Oceanogr* 2018; **64**:10783.
25. Ouyang X and Lee SY. Updated estimates of carbon accumulation rates in coastal marsh sediments. *Biogeosciences* 2014;**11**:5057-5071.
26. Chmura GL, Anisfeld SC and Cahoon DR *et al.* Global carbon sequestration in tidal, saline wetland soils. *Glob Biogeochem Cycles* 2003;**17**:1111.
27. Giri C, Ochieng E and Tieszen LL *et al.* Status and distribution of mangrove forests of the world using earth observation satellite data. *Global Ecol Biogeogr* 2011;**20**:154-159.
28. Chmura GL, Anisfeld SC and Cahoon DR *et al.* Global carbon sequestration in tidal, saline wetland soils. *Global Biogeochem Cycles* 2003;**17**:22-1.
29. Kusumaningtyas MA, Hutahaean AA and Fischer HW *et al.* Variability in the organic carbon stocks, sources, and accumulation rates of Indonesian mangrove ecosystems. *Estuarine Coast Shelf Sci* 2019;**218**:310-323.
30. Osland MJ, Feher LC and Griffith KT *et al.* Climatic controls on the global distribution, abundance, and species richness of mangrove forests. *Ecol Monogr* 2017;**87**:341-359.

31. Simard M, Fatoyinbo L and Smetanka C *et al.* Mangrove canopy height globally related to precipitation, temperature and cyclone frequency. *Nat Geosci* 2019;**12**:40-45.
32. Davidson EA and Janssens IA. Temperature sensitivity of soil carbon decomposition and feedbacks to climate change. *Nature* 2006;**440**:165-173.
33. Sanders CJ, Maher DT and Tait DR *et al.* Are global mangrove carbon stocks driven by rainfall? *J Geophys Res Biogeos* 2016; **121**: 2600-2609.
34. Ewel KC, Bourgeois JA and Cole TG *et al.* Variation in environmental characteristics and vegetation in high-rainfall mangrove forests, Kosrae, Micronesia. *Global Ecol Biogeogr Lett* 1998;**7**:49-56.
35. Castañeda-Moya E, Twilley RR and Rivera-Monroy VH. Allocation of biomass and net primary productivity of mangrove forests along environmental gradients in the Florida Coastal Everglades, USA. *Forest Ecol Manage* 2013;**307**:226-241.
36. Tan J, Jakob C and Rossow WB *et al.* Increases in tropical rainfall driven by changes in frequency of organized deep convection. *Nature* 2015;**519**:451-454. doi: 10.1038/nature14339.
37. Pickering MD, Horsburgh KJ and Blundell JR *et al.* The impact of future sea-level rise on the global tides. *Cont Shelf Res* 2017;**142**:50-68. doi: 10.1016/j.csr.2017.02.004.
38. Schuerch M, Spencer T and Temmerman S *et al.* Reply to “Global coastal wetland expansion under accelerated sea-level rise is unlikely”. *EarthArXiv* 2020. doi: 10.31223/osf.io/ycunb.
39. Kulp SA and Strauss BH. New elevation data triple estimates of global vulnerability to sea-level rise and coastal flooding. *Nat Commun* 2019;**10**:4844.
40. Dilley M, Chen RS and Deichmann U *et al.*; Natural Disaster Hotspots: A Global Risk Analysis. Washington, D.C.: World Bank, 2005.
41. Spencer T, Schuerch M and Nicholls RJ *et al.* Global coastal wetland change under sea-level rise and related stresses: The DIVA Wetland Change Model. *Global Planet Change* 2016;**139**:15-30.
42. McOwen CJ, Weatherdon LV and Bochove J-WV *et al.* A global map of saltmarshes. *Biodivers Data J* 2017; **5**: e11764.
43. Mendonça R, Müller RA and Clow D *et al.* Organic carbon burial in global lakes and reservoirs. *Nat Commun* 2017;**8**: 1694.
44. Duarte CM and Middelburg JJ Caraco N. Major role of marine vegetation on the oceanic carbon cycle. *Biogeosciences* 2005;**2**:1-8.
45. Quéré C, Andrew R and Friedlingstein P *et al.* Global Carbon Budget 2018. *Earth Syst Sci Data* 2018;**10**:2141-2194.
46. Breithaupt JL, Smoak JM and Smith TJ *et al.* Organic carbon burial rates in mangrove sediments: Strengthening the global budget. *Global Biogeochem Cycles* 2012;**26**: GB3011.
47. IPCC. *Climate Change 2014: Synthesis Report*, 2014.
48. Ouyang X and Lee SY. Connolly RM. The role of root decomposition in global mangrove and saltmarsh carbon budgets. *Earth Sci Rev* 2017;**166**:53-63.
49. Robertson AI and Aongi DM. Massive turnover rates of fine root detrital carbon in tropical Australian mangroves. *Oecologia* 2016;**180**:841-851.
50. Lamont K, Saintilan N and Kelleway JJ *et al.* Thirty-Year Repeat Measures of Mangrove Above- and Below-Ground Biomass Reveals Unexpectedly High Carbon Sequestration. *Ecosystems* 2020;**23**:370-382.

51. Santos IR, Maher DT and Larkin R *et al.* Carbon outwelling and outgassing vs. burial in an estuarine tidal creek surrounded by mangrove and saltmarsh wetlands. *Limnol Oceanogr* 2019;**64**:996-1013.
52. Pérez A, Libardoni BG and Sanders CJ. Factors influencing organic carbon accumulation in mangrove ecosystems. *Biol Lett* 2018;**14**: 0237.
53. Wang ZA, Kroeger KD, Ganju NK *et al.* Intertidal salt marshes as an important source of inorganic carbon to the coastal ocean. *Limnol Oceanogr* 2016;**61**:1916-1931.
54. Doughty CL, Langley JA and Walker WS *et al.* Mangrove Range Expansion Rapidly Increases Coastal Wetland Carbon Storage. *Estuarie Coast* 2015;**39**:385-396.
55. Saintilan N, Wilson NC and Rogers K *et al.* Mangrove expansion and salt marsh decline at mangrove poleward limits. *Global Change Biol* 2014;**20**:147-157.
56. Morris JT, Sundareshwar PV and Nietch CT *et al.* Responses of Coastal Wetlands to Rising Sea Level. *Ecology* 2002;**83**:2869-2877.
57. Törnqvist TrE, Cahoon DR and Day JW *et al.* Global coastal wetland expansion under accelerated sea-level rise is unlikely. *EarthArXiv* 2019. doi: 10.31223/osf.io/d2nhs.
58. Allen SE, Grimshaw HM and Parkinson JA *et al.* *Chemical analysis of ecological materials* 565, 1974.
59. Craft CB, Seneca ED and Broome SW. Loss on ignition and kjeldahl digestion for estimating organic carbon and total nitrogen in estuarine marsh soils: Calibration with dry combustion. *Estuaries* 1991;**14**:175-179.
60. Morris JT, Barber DC and Callaway JC *et al.* Contributions of organic and inorganic matter to sediment volume and accretion in tidal wetlands at steady state. *Earth Future* 2016;**4**:110-121.
61. Armentano TV and Woodwell GM. Sedimentation rates in a Long Island marsh determined by ²¹⁰Pb dating. *Limnol Oceanogr* 1975;**20**:452-456.
62. Delaune RD, Patrick WH, and Buresh RJ. Sedimentation rates determined by ¹³⁷Cs dating in a rapidly accreting salt marsh. *Nature* 1978;**275**:532-533.
63. Fick SE and Hijmans RJ. WorldClim 2: new 1-km spatial resolution climate surfaces for global land areas. *Int J Climatol* 2017;**37**:4302-4315.
64. Holgate SJ, Matthews A and Woodworth PL *et al.* New Data Systems and Products at the Permanent Service for Mean Sea Level. *J Coast Res* 2013;**29**:493-504.
65. Center for Hazards and Risk Research - CHRR - Columbia University, Center for International Earth Science Information Network - CIESIN - Columbia University, International Bank for Reconstruction and Development - The World Bank *et al.*; Global Cyclone Hazard Frequency and Distribution. Palisades, NY: NASA Socioeconomic Data and Applications Center (SEDAC), 2005.
66. Bates D, Mächler M and Bolker B *et al.* Fitting Linear Mixed-Effects Models Using lme4. *J Stat Softw* 2015;**67**:48.
67. Zhu K, Chiariello NR and Tobeck T *et al.* Nonlinear, interacting responses to climate limit grassland production under global change. *Proc Natl Acad Sci USA* 2016;**113**:10589-10594.
68. Nakagawa S and Schielzeth H. A general and simple method for obtaining R² from generalized linear mixed-effects models. *Methods Ecol Evol* 2013;**4**:133-142.
69. Reich PB, Luo Y and Bradford JB *et al.* Temperature drives global patterns in forest biomass distribution in leaves, stems, and roots. *Proc Natl Acad Sci USA* 2014;**111**:13721-13726.

70. Thrasher B, Maurer EP, McKellar C *et al.* Technical Note: Bias correcting climate model simulated daily temperature extremes with quantile mapping. *Hydrol Earth Syst Sci* 2012;**16**:3309-3314.
71. Taylor KE, Stouffer RJ and Meehl GA. An Overview of CMIP5 and the Experiment Design. *B Am Meteorol Soc* 2012;**93**:485-498.
72. van Vuuren DP, Edmonds J and Kainuma M *et al.* The representative concentration pathways: an overview. *Clim Change* 2011;**109**:5-31.
73. Moss RH, Edmonds JA and Hibbard KA *et al.* The next generation of scenarios for climate change research and assessment. *Nature* 2010;**463**:747-756.
74. Rasmussen DJ, Meinshausen M and Kopp RE. Probability-Weighted Ensembles of U.S. County-Level Climate Projections for Climate Risk Analysis. *J Appl Meteorol Climatol* 2016;**55**:2301-2322.
75. Meinshausen M, Raper SCB and Wigley TML. Emulating coupled atmosphere-ocean and carbon cycle models with a simpler model, MAGICC6 – Part 1: Model description and calibration. *Atmospheric Chem Phys* 2011;**11**:1417-1456.
76. Carleton T, Delgado M and Greenstone M *et al.*; Valuing the Global Mortality Consequences of Climate Change Accounting for Adaptation Costs and Benefits. 2019; doi: 10.2139/ssrn.3224365.
77. Kopp RE, Horton RM and Little CM *et al.* Probabilistic 21st and 22nd century sea-level projections at a global network of tide-gauge sites. *Earth Future* 2014;**2**:383-406.
78. R Core Team; R: A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing, 2016.

ORIGINAL UNEDITED MANUSCRIPT

Funding:

FW is funded by the National Natural Science Foundation of China (31670621,31870463), the Key Special Project for Introduced Talents Team of Southern Marine Science and Engineering Guangdong Laboratory (Guangzhou) (GML2019ZD0408), the Guangdong Natural Science Foundation for Distinguished Young Scholars (21050000107), the CAS Youth Innovation Promotion Association and the National Forestry and Grassland Administration Youth Talent Support Program (2020BJ003); ZL was funded by R & D program of Guangdong Provincial Department of Science and Technology (2018B030324003); CJS and IRS and most of the new field and laboratory data were supported by the Australian Research Council (DE160100443, DP150103286, DE140101733 and LE140100083); JT is funded by the NOAA National Estuarine Research Reserve Science Collaborative (NA09NOS4190153 and NA14NOS4190145).

Author contributions:

FW and CS conceived and designed the study together. FW collected the data from the literature, conducted the data analysis and modeling, and prepared the revision of this paper. CS collected the field survey and drafted the first version of the paper. JY provided the SMME-CMIP5 climate data. REK and JY provided the sea-level projections. MS provided data of the modeling of global wetland extents. All authors contributed to the writing and editing of the manuscript.

Competing interests:

Authors declare no competing interests.

Data and materials availability:

All the data used in this study was shown in results and supplementary information. The code and raw data will be available in Github after this paper is published.




Figure 1. The spatial distribution of tidal wetlands and the observed C accumulation rate (CAR) points. The right panel indicated the arithmetic average CAR at every 10-degree bands of latitude.

ORIGINAL UNEDITED MANUSCRIPT




Figure 2. National C accumulation rates worldwide. The 20 countries listed here where observations are available account for 62% global tidal wetland distribution. Diamond is the mean value, the box is interquartile range, error bar is the largest and smallest value within 1.5 times interquartile range above 75% and below 25%, respectively. Black points indicate outside values.




Figure 3. Projected global change in tidal wetland C sequestration. Results are displayed for moderate-emission scenario (RCP4.5) and high-emission scenario (RCP8.5) under the most restricted human adaptation scenario defined by Schuerch, Spencer [11] (i.e. population density threshold set as 5 people km⁻²; higher population density than this value would have no lateral accommodation space for tidal wetlands), giving the most conservative predicted increases in the global carbon accumulation rates found in this study. The colored shadings represent the 95% range (2.5%-97.5%) of the projected C distribution in global tidal wetlands. The solid lines denote the median of the distribution.




Figure 4. The projected 2100 spatial distribution of tidal wetland C sequestration changes. Relative changes in tidal wetland C sequestration are displayed for the RCP4.5 (a,c) and RCP8.5 (b,d) scenarios under two assumed human activity scenarios by Schuerch et al. [11]: inhibition of wetland inland migration in the regions with a population density (a,b) over 5 people km⁻² and in the regions with a population density (c,d) of over 300 people km⁻².

ORIGINAL UNEDITED MANUSCRIPT

Table 1. Top ten countries in median annual tidal wetland C sequestration under current conditions and projected scenarios in 2100. Tidal wetland C sequestrations are displayed in the moderate-emission RCP4.5 and high-emission RCP8.5 scenarios under two assumed human activity scenarios as defined by Schuerch et al. [11]: inhibition of wetland inland migration in the regions with a population density over 5 people km⁻² and in the regions with a population density of over 300 people km⁻².

Country	Current (C Tg)	RCP4.5 Pop. 300 (C Tg)	RCP8.5 Pop. 300 (C Tg)	RCP4.5 Pop. 5 (C Tg)	RCP8.5 Pop. 5 (C Tg)
Indonesia	14.7	19.22	19.10	15.06	10.35
Australia	6.86	12.71	26.63	12.25	25.29
USA	4.11	6.06	7.07	4.50	4.71
Brazil	3.26	4.24	5.51	3.83	4.36
Malaysia	2.01	3.54	3.28	3.23	2.56
Papua New Guinea	1.96	2.31	2.78	1.98	2.21
Mexico	1.56	1.85	7.43	0.97	0.73
Nigeria	1.33	1.03	0.61	0.99	0.15
China	1.24	1.87	3.64	1.82	3.45
Thailand	1.12	1.11	1.00	1.13	0.84
<i>Others</i>	<i>15.45</i>	<i>17.66</i>	<i>22.95</i>	<i>13.04</i>	<i>12.85</i>
Total	53.65	71.6	100	58.8	67.5

ORIGINAL UNEDITED MANUSCRIPT