

Sum Formulas For Generalized Tetranacci Numbers: Closed Forms of the Sum Formulas $\sum_{k=0}^n x^k W_k$ and $\sum_{k=1}^n x^k W_{-k}$

Yüksel Soykan

Department of Mathematics, Art and Science Faculty, Zonguldak Bülent Ecevit University, 67100, Zonguldak, Turkey

Email: yuksel_soykan@hotmail.com

Received: January 22, 2021; Accepted: February 12, 2021; Published: February 19, 2021

Copyright © 2021 by author(s) and Scitech Research Organisation (SRO).

This work is licensed under the Creative Commons Attribution International License (CC BY).

<http://creativecommons.org/licenses/by/4.0/>

Abstract

In this paper, closed forms of the sum formulas $\sum_{k=0}^n x^k W_k$ and $\sum_{k=1}^n x^k W_{-k}$ for generalized Tetranacci numbers are presented. As special cases, we give summation formulas of Tetranacci, Tetranacci-Lucas, and other fourth-order recurrence sequences.

2020 Mathematics Subject Classification. 11B37, 11B39, 11B83.

Keywords

Tetranacci numbers, Tetranacci-Lucas numbers, fourth order Pell numbers, sum formulas, summing formulas.

1. Introduction

There have been so many studies of the sequences of numbers in the literature which are defined recursively. Two of these type of sequences are the sequences of Tetranacci and Tetranacci-Lucas which are special case of generalized Tetranacci numbers. A generalized Tetranacci sequence

$$\{W_n\}_{n \geq 0} = \{W_n(W_0, W_1, W_2, W_3; r, s, t, u)\}_{n \geq 0}$$

is defined by the fourth-order recurrence relations

$$W_n = rW_{n-1} + sW_{n-2} + tW_{n-3} + uW_{n-4}, \quad (1)$$

with the initial values W_0, W_1, W_2, W_3 are arbitrary complex (or real) numbers not all being zero and r, s, t, u are real numbers.

This sequence has been studied by many authors and more detail can be found in the extensive literature dedicated to these sequences, see for example [5,8,9,11,30,32,33].

The sequence $\{W_n\}_{n \geq 0}$ can be extended to negative subscripts by defining

$$W_{-n} = -\frac{t}{u}W_{-(n-1)} - \frac{s}{u}W_{-(n-2)} - \frac{r}{u}W_{-(n-3)} + \frac{1}{u}W_{-(n-4)}$$

Cite this paper: Soykan, Y. (2021). Sum Formulas For Generalized Tetranacci Numbers: Closed Forms of the Sum Formulas $\sum_{k=0}^n x^k W_k$ and $\sum_{k=1}^n x^k W_{-k}$. Journal of Progressive Research in Mathematics, 18(1), 24-47. Retrieved from <http://scitecresearch.com/journals/index.php/jprm/article/view/2018>

for $n = 1, 2, 3, \dots$. Therefore, recurrence (1) holds for all integer n .

For some specific values of W_0, W_1, W_2, W_3 and r, s, t, u , it is worth presenting these special Tetranacci numbers in a table as a specific name. In literature, for example, the following names and notations (see Table 1) are used for the special cases of r, s, t, u and initial values.

In literature, for example, the following names and notations (see Table 1) are used for the special case of r, s, t, u and initial values.

Table 1. A few special case of generalized Tetranacci sequences.

No	Sequences (Numbers)	Notation	OEIS [12]	Ref.
1	Tetranacci	$\{M_n\} = \{W_n(0, 1, 1, 2; 1, 1, 1, 1)\}$	A000078	[26]
2	Tetranacci-Lucas	$\{R_n\} = \{W_n(4, 1, 3, 7; 1, 1, 1, 1)\}$	A073817	[26]
3	fourth order Pell	$\{P_n^{(4)}\} = \{W_n(0, 1, 2, 5; 2, 1, 1, 1)\}$	A103142	[27]
4	fourth order Pell-Lucas	$\{Q_n^{(4)}\} = \{W_n(4, 2, 6, 17; 2, 1, 1, 1)\}$	A331413	[[27]
5	modified fourth order Pell	$\{E_n^{(4)}\} = \{W_n(0, 1, 1, 3; 2, 1, 1, 1)\}$	A190139	[[27]
6	fourth order Jacobsthal	$\{J_n^{(4)}\} = \{W_n(0, 1, 1, 1; 1, 1, 1, 2)\}$	A007909	[28]
7	fourth order Jacobsthal-Lucas	$\{J_n^{(4)}\} = \{W_n(2, 1, 5, 10; 1, 1, 1, 2)\}$	A226309	[28]
8	modified fourth order Jacobsthal	$\{K_n^{(4)}\} = \{W_n(3, 1, 3, 10; 1, 1, 1, 2)\}$		[28]
9	fourth-order Jacobsthal Perrin	$\{Q_n^{(4)}\} = \{W_n(3, 0, 2, 8; 1, 1, 1, 2)\}$		[28]
10	adjusted fourth-order Jacobsthal	$\{S_n^{(4)}\} = \{W_n(0, 1, 1, 2; 1, 1, 1, 2)\}$		[28]
11	modified fourth-order Jacobsthal-Lucas	$\{R_n^{(4)}\} = \{W_n(4, 1, 3, 7; 1, 1, 1, 2)\}$		[28]
12	4-primes	$\{G_n\} = \{W_n(0, 0, 1, 2; 2, 3, 5, 7)\}$		[29]
13	Lucas 4-primes	$\{H_n\} = \{W_n(4, 2, 10, 41; 2, 3, 5, 7)\}$		[29]
14	modified 4-primes	$\{E_n\} = \{W_n(0, 0, 1, 1; 2, 3, 5, 7)\}$		[29]

Here OEIS stands for On-line Encyclopedia of Integer Sequences. For easy writing, from now on, we drop the superscripts from the sequences, for example we write J_n for $J_n^{(4)}$.

We present some works on summing formulas of the numbers in the following Table 2.

Table 2. A few special study of sum formulas.

Name of sequence	Papers which deal with summing formulas
Pell and Pell-Lucas	[1,3,31],[6,7]
Generalized Fibonacci	[4,13,14,15,16,17,19]
Generalized Tribonacci	[2,10,18]
Generalized Tetranacci	[20,25,32]
Generalized Pentanacci	[21,22]
Generalized Hexanacci	[23,24]

In this work, we investigate linear summation formulas of generalized Tetranacci numbers.

2 Linear Sum Formulas of Generalized Tetranacci Numbers with Positive Subscripts

The following theorem presents some linear summing formulas of generalized Tetranacci numbers with positive subscripts.

Theorem 1. *Let x be a real or complex numbers. For $n \geq 0$ we have the following formulas:*

(a) *If $rx + sx^2 + tx^3 + ux^4 - 1 \neq 0$, then*

$$\sum_{k=0}^n x^k W_k = \frac{\Theta_1(x)}{rx + sx^2 + tx^3 + ux^4 - 1}.$$

(b) If $r^2x + 2ux^2 - s^2x^2 + t^2x^3 - u^2x^4 + 2sx + 2rtx^2 - 2sux^3 - 1 \neq 0$ then

$$\sum_{k=0}^n x^k W_{2k} = \frac{\Theta_2(x)}{r^2x + 2ux^2 - s^2x^2 + t^2x^3 - u^2x^4 + 2sx + 2rtx^2 - 2sux^3 - 1}.$$

and

(c) If $r^2x + 2ux^2 - s^2x^2 + t^2x^3 - u^2x^4 + 2sx + 2rtx^2 - 2sux^3 - 1 \neq 0$ then

$$\sum_{k=0}^n x^k W_{2k+1} = \frac{\Theta_3(x)}{r^2x + 2ux^2 - s^2x^2 + t^2x^3 - u^2x^4 + 2sx + 2rtx^2 - 2sux^3 - 1},$$

where

$$\Theta_1(x) = x^{n+3}W_{n+3} - x^{n+2}(rx - 1)W_{n+2} - x^{n+1}(sx^2 + rx - 1)W_{n+1} + ux^{n+4}W_n - x^3W_3 + x^2(rx - 1)W_2 + x(sx^2 + rx - 1)W_1 + (tx^3 + sx^2 + rx - 1)W_0,$$

$$\Theta_2(x) = x^{n+1}(-ux^2 - sx + 1)W_{2n+2} + x^{n+2}(t + rs + rux)W_{2n+1} + x^{n+2}(u + t^2x - u^2x^2 + rt - sux)W_{2n} + ux^{n+2}(r + tx)W_{2n-1} - x^2(r + tx)W_3 + x(r^2x + ux^2 + sx + rtx^2 - 1)W_2 - x^2(t + rux - stx)W_1 + (r^2x + ux^2 - s^2x^2 + t^2x^3 + 2sx + 2rtx^2 - sux^3 - 1)W_0,$$

$$\Theta_3(x) = x^{n+1}(r + tx)W_{2n+2} + x^{n+1}(s - s^2x + t^2x^2 - u^2x^3 + ux - 2sux^2 + rtx)W_{2n+1} + x^{n+1}(t + rux - stx)W_{2n} - ux^{n+1}(ux^2 + sx - 1)W_{2n-1} + x(ux^2 + sx - 1)W_3 - x^2(t + rs + rux)W_2 + (r^2x + ux^2 - s^2x^2 + 2sx + rtx^2 - sux^3 - 1)W_1 - ux^2(r + tx)W_0.$$

Proof.

(a) Using the recurrence relation

$$W_n = rW_{n-1} + sW_{n-2} + tW_{n-3} + uW_{n-4}$$

i.e.

$$uW_{n-4} = W_n - rW_{n-1} - sW_{n-2} - tW_{n-3}$$

we obtain

$$\begin{aligned} ux^0W_0 &= x^0W_4 - rx^0W_3 - sx^0W_2 - tx^0W_1 \\ ux^1W_1 &= x^1W_5 - rx^1W_4 - sx^1W_3 - tx^1W_2 \\ ux^2W_2 &= x^2W_6 - rx^2W_5 - sx^2W_4 - tx^2W_3 \\ ux^3W_3 &= x^3W_7 - rx^3W_6 - sx^3W_5 - tx^3W_4 \\ &\vdots \\ ux^{n-3}W_{n-3} &= x^{n-3}W_{n+1} - rx^{n-3}W_n - sx^{n-3}W_{n-1} - tx^{n-3}W_{n-2} \\ ux^{n-2}W_{n-2} &= x^{n-2}W_{n+2} - rx^{n-2}W_{n+1} - sx^{n-2}W_n - tx^{n-2}W_{n-1} \\ ux^{n-1}W_{n-1} &= x^{n-1}W_{n+3} - rx^{n-1}W_{n+2} - sx^{n-1}W_{n+1} - tx^{n-1}W_n \\ ux^nW_n &= x^nW_{n+4} - rx^nW_{n+3} - sx^nW_{n+2} - tx^nW_{n+1} \end{aligned}$$

If we add the above equations side by side, we get (a).

(b) and (c) Using the recurrence relation

$$W_n = rW_{n-1} + sW_{n-2} + tW_{n-3} + uW_{n-4}$$

i.e.

$$rW_{n-1} = W_n - sW_{n-2} - tW_{n-3} - uW_{n-4}$$

we obtain

$$\begin{aligned}
rx^1W_3 &= x^1W_4 - sx^1W_2 - tx^1W_1 - ux^1W_0 \\
rx^2W_5 &= x^2W_6 - sx^2W_4 - tx^2W_3 - ux^2W_2 \\
rx^3W_7 &= x^3W_8 - sx^3W_6 - tx^3W_5 - ux^3W_4 \\
rx^4W_9 &= x^4W_{10} - sx^4W_8 - tx^4W_7 - ux^4W_6 \\
&\vdots \\
rx^{n-1}W_{2n-1} &= x^{n-1}W_{2n} - sx^{n-1}W_{2n-2} - tx^{n-1}W_{2n-3} - ux^{n-1}W_{2n-4} \\
rx^nW_{2n+1} &= x^nW_{2n+2} - sx^nW_{2n} - tx^nW_{2n-1} - ux^nW_{2n-2} \\
rx^{n+1}W_{2n+3} &= x^{n+1}W_{2n+4} - sx^{n+1}W_{2n+2} - tx^{n+1}W_{2n+1} - ux^{n+1}W_{2n}
\end{aligned}$$

Now, if we add the above equations side by side, we get

$$\begin{aligned}
r(-W_1 + \sum_{k=0}^n x^k W_{2k+1}) &= (x^n W_{2n+2} - W_2 - x^{-1} W_0 + \sum_{k=0}^n x^{k-1} W_{2k}) \\
&\quad -s(-W_0 + \sum_{k=0}^n x^k W_{2k}) - t(-x^{n+1} W_{2n+1} + \sum_{k=0}^n x^{k+1} W_{2k+1}) \\
&\quad -u(-x^{n+1} W_{2n} + \sum_{k=0}^n x^{k+1} W_{2k}).
\end{aligned} \tag{2}$$

Similarly, using the recurrence relation

$$W_n = rW_{n-1} + sW_{n-2} + tW_{n-3} + uW_{n-4}$$

i.e.

$$rW_{n-1} = W_n - sW_{n-2} - tW_{n-3} - uW_{n-4}$$

we write the following obvious equations;

$$\begin{aligned}
rx^1W_2 &= x^1W_3 - sx^1W_1 - tx^1W_0 - ux^1W_{-1} \\
rx^2W_4 &= x^2W_5 - sx^2W_3 - tx^2W_2 - ux^2W_1 \\
rx^3W_6 &= x^3W_7 - sx^3W_5 - tx^3W_4 - ux^3W_3 \\
rx^4W_8 &= x^4W_9 - sx^4W_7 - tx^4W_6 - ux^4W_5 \\
&\vdots \\
rx^{n-1}W_{2n-2} &= x^{n-1}W_{2n-1} - sx^{n-1}W_{2n-3} - tx^{n-1}W_{2n-4} - ux^{n-1}W_{2n-5} \\
rx^nW_{2n} &= x^nW_{2n+1} - sx^nW_{2n-1} - tx^nW_{2n-2} - ux^nW_{2n-3} \\
rx^{n+1}W_{2n+2} &= x^{n+1}W_{2n+3} - sx^{n+1}W_{2n+1} - tx^{n+1}W_{2n} - ux^{n+1}W_{2n-1} \\
rx^{n+2}W_{2n+4} &= x^{n+2}W_{2n+5} - sx^{n+2}W_{2n+3} - tx^{n+2}W_{2n+2} - ux^{n+2}W_{2n+1}
\end{aligned}$$

Now, if we add the above equations side by side, we obtain

$$\begin{aligned}
r(-W_0 + \sum_{k=0}^n x^k W_{2k}) &= (-W_1 + \sum_{k=0}^n x^k W_{2k+1}) - s(-x^{n+1} W_{2n+1} + \sum_{k=0}^n x^{k+1} W_{2k+1}) \\
&\quad -t(-x^{n+1} W_{2n} + \sum_{k=0}^n x^{k+1} W_{2k}) \\
&\quad -u(-x^{n+2} W_{2n+1} - x^{n+1} W_{2n-1} + x^1 W_{-1} + \sum_{k=0}^n x^{k+2} W_{2k+1})
\end{aligned}$$

Since

$$W_{-1} = -\frac{t}{u}W_0 - \frac{s}{u}W_1 - \frac{r}{u}W_2 + \frac{1}{u}W_3$$

we have

$$\begin{aligned} r(-W_0 + \sum_{k=0}^n x^k W_{2k}) &= (-W_1 + \sum_{k=0}^n x^k W_{2k+1}) - s(-x^{n+1}W_{2n+1} + \sum_{k=0}^n x^{k+1}W_{2k+1}) \\ &\quad - t(-x^{n+1}W_{2n} + \sum_{k=0}^n x^{k+1}W_{2k}) \\ &\quad - u(-x^{n+2}W_{2n+1} - x^{n+1}W_{2n-1} \\ &\quad + x^1(-\frac{t}{u}W_0 - \frac{s}{u}W_1 - \frac{r}{u}W_2 + \frac{1}{u}W_3) + \sum_{k=0}^n x^{k+2}W_{2k+1}). \end{aligned} \quad (3)$$

Then, solving the system (2)-(3), the required results of (b) and (c) follow. \square

Note that (a) of the above theorem can be written as follows:

$$\sum_{k=0}^n x^k W_k = \frac{g}{rx + sx^2 + tx^3 + ux^4 - 1}$$

where

$$g = x^{n+4}W_{n+4} - x^{n+3}(rx-1)W_{n+3} - x^{n+2}(sx^2+rx-1)W_{n+2} - x^{n+1}(tx^3+sx^2+rx-1)W_{n+1} - x^3W_3 + x^2(rx-1)W_2 + x(sx^2+rx-1)W_1 + (tx^3+sx^2+rx-1)W_0.$$

3 Special Cases

In this section, for the special cases of x , we present the closed form solutions (identities) of the sums $\sum_{k=0}^n x^k W_k$, $\sum_{k=0}^n x^k W_{2k}$ and $\sum_{k=0}^n x^k W_{2k+1}$ for the specific case of sequence $\{W_n\}$.

3.1 The case $x = 1$

In this subsection we consider the special case $x = 1$.

The case $x = 1$ of Theorem 1 is given in [20]. For the generalized 4-primes sequence case ($x = 1, r = 2, s = 3, t = 5, u = 7$), see [29].

We only consider the case $x = 1, r = 1, s = 1, t = 1, u = 2$ (which is not considered in [20]).

Observe that setting $x = 1, r = 1, s = 1, t = 1, u = 2$ (i.e. for the generalized fourth order Jacobsthal sequence case) in Theorem 1 (b),(c) makes the right hand side of the sum formulas to be an indeterminate form. Application of L'Hospital rule however provides the evaluation of the sum formulas.

Theorem 2. *If $r = 1, s = 1, t = 1, u = 2$ then for $n \geq 0$ we have the following formulas:*

(a) $\sum_{k=0}^n W_k = \frac{1}{4}(W_{n+3} - W_{n+1} + 2W_n - W_3 + W_1 + 2W_0).$

(b) $\sum_{k=0}^n W_{2k} = \frac{1}{12}((2n+7)W_{2n+2} - 2(2n+5)W_{2n+1} + (2n+13)W_{2n} - 2(2n+5)W_{2n-1} + 5W_3 - 12W_2 + 5W_1 - 6W_0).$

(c) $\sum_{k=0}^n W_{2k+1} = \frac{1}{12}(-(2n+3)W_{2n+2} + 4(n+5)W_{2n+1} - (2n+3)W_{2n} + 2(2n+7)W_{2n-1} - 7W_3 + 10W_2 - W_1 + 10W_0).$

Proof.

(a) We use Theorem 1 (a). If we set $x = 1, r = 1, s = 1, t = 1, u = 2$ in Theorem 1 (a) we get (a).

(b) We use Theorem 1 (b). If we set $r = 1, s = 1, t = 1, u = 2$ in Theorem 1 (b) then we have

$$\sum_{k=0}^n x^k W_{2k} = \frac{g_1(x)}{-4x^4 - 3x^3 + 5x^2 + 3x - 1}$$

where

$$g_1(x) = -(2x^2 + x - 1)x^{n+1}W_{2n+2} + (2x+2)x^{n+2}W_{2n+1} - (4x^2 + x - 3)x^{n+2}W_{2n} + 2(x+1)x^{n+2}W_{2n-1} + x(3x^2 + 2x - 1)W_2 - (x+1)x^2W_1 + (-x^3 + 3x^2 + 3x - 1)W_0 - (x+1)x^2W_3.$$

For $x = 1$, the right hand side of the above sum formula is an indeterminate form. Now, we can use L'Hospital rule. Then we get (b) using

$$\sum_{k=0}^n W_{2k} = \frac{\frac{d}{dx}(g_1(x))}{\frac{d}{dx}(-4x^4 - 3x^3 + 5x^2 + 3x - 1)} \Big|_{x=1}.$$

(c) We use Theorem 1 (c). If we set $r = 1, s = 1, t = 1, u = 2$ in Theorem 1 (c) then we have

$$\sum_{k=0}^n x^k W_{2k+1} = \frac{g_2(x)}{-4x^4 - 3x^3 + 5x^2 + 3x - 1}$$

where

$$g_2(x) = (x+1)x^{n+1}W_{2n+2} - (4x^3 + 3x^2 - 2x - 1)x^{n+1}W_{2n+1} + (x+1)x^{n+1}W_{2n} - 2(2x^2 + x - 1)x^{n+1}W_{2n-1} + x(2x^2 + x - 1)W_3 - (2x+2)x^2W_2 + (-2x^3 + 2x^2 + 3x - 1)W_1 - 2(x+1)x^2W_0.$$

For $x = 1$, the right hand side of the above sum formula is an indeterminate form. Now, we can use L'Hospital rule. Then we get (c) using

$$\sum_{k=0}^n W_{2k+1} = \frac{\frac{d}{dx}(g_2(x))}{\frac{d}{dx}(-4x^4 - 3x^3 + 5x^2 + 3x - 1)} \Big|_{x=1}.$$

□

Taking $W_n = J_n$ with $J_0 = 0, J_1 = 1, J_2 = 1, J_3 = 1$ in the last theorem, we have the following corollary which presents linear sum formula of fourth-order Jacobsthal numbers.

Corollary 1. For $n \geq 0$, fourth order Jacobsthal numbers have the following property:

(a) $\sum_{k=0}^n J_k = \frac{1}{4}(J_{n+3} - J_{n+1} + 2J_n).$

(b) $\sum_{k=0}^n J_{2k} = \frac{1}{12}((2n+7)J_{2n+2} - 2(2n+5)J_{2n+1} + (2n+13)J_{2n} - 2(2n+5)J_{2n-1} - 2).$

(c) $\sum_{k=0}^n J_{2k+1} = \frac{1}{12}(-(2n+3)J_{2n+2} + 4(n+5)J_{2n+1} - (2n+3)J_{2n} + 2(2n+7)J_{2n-1} + 2).$

From the last theorem, we have the following corollary which gives linear sum formula of fourth order Jacobsthal-Lucas numbers (take $W_n = j_n$ with $j_0 = 2, j_1 = 1, j_2 = 5, j_3 = 10$).

Corollary 2. For $n \geq 0$, fourth order Jacobsthal-Lucas numbers have the following property:

(a) $\sum_{k=0}^n j_k = \frac{1}{4}(j_{n+3} - j_{n+1} + 2j_n - 5).$

(b) $\sum_{k=0}^n j_{2k} = \frac{1}{12}((2n+7)j_{2n+2} - 2(2n+5)j_{2n+1} + (2n+13)j_{2n} - 2(2n+5)j_{2n-1} - 17).$

(c) $\sum_{k=0}^n j_{2k+1} = \frac{1}{12}(-(2n+3)j_{2n+2} + 4(n+5)j_{2n+1} - (2n+3)j_{2n} + 2(2n+7)j_{2n-1} - 1).$

Taking $W_n = K_n$ with $K_0 = 3, K_1 = 1, K_2 = 3, K_3 = 10$ in the last theorem, we have the following corollary which presents linear sum formula of modified fourth order Jacobsthal numbers.

Corollary 3. For $n \geq 0$, modified fourth order Jacobsthal numbers have the following property:

- (a) $\sum_{k=0}^n K_k = \frac{1}{4}(K_{n+3} - K_{n+1} + 2K_n - 3)$.
- (b) $\sum_{k=0}^n K_{2k} = \frac{1}{12}((2n+7)K_{2n+2} - 2(2n+5)K_{2n+1} + (2n+13)K_{2n} - 2(2n+5)K_{2n-1} + 1)$.
- (c) $\sum_{k=0}^n K_{2k+1} = \frac{1}{12}(-(2n+3)K_{2n+2} + 4(n+5)K_{2n+1} - (2n+3)K_{2n} + 2(2n+7)K_{2n-1} - 11)$.

From the last theorem, we have the following corollary which gives linear sum formula of fourth-order Jacobsthal Perrin numbers (take $W_n = Q_n$ with $Q_0 = 3, Q_1 = 0, Q_2 = 2, Q_3 = 8$).

Corollary 4. For $n \geq 0$, fourth-order Jacobsthal Perrin numbers have the following property:

- (a) $\sum_{k=0}^n Q_k = \frac{1}{4}(Q_{n+3} - Q_{n+1} + 2Q_n - 2)$.
- (b) $\sum_{k=0}^n Q_{2k} = \frac{1}{12}((2n+7)Q_{2n+2} - 2(2n+5)Q_{2n+1} + (2n+13)Q_{2n} - 2(2n+5)Q_{2n-1} - 2)$.
- (c) $\sum_{k=0}^n Q_{2k+1} = \frac{1}{12}(-(2n+3)Q_{2n+2} + 4(n+5)Q_{2n+1} - (2n+3)Q_{2n} + 2(2n+7)Q_{2n-1} - 6)$.

Taking $W_n = S_n$ with $S_0 = 0, S_1 = 1, S_2 = 1, S_3 = 2$ in the theorem, we have the following corollary which presents linear sum formula of adjusted fourth-order Jacobsthal numbers.

Corollary 5. For $n \geq 0$, adjusted fourth-order Jacobsthal numbers have the following property:

- (a) $\sum_{k=0}^n S_k = \frac{1}{4}(S_{n+3} - S_{n+1} + 2S_n - 1)$.
- (b) $\sum_{k=0}^n S_{2k} = \frac{1}{12}((2n+7)S_{2n+2} - 2(2n+5)S_{2n+1} + (2n+13)S_{2n} - 2(2n+5)S_{2n-1} + 3)$.
- (c) $\sum_{k=0}^n S_{2k+1} = \frac{1}{12}(-(2n+3)S_{2n+2} + 4(n+5)S_{2n+1} - (2n+3)S_{2n} + 2(2n+7)S_{2n-1} - 5)$.

From the last theorem, we have the following corollary which gives linear sum formula of modified fourth-order Jacobsthal-Lucas numbers (take $W_n = R_n$ with $R_0 = 4, R_1 = 1, R_2 = 3, R_3 = 7$).

Corollary 6. For $n \geq 0$, modified fourth-order Jacobsthal-Lucas numbers have the following property:

- (a) $\sum_{k=0}^n R_k = \frac{1}{4}(R_{n+3} - R_{n+1} + 2R_n + 2)$.
- (b) $\sum_{k=0}^n R_{2k} = \frac{1}{12}((2n+7)R_{2n+2} - 2(2n+5)R_{2n+1} + (2n+13)R_{2n} - 2(2n+5)R_{2n-1} - 20)$.
- (c) $\sum_{k=0}^n R_{2k+1} = \frac{1}{12}(-(2n+3)R_{2n+2} + 4(n+5)R_{2n+1} - (2n+3)R_{2n} + 2(2n+7)R_{2n-1} + 20)$.

3.2 The case $x = -1$

In this subsection we consider the special case $x = -1$.

In this section, we present the closed form solutions (identities) of the sums $\sum_{k=0}^n (-1)^k W_k$, $\sum_{k=0}^n (-1)^k W_{2k}$ and $\sum_{k=0}^n (-1)^k W_{2k+1}$ for the specific case of the sequence $\{W_n\}$.

Taking $x = -1, r = s = t = u = 1$ in Theorem 1 (a), (b) and (c), we obtain the following proposition.

Proposition 3. If $x = -1, r = s = t = u = 1$ then for $n \geq 0$ we have the following formulas:

- (a) $\sum_{k=0}^n (-1)^k W_k = (-1)^n (W_{n+3} - 2W_{n+2} + W_{n+1} - W_n) - W_3 + 2W_2 - W_1 + 2W_0$.
- (b) $\sum_{k=0}^n (-1)^k W_{2k} = (-1)^n (W_{2n+2} - W_{2n+1} - W_{2n}) - W_2 + W_1 + 2W_0$.
- (c) $\sum_{k=0}^n (-1)^k W_{2k+1} = (-1)^n (W_{2n} + W_{2n-1}) - W_3 + W_2 + 2W_1$.

From the above proposition, we have the following corollary which gives linear sum formulas of Tetranacci numbers (take $W_n = M_n$ with $M_0 = 0, M_1 = 1, M_2 = 1, M_3 = 2$).

Corollary 7. For $n \geq 0$, Tetranacci numbers have the following properties.

- (a) $\sum_{k=0}^n (-1)^k M_k = (-1)^n (M_{n+3} - 2M_{n+2} + M_{n+1} - M_n) - 1.$
 (b) $\sum_{k=0}^n (-1)^k M_{2k} = (-1)^n (M_{2n+2} - M_{2n+1} - M_{2n}).$
 (c) $\sum_{k=0}^n (-1)^k M_{2k+1} = (-1)^n (M_{2n} + M_{2n-1}) + 1.$

Taking $W_n = R_n$ with $R_0 = 4, R_1 = 1, R_2 = 3, R_3 = 7$ in the above proposition, we have the following corollary which presents linear sum formulas of Tetranacci-Lucas numbers.

Corollary 8. *For $n \geq 0$, Tetranacci-Lucas numbers have the following properties.*

- (a) $\sum_{k=0}^n (-1)^k R_k = (-1)^n (R_{n+3} - 2R_{n+2} + R_{n+1} - R_n) + 6.$
 (b) $\sum_{k=0}^n (-1)^k R_{2k} = (-1)^n (R_{2n+2} - R_{2n+1} - R_{2n}) + 6.$
 (c) $\sum_{k=0}^n (-1)^k R_{2k+1} = (-1)^n (R_{2n} + R_{2n-1}) - 2.$

Taking $x = -1, r = 2, s = t = u = 1$ in Theorem 1 (a), (b) and (c), we obtain the following proposition.

Proposition 4. *If $x = -1, r = 2, s = t = u = 1$ then for $n \geq 0$ we have the following formulas:*

- (a) $\sum_{k=0}^n (-1)^k W_k = \frac{1}{2}((-1)^n (W_{n+3} - 3W_{n+2} + 2W_{n+1} - W_n) - W_3 + 3W_2 - 2W_1 + 3W_0).$
 (b) $\sum_{k=0}^n (-1)^k W_{2k} = \frac{1}{2}((-1)^n (W_{2n+2} - W_{2n+1} - 2W_{2n} - W_{2n-1}) + W_3 - 3W_2 + 3W_0).$
 (c) $\sum_{k=0}^n (-1)^k W_{2k+1} = \frac{1}{2}((-1)^n (W_{2n+2} - W_{2n+1} + W_{2n-1}) - W_3 + W_2 + 4W_1 + W_0).$

From the last proposition, we have the following corollary which gives linear sum formulas of fourth-order Pell numbers (take $W_n = P_n$ with $P_0 = 0, P_1 = 1, P_2 = 2, P_3 = 5$).

Corollary 9. *For $n \geq 0$, fourth-order Pell numbers have the following properties:*

- (a) $\sum_{k=0}^n (-1)^k P_k = \frac{1}{2}((-1)^n (P_{n+3} - 3P_{n+2} + 2P_{n+1} - P_n) - 1).$
 (b) $\sum_{k=0}^n (-1)^k P_{2k} = \frac{1}{2}((-1)^n (P_{2n+2} - P_{2n+1} - 2P_{2n} - P_{2n-1}) - 1).$
 (c) $\sum_{k=0}^n (-1)^k P_{2k+1} = \frac{1}{2}((-1)^n (P_{2n+2} - P_{2n+1} + P_{2n-1}) + 1).$

Taking $W_n = Q_n$ with $Q_0 = 4, Q_1 = 2, Q_2 = 6, Q_3 = 17$ in the last proposition, we have the following corollary which presents linear sum formulas of fourth-order Pell-Lucas numbers.

Corollary 10. *For $n \geq 0$, fourth-order Pell-Lucas numbers have the following properties:*

- (a) $\sum_{k=0}^n (-1)^k Q_k = \frac{1}{2}((-1)^n (Q_{n+3} - 3Q_{n+2} + 2Q_{n+1} - Q_n) + 9).$
 (b) $\sum_{k=0}^n (-1)^k Q_{2k} = \frac{1}{2}((-1)^n (Q_{2n+2} - Q_{2n+1} - 2Q_{2n} - Q_{2n-1}) + 11).$
 (c) $\sum_{k=0}^n (-1)^k Q_{2k+1} = \frac{1}{2}((-1)^n (Q_{2n+2} - Q_{2n+1} + Q_{2n-1}) + 1).$

Observe that setting $x = -1, r = 1, s = 1, t = 1, u = 2$ (i.e. for the generalized fourth order Jacobsthal case) in Theorem 1 (a), (b) and (c), makes the right hand side of the sum formulas to be an indeterminate form. Application of L'Hospital rule however provides the evaluation of the sum formulas.

Theorem 5. *If $r = 1, s = 1, t = 1, u = 2$ then for $n \geq 0$ we have the following formulas:*

- (a) $\sum_{k=0}^n (-1)^k W_k = \frac{1}{6}((-1)^n (-(n+3)W_{n+3} + (2n+5)W_{n+2} - nW_{n+1} + 2(n+4)W_n) + 3W_3 - 5W_2 - 2W_0).$
 (b) $\sum_{k=0}^n (-1)^k W_{2k} = \frac{1}{10}((-1)^n (-(3n+5)W_{2n+2} + 2(n+2)W_{2n+1} + (7n+18)W_{2n} + 2(n+2)W_{2n-1}) - 2W_3 + 7W_2 - 2W_1 - 6W_0).$

$$(c) \sum_{k=0}^n (-1)^k W_{2k+1} = \frac{1}{10} (-(-1)^n ((n+1)W_{2n+2} - (4n+13)W_{2n+1} + (n+1)W_{2n} + 2(3n+5)W_{2n-1}) + 5W_3 - 8W_1 - 4W_2 - 4W_0).$$

Proof.

(a) We use Theorem 1 (a). If we set $r = 1, s = 1, t = 1, u = 2$ in Theorem 1 (a) then we have

$$\sum_{k=0}^n x^k W_k = \frac{g_3(x)}{2x^4 + x^3 + x^2 + x - 1}$$

where

$$g_3(x) = x^{n+3}W_{n+3} - (x-1)x^{n+2}W_{n+2} - (x^2+x-1)x^{n+1}W_{n+1} + 2x^{n+4}W_n - x^3W_3 + x^2(x-1)W_2 + x(x^2+x-1)W_1 + (x^3+x^2+x-1)W_0.$$

For $x = -1$, the right hand side of the above sum formula is an indeterminate form. Now, we can use L'Hospital rule. Then we get (b) using

$$\sum_{k=0}^n (-1)^k W_k = \frac{\frac{d}{dx}(g_3(x))}{\frac{d}{dx}(2x^4 + x^3 + x^2 + x - 1)} \Big|_{x=-1}.$$

(b) We use Theorem 1 (b). If we set $r = 1, s = 1, t = 1, u = 2$ in Theorem 1 (b) then we have

$$\sum_{k=0}^n x^k W_{2k} = \frac{g_4(x)}{-4x^4 - 3x^3 + 5x^2 + 3x - 1}$$

where

$$g_4(x) = -(2x^2+x-1)x^{n+1}W_{2n+2} + (2x+2)x^{n+2}W_{2n+1} - (4x^2+x-3)x^{n+2}W_{2n} + 2(x+1)x^{n+2}W_{2n-1} - (x+1)x^2W_3 + x(3x^2+2x-1)W_2 - (x+1)x^2W_1 + (-x^3+3x^2+3x-1)W_0.$$

For $x = -1$, the right hand side of the above sum formula is an indeterminate form. Now, we can use L'Hospital rule (twice). Then we get (b) using

$$\sum_{k=0}^n (-1)^k W_{2k} = \frac{\frac{d^2}{dx^2}(g_4(x))}{\frac{d^2}{dx^2}(-4x^4 - 3x^3 + 5x^2 + 3x - 1)} \Big|_{x=-1}.$$

(c) We use Theorem 1 (c). If we set $r = 1, s = 1, t = 1, u = 2$ in Theorem 1 (c) then we have

$$\sum_{k=0}^n x^k W_{2k+1} = \frac{g_5(x)}{-4x^4 - 3x^3 + 5x^2 + 3x - 1}$$

where

$$g_5(x) = (x+1)x^{n+1}W_{2n+2} - (4x^3+3x^2-2x-1)x^{n+1}W_{2n+1} + (x+1)x^{n+1}W_{2n} - 2(2x^2+x-1)x^{n+1}W_{2n-1} + x(2x^2+x-1)W_3 - (2x+2)x^2W_2 + (-2x^3+2x^2+3x-1)W_1 - 2(x+1)x^2W_0.$$

For $x = -1$, the right hand side of the above sum formula is an indeterminate form. Now, we can use L'Hospital rule (twice). Then we get (c) using

$$\sum_{k=0}^n (-1)^k W_{2k+1} = \frac{\frac{d^2}{dx^2}(g_5(x))}{\frac{d^2}{dx^2}(-4x^4 - 3x^3 + 5x^2 + 3x - 1)} \Big|_{x=-1}.$$

□

Taking $W_n = J_n$ with $J_0 = 0, J_1 = 1, J_2 = 1, J_3 = 1$ in the last theorem, we have the following corollary which presents linear sum formula of fourth-order Jacobsthal numbers.

Corollary 11. *For $n \geq 0$, fourth order Jacobsthal numbers have the following property:*

- (a) $\sum_{k=0}^n (-1)^k J_k = \frac{1}{6}((-1)^n (-(n+3)J_{n+3} + (2n+5)J_{n+2} - nJ_{n+1} + 2(n+4)J_n) - 2)$.
- (b) $\sum_{k=0}^n (-1)^k J_{2k} = \frac{1}{10}((-1)^n (-(3n+5)J_{2n+2} + 2(n+2)J_{2n+1} + (7n+18)J_{2n} + 2(n+2)J_{2n-1}) + 3)$.
- (c) $\sum_{k=0}^n (-1)^k J_{2k+1} = \frac{1}{10}(-(-1)^n ((n+1)J_{2n+2} - (4n+13)J_{2n+1} + (n+1)J_{2n} + 2(3n+5)J_{2n-1}) - 7)$.

From the last theorem, we have the following corollary which gives linear sum formula of fourth order Jacobsthal-Lucas numbers (take $W_n = j_n$ with $j_0 = 2, j_1 = 1, j_2 = 5, j_3 = 10$).

Corollary 12. *For $n \geq 0$, fourth order Jacobsthal-Lucas numbers have the following property:*

- (a) $\sum_{k=0}^n (-1)^k j_k = \frac{1}{6}((-1)^n (-(n+3)j_{n+3} + (2n+5)j_{n+2} - nj_{n+1} + 2(n+4)j_n) + 1)$.
- (b) $\sum_{k=0}^n (-1)^k j_{2k} = \frac{1}{10}((-1)^n (-(3n+5)j_{2n+2} + 2(n+2)j_{2n+1} + (7n+18)j_{2n} + 2(n+2)j_{2n-1}) + 1)$.
- (c) $\sum_{k=0}^n (-1)^k j_{2k+1} = \frac{1}{10}(-(-1)^n ((n+1)j_{2n+2} - (4n+13)j_{2n+1} + (n+1)j_{2n} + 2(3n+5)j_{2n-1}) + 14)$.

Taking $W_n = K_n$ with $K_0 = 3, K_1 = 1, K_2 = 3, K_3 = 10$ in the last theorem, we have the following corollary which presents linear sum formula of modified fourth order Jacobsthal numbers.

Corollary 13. *For $n \geq 0$, modified fourth order Jacobsthal numbers have the following property:*

- (a) $\sum_{k=0}^n (-1)^k K_k = \frac{1}{6}((-1)^n (-(n+3)K_{n+3} + (2n+5)K_{n+2} - nK_{n+1} + 2(n+4)K_n) + 9)$.
- (b) $\sum_{k=0}^n (-1)^k K_{2k} = \frac{1}{10}((-1)^n (-(3n+5)K_{2n+2} + 2(n+2)K_{2n+1} + (7n+18)K_{2n} + 2(n+2)K_{2n-1}) - 19)$.
- (c) $\sum_{k=0}^n (-1)^k K_{2k+1} = \frac{1}{10}(-(-1)^n ((n+1)K_{2n+2} - (4n+13)K_{2n+1} + (n+1)K_{2n} + 2(3n+5)K_{2n-1}) + 18)$.

From the last theorem, we have the following corollary which gives linear sum formula of fourth-order Jacobsthal Perrin numbers (take $W_n = Q_n$ with $Q_0 = 3, Q_1 = 0, Q_2 = 2, Q_3 = 8$).

Corollary 14. *For $n \geq 0$, fourth-order Jacobsthal Perrin numbers have the following property:*

- (a) $\sum_{k=0}^n (-1)^k Q_k = \frac{1}{6}((-1)^n (-(n+3)Q_{n+3} + (2n+5)Q_{n+2} - nQ_{n+1} + 2(n+4)Q_n) + 8)$.
- (b) $\sum_{k=0}^n (-1)^k Q_{2k} = \frac{1}{10}((-1)^n (-(3n+5)Q_{2n+2} + 2(n+2)Q_{2n+1} + (7n+18)Q_{2n} + 2(n+2)Q_{2n-1}) - 20)$.
- (c) $\sum_{k=0}^n (-1)^k Q_{2k+1} = \frac{1}{10}(-(-1)^n ((n+1)Q_{2n+2} - (4n+13)Q_{2n+1} + (n+1)Q_{2n} + 2(3n+5)Q_{2n-1}) + 20)$.

Taking $W_n = S_n$ with $S_0 = 0, S_1 = 1, S_2 = 1, S_3 = 2$ in the theorem, we have the following corollary which presents linear sum formula of adjusted fourth-order Jacobsthal numbers.

Corollary 15. *For $n \geq 0$, adjusted fourth-order Jacobsthal numbers have the following property:*

- (a) $\sum_{k=0}^n (-1)^k S_k = \frac{1}{6}((-1)^n (-(n+3)S_{n+3} + (2n+5)S_{n+2} - nS_{n+1} + 2(n+4)S_n) + 1)$.
- (b) $\sum_{k=0}^n (-1)^k S_{2k} = \frac{1}{10}((-1)^n (-(3n+5)S_{2n+2} + 2(n+2)S_{2n+1} + (7n+18)S_{2n} + 2(n+2)S_{2n-1}) + 1)$.
- (c) $\sum_{k=0}^n (-1)^k S_{2k+1} = \frac{1}{10}(-(-1)^n ((n+1)S_{2n+2} - (4n+13)S_{2n+1} + (n+1)S_{2n} + 2(3n+5)S_{2n-1}) - 2)$.

From the last theorem, we have the following corollary which gives linear sum formula of modified fourth-order Jacobsthal-Lucas numbers (take $W_n = R_n$ with $R_0 = 4, R_1 = 1, R_2 = 3, R_3 = 7$).

Corollary 16. *For $n \geq 0$, modified fourth-order Jacobsthal-Lucas numbers have the following property:*

- (a) $\sum_{k=0}^n (-1)^k R_k = \frac{1}{6}((-1)^n (-(n+3)R_{n+3} + (2n+5)R_{n+2} - nR_{n+1} + 2(n+4)R_n) - 2)$.
- (b) $\sum_{k=0}^n (-1)^k R_{2k} = \frac{1}{10}((-1)^n (-(3n+5)R_{2n+2} + 2(n+2)R_{2n+1} + (7n+18)R_{2n} + 2(n+2)R_{2n-1}) - 19)$.
- (c) $\sum_{k=0}^n (-1)^k R_{2k+1} = \frac{1}{10}(-(-1)^n ((n+1)R_{2n+2} - (4n+13)R_{2n+1} + (n+1)R_{2n} + 2(3n+5)R_{2n-1}) - 1)$.

Taking $x = -1, r = 2, s = 3, t = 5, u = 7$ in Theorem 1 (a), (b) and (c), we obtain the following proposition.

Proposition 6. *If $r = 2, s = 3, t = 5, u = 7$ then for $n \geq 0$ we have the following formulas:*

- (a) $\sum_{k=0}^n (-1)^k W_k = \frac{1}{2}((-1)^n (-W_{n+3} + 3W_{n+2} + 7W_n) + W_3 - 3W_2 - 5W_0)$.
- (b) $\sum_{k=0}^n (-1)^k W_{2k} = \frac{1}{6}((-1)^n (-W_{2n+2} + W_{2n+1} + 12W_{2n} + 7W_{2n-1}) - W_3 + 3W_2 + 2W_1 - W_0)$.
- (c) $\sum_{k=0}^n (-1)^k W_{2k+1} = \frac{1}{6}((-1)^n (9W_{2n+1} - W_{2n+2} + 2W_{2n} - 7W_{2n-1}) + W_3 - W_2 - 6W_1 - 7W_0)$.

From the last proposition, we have the following corollary which gives linear sum formulas of 4-primes numbers (take $W_n = G_n$ with $G_0 = 0, G_1 = 0, G_2 = 1, G_3 = 2$).

Corollary 17. *For $n \geq 0$, 4-primes numbers have the following properties:*

- (a) $\sum_{k=0}^n (-1)^k G_k = \frac{1}{2}((-1)^n (-G_{n+3} + 3G_{n+2} + 7G_n) - 1)$.
- (b) $\sum_{k=0}^n (-1)^k G_{2k} = \frac{1}{6}((-1)^n (-G_{2n+2} + G_{2n+1} + 12G_{2n} + 7G_{2n-1}) + 1)$.
- (c) $\sum_{k=0}^n (-1)^k G_{2k+1} = \frac{1}{6}((-1)^n (9G_{2n+1} - G_{2n+2} + 2G_{2n} - 7G_{2n-1}) + 1)$.

Taking $W_n = H_n$ with $H_0 = 4, H_1 = 2, H_2 = 10, H_3 = 41$ in the last proposition, we have the following corollary which presents linear sum formulas of Lucas 4-primes numbers.

Corollary 18. *For $n \geq 0$, Lucas 4-primes numbers have the following properties:*

- (a) $\sum_{k=0}^n (-1)^k H_k = \frac{1}{2}((-1)^n (-H_{n+3} + 3H_{n+2} + 7H_n) - 9)$.
- (b) $\sum_{k=0}^n (-1)^k H_{2k} = \frac{1}{6}((-1)^n (-H_{2n+2} + H_{2n+1} + 12H_{2n} + 7H_{2n-1}) - 11)$.
- (c) $\sum_{k=0}^n (-1)^k H_{2k+1} = \frac{1}{6}((-1)^n (9H_{2n+1} - H_{2n+2} + 2H_{2n} - 7H_{2n-1}) - 9)$.

From the last proposition, we have the following corollary which gives linear sum formulas of modified 4-primes numbers (take $W_n = E_n$ with $E_0 = 0, E_1 = 0, E_2 = 1, E_3 = 1$).

Corollary 19. *For $n \geq 0$, modified 4-primes numbers have the following properties:*

- (a) $\sum_{k=0}^n (-1)^k E_k = \frac{1}{2}((-1)^n (-E_{n+3} + 3E_{n+2} + 7E_n) - 2)$.
- (b) $\sum_{k=0}^n (-1)^k E_{2k} = \frac{1}{6}((-1)^n (-E_{2n+2} + E_{2n+1} + 12E_{2n} + 7E_{2n-1}) + 2)$.
- (c) $\sum_{k=0}^n (-1)^k E_{2k+1} = \frac{1}{6}((-1)^n (9E_{2n+1} - E_{2n+2} + 2E_{2n} - 7E_{2n-1}))$.

3.3 The case $x = i$

In this subsection we consider the special case $x = i$. Taking $x = i, r = s = t = u = 1$ in Theorem 1 (a), (b) and (c), we obtain the following proposition.

Proposition 7. *If $x = i, r = s = t = u = 1$ then for $n \geq 0$ we have the following formulas:*

- (a) $\sum_{k=0}^n i^k W_k = i^n (iW_{n+3} + (1-i)W_{n+2} - (1+2i)W_{n+1} - W_n) - iW_3 - (1-i)W_2 + (1+2i)W_1 + 2W_0.$
- (b) $\sum_{k=0}^n i^k W_{2k} = \frac{(5+4i)}{41} (i^n (- (1+2i)W_{2n+2} + (2+i)W_{2n+1} + 3W_{2n} + (1+i)W_{2n-1}) - (1+i)W_3 + (2+3i)W_2 - W_1 + (3-3i)W_0).$
- (c) $\sum_{k=0}^n i^k W_{2k+1} = \frac{(5+4i)}{41} (i^n ((1-i)W_{2n+2} + (2-2i)W_{2n+1} - iW_{2n} - (1+2i)W_{2n-1}) + (1+2i)W_3 - (2+i)W_2 + (2-4i)W_1 - (1+i)W_0).$

From the above proposition, we have the following corollary which gives linear sum formulas of Tetranacci numbers (take $W_n = M_n$ with $M_0 = 0, M_1 = 1, M_2 = 1, M_3 = 2$).

Corollary 20. *For $n \geq 0$, Tetranacci numbers have the following properties.*

- (a) $\sum_{k=0}^n i^k M_k = i^n (iM_{n+3} + (1-i)M_{n+2} - (1+2i)M_{n+1} - M_n) + i.$
- (b) $\sum_{k=0}^n i^k M_{2k} = \frac{(5+4i)}{41} (i^n (- (1+2i)M_{2n+2} + (2+i)M_{2n+1} + 3M_{2n} + (1+i)M_{2n-1}) + (-1+i)).$
- (c) $\sum_{k=0}^n i^k M_{2k+1} = \frac{(5+4i)}{41} (i^n ((1-i)M_{2n+2} + (2-2i)M_{2n+1} - iM_{2n} - (1+2i)M_{2n-1}) + (2-i)).$

Taking $W_n = R_n$ with $R_0 = 4, R_1 = 1, R_2 = 3, R_3 = 7$ in the above proposition, we have the following corollary which presents linear sum formulas of Tetranacci-Lucas numbers.

Corollary 21. *For $n \geq 0$, Tetranacci-Lucas numbers have the following properties.*

- (a) $\sum_{k=0}^n i^k R_k = i^n (iR_{n+3} + (1-i)R_{n+2} - (1+2i)R_{n+1} - R_n) + (6-2i).$
- (b) $\sum_{k=0}^n i^k R_{2k} = \frac{(5+4i)}{41} (i^n (- (1+2i)R_{2n+2} + (2+i)R_{2n+1} + 3R_{2n} + (1+i)R_{2n-1}) + 10(1-i)).$
- (c) $\sum_{k=0}^n i^k R_{2k+1} = \frac{(5+4i)}{41} (i^n ((1-i)R_{2n+2} + (2-2i)R_{2n+1} - iR_{2n} - (1+2i)R_{2n-1}) + (-1+3i)).$

Corresponding sums of the other fourth order generalized Tetranacci numbers can be calculated similarly.

4 Linear Sum Formulas of Generalized Tetranacci Numbers with Negative Subscripts

The following theorem present some linear summing formulas of generalized Tetranacci numbers with negative subscripts.

Theorem 8. *Let x be a real or complex numbers. For $n \geq 1$ we have the following formulas:*

- (a) *If $rx^3 + sx^2 + tx + u - x^4 \neq 0$, then*

$$\sum_{k=1}^n x^k W_{-k} = \frac{\Theta_4(x)}{rx^3 + sx^2 + tx + u - x^4}$$

- (b) *If $2sx^3 + t^2x + 2ux^2 + r^2x^3 - s^2x^2 - u^2 - x^4 + 2rtx^2 - 2sux \neq 0$ then*

$$\sum_{k=1}^n x^k W_{-2k} = \frac{x\Theta_5(x)}{2sx^3 + t^2x + 2ux^2 + r^2x^3 - s^2x^2 - u^2 - x^4 + 2rtx^2 - 2sux}$$

(c) If $2sx^3 + t^2x + 2ux^2 + r^2x^3 - s^2x^2 - u^2 - x^4 + 2rtx^2 - 2sux \neq 0$ then

$$\sum_{k=1}^n x^k W_{-2k+1} = \frac{x\Theta_6(x)}{2sx^3 + t^2x + 2ux^2 + r^2x^3 - s^2x^2 - u^2 - x^4 + 2rtx^2 - 2sux}$$

where

$$\Theta_4(x) = -x^{n+1}W_{-n+3} + x^{n+1}(r-x)W_{-n+2} + x^{n+1}(s+rx-x^2)W_{-n+1} + x^{n+1}(t+rx^2+sx-x^3)W_{-n} + xW_3 - x(r-x)W_2 + x(-s-rx+x^2)W_1 + x(-t-rx^2-sx+x^3)W_0,$$

$$\Theta_5(x) = x^n(u+sx-x^2)W_{-2n+2} - x^n(ru+tx+rsx)W_{-2n+1} + x^n(2sx^2-s^2x+r^2x^2-su+ux-x^3+rtx)W_{-2n} - ux^n(t+rx)W_{-2n-1} + (t+rx)W_3 + (-u-r^2x-rt-sx+x^2)W_2 + (ru-st+tx)W_1 - (2sx^2-s^2x+r^2x^2-su+ux+t^2-x^3+2rtx)W_0,$$

$$\Theta_6(x) = -x^{n+1}(t+rx)W_{-2n+2} + x^{n+1}(u+r^2x+rt+sx-x^2)W_{-2n+1} - x^{n+1}(ru-st+tx)W_{-2n} + ux^n(u+sx-x^2)W_{-2n-1} + (-u-sx+x^2)W_3 + (ru+tx+rsx)W_2 + (-2sx^2+s^2x-r^2x^2+su-ux+x^3-rtx)W_1 + u(t+rx)W_0.$$

Proof.

(a) Using the recurrence relation

$$W_{-n+4} = rW_{-n+3} + sW_{-n+2} + tW_{-n+1} + uW_{-n}$$

i.e.

$$uW_{-n} = W_{-n+4} - rW_{-n+3} - sW_{-n+2} - tW_{-n+1}$$

we obtain

$$\begin{aligned} ux^n W_{-n} &= x^n W_{-n+4} - rx^n W_{-n+3} - sx^n W_{-n+2} - tx^n W_{-n+1} \\ ux^{n-1} W_{-n+1} &= x^{n-1} W_{-n+5} - rx^{n-1} W_{-n+4} - sx^{n-1} W_{-n+3} - tx^{n-1} W_{-n+2} \\ ux^{n-2} W_{-n+2} &= x^{n-2} W_{-n+6} - rx^{n-2} W_{-n+5} - sx^{n-2} W_{-n+4} - tx^{n-2} W_{-n+3} \\ &\vdots \\ ux^4 W_{-4} &= x^4 W_0 - rx^4 W_{-1} - sx^4 W_{-2} - tx^4 W_{-3} \\ ux^3 W_{-3} &= x^3 W_1 - rx^3 W_0 - sx^3 W_{-1} - tx^3 W_{-2} \\ ux^2 W_{-2} &= x^2 W_2 - rx^2 W_1 - sx^2 W_0 - tx^2 W_{-1} \\ ux^1 W_{-1} &= x^1 W_3 - rx^1 W_2 - sx^1 W_1 - tx^1 W_0 \end{aligned}$$

If we add the above equations side by side, we obtain

$$\begin{aligned} u\left(\sum_{k=1}^n x^k W_{-k}\right) &= (-x^{n+1}W_{-n+3} - x^{n+2}W_{-n+2} - x^{n+3}W_{-n+1} - x^{n+4}W_{-n} \\ &\quad + x^1W_3 + x^2W_2 + x^3W_1 + x^4W_0 + x^4 \sum_{k=1}^n x^k W_{-k}) \\ &\quad - r(-x^{n+1}W_{-n+2} - x^{n+2}W_{-n+1} - x^{n+3}W_{-n} + x^1W_2 + x^2W_1 + x^3W_0 + x^3 \sum_{k=1}^n x^k W_{-k}) \\ &\quad - s(-x^{n+1}W_{-n+1} - x^{n+2}W_{-n} + x^1W_1 + x^2W_0 + x^2 \sum_{k=1}^n x^k W_{-k}) \\ &\quad - t(-x^{n+1}W_{-n} + x^1W_0 + x^1 \sum_{k=1}^n x^k W_{-k}) \end{aligned}$$

From the last equation we get (a).

(b) and (c) Using the recurrence relation

$$W_{-n+4} = rW_{-n+3} + sW_{-n+2} + tW_{-n+1} + uW_{-n}$$

i.e.

$$tW_{-n+1} = W_{-n+4} - rW_{-n+3} - sW_{-n+2} - uW_{-n}$$

we obtain

$$\begin{aligned} tx^n W_{-2n+1} &= x^n W_{-2n+4} - rx^n W_{-2n+3} - sx^n W_{-2n+2} - ux^n W_{-2n} \\ tx^{n-1} W_{-2n+3} &= x^{n-1} W_{-2n+6} - rx^{n-1} W_{-2n+5} - sx^{n-1} W_{-2n+4} - ux^{n-1} W_{-2n+2} \\ tx^{n-2} W_{-2n+5} &= x^{n-2} W_{-2n+8} - rx^{n-2} W_{-2n+7} - sx^{n-2} W_{-2n+6} - ux^{n-2} W_{-2n+4} \\ tx^{n-3} W_{-2n+7} &= x^{n-3} W_{-2n+10} - rx^{n-3} W_{-2n+9} - sx^{n-3} W_{-2n+8} - ux^{n-3} W_{-2n+6} \\ &\vdots \\ tx^3 W_{-5} &= x^3 W_{-2} - rx^3 W_{-3} - sx^3 W_{-4} - ux^3 W_{-6} \\ tx^2 W_{-3} &= x^2 W_0 - rx^2 W_{-1} - sx^2 W_{-2} - ux^2 W_{-4} \\ tx^1 W_{-1} &= x^1 W_2 - rx^1 W_1 - sx^1 W_0 - ux^1 W_{-2}. \end{aligned}$$

If we add the above equations side by side, we get

$$\begin{aligned} t \sum_{k=1}^n x^k W_{-2k+1} &= (-x^{n+1} W_{-2n+2} - x^{n+2} W_{-2n} + x^2 W_0 + x^1 W_2 + \sum_{k=1}^n x^{k+2} W_{-2k}) \\ &\quad - r(-x^{n+1} W_{-2n+1} + x^1 W_1 + \sum_{k=1}^n x^{k+1} W_{-2k+1}) \\ &\quad - s(-x^{n+1} W_{-2n} + x^1 W_0 + \sum_{k=1}^n x^{k+1} W_{-2k}) - u(\sum_{k=1}^n x^k W_{-2k}). \end{aligned} \quad (4)$$

Similarly, using the recurrence relation

$$W_{-n+4} = rW_{-n+3} + sW_{-n+2} + tW_{-n+1} + uW_{-n}$$

i.e.

$$tW_{-n} = W_{-n+3} - rW_{-n+2} - sW_{-n+1} - uW_{-n-1}$$

we obtain

$$\begin{aligned} tx^n W_{-2n} &= x^n W_{-2n+3} - rx^n W_{-2n+2} - sx^n W_{-2n+1} - ux^n W_{-2n-1} \\ tx^{n-1} W_{-2n+2} &= x^{n-1} W_{-2n+5} - rx^{n-1} W_{-2n+4} - sx^{n-1} W_{-2n+3} - ux^{n-1} W_{-2n+1} \\ tx^{n-2} W_{-2n+4} &= x^{n-2} W_{-2n+7} - rx^{n-2} W_{-2n+6} - sx^{n-2} W_{-2n+5} - ux^{n-2} W_{-2n+3} \\ tx^{n-3} W_{-2n+6} &= x^{n-3} W_{-2n+9} - rx^{n-3} W_{-2n+8} - sx^{n-3} W_{-2n+7} - ux^{n-3} W_{-2n+5} \\ &\vdots \\ tx^4 W_{-8} &= x^4 W_{-5} - rx^4 W_{-6} - sx^4 W_{-7} - ux^4 W_{-9} \\ tx^3 W_{-6} &= x^3 W_{-3} - rx^3 W_{-4} - sx^3 W_{-5} - ux^3 W_{-7} \\ tx^2 W_{-4} &= x^2 W_{-1} - rx^2 W_{-2} - sx^2 W_{-3} - ux^2 W_{-5} \\ tx^1 W_{-2} &= x^1 W_1 - rx^1 W_0 - sx^1 W_{-1} - ux^1 W_{-3}. \end{aligned}$$

If we add the above equations side by side, we get

$$\begin{aligned} t \sum_{k=1}^n x^k W_{-2k} &= (-x^{n+1} W_{-2n+1} + x^1 W_1 + \sum_{k=1}^n x^{k+1} W_{-2k+1}) \\ &\quad -r(-x^{n+1} W_{-2n} + x^1 W_0 + \sum_{k=1}^n x^{k+1} W_{-2k}) \\ &\quad -s(\sum_{k=1}^n x^k W_{-2k+1}) - u(x^n W_{-2n-1} - x^0 W_{-1} + \sum_{k=1}^n x^{k-1} W_{-2k+1}). \end{aligned}$$

Since

$$W_{-1} = -\frac{t}{u} W_0 - \frac{s}{u} W_1 - \frac{r}{u} W_2 + \frac{1}{u} W_3$$

it follows that

$$\begin{aligned} t \sum_{k=1}^n x^k W_{-2k} &= (-x^{n+1} W_{-2n+1} + x^1 W_1 + \sum_{k=1}^n x^{k+1} W_{-2k+1}) \\ &\quad -r(-x^{n+1} W_{-2n} + x^1 W_0 + \sum_{k=1}^n x^{k+1} W_{-2k}) - s(\sum_{k=1}^n x^k W_{-2k+1}) \\ &\quad -u(x^n W_{-2n-1} - x^0 (-\frac{t}{u} W_0 - \frac{s}{u} W_1 - \frac{r}{u} W_2 + \frac{1}{u} W_3) + \sum_{k=1}^n x^{k-1} W_{-2k+1}). \end{aligned} \quad (5)$$

Then, solving system (4)-(5) the required results of (b) and (c) follow. \square

5 Specific Cases

In this section, for the specific cases of x , we present the closed form solutions (identities) of the sums $\sum_{k=1}^n x^k W_{-k}$, $\sum_{k=1}^n x^k W_{-2k}$ and $\sum_{k=0}^n x^k W_{-2k+1}$ for the specific case of sequence $\{W_n\}$.

5.1 The case $x = 1$

In this subsection we consider the special case $x = 1$.

The case $x = 1$ of Theorem 8 is given in [20]. For the generalized 4-primes sequence case ($x = 1, r = 2, s = 3, t = 5, u = 7$), see [29].

We only consider the cases $x = 1, r = 1, s = 1, t = 1, u = 2$ (which is not considered in [20]).

Observe that setting $x = 1, r = 1, s = 1, t = 1, u = 2$ (i.e. for the generalized fourth order Jacobsthal case) in Theorem 8 (a),(b),(c) makes the right hand side of the sum formulas to be an indeterminate form. Application of L'Hospital rule however provides the evaluation of the sum formulas.

Taking $r = 1, s = 1, t = 1, u = 2$ in Theorem 8, we obtain the following theorem.

Theorem 9. *If $r = 1, s = 1, t = 1, u = 2$ then for $n \geq 1$ we have the following formulas:*

- (a) $\sum_{k=1}^n W_{-k} = \frac{1}{4}(-W_{-n+3} + W_{-n+1} + 2W_{-n} + W_3 - W_1 - 2W_0)$.
- (b) $\sum_{k=1}^n W_{-2k} = \frac{1}{12}((2n+1)W_{-2n+2} - 2(2n+3)W_{-2n+1} + (2n+7)W_{-2n} - 2(2n+3)W_{-2n-1} + 3W_3 - 4W_2 + 3W_1 - 10W_0)$.
- (c) $\sum_{k=1}^n W_{-2k+1} = \frac{1}{12}(-(2n+5)W_{-2n+2} + 4(n+2)W_{-2n+1} - (2n+5)W_{-2n} + 2(2n+1)W_{-2n-1} - W_3 + 6W_2 - 7W_1 + 6W_0)$.

Proof.

- (a) We use Theorem 8 (a). If we set $r = 1, s = 1, t = 1, u = 2$ in Theorem 8 (a) we get (a).
 (b) We use Theorem 8 (b). If we set $r = 1, s = 1, t = 1, u = 2$ in Theorem 8 (b) then we have

$$\sum_{k=1}^n x^k W_{-2k} = \frac{g_6(x)}{-x^4 + 3x^3 + 5x^2 - 3x - 4}$$

where

$$g_6(x) = (-x^2 + x + 2)x^{n+1}W_{-2n+2} - (2x + 2)x^{n+1}W_{-2n+1} + (-x^3 + 3x^2 + 2x - 2)x^{n+1}W_{-2n} - 2(x + 1)x^{n+1}W_{-2n-1} + x(x + 1)W_3 - x(-x^2 + 2x + 3)W_2 + x(x + 1)W_1 - x(-x^3 + 3x^2 + 3x - 1)W_0.$$

For $x = 1$, the right hand side of the above sum formula is an indeterminate form. Now, we can use L'Hospital rule. Then we get (b) using

$$\sum_{k=1}^n W_{-2k} = \frac{\frac{d}{dx}(g_6(x))}{\frac{d}{dx}(-x^4 + 3x^3 + 5x^2 - 3x - 4)} \Big|_{x=1}.$$

- (c) We use Theorem 8 (c). If we set $r = 1, s = 1, t = 1, u = 2$ in Theorem 8 (c) then we have

$$\sum_{k=1}^n x^k W_{-2k+1} = \frac{g_7(x)}{-x^4 + 3x^3 + 5x^2 - 3x - 4}$$

where

$$g_7(x) = -(x + 1)x^{n+2}W_{-2n+2} + (-x^2 + 2x + 3)x^{n+2}W_{-2n+1} - (x + 1)x^{n+2}W_{-2n} + 2(-x^2 + x + 2)x^{n+1}W_{-2n-1} - x(-x^2 + x + 2)W_3 + x(2x + 2)W_2 - x(-x^3 + 3x^2 + 2x - 2)W_1 + 2xW_0(x + 1).$$

For $x = 1$, the right hand side of the above sum formula is an indeterminate form. Now, we can use L'Hospital rule. Then we get (c) using

$$\sum_{k=1}^n W_{-2k+1} = \frac{\frac{d}{dx}(g_7(x))}{\frac{d}{dx}(-x^4 + 3x^3 + 5x^2 - 3x - 4)} \Big|_{x=1}.$$

□

Taking $W_n = J_n$ with $J_0 = 0, J_1 = 1, J_2 = 1, J_3 = 1$ in the last theorem, we have the following corollary which presents linear sum formula of fourth-order Jacobsthal numbers.

Corollary 22. For $n \geq 1$, fourth order Jacobsthal numbers have the following property

- (a) $\sum_{k=1}^n J_{-k} = \frac{1}{4}(-J_{-n+3} + J_{-n+1} + 2J_{-n})$.
 (b) $\sum_{k=1}^n J_{-2k} = \frac{1}{12}((2n + 1)J_{-2n+2} - 2(2n + 3)J_{-2n+1} + (2n + 7)J_{-2n} - 2(2n + 3)J_{-2n-1} + 2)$.
 (c) $\sum_{k=1}^n J_{-2k+1} = \frac{1}{12}(-(2n + 5)J_{-2n+2} + 4(n + 2)J_{-2n+1} - (2n + 5)J_{-2n} + 2(2n + 1)J_{-2n-1} - 2)$.

From the last theorem, we have the following corollary which gives linear sum formulas of fourth order Jacobsthal-Lucas numbers (take $W_n = j_n$ with $j_0 = 2, j_1 = 1, j_2 = 5, j_3 = 10$).

Corollary 23. For $n \geq 1$, fourth order Jacobsthal-Lucas numbers have the following property

- (a) $\sum_{k=1}^n j_{-k} = \frac{1}{4}(-j_{-n+3} + j_{-n+1} + 2j_{-n} + 5)$.
 (b) $\sum_{k=1}^n j_{-2k} = \frac{1}{12}((2n + 1)j_{-2n+2} - 2(2n + 3)j_{-2n+1} + (2n + 7)j_{-2n} - 2(2n + 3)j_{-2n-1} - 7)$.
 (c) $\sum_{k=1}^n j_{-2k+1} = \frac{1}{12}(-(2n + 5)j_{-2n+2} + 4(n + 2)j_{-2n+1} - (2n + 5)j_{-2n} + 2(2n + 1)j_{-2n-1} + 25)$.

Taking $W_n = K_n$ with $K_0 = 3, K_1 = 1, K_2 = 3, K_3 = 10$ in the last theorem, we have the following corollary which presents linear sum formula of modified fourth order Jacobsthal numbers.

Corollary 24. *For $n \geq 1$, modified fourth order Jacobsthal numbers have the following property:*

- (a) $\sum_{k=1}^n K_{-k} = \frac{1}{4}(-K_{-n+3} + K_{-n+1} + 2K_{-n} + 3)$.
- (b) $\sum_{k=1}^n K_{-2k} = \frac{1}{12}((2n+1)K_{-2n+2} - 2(2n+3)K_{-2n+1} + (2n+7)K_{-2n} - 2(2n+3)K_{-2n-1} - 9)$.
- (c) $\sum_{k=1}^n K_{-2k+1} = \frac{1}{12}(-(2n+5)K_{-2n+2} + 4(n+2)K_{-2n+1} - (2n+5)K_{-2n} + 2(2n+1)K_{-2n-1} + 19)$.

From the last theorem, we have the following corollary which gives linear sum formula of fourth-order Jacobsthal Perrin numbers (take $W_n = Q_n$ with $Q_0 = 3, Q_1 = 0, Q_2 = 2, Q_3 = 8$).

Corollary 25. *For $n \geq 1$, fourth-order Jacobsthal Perrin numbers have the following property:*

- (a) $\sum_{k=1}^n Q_{-k} = \frac{1}{4}(-Q_{-n+3} + Q_{-n+1} + 2Q_{-n} + 2)$.
- (b) $\sum_{k=1}^n Q_{-2k} = \frac{1}{12}((2n+1)Q_{-2n+2} - 2(2n+3)Q_{-2n+1} + (2n+7)Q_{-2n} - 2(2n+3)Q_{-2n-1} - 14)$.
- (c) $\sum_{k=1}^n Q_{-2k+1} = \frac{1}{12}(-(2n+5)Q_{-2n+2} + 4(n+2)Q_{-2n+1} - (2n+5)Q_{-2n} + 2(2n+1)Q_{-2n-1} + 22)$.

Taking $W_n = S_n$ with $S_0 = 0, S_1 = 1, S_2 = 1, S_3 = 2$ in the last theorem, we have the following corollary which presents linear sum formula of adjusted fourth-order Jacobsthal numbers.

Corollary 26. *For $n \geq 1$, adjusted fourth-order Jacobsthal numbers have the following property:*

- (a) $\sum_{k=1}^n S_{-k} = \frac{1}{4}(-S_{-n+3} + S_{-n+1} + 2S_{-n} + 1)$.
- (b) $\sum_{k=1}^n S_{-2k} = \frac{1}{12}((2n+1)S_{-2n+2} - 2(2n+3)S_{-2n+1} + (2n+7)S_{-2n} - 2(2n+3)S_{-2n-1} - 1)$.
- (c) $\sum_{k=1}^n S_{-2k+1} = \frac{1}{12}(-(2n+5)S_{-2n+2} + 4(n+2)S_{-2n+1} - (2n+5)S_{-2n} + 2(2n+1)S_{-2n-1} - 3)$.

From the last theorem, we have the following corollary which gives linear sum formula of modified fourth-order Jacobsthal-Lucas numbers (take $W_n = R_n$ with $R_0 = 4, R_1 = 1, R_2 = 3, R_3 = 7$).

Corollary 27. *For $n \geq 1$, modified fourth-order Jacobsthal-Lucas numbers have the following property:*

- (a) $\sum_{k=1}^n R_{-k} = \frac{1}{4}(-R_{-n+3} + R_{-n+1} + 2R_{-n} - 2)$.
- (b) $\sum_{k=1}^n R_{-2k} = \frac{1}{12}((2n+1)R_{-2n+2} - 2(2n+3)R_{-2n+1} + (2n+7)R_{-2n} - 2(2n+3)R_{-2n-1} - 28)$.
- (c) $\sum_{k=1}^n R_{-2k+1} = \frac{1}{12}(-(2n+5)R_{-2n+2} + 4(n+2)R_{-2n+1} - (2n+5)R_{-2n} + 2(2n+1)R_{-2n-1} + 28)$.

5.2 The case $x = -1$

In this subsection we consider the special case $x = -1$.

Taking $x = -1, r = s = t = u = 1$ in Theorem 8 (a) and (b) (or (c)), we obtain the following proposition.

Proposition 10. *If $r = s = t = u = 1$ then for $n \geq 1$ we have the following formulas:*

- (a) $\sum_{k=1}^n (-1)^k W_{-k} = (-1)^n (-W_{-n+3} + 2W_{-n+2} - W_{-n+1} + 2W_{-n}) + W_3 - 2W_2 + W_1 - 2W_0$.
- (b) $\sum_{k=1}^n (-1)^k W_{-2k} = (-1)^n (-W_{-2n+2} + W_{-2n+1} + 2W_{-2n}) + W_2 - W_1 - 2W_0$.
- (c) $\sum_{k=1}^n (-1)^k W_{-2k+1} = (-1)^n (W_{-2n+1} - W_{-2n} - W_{-2n-1}) + W_3 - W_2 - 2W_1$.

From the above proposition, we have the following corollary which gives linear sum formulas of Tetranacci numbers (take $W_n = M_n$ with $M_0 = 0, M_1 = 1, M_2 = 1, M_3 = 2$).

Corollary 28. For $n \geq 1$, Tetranacci numbers have the following properties.

- (a) $\sum_{k=1}^n (-1)^k M_{-k} = (-1)^n (-M_{-n+3} + 2M_{-n+2} - M_{-n+1} + 2M_{-n}) + 1.$
- (b) $\sum_{k=1}^n (-1)^k M_{-2k} = (-1)^n (-M_{-2n+2} + M_{-2n+1} + 2M_{-2n}).$
- (c) $\sum_{k=1}^n (-1)^k M_{-2k+1} = (-1)^n (M_{-2n+1} - M_{-2n} - M_{-2n-1}) - 1.$

Taking $W_n = R_n$ with $R_0 = 4, R_1 = 1, R_2 = 3, R_3 = 7$ in the above proposition, we have the following corollary which presents linear sum formulas of Tetranacci-Lucas numbers.

Corollary 29. For $n \geq 1$, Tetranacci-Lucas numbers have the following properties.

- (a) $\sum_{k=1}^n (-1)^k R_{-k} = (-1)^n (-R_{-n+3} + 2R_{-n+2} - R_{-n+1} + 2R_{-n}) - 6.$
- (b) $\sum_{k=1}^n (-1)^k R_{-2k} = (-1)^n (-R_{-2n+2} + R_{-2n+1} + 2R_{-2n}) - 6.$
- (c) $\sum_{k=1}^n (-1)^k R_{-2k+1} = (-1)^n (R_{-2n+1} - R_{-2n} - R_{-2n-1}) + 2.$

Taking $x = -1, r = 2, s = t = u = 1$ in Theorem 8 (a) and (b) (or (c)), we obtain the following proposition.

Proposition 11. If $r = 2, s = t = u = 1$ then for $n \geq 1$ we have the following formulas:

- (a) $\sum_{k=1}^n (-1)^k W_{-k} = \frac{1}{2}((-1)^n (-W_{-n+3} + 3W_{-n+2} - 2W_{-n+1} + 3W_{-n}) + W_3 - 3W_2 + 2W_1 - 3W_0).$
- (b) $\sum_{k=1}^n (-1)^k W_{-2k} = \frac{1}{2}((-1)^n (-W_{-2n+2} + W_{-2n+1} + 4W_{-2n} + W_{-2n-1}) - W_3 + 3W_2 - 3W_0).$
- (c) $\sum_{k=1}^n (-1)^k W_{-2k+1} = \frac{1}{2}((-1)^n (-W_{-2n+2} + 3W_{-2n+1} - W_{-2n-1}) + W_3 - W_2 - 4W_1 - W_0).$

From the last proposition, we have the following corollary which gives linear sum formulas of fourth-order Pell numbers (take $W_n = P_n$ with $P_0 = 0, P_1 = 1, P_2 = 2, P_3 = 5$).

Corollary 30. For $n \geq 1$, fourth-order Pell numbers have the following properties:

- (a) $\sum_{k=1}^n (-1)^k P_{-k} = \frac{1}{2}((-1)^n (-P_{-n+3} + 3P_{-n+2} - 2P_{-n+1} + 3P_{-n}) + 1).$
- (b) $\sum_{k=1}^n (-1)^k P_{-2k} = \frac{1}{2}((-1)^n (-P_{-2n+2} + P_{-2n+1} + 4P_{-2n} + P_{-2n-1}) + 1).$
- (c) $\sum_{k=1}^n (-1)^k P_{-2k+1} = \frac{1}{2}((-1)^n (-P_{-2n+2} + 3P_{-2n+1} - P_{-2n-1}) - 1).$

Taking $W_n = Q_n$ with $Q_0 = 4, Q_1 = 2, Q_2 = 6, Q_3 = 17$ in the last proposition, we have the following corollary which presents linear sum formulas of fourth-order Pell-Lucas numbers.

Corollary 31. For $n \geq 1$, fourth-order Pell-Lucas numbers have the following properties:

- (a) $\sum_{k=1}^n (-1)^k Q_{-k} = \frac{1}{2}((-1)^n (-Q_{-n+3} + 3Q_{-n+2} - 2Q_{-n+1} + 3Q_{-n}) - 9).$
- (b) $\sum_{k=1}^n (-1)^k Q_{-2k} = \frac{1}{2}((-1)^n (-Q_{-2n+2} + Q_{-2n+1} + 4Q_{-2n} + Q_{-2n-1}) - 11).$
- (c) $\sum_{k=1}^n (-1)^k Q_{-2k+1} = \frac{1}{2}((-1)^n (-Q_{-2n+2} + 3Q_{-2n+1} - Q_{-2n-1}) - 1).$

Observe that setting $x = -1, r = 1, s = 1, t = 1, u = 2$ (i.e. for the generalized fourth order Jacobsthal case) in Theorem 8 (a),(b),(c) makes the right hand side of the sum formulas to be an indeterminate form. Application of L'Hospital rule however provides the evaluation of the sum formulas.

Taking $r = 1, s = 1, t = 1, u = 2$ in Theorem 8, we obtain the following theorem.

Theorem 12. If $r = 1, s = 1, t = 1, u = 2$ then for $n \geq 1$ we have the following formulas:

- (a) $\sum_{k=1}^n (-1)^k W_{-k} = \frac{1}{6}((-1)^n (-(n+1)W_{-n+3} + (2n+3)W_{-n+2} - (n+4)W_{-n+1} + 2(n+3)W_{-n}) + W_3 - 3W_2 + 4W_1 - 6W_0).$
- (b) $\sum_{k=1}^n (-1)^k W_{-2k} = \frac{1}{10}((-1)^n (-(3n+4)W_{-2n+2} + 2(n+1)W_{-2n+1} + (7n+13)W_{-2n} + 2(n+1)W_{-2n-1}) - W_3 + 5W_2 - W_1 - 12W_0).$
- (c) $\sum_{k=1}^n (-1)^k W_{-2k+1} = \frac{1}{10}((-1)^n (-(n+2)W_{-2n+2} + (4n+9)W_{-2n+1} - (n+2)W_{-2n} - 2(3n+4)W_{-2n-1}) + 4W_3 - 13W_1 - 2W_2 - 2W_0).$

Proof.

- (a) We use Theorem 8 (a). If we set $r = 1, s = 1, t = 1, u = 2$ in Theorem 8 (a) then we have

$$\sum_{k=1}^n x^k W_{-k} = \frac{g_8(x)}{-x^4 + x^3 + x^2 + x + 2}$$

where

$$g_8(x) = xW_3 + x(x-1)W_2 - x(-x^2 + x + 1)W_1 - x(-x^3 + x^2 + x + 1)W_0 - x^{n+1}W_{-n+3} - (x-1)x^{n+1}W_{-n+2} - (-x^2 + x + 1)x^{n+1}W_{-n+1} + (-x^3 + x^2 + x + 1)x^{n+1}W_{-n}.$$

For $x = -1$, the right hand side of the above sum formula is an indeterminate form. Now, we can use L'Hospital rule. Then we get (a) using

$$\sum_{k=1}^n (-1)^k W_{-k} = \frac{\frac{d}{dx}(g_8(x))}{\frac{d}{dx}(-x^4 + x^3 + x^2 + x + 2)} \Big|_{x=-1}.$$

- (b) We use Theorem 8 (b). If we set $r = 1, s = 1, t = 1, u = 2$ in Theorem 8 (b) then we have

$$\sum_{k=1}^n x^k W_{-2k} = \frac{g_9(x)}{-x^4 + 3x^3 + 5x^2 - 3x - 4}$$

where

$$g_9(x) = (-x^2 + x + 2)x^{n+1}W_{-2n+2} - (2x + 2)x^{n+1}W_{-2n+1} + (-x^3 + 3x^2 + 2x - 2)x^{n+1}W_{-2n} - 2(x + 1)x^{n+1}W_{-2n-1} + x(x + 1)W_3 - x(-x^2 + 2x + 3)W_2 + x(x + 1)W_1 - x(-x^3 + 3x^2 + 3x - 1)W_0.$$

For $x = -1$, the right hand side of the above sum formula is an indeterminate form. Now, we can use L'Hospital rule (twice). Then we get (b) using

$$\sum_{k=1}^n (-1)^k W_{-2k} = \frac{\frac{d^2}{dx^2}(g_9(x))}{\frac{d^2}{dx^2}(-x^4 + 3x^3 + 5x^2 - 3x - 4)} \Big|_{x=-1}.$$

- (c) We use Theorem 8 (c). If we set $r = 1, s = 1, t = 1, u = 2$ in Theorem 8 (c) then we have

$$\sum_{k=1}^n x^k W_{-2k+1} = \frac{g_{10}(x)}{-x^4 + 3x^3 + 5x^2 - 3x - 4}$$

where

$$g_{10}(x) = -(x + 1)x^{n+2}W_{-2n+2} + (-x^2 + 2x + 3)x^{n+2}W_{-2n+1} - (x + 1)x^{n+2}W_{-2n} + 2(-x^2 + x + 2)x^{n+1}W_{-2n-1} - x(-x^2 + x + 2)W_3 + x(2x + 2)W_2 - x(-x^3 + 3x^2 + 2x - 2)W_1 + 2x(x + 1)W_0.$$

For $x = -1$, the right hand side of the above sum formula is an indeterminate form. Now, we can use L'Hospital rule (twice). Then we get (c) using

$$\sum_{k=1}^n (-1)^k W_{-2k+1} = \frac{\frac{d^2}{dx^2}(g_{10}(x))}{\frac{d^2}{dx^2}(-x^4 + 3x^3 + 5x^2 - 3x - 4)} \Big|_{x=-1}.$$

□

Taking $W_n = J_n$ with $J_0 = 0, J_1 = 1, J_2 = 1, J_3 = 1$ in the last proposition, we have the following corollary which presents linear sum formula of fourth-order Jacobsthal numbers.

Corollary 32. *For $n \geq 1$, fourth order Jacobsthal numbers have the following property*

- (a) $\sum_{k=1}^n (-1)^k J_{-k} = \frac{1}{6}((-1)^n (-(n+1)J_{-n+3} + (2n+3)J_{-n+2} - (n+4)J_{-n+1} + 2(n+3)J_{-n}) + 2)$.
- (b) $\sum_{k=1}^n (-1)^k J_{-2k} = \frac{1}{10}((-1)^n (-(3n+4)J_{-2n+2} + 2(n+1)J_{-2n+1} + (7n+13)J_{-2n} + 2(n+1)J_{-2n-1}) + 3)$.
- (c) $\sum_{k=1}^n (-1)^k J_{-2k+1} = \frac{1}{10}((-1)^n (-(n+2)J_{-2n+2} + (4n+9)J_{-2n+1} - (n+2)J_{-2n} - 2(3n+4)J_{-2n-1}) - 11)$.

From the last proposition, we have the following corollary which gives linear sum formulas of fourth order Jacobsthal-Lucas numbers (take $W_n = j_n$ with $j_0 = 2, j_1 = 1, j_2 = 5, j_3 = 10$).

Corollary 33. *For $n \geq 1$, fourth order Jacobsthal-Lucas numbers have the following property*

- (a) $\sum_{k=1}^n (-1)^k j_{-k} = \frac{1}{6}((-1)^n (-(n+1)j_{-n+3} + (2n+3)j_{-n+2} - (n+4)j_{-n+1} + 2(n+3)j_{-n}) - 13)$.
- (b) $\sum_{k=1}^n (-1)^k j_{-2k} = \frac{1}{10}((-1)^n (-(3n+4)j_{-2n+2} + 2(n+1)j_{-2n+1} + (7n+13)j_{-2n} + 2(n+1)j_{-2n-1}) - 10)$.
- (c) $\sum_{k=1}^n (-1)^k j_{-2k+1} = \frac{1}{10}((-1)^n (-(n+2)j_{-2n+2} + (4n+9)j_{-2n+1} - (n+2)j_{-2n} - 2(3n+4)j_{-2n-1}) + 13)$.

Taking $W_n = K_n$ with $K_0 = 3, K_1 = 1, K_2 = 3, K_3 = 10$ in the last proposition, we have the following corollary which presents linear sum formula of modified fourth order Jacobsthal numbers.

Corollary 34. *For $n \geq 1$, modified fourth order Jacobsthal numbers have the following property:*

- (a) $\sum_{k=1}^n (-1)^k K_{-k} = \frac{1}{6}((-1)^n (-(n+1)K_{-n+3} + (2n+3)K_{-n+2} - (n+4)K_{-n+1} + 2(n+3)K_{-n}) - 13)$.
- (b) $\sum_{k=1}^n (-1)^k K_{-2k} = \frac{1}{10}((-1)^n (-(3n+4)K_{-2n+2} + 2(n+1)K_{-2n+1} + (7n+13)K_{-2n} + 2(n+1)K_{-2n-1}) - 32)$.
- (c) $\sum_{k=1}^n (-1)^k K_{-2k+1} = \frac{1}{10}((-1)^n (-(n+2)K_{-2n+2} + (4n+9)K_{-2n+1} - (n+2)K_{-2n} - 2(3n+4)K_{-2n-1}) + 15)$.

From the last proposition, we have the following corollary which gives linear sum formula of fourth-order Jacobsthal Perrin numbers (take $W_n = Q_n$ with $Q_0 = 3, Q_1 = 0, Q_2 = 2, Q_3 = 8$).

Corollary 35. *For $n \geq 1$, fourth-order Jacobsthal Perrin numbers have the following property:*

- (a) $\sum_{k=1}^n (-1)^k Q_{-k} = \frac{1}{6}((-1)^n (-(n+1)Q_{-n+3} + (2n+3)Q_{-n+2} - (n+4)Q_{-n+1} + 2(n+3)Q_{-n}) - 16)$.
- (b) $\sum_{k=1}^n (-1)^k Q_{-2k} = \frac{1}{10}((-1)^n (-(3n+4)Q_{-2n+2} + 2(n+1)Q_{-2n+1} + (7n+13)Q_{-2n} + 2(n+1)Q_{-2n-1}) - 34)$.
- (c) $\sum_{k=1}^n (-1)^k Q_{-2k+1} = \frac{1}{10}((-1)^n (-(n+2)Q_{-2n+2} + (4n+9)Q_{-2n+1} - (n+2)Q_{-2n} - 2(3n+4)Q_{-2n-1}) + 22)$.

Taking $W_n = S_n$ with $S_0 = 0, S_1 = 1, S_2 = 1, S_3 = 2$ in the last proposition, we have the following corollary which presents linear sum formula of adjusted fourth-order Jacobsthal numbers.

Corollary 36. *For $n \geq 1$, adjusted fourth-order Jacobsthal numbers have the following property:*

- (a) $\sum_{k=1}^n (-1)^k S_{-k} = \frac{1}{6}((-1)^n (-(n+1)S_{-n+3} + (2n+3)S_{-n+2} - (n+4)S_{-n+1} + 2(n+3)S_{-n}) + 3)$.
- (b) $\sum_{k=1}^n (-1)^k S_{-2k} = \frac{1}{10}((-1)^n (-(3n+4)S_{-2n+2} + 2(n+1)S_{-2n+1} + (7n+13)S_{-2n} + 2(n+1)S_{-2n-1}) + 2)$.
- (c) $\sum_{k=1}^n (-1)^k S_{-2k+1} = \frac{1}{10}((-1)^n (-(n+2)S_{-2n+2} + (4n+9)S_{-2n+1} - (n+2)S_{-2n} - 2(3n+4)S_{-2n-1}) - 7)$.

From the last proposition, we have the following corollary which gives linear sum formula of modified fourth-order Jacobsthal-Lucas numbers (take $W_n = R_n$ with $R_0 = 4, R_1 = 1, R_2 = 3, R_3 = 7$).

Corollary 37. For $n \geq 1$, modified fourth-order Jacobsthal-Lucas numbers have the following property:

- (a) $\sum_{k=1}^n (-1)^k R_{-k} = \frac{1}{6}((-1)^n (-(n+1)R_{-n+3} + (2n+3)R_{-n+2} - (n+4)R_{-n+1} + 2(n+3)R_{-n}) - 22)$.
- (b) $\sum_{k=1}^n (-1)^k R_{-2k} = \frac{1}{10}((-1)^n (-(3n+4)R_{-2n+2} + 2(n+1)R_{-2n+1} + (7n+13)R_{-2n} + 2(n+1)R_{-2n-1}) - 41)$.
- (c) $\sum_{k=1}^n (-1)^k R_{-2k+1} = \frac{1}{10}((-1)^n (-(n+2)R_{-2n+2} + (4n+9)R_{-2n+1} - (n+2)R_{-2n} - 2(3n+4)R_{-2n-1}) + 1)$.

Taking $x = -1, r = 2, s = 3, t = 5, u = 7$ in Theorem 8 (a), (b) and (c), we obtain the following proposition.

Proposition 13. If $r = 2, s = 3, t = 5, u = 7$ then for $n \geq 1$ we have the following formulas:

- (a) $\sum_{k=1}^n (-1)^k W_{-k} = \frac{1}{2}((-1)^n (W_{-n+3} - 3W_{-n+2} - 5W_{-n}) - W_3 + 3W_2 + 5W_0)$.
- (b) $\sum_{k=1}^n (-1)^k W_{-2k} = \frac{1}{6}((-1)^n (W_{-2n+2} - W_{-2n+1} - 6W_{-2n} - 7W_{-2n-1}) + W_3 - 3W_2 - 2W_1 + W_0)$.
- (c) $\sum_{k=1}^n (-1)^k W_{-2k+1} = \frac{1}{6}((-1)^n (W_{-2n+2} - 3W_{-2n+1} - 2W_{-2n} + 7W_{-2n-1}) - W_3 + W_2 + 6W_1 + 7W_0)$.

From the last proposition, we have the following corollary which gives linear sum formulas of 4-primes numbers (take $W_n = G_n$ with $G_0 = 0, G_1 = 0, G_2 = 1, G_3 = 2$).

Corollary 38. For $n \geq 1$, 4-primes numbers have the following properties:

- (a) $\sum_{k=1}^n (-1)^k G_{-k} = \frac{1}{2}((-1)^n (G_{-n+3} - 3G_{-n+2} - 5G_{-n}) + 1)$.
- (b) $\sum_{k=1}^n (-1)^k G_{-2k} = \frac{1}{6}((-1)^n (G_{-2n+2} - G_{-2n+1} - 6G_{-2n} - 7G_{-2n-1}) - 1)$.
- (c) $\sum_{k=1}^n (-1)^k G_{-2k+1} = \frac{1}{6}((-1)^n (G_{-2n+2} - 3G_{-2n+1} - 2G_{-2n} + 7G_{-2n-1}) - 1)$.

Taking $W_n = H_n$ with $H_0 = 4, H_1 = 2, H_2 = 10, H_3 = 41$ in the last proposition, we have the following corollary which presents linear sum formulas of Lucas 4-primes numbers.

Corollary 39. For $n \geq 1$, Lucas 4-primes numbers have the following properties:

- (a) $\sum_{k=1}^n (-1)^k H_{-k} = \frac{1}{2}((-1)^n (H_{-n+3} - 3H_{-n+2} - 5H_{-n}) + 9)$.
- (b) $\sum_{k=1}^n (-1)^k H_{-2k} = \frac{1}{6}((-1)^n (H_{-2n+2} - H_{-2n+1} - 6H_{-2n} - 7H_{-2n-1}) + 11)$.
- (c) $\sum_{k=1}^n (-1)^k H_{-2k+1} = \frac{1}{6}((-1)^n (H_{-2n+2} - 3H_{-2n+1} - 2H_{-2n} + 7H_{-2n-1}) + 9)$.

From the last proposition, we have the following corollary which gives linear sum formulas of modified 4-primes numbers (take $W_n = E_n$ with $E_0 = 0, E_1 = 0, E_2 = 1, E_3 = 1$).

Corollary 40. For $n \geq 1$, modified 4-primes numbers have the following properties:

- (a) $\sum_{k=1}^n (-1)^k E_{-k} = \frac{1}{2}((-1)^n (E_{-n+3} - 3E_{-n+2} - 5E_{-n}) + 2)$.
- (b) $\sum_{k=1}^n (-1)^k E_{-2k} = \frac{1}{6}((-1)^n (E_{-2n+2} - E_{-2n+1} - 6E_{-2n} - 7E_{-2n-1}) - 2)$.
- (c) $\sum_{k=1}^n (-1)^k E_{-2k+1} = \frac{1}{6}((-1)^n (E_{-2n+2} - 3E_{-2n+1} - 2E_{-2n} + 7E_{-2n-1}))$.

5.3 The case $x = i$

In this subsection we consider the special case $x = i$. Taking $r = s = t = u = 1$ in Theorem 8, we obtain the following proposition.

Proposition 14. *If $r = s = t = u = 1$ then for $n \geq 1$ we have the following formulas:*

- (a) $\sum_{k=1}^n i^k W_{-k} = i(i^n(W_{-n+3} - (1-i)W_{-n+2} - (2+i)W_{-n+1} - 2iW_{-n}) - W_3 + (1-i)W_2 + (2+i)W_1 + 2iW_0)$.
- (b) $\sum_{k=1}^n i^k W_{-2k} = \frac{(4+5i)}{41}(i^n(-(2+i)W_{-2n+2} + (1+2i)W_{-2n+1} + (4-2i)W_{-2n} + (1+i)W_{-2n-1}) - (1+i)W_3 + (3+2i)W_2 - iW_1 - (3-3i)W_0)$.
- (c) $\sum_{k=1}^n i^k W_{-2k+1} = \frac{(4+5i)}{41}(i^n(-(1-i)W_{-2n+2} + (2-3i)W_{-2n+1} - W_{-2n} - (2+i)W_{-2n-1}) + (2+i)W_3 - (1+2i)W_2 - (4-2i)W_1 - (1+i)W_0)$.

From the above proposition, we have the following corollary which gives linear sum formulas of Tetranacci numbers (take $W_n = M_n$ with $M_0 = 0, M_1 = 1, M_2 = 1, M_3 = 2$).

Corollary 41. *For $n \geq 1$, Tetranacci numbers have the following properties.*

- (a) $\sum_{k=1}^n i^k M_{-k} = i(i^n(M_{-n+3} - (1-i)M_{-n+2} - (2+i)M_{-n+1} - 2iM_{-n}) + 1)$.
- (b) $\sum_{k=1}^n i^k M_{-2k} = \frac{(4+5i)}{41}(i^n(-(2+i)M_{-2n+2} + (1+2i)M_{-2n+1} + (4-2i)M_{-2n} + (1+i)M_{-2n-1}) + (1-i))$.
- (c) $\sum_{k=1}^n i^k M_{-2k+1} = \frac{(4+5i)}{41}(i^n(-(1-i)M_{-2n+2} + (2-3i)M_{-2n+1} - M_{-2n} - (2+i)M_{-2n-1}) + (-1+2i))$.

Taking $W_n = R_n$ with $R_0 = 4, R_1 = 1, R_2 = 3, R_3 = 7$ in the above proposition, we have the following corollary which presents linear sum formulas of Tetranacci-Lucas numbers.

Corollary 42. *For $n \geq 1$, Tetranacci-Lucas numbers have the following properties.*

- (a) $\sum_{k=1}^n i^k R_{-k} = i(i^n(R_{-n+3} - (1-i)R_{-n+2} - (2+i)R_{-n+1} - 2iR_{-n}) + 2(-1+3i))$.
- (b) $\sum_{k=1}^n i^k R_{-2k} = \frac{(4+5i)}{41}(i^n(-(2+i)R_{-2n+2} + (1+2i)R_{-2n+1} + (4-2i)R_{-2n} + (1+i)R_{-2n-1}) + 10(-1+i))$.
- (c) $\sum_{k=1}^n i^k R_{-2k+1} = \frac{(4+5i)}{41}(i^n(-(1-i)R_{-2n+2} + (2-3i)R_{-2n+1} - R_{-2n} - (2+i)R_{-2n-1}) + (3-i))$.

Corresponding sums of the other fourth order generalized Tetranacci numbers can be calculated similarly.

References

- [1] Akbulak, M., Öteleş, A., On the sum of Pell and Jacobsthal numbers by matrix method, Bulletin of the Iranian Mathematical Society, 40 (4), 1017-1025, 2014.
- [2] Frontczak, R., Sums of Tribonacci and Tribonacci-Lucas Numbers, International Journal of Mathematical Analysis, 12 (1), 19-24, 2018.
- [3] Gökbaş, H., Köse, H., Some Sum Formulas for Products of Pell and Pell-Lucas Numbers, Int. J. Adv. Appl. Math. and Mech. 4(4), 1-4, 2017.
- [4] Hansen., R.T., General Identities for Linear Fibonacci and Lucas Summations, Fibonacci Quarterly, 16(2), 121-28, 1978.

- [5] Hathiwala, G. S., Shah, D. V., Binet–Type Formula For The Sequence of Tetranacci Numbers by Alternate Methods, *Mathematical Journal of Interdisciplinary Sciences* 6 (1), 37–48, 2017.
- [6] Koshy, T., *Fibonacci and Lucas Numbers with Applications*, A Wiley-Interscience Publication, New York, 2001.
- [7] Koshy, T., *Pell and Pell-Lucas Numbers with Applications*, Springer, New York, 2014.
- [8] Melham, R. S., Some Analogs of the Identity $F_n^2 + F_{n+1}^2 = F_{2n+1}^2$, *Fibonacci Quarterly*, 305-311, 1999.
- [9] Natividad, L. R., On Solving Fibonacci-Like Sequences of Fourth, Fifth and Sixth Order, *International Journal of Mathematics and Computing*, 3 (2), 2013.
- [10] Parpar, T., k'ncü Mertebeden Rekürans Bağıntısının Özellikleri ve Bazı Uygulamaları, Selçuk Üniversitesi, Fen Bilimleri Enstitüsü, Yüksek Lisans Tezi, 2011.
- [11] Singh, B., Bhadouria, P., Sikhwal, O., Sisodiya, K., A Formula for Tetranacci-Like Sequence, *Gen. Math. Notes*, 20 (2), 136-141, 2014.
- [12] Sloane, N.J.A., The on-line encyclopedia of integer sequences. Available: <http://oeis.org/>
- [13] Soykan, Y., On Summing Formulas For Generalized Fibonacci and Gaussian Generalized Fibonacci Numbers, *Advances in Research*, 20(2), 1-15, 2019.
- [14] Soykan, Y., Corrigendum: On Summing Formulas for Generalized Fibonacci and Gaussian Generalized Fibonacci Numbers, *Advances in Research*, 21(10), 66-82, 2020. DOI: 10.9734/AIR/2020/v21i1030253
- [15] Soykan, Y., On Summing Formulas for Horadam Numbers, *Asian Journal of Advanced Research and Reports* 8(1): 45-61, 2020, DOI: 10.9734/AJARR/2020/v8i130192.
- [16] Soykan, Y., Generalized Fibonacci Numbers: Sum Formulas, *Journal of Advances in Mathematics and Computer Science*, 35(1), 89-104, 2020, DOI: 10.9734/JAMCS/2020/v35i130241.
- [17] Soykan Y., Generalized Tribonacci Numbers: Summing Formulas, *Int. J. Adv. Appl. Math. and Mech.* 7(3), 57-76, 2020.
- [18] Soykan, Y., Summing Formulas For Generalized Tribonacci Numbers, *Universal Journal of Mathematics and Applications*, 3(1), 1-11, 2020. ISSN 2619-9653, DOI: <https://doi.org/10.32323/ujma.637876>
- [19] Soykan, Y., On Sum Formulas for Generalized Tribonacci Sequence, *Journal of Scientific Research & Reports*, 26(7), 27-52, 2020. ISSN: 2320-0227, DOI: 10.9734/JSRR/2020/v26i730283
- [20] Soykan, Y., Summation Formulas For Generalized Tetranacci Numbers, *Asian Journal of Advanced Research and Reports*, 7(2), 1-12, 2019. doi.org/10.9734/ajarr/2019/v7i230170.
- [21] Soykan, Y., Sum Formulas For Generalized Fifth-Order Linear Recurrence Sequences, *Journal of Advances in Mathematics and Computer Science*, 34(5), 1-14, 2019; Article no.JAMCS.53303, ISSN: 2456-9968, DOI: 10.9734/JAMCS/2019/v34i530224.
- [22] Soykan, Y., Linear Summing Formulas of Generalized Pentanacci and Gaussian Generalized Pentanacci Numbers, *Journal of Advanced in Mathematics and Computer Science*, 33(3): 1-14, 2019.
- [23] Soykan, Y., On Summing Formulas of Generalized Hexanacci and Gaussian Generalized Hexanacci Numbers, *Asian Research Journal of Mathematics*, 14(4), 1-14, 2019; Article no.ARJOM.50727.
- [24] Soykan, Y., A Study On Sum Formulas of Generalized Sixth-Order Linear Recurrence Sequences, *Asian Journal of Advanced Research and Reports*, 14(2), 36-48, 2020. DOI: 10.9734/AJARR/2020/v14i230329

-
- [25] Soykan, Y., Matrix Sequences of Tribonacci and Tribonacci-Lucas Numbers, *Communications in Mathematics and Applications*, 11(2), 281-295, 2020. DOI: 10.26713/cma.v11i2.1102
- [26] Soykan, Y., Gaussian Generalized Tetranacci Numbers, *Journal of Advances in Mathematics and Computer Science*, 31(3): 1-21, Article no.JAMCS.48063, 2019.
- [27] Soykan, Y., A Study of Generalized Fourth-Order Pell Sequences, *Journal of Scientific Research and Reports*, 25(1-2), 1-18, 2019.
- [28] Polath, E.E., Soykan, Y., A Study on Generalized Fourth-Order Jacobsthal Sequences, Submitted.
- [29] Soykan, Y., On Generalized 4-primes Numbers, *Int. J. Adv. Appl. Math. and Mech.* 7(4), 20-33, 2020.
- [30] Soykan, Y., Properties of Generalized (r,s,t,u) -Numbers, *Earthline Journal of Mathematical Sciences*, 5(2), 297-327, 2021. <https://doi.org/10.34198/ejms.5221.297327>
- [31] Öteleş, A., Akbulak, M., A Note on Generalized k -Pell Numbers and Their Determinantal Representation, *Journal of Analysis and Number Theory*, 4(2), 153-158, 2016.
- [32] Waddill, M. E., The Tetranacci Sequence and Generalizations, *Fibonacci Quarterly*, 9-20, 1992.
- [33] Waddill, Another Generalized Fibonacci Sequence, M. E., *Fibonacci Quarterly*, 5 (3), 209-227, 1967.