

## Effect of production system on fatty acid composition in subcutaneous adipose tissue of Ile de France lambs

Martin Janíček<sup>1</sup>, Milan Margetín<sup>1,2</sup>, Klára Vavrišínová<sup>1</sup>, Katarína Hozáková<sup>1</sup>

<sup>1</sup> Department of Animal Husbandry, Faculty of Agrobiolgy and Food Resources, Slovak University of Agriculture in Nitra

<sup>2</sup> NPPC - Research Institute for Animal Production in Nitra



Licensed under a Creative Commons Attribution 4.0 International License



A study was conducted on the effect of two different lamb production systems on the fatty acid (FA) composition of subcutaneous fat of 40 purebred Ile de France lambs. In the first production system, ewes and lambs grazed on pasture without concentrate (GS), whereas in the second production system, ewes and lambs were housed indoor and fed silage/hay and concentrate (IS). Twenty lambs (7 females and 13 males) were included in each group. Lambs were slaughtered at  $28.75 \pm 2.76$  kg. The FA were determined by gas chromatography and analysed through ANOVA by considering the following fixed effects: production system, sex and the interaction between production system and sex. Subcutaneous fat of GS lambs had greater proportion of C18:3n-3 ( $P < 0.001$ ), C22:5n-3 ( $P < 0.05$ ) and C22:6n-3 ( $P < 0.05$ ) than IS lambs, which resulted in a higher sum of n-3 polyunsaturated FA in GS compared to IS lambs (2.00 vs. 1.15 g/100 g FAME,  $P < 0.001$ ). Moreover, subcutaneous fat of GS lambs had greater proportion of c9,t11-C18:2 ( $P < 0.001$ ) and sum of detected conjugated linoleic acid isomers than IS lambs (2.21 vs. 0.67 g/100 g FAME,  $P < 0.001$ ). Females had significantly greater proportion of C18:2n-6 ( $P < 0.05$ ) and C18:3n-6 ( $P < 0.001$ ) than males. We can conclude that the GS system where lambs are raised under grazing conditions may provide carcasses with a more acceptable subcutaneous fat, as far as a human health and nutrition perspective is concerned.

**Keywords:** lamb, production system, subcutaneous fat, fatty acid composition

### 1 Introduction

The quality of food, especially in regard to its benefits for human health, plays an important role in affecting customer purchase decision. In recent years, a discussion has emerged surrounding both the positive and negative effects of red meat consumption, including lamb meat consumption (McAfee et al., 2010, Corpet, 2011). The average consumption of lean red meat, which is considered part of a balanced diet, evidently does not increase the risk of cardiovascular disease and colon cancer. It presents a positive effect on the intake of essential nutrients and a spectrum of beneficial fatty acids (FA; Nudda et al., 2009, Nudda et al., 2011, Binnie et al., 2014).

The proportion of FA may differ depending on the location of adipose tissue in the lamb's carcass. Potkanski et al. (2002) and Wood et al. (2008) reported that intramuscular fat has a more beneficial proportion of healthy FA than subcutaneous fat. As reported by Díaz et al. (2005), the proportion of FA in sheep meat is affected primarily by diet, as well as factors such as age, sex, breed and production system. Vasta et al. (2012) reported that pasture grazing is the best feeding system for lambs because the meat is much juicier and tastier, whilst having a higher proportion of beneficial FA for human health. Previous studies indicated higher proportion of n-3 polyunsaturated FA (PUFA) in the tissue of lambs from pasture than from concentrate-fed (Santos-Silva et al., 2002, Cividini et al., 2008, Araba et al., 2009, Popova, 2014, Karaca et al., 2016).

The demand for light lambs in the Mediterranean countries is largely affecting Slovak lamb meat production. The last decade has been characterized by light lamb's price fall. A large number of

\* **Corresponding Author:** Martin Janíček. Slovak University of Agriculture in Nitra, Faculty of Agrobiolgy and Food Resources, Tr. A. Hlinku 2, 949 76 Nitra, Slovakia; E-mail: [martin.janicek@uniag.sk](mailto:martin.janicek@uniag.sk). ORCID: <https://orcid.org/0000-0002-3989-3833>

stakeholders have transitioned from the dairy to the meat production system, due to lower production costs. There are two semi-intensive systems that are preferable for heavy carcass lamb production; the grazing system with ewes not receiving concentrate and the indoor system with ewes fed meadow hay or alfalfa silage, with unrestricted access to concentrate for lambs. The grazing system is more convenient in the foothill pastures and the indoor system in the areas rich in arable land. There is an opportunity for heavy lambs in Slovakia, due to growing demands from cultural minorities in Europe that tend to prefer lambs of a higher weight.

For this reason, it is necessary to study the quality of heavy lambs from Slovak production systems. The aim of this study was to investigate the effect of grazing and indoor systems on the FA composition of subcutaneous fat in Ile de France heavy lambs.

## 2 Material and methods

### 2.1 Experimental design and animal management

Forty Ile de France heavy lambs were included in the trial. Two experimental groups, namely the indoor system (IS) and the grazing system (GS) were formed, each consisting of 7 female and 13 male lambs. The IS consisted of lambs housed with ewes and allowed to suckle milk *ad libitum*. Lambs received an access to the separate area two weeks after birth, where lamb starter (crude protein: min 19.0%; crude fiber: max 9.0%; crude fat: max 6.0%; ash: max 10.0%; calcium: min 1.12%; phosphorus: min 0.61%; sodium: min 0.2%) and hay were offered *ad libitum*. Moreover, until 60 days of age, lambs were fed 0.2 kg mixture of lucerne and maize silage (ratio 2:1) per capita per day. The ewes were fed 1 kg maize silage, 1 kg lucerne silage, 2 kg meadow hay and 0.2 kg of concentrate per capita per day. Lambs in the GS were born indoor and shortly after parturition they were moved with ewes to pasture. The pasture was natural and rich in *Lolium perenne*, *Lolium multiflorum*, *Trifolium repens*, *Trifolium pretense*, *Poa pratensis* and *Festuca rubra*.

The slaughter of lambs was carried out in the authorised slaughterhouse of Slovak University of Agriculture in Nitra. The average age at slaughter was  $107 \pm 8$  days, and the average live weight was  $29.23 \pm 8.20$  kg for GS and  $32.2 \pm 2.62$  kg for IS. The average daily gain was  $0.242 \pm 0.034$  kg for GS and  $0.264 \pm 0.029$  kg for IS. Animal care, management and slaughter procedures followed the EU Directive 2010/63/EU on the protection of animals used in scientific experiments.

### 2.2 Fatty acid analysis

Two g of subcutaneous fat were taken from the tailhead area of every lamb to determine the FA profile of subcutaneous fat. Samples were collected 24 h after slaughter and carcass chilling. Capillary gas chromatography was used to determine the proportion of the individual FA and FA isomers. The fatty acid methyl esters (FAME) preparation and gas chromatography determination are described in Margetin et al. (2018).

The FA profile of subcutaneous fat was detected in grams of individual FAME per 100 g of total detected FAME. The average relative standard deviation of analysed FAME with content  $>0.5$  g/100 g was 1.1% for the entire analytical procedure and five replicated samples.

### 2.3 Statistical analysis

The experimental data was analysed by ANOVA using the GLM procedure of the software SAS (SAS Institute Inc., Cary, NC, USA). The linear model included the fixed effects of production system, sex and the interaction between production system and sex. Least squares means were compared by applying Scheffe's test at the significance limits:  $P < 0.05$ ,  $P < 0.01$  and  $P < 0.001$ .

## 3 Results and discussion

The composition of important FA identified in subcutaneous fat which are believed to affect human health, are reported in Table 1. Saturated FA (SFA) are formed from hydrolysed unsaturated FA (UFA) and this process depends on the biohydrogenation activity in rumen (Zervas and Tsiplakou, 2011). The ewes' maternal milk is rich in lauric acid (C12:0) and myristic acid (C14:0); these FA are mostly provided through the diet used to feed lambs, including dietary fat (Juárez et al., 2008). As previously reported by Santos-Silva et al. (2002), lambs raised under a pasture system showed higher C14:0 in intramuscular fat in relation to concentrate-fed lambs. The authors interpreted this result as consequence of the greater C14:0 proportion in maternal milk compared to concentrate-fed diet. In the

current study, lambs from both groups had unlimited access to maternal milk and no effect on the proportion of C12:0 and C14:0 was observed. The effect of different production systems significantly affected the proportion of palmitic acid (C16:0; 28.45 vs. 22.67 g/100 g FAME,  $P < 0.001$ ) and margaric acid (C17:0; 1.47 vs. 1.28 g/100 g FAME,  $P < 0.01$ ) and the total SFA (60.20 vs. 55.93 g/100 g FAME,  $P < 0.001$ ). The proportion of FA groups and their ratios are shown in Table 2. Mora et al. (2016) and Leão et al. (2011) found a similar concentration of C16:0 in subcutaneous fat of lambs finished by concentrates. Díaz et al. (2002) reported similar results concerning the proportion of C16:0 in subcutaneous loin fat, yet the concentration of the total SFA was higher in pastured lambs (59.37 vs. 57.16% of FA,  $P < 0.01$ ). A potential explanation could be that the energy intake of IS lambs was higher because the IS lambs' feed was richer in energy. Thus, the *de novo* synthesis of C16:0 was also higher. The GS lambs had higher proportion of SFA only for stearic acid (C18:0; 17.86 vs. 14.96 g/100 g FAME,  $P < 0.01$ ), which is one of the primary FA in meat of ruminants. It does not have an effect on cholesterol concentration in the bloodstream because it is desaturated and converted to oleic acid (c9-C18:1; Williams, 2000). The results of C18:0 were similar to some earlier investigations (Díaz et al., 2002, Guler et al., 2011, Karaca et al., 2016). It is likely that the ewes' milk fat consumed by GS lambs had a higher proportion of C18:1, which is biohydrogenated to C18:0 by rumen microbiota. That could be affected by the nature of dietary lipids, as well as ewes' rumen FA metabolism and FA synthesis in the intestinal, adipose tissue and mammary gland of ewes.

Proportion of palmitoleic acid (c9-C16:1), n-7 monounsaturated FA (MUFA), was higher in GS than IS group (0.64 vs. 0.48 g/100 g FAME,  $P < 0.001$ ). Karaca et al. (2016) reported different findings, i.e. pastured lambs had a lower proportion of c9-C16:1 than intensively fattened lambs (3.58 vs. 5.24% of FA,  $P < 0.001$ ). Proportion of c9-C18:1 was significantly higher in IS than GS (27.00 vs. 25.52 g/100 g FAME,  $P < 0.05$ ). The GS group also had a higher proportion of elaidic acid (t9-C18:1) than the IS group (0.33 vs. 0.22 g/100 g FAME,  $P < 0.001$ ).

**Table 1** Fatty acid composition (g/100 g FAME) of subcutaneous fat of lambs according to the production system (GS – grazing system, IS – indoor system) and sex (male and female)

Fatty acid	Production system (PS)		P-value	Sex		P-value	P-value PS x Sex
	GS	IS		Male	Female		
C12:0	1.22	1.19	0.833	1.35	1.05	0.027	0.194
C14:0	8.38	9.38	0.101	9.39	8.37	0.091	0.096
C16:0	22.67	28.45	<0.001	26.20	24.92	0.032	0.226
c9-C16:1	0.64	0.48	<0.001	0.57	0.55	0.166	0.196
C17:0	1.28	1.47	0.001	1.34	1.41	0.180	0.627
C18:0	17.86	14.96	0.002	16.40	16.43	0.976	0.730
c9-C18:1	25.52	27.00	0.030	25.62	26.91	0.055	0.017
t9-C18:1	0.33	0.22	<0.001	0.27	0.28	0.595	0.105
t11-C18:1	4.04	1.18	<0.001	2.60	2.62	0.777	0.341
C18:2n-6	2.84	2.26	<0.001	2.41	2.68	0.040	0.931
c9,t11-C18:2	1.96	0.60	<0.001	1.23	1.33	0.374	0.793
C18:3n-6	0.02	0.03	<0.001	0.01	0.03	<0.001	0.402
C18:3n-3	1.38	0.84	<0.001	1.15	1.07	0.243	0.962
C20:4n-6	0.29	0.10	0.176	0.08	0.30	0.104	0.143
C20:5n-3	0.08	0.04	0.117	0.03	0.09	0.090	0.088
C22:5n-3	0.36	0.19	0.051	0.20	0.35	0.103	0.092
C22:6n-3	0.09	0.04	0.050	0.05	0.09	0.090	0.070

The proportion of vaccenic acid (t11-C18:1) and rumenic acid (c9,t11-C18:2), the latter being the important isomer of conjugated linoleic acid (CLA), were significantly higher in GS than IS (4.04 vs. 1.18 and 1.96 vs. 0.60 g/100 g FAME, respectively,  $P < 0.001$ ). Nuernberg et al. (2008) did not find any differences in the proportion of t11-C18:1 and c9,t11-C18:2 in subcutaneous fat of pastured and intensive reared lambs. Di Memmo (2014) reported a high proportion of total CLA in the meat of lambs which were fed their mother's milk at pasture. We found a similar effect on the total CLA, when the proportion was more than threefold higher in GS than IS (2.21 vs. 0.67 g/100 g FAME,  $P < 0.001$ , Table 2). The higher proportion of CLA in subcutaneous fat of GS than IS is in agreement with studies focused on differences in intramuscular fat (Auroseau et al., 2004, Scerra et al., 2011, Popova, 2014). The difference of MUFA between GS and IS was not significant. Pastured lambs exhibited

higher proportion of both essential PUFA, linoleic acid (C18:2n-6) and  $\alpha$ -linolenic acid (C18:3n-3) in subcutaneous fat compared to the IS lambs (2.84 vs. 2.26 and 1.38 vs. 0.84 g/100 g FAME, respectively,  $P < 0.001$ ). In agreement with results of Karaca et al. (2016), the proportion of C18:2n-6 was higher in the IS lambs.

The second important n-6 FA, i.e. the  $\gamma$ -linolenic acid (C18:3n-6), was higher in IS than GS lambs (0.03 vs. 0.02 g/100 g FAME,  $P < 0.001$ ). However, these values are relatively low. The essential C18:3n-3 plays an important role in the production of the n-3 long chain PUFA, resulting in a positive effect on higher proportions of docosapentaenoic acid (C22:5 n-3) and docosahexaenoic acid (C22:6n-3) in the GS than IS lambs (0.36 vs 0.19 and 0.09 and 0.04 g/100 g FAME, respectively,  $P < 0.05$ ). The results of n-6 and n-3 acids obtained in this study were similar to those reported by Karaca et al. (2016). The proportion of total n-6 PUFA and n-3 PUFA was higher in the GS than in the IS lambs (3.18 vs. 2.42 and 2.00 vs. 1.15 g/100 g FAME, respectively,  $P < 0.001$ ). This affected total PUFA, which was higher for the GS than the IS lambs (9.17 vs. 5.55 g/100 g FAME,  $P < 0.001$ ). Value of the important n-6/n-3 ratio was more suitable in subcutaneous fat of the GS lambs, even if in both groups the values did not exceed the recommended standard 4:1 (1.61 vs. 2.15,  $P < 0.001$ ). According to Simopoulos (2016), increasing both the proportion of n-6 FA and n-6/n-3 ratio in phospholipid membrane of red blood cells, can cause an increased risk of obesity. On the other hand, a high proportion of n-3 FA in the phospholipid membrane of red blood cells can reduce this risk.

**Table 2** Groups of and ratios between fatty acids (g/100 g FAME) of subcutaneous fat of lambs according to the production system (GS – grazing system, IS – indoor system) and sex (male and female)

Fatty acid	Production system (PS)		P-value	Sex		P-value	P-value PS x Sex
	GS	IS		Male	Female		
SFA <sup>a</sup>	55.93	60.20	<0.001	59.43	56.70	0.006	0.071
MUFA <sup>b</sup>	34.90	34.08	0.337	33.74	35.25	0.085	0.027
PUFA <sup>c</sup>	9.17	5.55	<0.001	6.89	7.82	0.010	0.152
CLA <sup>d</sup>	2.21	0.67	<0.001	1.38	1.49	0.361	0.848
n-6 PUFA <sup>e</sup>	3.18	2.42	0.001	2.54	3.06	0.025	0.320
n-3 PUFA <sup>f</sup>	2.00	1.15	<0.001	1.48	1.67	0.200	0.063
PUFA/SFA	0.17	0.09	<0.001	0.12	0.14	0.007	0.319
n-6/n-3	1.61	2.15	<0.001	1.76	2.00	0.041	0.063
LA <sup>g</sup> /ALA <sup>h</sup>	2.15	2.77	0.003	2.17	2.75	0.006	0.627

<sup>a</sup> sum of saturated fatty acids: C8:0 + C10:0 + C11:0 + C12:0 + C13:0 + iso-C14:0 + C14:0 + iso-C15:0 + anteiso-C15:0 + C15:0 + iso-C16:0 + C16:0 + iso-C17:0 + anteiso-C17:0 + C17:0 + iso-C18:0 + C18:0 + C19:0 + C20:0 + C21:0 + C22:0

<sup>b</sup> sum of monounsaturated fatty acids: C12:1 + C14:1 + t-C16:1 + c-C16:1 + c9-C16:1 + C17:1 + t6/t7/t8-C18:1 + t9-C18:1 + t10-C18:1 + t11-C18:1 + t12-C18:1 + c9-C18:1 + (t15-C18:1 + c11-C18:1) + c12-C18:1 + c13-C18:1 + (c14-C18:1 + t9, t12-C18:2/2) + c15-C18:1 + (C18:2/2 + C19:1) + C20:1

<sup>c</sup> sum of polyunsaturated fatty acids: (c14-C18:1 + t9, t12-C18:2/2) + c9, t13-C18:2 + (t8, c13-C18:2 + c9, t12-C18:2) + (t9, c12-C18:2 + t11, c15-C18:2) + C18:2n-6 + c9, c15-C18:2 + c12, c15-C18:2 + cc-C18:2 + C18:3 n-6 + (C18:2 + C19:1/2) + cyklo + t9, c12, c15-C18:3 + C18:3n-3 + c9, t11-C18:2 + ct-CLA + cc-CLA + tc-CLA + tt-CLA + C18:3 + C20:2 + C20:3n-9 + C20:3n-6 + C20:4n-6 + C20:3n-3 + C20:4n-3 + C20:5n-3 + furyl C22 + C22:4n-3 + C22:5n-3 + C22:6n-3

<sup>d</sup> sum of n-6 PUFA: C18:2n-6 + C18:3n-6 + C20:3n-6 + C20:4n-6

<sup>e</sup> sum of n-3 PUFA: C18:3n-3 + C20:3n-3 + C20:4n-3 + C20:5n-3 + C22:4n-3 + C22:5n-3 + C22:6n-3

<sup>f</sup> sum of conjugated linoleic acid isomers: c9, t11-C18:2 + ct-CLA + cc-CLA + tc-CLA + tt-CLA

<sup>g</sup> linoleic acid

<sup>h</sup>  $\alpha$ -linolenic acid

The sex effect significantly explained the variation of C12:0 and C16:0; indeed, the proportion of both FA was higher in subcutaneous fat of males than females (1.35 vs. 1.05 and 26.20 vs. 24.92 g/100 g FAME,  $P < 0.001$ ), whereas females had significantly higher proportion of C18:2n-6 (2.68 vs. 2.41 g/100 g FAME,  $P < 0.05$ ) and C18:3n-6 (0.03 vs. 0.01 g/100 g FAME,  $P < 0.001$ ) than males (Table 1). Males had a lower n-6/n-3 ratio than females (1.76 vs. 2.00,  $P < 0.05$ , Table 2). The production system

by sex interaction was significant for c9-C18:1 ( $P=0.017$ ), which possibly affected the sum of MUFA ( $P=0.027$ ). Interaction effect in proportion of healthy FA was not significant. Because of that, our sample data did not provide enough evidence to conclude that the relationship between production system and sex has positive effect on FA composition. To the best of authors' knowledge, investigations on the interaction effect between grazing/indoor system and sex have not been published yet. Therefore, appropriate comparison with the literature could not be carried out.

#### 4 Conclusions

Results of this study indicated that the production system affected the FA profile of subcutaneous fat in Ile de France heavy lambs. The muscular lipids of lambs raised on pasture exhibited higher proportions of n-3 FA, in particular  $\alpha$ -linolenic acid, docosapentaenoic acid, docosahexaenoic acid and CLA, alongside a preferable n-6/n-3 ratio, which are beneficial for human health. Therefore, the pasture grazed lambs may be considered as a better nutritional source for humans than lambs raised in indoor systems.

#### Acknowledgments

The present study was supported by the Faculty of Agriculture and Food Resources found (GA FAPZ No. 2/2019) and by the Grant of Ministry of Education, Science, Research and Sport of the Slovak Republic (KEGA No. 015SPU-4/2019).

#### References

- Ackman, R. G. (2002). The gas chromatograph in practical analyses of common and uncommon fatty acids for the 21st century. *Analytica Chimica Acta*, 465(1-2), 175-192. [https://doi.org/10.1016/S0003-2670\(02\)00098-3](https://doi.org/10.1016/S0003-2670(02)00098-3)
- Araba, A., Bouarour, M., Bas, P., Morand-Fehr, P., El Aich, A., Kabbali, A. (2009). Performance carcass characteristics and meat quality of Timahdite-breed lambs finished on pasture or on hay and concentrate. *Options Méditerranéennes : Série A. Séminaires Méditerranéens*, 85, 465-469.
- Auroseau, B., Bauchart, D., Calichon, E., Micol, D., Priolo, A. (2004). Effect of grass and concentrate feeding systems and role of growth on triglyceride and phospholipid and their fatty acids in the M. Longissimus thoracis of lambs. *Meat Science*, 66(3), 531-541. [https://doi.org/10.1016/S0309-1740\(03\)00156-6](https://doi.org/10.1016/S0309-1740(03)00156-6)
- Binnie, M. A., Barlow, K., Johnson, V., Harrison, C. (2014). Red meats: Time for a paradigm shift in dietary advice. *Meat Science*, 98(3), 445-451. <https://doi.org/10.1016/j.meatsci.2014.06.024>
- Cividini, A., Levart, A., Zgur, S. (2008). Fatty acid composition as affected by production system, waning and sex. *Acta agriculturae Slovenica*, 2, 47-52.
- Corpet, D. E. (2011). Red meat and colon cancer: Should we become vegetarians, or can we make meat safer? *Meat Science*, 89(3), 310-316. <https://doi.org/10.1016/j.meatsci.2011.04.009>
- Díaz, M. T., Velasco, S., Cañeque, V., Lauzurica, S., Ruiz de Huidobro, F., Pérez, C., González, J., Manzanares, C. (2002). Use of concentrate or pasture for fattening lambs and its effect on carcass and meat quality. *Small Ruminant Research*, 43, 257-268. [https://doi.org/10.1016/S0921-4488\(02\)00016-0](https://doi.org/10.1016/S0921-4488(02)00016-0)
- Díaz, M. T., Álvarez, I., La Fuente, J., Sañudo, C., Campo, M. M., Oliver, M. A. Cañeque, V. (2005). Fatty acid composition of meat from typical lamb production systems of Spain, United Kingdom, Germany and Uruguay. *Meat Science*, 71(2), 256-263. <https://doi.org/10.1016/j.meatsci.2005.03.020>
- Di Memmo, D. (2015). *Influence of multiple injections of vitamin E on quality traits and oxidative stability of lamb meat*. Doctorate thesis. Campobasso : University of Molise. 130 p.
- Guler, G. O., Aktumsek, A., and Karabacak, A. (2011). Effect of Feeding Regime on Fatty Acid Composition of Longissimus dorsi Muscle and Subcutaneous Adipose Tissue of Akkaraman Lambs, *Kafkas Universitesi Veteriner Fakultesi Dergisi*, 17, 885-892. <https://doi.org/10.9775/kvfd.2011.4495>
- Juárez, M., Horcada, A., Alcalde M. J., Valera, M., Mullen, A. M., Molina, A. (2008) Estimation of factors influencing fatty acid profiles in light lambs. *Meat Science*, 79(2), 203-210. <https://doi.org/10.1016/j.meatsci.2007.08.014>
- Karaca, S., Yilmaz, A., Kor, A., Bingöl, M., Cavidoglu, I., Ser, G. (2016). The effect of feeding system on slaughter-carcass characteristics, meat quality, and fatty acid composition of lambs. *Archives Animal Breeding*, 59, 121-129. <https://doi.org/10.5194/aab-59-121-2016>
- Leão, A. G., Silva Sobrinho, A. G., Moreno, G. M. B., Souza, H. B. A., Perez, H. L., Loureiro, C. M. B. (2011). Características nutricionais da carne de cordeiros terminados com dietas contendo cana-deaçúcar ou silagem de

milho e dois níveis de concentrado. *Revista Brasileira de Zootecnia*, 40(5), 1072-1079.  
<https://doi.org/10.1590/S1516-35982011000500019>

Margetin, M., Oravcová, M., Margetinová, J., Kubinec, R. (2018). Fatty Acids in Intramuscular Fat of Ile De France Lambs in Two Different Production Systems. *Archives Animal Breeding*, 61(4), 395-403.  
<https://doi.org/10.5194/aab-61-395-2018>

McAfee, A. J., McSorley, E. M., Cuskelly, G. J., Moss, B. W., Wallace, J. M., Bonham, M. P., Fearon, A. M. (2010). Red meat consumption: an overview of the risk and benefits. *Meat Science*, 84(1), 1-13.  
<https://doi.org/10.1016/j.meatsci.2009.08.029>

Mora, N. H. A. P., De Macedo, F. A. F., Feihrmann, A. C., Possamai, A. P. S., Torres, M. G., Mexia, A. A. (2016). Lipid composition and sensory traits of meat from Pantaneiro lambs slaughtered with different subcutaneous fat thickness. *Acta Scientiarum. Technology*, 38(2), 145-151.  
<https://doi.org/10.4025/actascitechnol.v28i2.28341>

Nudda, A., Mele, M., Serra, A., Manca, M. G., Boe, R., Secchiari, P. (2009). Comparison of fatty acid profile in lamb meat and baby food based on lamb meat. *Italian Journal of Animal Science*, 8(2), 525-527.  
<https://doi.org/10.4081/ijas.2009.s2.525>

Nudda, A., McGuire, M. K., Battacone, G., Manca, M. G., Boe, R., Pulina, G. (2011). Documentation of fatty acid profiles in lamb meat and lamb-based infant foods. *Journal of Food Science*, 76(2), 43-47.  
<https://doi.org/10.1111/j.1750-3841.2010.02027.x>

Nuernberg, K., Fischer, A., Nuernberg, G., Ender, K., Dannenberger, D. (2008). Meat quality and fatty acid composition of lipids in muscle and fatty tissue of Skudde lambs fed grass versus concentrate. *Small Ruminant Research*, 74(1-3), 279-283. <https://doi.org/10.1016/j.smallrumres.2007.07.009>

Popova, T. (2014). Fatty acid composition of Longissimus dorsi and Semimembranosus muscles during storage in lambs reared indoors and on pasture. *Emirates Journal of Food and Agriculture*, 26(3), 302-308.  
<https://doi.org/10.9755/ejfa.v26i3.16771>

Potkanski, A., Čermák, B., Szumacher-Strabel, M., Kowalczyk, J., Cieslak, A. (2002). Effects of different amounts and types of fat on fatty acid composition of fat deposit in lambs. *Czech Journal Animal Science*, 47(2), 72-75.

Santos-Silva, J., Bessa, R. J. B., Santos-Silva, F. (2002). Effect of genotype, feeding system and slaughter weight on the quality of light lambs: II. Fatty acid composition of meat. *Livestock Production Science*, 77(2-3), 187-194. [https://doi.org/10.1016/S0301-6226\(02\)00059-3](https://doi.org/10.1016/S0301-6226(02)00059-3)

Scerra, M., Luciano, G., Caparra, P., Foti, F., Cilione, C., Giorgi, A., Scerra, V. (2011). Influence of stall finishing duration of Italian Merino lambs, raised on pasture on intramuscular fatty acid composition. *Meat Science*, 89(2), 238-242. <https://doi.org/10.1016/j.meatsci.2011.04.012>

Simopoulos, A. P. (2016). An increase in the omega-6/omega-3 fatty acid ratio increases the risk for obesity. *Nutrients*, 128(8), 1-17. <https://doi.org/10.3390.Fnu8030128>

Vasta, V., D'Alessandro, A. G., Priolo, A., Petrotos, K., Martemucci, G. (2012). Volatile compound profile of ewe's milk and meat of their suckling lambs in relation to pasture vs. indoor feeding system. *Small Ruminant Research*, 105(1-3), 16-21. <https://doi.org/10.1016/j.smallrumres.2012.02.010>

Willians, C. M. (2000). Dietary fatty acids human health. *Nutrition Animal*, 49(3), 165-180.  
<https://doi.org/10.1051/animres:2000116>

Wood, J. D., Enser, M., Fisher, A. V., Nute, G. R., Sheard, P. R., Richardson, R. I., Hughes, S. I., Whittington, F. M. (2008). Fat deposition, fatty acid composition and meat quality: A review. *Meat Science*, 78(4), 343-358.  
<https://doi.org/10.1016/j.meatsci.2007.07.019>

Zervas, G., Tsiplakou, E. (2011). The effect of feeding systems on the characteristics of products from small ruminants. *Small Ruminant Research*, 101, 140-149. <https://doi.org/10.1016/j.smallrumres.2011.09.034>