



# Termites host specific fungal communities that differ from those in their ambient environments

Author	Tomas Vetrovsky, Patrik Soukup, Petr Stiblik, Katerina Votypkova, Amrita Chakraborty, Inaki Odriozola Larranaga, David Sillam-Dusses, Nathan Lo, Thomas Bourguignon, Petr Baldrian, Jan Sobotnik, Miroslav Kolarik
journal or publication title	Fungal Ecology
volume	48
page range	100991
year	2020-09-29
Publisher	Elsevier Ltd British Mycological Society
Rights	This article/chapter was published in Fungal Ecology, Vetrovsky, Tomas Soukup, Patrik Stiblik, Petr Votypkova, Katerina Chakraborty, Amrita Larranaga, Inaki Odriozola Sillam-Dusses, David Lo, Nathan Bourguignon, Thomas Baldrian, Petr Sobotnik, Jan Kolarik, Miroslav, Termites host specific fungal communities that differ from those in their ambient environments, 100991, (C) Elsevier Ltd and British Mycological Society 2020.
Author's flag	author
URL	<a href="http://id.nii.ac.jp/1394/00001668/">http://id.nii.ac.jp/1394/00001668/</a>

doi: info:doi/10.1016/j.funeco.2020.100991

**Title: Termites host specific fungal communities that differ from those in their ambient environments**

**Running title: Termite body mycobiome**

**Authors:** Tomáš Větrovský<sup>a,b</sup>, Patrik Soukup<sup>a</sup>, Petr Stiblík<sup>a</sup>, Kateřina Votýpková<sup>a</sup>, Amrita Chakraborty<sup>a</sup>, Iñaki Odriozola Larrañaga<sup>b</sup>, David Sillam-Dussès<sup>c</sup>, Nathan Lo<sup>d</sup>, Thomas Bourguignon<sup>a,e</sup>, Petr Baldrian<sup>b</sup>, Jan Šobotník<sup>a</sup>, Miroslav Kolařík<sup>f,g</sup>

Addresses:

[a] Faculty of Forestry and Wood Sciences, Czech University of Life Sciences, Kamýcká 129, 165 21 Prague 6 Suchbát, Czech Republic

[b] Laboratory of Environmental Microbiology, Institute of Microbiology of the CAS, Vídeňská 1083, 14220 Praha 4, Czech Republic

[c] Laboratory of Experimental and Comparative Ethology, EA44443, University Paris 13, Sorbonne Paris Cité, 99 Avenue Jean-Baptiste Clément, 93430 Villetaneuse, France

[d] School of Life and Environmental Sciences, University of Sydney, Sydney, NSW 2006, Australia

[e] Okinawa Institute of Science & Technology Graduate University, 1919–1 Tancha, Onna-son, Okinawa 904–0495, Japan

[f] Laboratory of Fungal Genetics and Metabolism, Institute of Microbiology of the CAS, Vídeňská 1083, 14220 Praha 4, Czech Republic

[g] Department of Botany, Faculty of Science, Charles University, Benátská 2, 128 01, Prague 2, Czech Republic

**Corresponding author:** Miroslav Kolařík, fax: +420 241 062 347, phone: (+420) 29644 2332, E-mail: [mkolařík@biomed.cas.cz](mailto:mkolařík@biomed.cas.cz); Jan Šobotník, phone: (+420) 22438 3776, E-mail: [sobotnik@fld.czu.cz](mailto:sobotnik@fld.czu.cz)

## **Abstract**

Termites are important plant biomass decomposers. Their digestive activity typically relies on prokaryotes and protozoa present in their guts. In some cases, such as in fungus-growing termites, digestion also relies on ectosymbiosis with specific fungal taxa. To date, the mycobiome of termites has yet to be investigated in detail. We evaluated the specificity of whole-termite associated fungal communities in three wood-feeding termite species. We showed that the whole-termite fungal community spectra are stable over diverse environments, regardless of the host species, and differ markedly from the wood in which they nest. The core mycobiome is similar to that found in other ecologically related insects and consists of a narrow spectrum of common filamentous fungi and yeasts, known for their stress tolerance and their ability to decompose plant biomass. The observed patterns suggest that a number of fungal strains may have a symbiotic relationship with termites, and our results set the stage for future investigations into the interactions between fungi, termites, and their other gut microbiota.

**Keywords:** mycobiome; symbiosis; *Coptotermes*; *Heterotermes*; *Nasutitermes*; yeast; moulds

## Introduction

Dead plant materials are mostly made of lignocellulose, the most common polymer on Earth, which relatively few metazoan taxa are able to significantly decompose (Lo et al., 2003). All key taxa consuming dead plant tissues, such as ruminants, earthworms and insects, largely rely upon a rich microbial consortium, which possess the necessary metabolic pathways for lignocellulose decomposition (Watanabe and Tokuda 2010; Brune and Dietrich 2015). Termites process this cellulose far more efficiently than other decomposers (Brune 2014) and their dominance in tropical ecosystems makes them key players at a global scale (Bignell and Eggleton 2000; Bar-On et al., 2018; Griffiths et al., 2019). While early branching termite lineages (“lower” termites) feed exclusively on wood or grasses, the “higher” termites (i.e. the crown family Termitidae) consume a variety of plant materials irrespective of decomposition status, and a majority of these taxa are soil-feeders (Jouquet et al., 2006; Krishna et al., 2013). To digest cellulose, termites rely on their own endogenous cellulases (Watanabe et al. 1998), in combination with microbial cellulases in their guts. Lower termites depend primarily on flagellate protozoa, with some contribution by prokaryotes, while “higher” termites lack cellulolytic flagellates completely and depend on bacteria and archaea for cellulose decomposition (Brune and Ohkuma 2011; Hongoh 2011; Brune and Deitrich 2015).

In herbivorous or detritivorous insects, both prokaryotes and fungi are generally thought to form core taxa of the gut microbiome, despite the fact that most studies have considered prokaryotes only (Gurung et al., 2019; Ravenscraft et al., 2019). Both types of microorganism can act as nutritional symbionts assisting with digestion, detoxification and essential nutrients synthesis, or as protective symbionts (Dillon and Dillon 2004; Gurung et al., 2019). Other roles, such as the effect on the host cells physiology and interactions with other microbes can be expected, as is known in mammals (Lai et al., 2018). The core gut mycobiome of wood feeding insects covers a relatively narrow set of ubiquitous yeasts and filamentous fungi such as *Candida*, *Mucor*, *Aspergillus*, *Penicillium*, *Alternaria* or *Trichoderma* (Pérez et al., 2003; Rojas-Jiménez and Hernández 2015; Ziganshina et al. 2018) and the same taxon spectrum is reported not only in other insects (Moraes et al., 2001; Fredensborg et al., 2020), but also in mammals (Lai et al., 2018).

Associations between termites and fungi have so far been considered in two categories: firstly, interactions that affect the discovery and consumption of food or its nutrient value, but which fall short of mutualism; secondly, the cultivation of fungus-combs (*Termitomyces* spp.) by fungus farming Macrotermitinae (Lenz et al., 1991; Rouland-Lefèvre 2000). However, apart from a few studies reporting common yeasts and filamentous fungi, found in termite guts (Prillinger et al., 1996; Prillinger and König 2006), the fungi associated with termites are yet to be systematically investigated. The only studies which have compared the microbiota of termite guts and termite ambient environments showed that fungal assemblages of guts differ markedly from nest walls or food nodules in litter and humus feeding termites (Menezes et al., 2018; Moreira et al., 2018). Based on the current

knowledge of insect microbiomes, we hypothesised that fungi, which are a neglected part of the termite gut microbiome, form predictable communities and have stable interactions with their hosts.

We compared the specificity of body associated fungal communities (i.e. fungi in gut and on exoskeleton) in three ecologically similar species, *Heterotermes tenuis*, *Coptotermes testaceus* (both lower termites, Rhinotermitidae) and *Nasutitermes octopilis* (higher termites, Termitidae: Nasutitermitinae), which can be simultaneously collected from the same large wood item. We examined the mycobiomes of whole termite bodies as a proxy for termite gut mycobiomes, which enabled us to analyse large sample sizes, necessary for statistical testing. We hypothesized that fungal communities are similar in termites with a similar diet, and more alike in the genera *Heterotermes* and *Coptotermes* compared to *Nasutitermes*, as *Coptotermes* is nested within the genus *Heterotermes* (Bourguignon et al., 2016, Buček et al., 2019). We examined fungal communities using high-throughput sequencing of ITS2 metabarcodes of termite bodies, their food source (narrow termite galleries), and intact control wood near to areas where termites were feeding. The patterns described below are based on repeated samples from the same log, usually of multiple species from the same trunk, which allowed us to test for termite species and colony-level specificity of the associations.

## Material and Methods

### *Study site and sampling*

The samples were collected in November 2014 in Nouragues Nature reserve (French Guiana; N 04°05', W 52°41'). Large wood items were inspected for the presence of two “lower” termite species, *Coptotermes testaceus* (Rhinotermitidae) with a preference to sound white wood, *Heterotermes tenuis* (Rhinotermitidae) preferring red-rot wood, and one “higher” termite species, *Nasutitermes octopilis*, (Termitidae: Nasutitermitinae) having no clear specialisation to the wood-decomposition degree.

A single sample set comprised of three samples: (1) 10 workers from a single foraging party, (2) their feeding substrate (approx. 1 cm<sup>3</sup> piece of wood containing gallery) and (3) the control sample (approx. 1 cm<sup>3</sup> of wood roughly 10 cm away from the closest termite gallery) (Fig. 1). Two or three sample sets, collected 1 m away from each other, were taken from the single wood log. Visually healthy workers were collected and narrow termite galleries with minimal amounts of frass were selected. Samples were firstly stored in RNAlater® solution at -20 °C within 12 h following collection, and shipped to Prague where they were stored at -80 °C until DNA extraction. In total, 82 samples sets (*Coptotermes*: n = 28, *Heterotermes*: n = 31, *Nasutitermes*: n = 23) originated from 23 trunks were studied. Storage in RNA later ® solution caused hardening of termite bodies preventing gut dissection. Thus, as extraction of the intact intestine was impossible, we used whole termite bodies as a proxy for the study of intestinal microbiota.

### *DNA extraction and PCR amplification*

Total DNA was extracted using Macherey-Nagel NucleoSpin® Soil kit with following modifications. Each termite sample was homogenized together in 500 µL of SL1 Lysis buffer, 100 µL of SX enhancer buffer and two sterilized steel beads (3 mm diameter) using a Mixer Mill MM 400 for 2 min, set on 30 Hz. Sample lysis by using a vortex was shortened to 2 min. The wood samples were mechanically crushed to small pieces, placed in a 2 mL tube with a five steel beads, frozen in liquid nitrogen for 1 min and grinded in Mixer mill Retsch MM 400 for 10 min at 30 Hz. 550 µL SL2 of extraction buffer was added to homogenized material and the grinding was repeated once more. Sample lysis was extended to 10 min.

PCR amplification of the fungal ITS2 region from DNA was performed using gITS7 (50-GTGARTCATCGARTCTTTG-30) and ITS4 (50-TCCTCCGCTTATTGATATGC-30) (Ihrmark et al., 2012; Tedersoo et al., 2015), each of them was barcoded in three PCR reactions per sample. The PCR reactions contained 2.5 µL of 10× buffer for DyNAzyme II DNA Polymerase, 0.75 µL of bovine serum albumin (20 mg/mL), 1 µL of each primer (0.01 mM), 0.5 µL of PCR Nucleotide Mix (10 mM each), 0.75 µL of polymerase (2 U/µL DyNAzyme II DNA polymerase), and 1 µL of template DNA.

PCR was performed by using an Eppendorf Mastercycler® (Eppendorf AG, Hamburg, Germany) nexus cycler. The PCR cycling parameters were 94 °C – 5 min (1 cycle), 94 °C – 45 sec, 56 °C – 35 sec, 72 °C – 30 sec (40 cycles), final extension at 72 °C – 10 min. PCR triplicates were combined and purified using MinElute PCR Purification Kit (Qiagen GmbH, Hilden, Germany) according to provided manual and eluted in 20 µL. Paired-ends amplicon reads were sequenced on Illumina MiSeq sequencer (Illumina Inc., USA) using V2 chemistry producing 2 × 250 bp output.

### *Data processing*

Raw fungal ITS paired-end sequences were joined using fastq-join software (Aronesty 2011) and demultiplexed, filtered and trimmed using the pipeline SEED 2 (version 2.1.05) (Vetrovsky et al., 2018). Low-quality sequences (mean Phred quality score < 30) and all sequences with mismatches in barcodes were removed from the dataset. After the quality filtering, all fungal sequences were extracted from the joined sequences using ITSx (v 1.0.11) (Bengtsson-Palme et al., 2013) to acquire complete ITS2 region. All ITS2 sequences shorter than 40 bp were discarded, yielding a dataset of 3 967 992 fungal ITS2 sequences (length distribution 40-395 bp, avg. 175 bp). The dataset was clustered into operational taxonomic units (OTUs) using UPARSE implementation in USEARCH version 8.1.1861 (Edgar 2013) with 97% similarity threshold (109 476 fungal chimeric sequences were excluded during this step). A total of 10 742 fungal OTUs (without singletons) were obtained during the clustering step. To reduce the influence of contaminations and minimize the effects of barcode hopping all OTUs with up to 4 reads were discarded, what resulted in 2857 OTUs used for further analysis.

The most abundant sequence from each cluster was used as a representative sequence for taxonomic classification. Fungal sequences were classified based on BLAST best hit against UNITE database, version 7.2 (Koljalg et al., 2013). The functional guild of each fungal OTU was assigned based on the FUNGuild database (Nguyen et al., 2016). For alpha diversity estimation, all fungal samples were resampled to 909 sequences. Diversity indices were estimated using SEED 2 version 2.1.05. The abundances of sequence reads were plotted on the phylogenetic tree constructed using NCBI molecular data via phyloT (Letunic 2015) and the iTol visualisation tool (Letunic and Bork 2019). Data were deposited in the MG-RAST database under accession number mgp91984 and in NCBI Sequence Read Archive (SRA) under BioProject accession number PRJNA639228. Processes data (extracted ITS2 reads) were deposited in the GlobalFungi Database (<https://globalfungi.com>, Větrovský et al. 2020).

### *Statistical analysis*

To test the null hypothesis of no difference between termite body, gallery and wood (control) fungal community composition, PERMANOVA analysis (Anderson 2001) was performed with `adonis()` function of *vegan* package (Oksanen et al., 2018) in R (R Core Team, 2018). Euclidean distance on Hellinger-transformed fungal composition (i.e. Hellinger distance matrix) (Legendre and Gallagher 2001) was used as response matrix, and sample type (body, gallery or wood) was used as fixed explanatory variable. Since observations were paired within triplets (the three sample types were sampled in each triplet), which, in turn, were nested in logs, the permutations were constrained to occur within triplets, using the variable triplet as blocking factor (or strata). To visualize the results, non-metric multidimensional scaling (NMDS) was performed in two different ways. In the first way, raw community data was ordinated by their fungal composition. This NMDS plot shows all the variability in the dataset. In the second way, community data was first regressed against triplet and log effects (i.e. the effect of spatial variability due to the experimental design was removed from the data) and, then, the residualized distance matrix was ordinated using NMDS as suggested by Anderson et al. (2017). This plot shows the variability in the dataset, once the effect of triplet and log has been taken into account.

To test the null hypothesis of no effect of sample type and termite species on fungal diversity (measured with Chao 1 index, Shannon-Wiener diversity index, and Pielou's evenness), linear mixed effect models were fitted using the function `lme()` from the R package *nlme* (Pinheiro et al., 2018). The interaction between termite species and sample type was fitted as the fixed part of the model, and, a random structure of the form  $\sim 1|triplet/log$  was included in each model to account for the fact that measurements were grouped in triplets, which, in turn, were nested in logs. Tukey *post-hoc* tests were performed using the function `lsmeans()` of the R package *lsmeans* (Lenth 2016).

To identify the fungal OTUs contributing to the separation between termite bodies, galleries and wood, partial redundancy analysis (partial RDA) was used (Legendre and Legendre 2012) for each termite species separately. Separating the communities by termite species allowed checking whether similar OTUs contribute to the separation between sample types in the three termite species. In each RDA, Hellinger-transformed fungal OTU composition was used as response matrix, sample type was used as fixed explanatory factor, and the analysis was conditioned with the effect of the log and triplet. 1% of the OTUs with highest loadings to the ordination axes RDA1 and RDA2 in the three partial RDAs were depicted in triplots (Legendre and Legendre 2012).

Lastly, to test and quantify the effect of termite species and log identity on fungal mycobiome composition, variation partitioning was performed based on RDA (Legendre and Legendre 2012). Variation in Hellinger-transformed fungal OTU composition of termite bodies and galleries was partitioned in the effect of termite species and log identity. Since the number of body and gallery samples per species was not equal, *Coptotermes* and *Heterotermes* were randomly subsampled to balance the design, which makes the hypothesis testing more robust to the presence of heterogeneous group dispersions (Anderson and Walsh 2013). The partial effect of each fraction (i.e. the effect of a fraction –e.g. species– once the effect of the other fraction –e.g. log identity– has been taken into account) was tested using a permutation test in partial RDA results.

## Results

### *Fungal diversity*

The diversity of fungal OTUs was significantly higher in termite bodies of all three species than in their galleries and intact wood, and was also significantly different between termite species. The estimated number of OTUs (Chao-1 estimate) in termite body samples, counted from the resampled dataset, ranged from 26 to 221 with an average 92-101 per species. Estimated OTU numbers and diversity indices were at least two times lower in termite galleries and in control wood. The fungal communities from termite bodies were significantly more even than termite galleries and control wood samples (Fig. 2).

### *Fungal community composition*

The wood control and galleries were dominated by Basidiomycota followed by Ascomycota while there was an obvious shift to the dominance of Ascomycota over Basidiomycota in the termite bodies, with the addition of Mucoromycotina and Chytridiomycota members (Fig. 3A). A significant diversity of fungal taxa unidentified at the phylum level was recovered for all three treatments. At the finer taxonomic scale, 25 fungal orders were most abundant (Fig. 3B, 4). Among the most abundant orders, Mucoromycotina GS23 (artificial group, see Figure 3 for definition), Eurotiales, Hypocreales, Ophiostomatales and Saccharomycetales were typically associated with termite bodies, whereas



Chaetosphaeriales, Auriculariales and partially also Corticiales were associated with wood and galleries. Wood was also marked by the high abundance of Polyporales (Fig. 3, 4, Supplementary Table 1). Finally, the members of the order Hymenochaetales were abundant in all variants. The majority of the fungal taxa identified to the ecological guild were predicted to be saprotrophs, combination of mixed trophic modes (mostly saprotrophs and pathotrophs) and pathotrophs. Saprotrophs and pathotrophs were more abundant in termite bodies (frequency of reads in saprotrophs - 39-23%; pathotrophs - 5-11 %) than in galleries and wood (22-11%; 0.1 -3%). Those taxa belonged mostly to plant pathogens, with the small fraction of insect pathogenic fungi (0.05-0.25%) dominated by *Metarhizium* spp. and *Lecanicillium* spp. (Supplementary table 2).

Multivariate analysis of the raw OTU dataset did not clearly separate samples by their types, but showed that the intestinal mycobiota of all three termite species is rather homogenous and similar, in comparison to the very heterogeneous communities colonizing their galleries and wood controls (k = 3 dimensions, final stress = 0.24, Fig. 5A). By contrast, once the spatial variability due to the experimental design (i.e. the effect of log and triplet identity) is removed (k = 3 dimensions, final stress = 0.25, Fig. 5B), body samples clearly separate from galleries and controls. The NMDS revealed a high stress value indicating that 2D graphical representation only roughly corresponds with the underlying data. However, the observed patterns were confirmed by the PERMANOVA analysis which showed that fungal communities from termite bodies were significantly different from galleries and controls (permutations = 999, *P*-value = 0.001).

Constrained RDA analysis with the removed effect of the sampling design revealed a clear separation of samples based on their type in all three termite species. The first axis of RDA (RDA1) separates termite bodies from galleries and controls, whereas the second axis (RDA2) separates galleries from controls (Fig. 6). As opposed to the unconstrained ordination (NMDS, Fig. 5), the constrained ordination (Fig. 6) distinguishes between the fungal compositions of galleries and controls. OTU 12, 20 and 34 are consistently positively associated with termite bodies in all three species. A further 13 OTUs are associated with two termite species (Table 1). The fungal genera linked with termite bodies (i.e. with high negative RDA1 axis loadings, Fig. 6), in all three termite species includes a narrow spectrum of filamentous ascomycetes (*Trichoderma*, *Penicillium*, *Scytalidium*, *Hawksworthiomyces*, *Lasiodiplodia*), a few basidiomycete genera (*Malassezia*, *Phlebia*, *Hyphodontia*, *Corticium*, *Wrightoporia* etc.), a single but abundant taxon from Mucoromycotina and a chytrid species from the genus *Spizellomyces* specifically associated with *Coptotermes* and *Heterotermes* (Table 1, Supplementary Table 3). Fungal genera linked with galleries include mostly wood saprobes from Basidiomycota (*Resinicium*, *Hyphodontia*, and unidentified genera), the very abundant genus *Chaetosphaeria*, and other wood inhabiting ascomycetes (*Pseudolachnella*, *Orbilia*, *Calonectria*, etc.). Genera linked with wood were *Auricularia*, *Porothelium* and numerous, mostly unidentified, genera of Polyporales, Auriculariales and Agaricomycetes but also various wood rotting

ascomycota (*Hypoxylon*, *Kretzschmaria*, *Camarops*, *Cordana*, *Chaetosphaeria*) (Fig. 6, Supplementary table 3).

The fungal community composition of termite bodies and galleries was significantly affected by both termite species and log identity. Total explained variation in gallery mycobiomes (0.118) was more than the half (0.261) of the explained variation in body mycobiomes. Accordingly, the variations explained by only termite species, only log and the shared fraction (i.e. the fraction that cannot be clearly attributed to either species or log) were more than double in termite bodies compared with galleries (Fig. 7).

## Discussion

### *Termite associated mycobiome*

Fungal communities of galleries and intact overlapped and differed from termite communities, which also overlapped with each other (Figs 3-6). The termite mycobiota is likely to be a mix of fungi present on insect cuticular tissues (mostly from the mouthparts and pathogenic fungi present on the body surface), fungi present in the gut, and possibly fungi present internally in other organs or in the haemolymph. In our study, the fraction of insect pathogenic fungi was higher in termites than in their galleries, but their overall abundance was very low and did not contribute to the separation of the studied sample types (Table 1, Supplementary Table 2). Intestinal fungi appear to dominate the termite mycobiome. Whether fungi occur in other internal organs (i.e. haemolymph, gonad rudiments) is unclear. The presence of fungi on termite exterior cuticles could potentially reduce the differences between the termite and gallery communities, due to the fact that termites are in close contact with their galleries. Despite this limitation, we found statistically significant differences between both communities.

In our study, representatives of the Saccharomycetales, Malasseziales, Eurotiales, Hypocreales and Mucoromycota common in whole termites, and much less frequent in galleries and wood (Fig. 3, Supplementary Table 1), can be considered as typical members of termite mycobiome. Two previous studies quantified termite associated fungi using ITS metabarcoding. They found Eurotiales, Trichosphaeriales and Pleosporales (Menezes et al. 2018) together with Hypocreales (Moreira et al. 2018) are associated with guts and much less abundant in surrounding environments. This is in line with our results, including the presence of Trichosphaeriales and Pleosporales, which were rare in our study, but typically present in termite bodies (Fig. 3, Supplementary Table 1).

Yeasts, i.e. species from Saccharomycetales, Malasseziales and Trichosporonales, are the best studied fungi in “lower” termites (Prillinger et al., 1996; Prillinger and König 2006) and the insect gut in general (Blackwell 2017; Stefanini 2018). Genera frequently found in our study, *Candida*, *Debaryomyces*, *Pichia*, *Cryptococcus*, and *Trichosporon*, are known as typical termite gut inhabitants (Prillinger et al., 1996; Prillinger and König 2006). At the species level (i.e. OTUs with  $\geq 99$  %

similarity) we identified several taxa already known as intestinal symbionts of various insects (e.g. *Candida haemulonis*, *C. parapsilosis* (Suh et al., 2007; Bozic et al., 2017), *C. elateridarum* (Suh and Blackwell 2004), *Malassezia restricta* (Zhang et al., 2003), *Metschnikowia pulcherrima* (Woolfolk and Inglis 2004) and *Trichosporon insectorum* (Fuentefria et al., 2008) (Supplementary Table 1). Surprisingly, yeasts (with the exception of *Malassezia*) did not contribute to the statistical separation of gut and gallery associated fungal communities, when the effect of sampling design was removed (Fig. 6). This was partially due to the high inter-sample variability of yeast communities, but also because of their consistent occurrence (although in very low abundances) in the galleries.

The statistical separation of the whole termite mycobiota in our study was mostly due to the differences among representatives of ubiquitous genera of plant endophytes and saprobes including *Mucoromycotina* spp., *Trichoderma*, *Hawksworthiomyces* and *Penicillium* (Table 1). Data on termite gut associated filamentous fungi are scarce (for review see König et al., 2006; Prillinger and König 2006). The genera *Trichoderma*, *Penicillium*, *Aspergillus* and *Alternaria* (Hendee 1935; Rajagopal et al., 1979, 1981; Varna et al., 1994; Jayasimha and Henderson 2007), together with numerous *Mucoromycotina* spp. (Zoberi et al. 1990), were already reported from termite guts, which corresponds to our results (Table 1). Interestingly, a similar spectrum of genera (i.e. *Penicillium*, *Trichoderma*, *Fusarium*, *Cladosporium*, *Aspergillus*, *Rhizopus*, and *Mucor*) is also present in the guts of other plant biomass feeders such as isopods (Kayang et al., 1996), *Tenebrio molitor* (Fredensborg et al., 2020), wood feeding beetles (Rojas-Jiménez and Hernández 2015; Mohammed et al., 2018; Ziganshina et al., 2018), cockroaches (Salehzadeh et al., 2007), bark beetles (Perez et al., 2003) and grasshoppers (Idowu et al., 2009). It is also worth mentioning the abundant presence of *Spizellomyces* sp., identified in more than 50% of *C. testaceus* and *H. tenuis* samples, but absent in *N. octopilis*. *Spizellomyces* is a genus of zoosporic fungi living in soil, or as plant pollen, or fungal parasites (Wakefield et al., 2010), and its association with termites calls for further studies.

Although previous works studied the presence of various fungi in termites (Prillinger et al., 1996; König et al., 2006; Jayasimha and Henderson 2007; Santana et al. 2015; Menezes et al. 2018; Moreira et al. 2018), our study is the first large and systematic comparison targeting the fungal communities using a statistically robust dataset of termites, their environments, and their feeding substrates. In agreement with previous studies (König et al., 2006; Prillinger and König 2006), our results indicate that the termite intestine is inhabited by ubiquitous environmental fungi. We showed that the termite associated community is distinct and relatively homogenous and stable over diverse environments and termite species, compared with termite galleries and control wood. In addition, termite galleries represent a specific habitat which significantly differs from wood in fungal community composition. This is similar to results found in studies of humus and litter feeding termites, where intestinal fungal communities differed substantially from communities of feeding

nodules and/or nest walls (Menezes et al. 2018, Moreira et al. 2018) suggesting that termites possessed a host-defined intestinal mycobiome.

Strong environmental filtering appears to allow a relatively small number of fungi to grow freely and persist inside termites, across different species, after being taken up from the environment (soil galleries used for foraging, or the wood upon which termites feed). Vertical transmission of fungi by termites may also occur, although our results do not provide a clear answer on this. Both modes of symbiont acquisition, or their combination, can result in the observed stability of the intestinal communities across various collection dates, termite populations and species. In addition, the galleries themselves host specific fungal communities, which are more similar to intact wood and less affected by the termite species that form the galleries (Fig. 6, 7).

Interestingly, the effect of termite species on fungal community composition in whole termites was very low (Fig. 7), which shows that different termites shape their fungal communities in a similar way. This is in contrast to patterns found in bacteria (Colman et al., 2012; Bourguignon et al., 2018; Chouvenec et al., 2018; Menezes et al., 2018; Moreira et al., 2018), which are more host specific. This is partly explained by the fact that many termite-associated bacteria are highly co-evolved vertically transmitted obligate symbionts (of termites or associated protists), whereas most identified fungi are presumably facultative associates, frequently existing as environmental fungi. Higher OTU diversity in whole termites in comparison to galleries and control wood is another feature constantly shared among different termite species (Fig. 2). This pattern is expected if we consider that the intestine itself is highly compartmentalised, which results in an increase in microbial diversity (Mikaelyan et al., 2017).

The core mycobiome of the termite gut is composed of plant biomass decomposers (see below), which are stress tolerant, fast growing and sporulating. It is likely that they are pre-disposed to live in the environment of the termite gut, which is characterized by harsh microaerobic conditions, steep gradients of oxygen and hydrogen, and activity of strong hydrolytic enzymes (König et al., 2006). Furthermore, it is possible that such features allow these fungi to live not only in termites, but also in taxonomically distant insect plant biomass feeders. The apparent stability of the fungal community between different termites could be considered evidence for symbiosis. Although it appears likely that the fungi we identified are able to live and grow in the termite gut, it is also possible that the origin of some strains is from the digested material, but were nevertheless picked up using our methods.

#### *Ecological role of gut associated fungi*

Both the presence and the ecological role of fungi in the termite gut have been poorly studied so far. However, fungi have generally not been considered as an important part of the termite holobiont (Slaytor 1992; Brune and Dietrich 2015; Peterson and Scharf 2016) and their presence is usually

ignored. It has been shown that the intestinal fungi were not essential for *Nasutitermes exitiosus* survival in the lab experiments (Eutick et al., 1978), but similar studies in other termite species are needed to confirm their facultative status. More insight into their ecological functions has been provided with the transcriptome data. The fungal contigs from nymphoid neotenic (i.e. the developmental stage fed mostly by proteinaceous labial gland secretions) intestinomes in *Reticulitermes* spp. represent 10.2% of the fraction of non-termite origin (Dedeine et al., 2015). Another study showed that 18% of all carbohydrate-active enzymes in *Coptotermes formosanus* transcriptomes were of fungal origin, similar to bacteria (24%) but not to protists (6% only) (Zhang et al., 2012). In *Reticulitermes flavipes* symbiont libraries (gut content only), fungi represent 7% of the non-animal fraction of the reads (protists 71%, prokaryotes 21%) (Tartar et al., 2009). Little is known concerning fungal gut biomass. In the termites *Zootermopsis angusticollis* and *Neotermes castaneus*,  $10^7$ - $10^8$  yeast cells per millilitre of gut content were found, which is comparable to the number of flagellates, and similar, or two orders of magnitude lower, than the numbers found in bacteria (König et al., 2006, Prillinger and König 2006). This suggests that fungi may actively proliferate in termite guts and they might be an important part of the lignocellulolytic machinery as proposed by Zhang et al. (2012).

Ecologically, yeast are typical inhabitants of the insect gut, including termites (Blackwell 2017; Stefanini 2018), and they can extracellularly decompose cellulose, hemicellulose and xylans, thus contributing to wood digestion (Prillinger and König 2006; Schäfer et al., 1996). Interestingly, the dominant fungal strains identified in our study, especially *Trichoderma* and *Penicillium*, are well known for their ability to degrade cellulose, hemicellulose, and lignin, and are often used in biotechnology (de França Passos et al., 2018). Significant lignocellulosic activities have also been reported in *Phlebia*, *Hyphodontia*, *Scytalidium* (Eriksson et al., 2012), *Hawksworthiomyces* (De Beer et al., 2016) and *Lasiodiplodia* (Félix et al., 2018). Such strong enzymatic activities were shown *in vitro* directly in the strains from termites (Tarayre et al., 2015). This information, together with published transcriptomic data, reinforce the idea that, in termites, fungi may contribute to the degradation of lignocellulose and hemicellulose (Tartar et al., 2009). In addition, detoxification ability, which is well known in fungi, was also found in the yeasts from termites (Molnar et al., 2004) and therefore toxin degradation could be another important role of the intestinal fungal symbionts. However, further characterisation of the real contribution of fungi to food-processing in termites still remains to be undertaken.

## Acknowledgements

We are grateful to the Agence Nationale de la Recherche Investissement d'Avenir for the Nouragues travel grant which allows us to collect the samples in French Guiana, to the Czech Science Foundation

(project no. 16-05318S), and Czech University of Life Sciences Prague (project CIGA 20184303). NL is supported by an Australian Research Council Future Fellowship.

## Reference

- Anderson, M.J., 2001. A new method for non-parametric multivariate analysis of variance. *Austral Ecol.* 26, 32-46. <https://doi.org/10.1111/j.1442-9993.2001.01070.pp.x>
- Anderson, M.J., Walsh, D.C., 2013. PERMANOVA, ANOSIM, and the Mantel test in the face of heterogeneous dispersions: what null hypothesis are you testing? *Ecol. Monogr.* 83, 557-574. <https://doi.org/10.1890/12-2010.1>
- Anderson, T.R., Pond, DW, Mayor DJ, 2017. The role of microbes in the nutrition of detritivorous invertebrates: A stoichiometric analysis. *Front. Microbiol.* 7, 2113. <https://doi.org/10.3389/fmicb.2016.02113>
- Aronesty, E., 2011. ea-utils: Command-line tools for processing biological sequencing data, 1.1.2 ed. Expression Analysis. Durham, NC. Available online at: <http://code.google.com/p/ea-utils>
- Bar-On YM, Phillips R, Milo R, 2018. The biomass distribution on Earth. *Proc. Nat. Acad. Sci.* 115, 6506-6511. <https://doi.org/10.1073/pnas.1711842115>
- Bengtsson-Palme, J., Ryberg, M., Hartmann, M., Branco, S., Wang, Z., Godhe, A., De Wit, P., Sanchez-Garcia, M., Ebersberger, I., de Sousa, F., Amend, A.S., Jumpponen, A., Unterseher, M., Kristiansson, E., Abarenkov, K., Bertrand, Y.J.K., Sanli, K., Eriksson, K.M., Vik, U., Veldre, V., Nilsson, R.H., 2013. Improved software detection and extraction of ITS1 and ITS2 from ribosomal ITS sequences of fungi and other eukaryotes for analysis of environmental sequencing data. *Methods Ecol. Evol.* 4, 914-919. <https://doi.org/10.1111/2041-210X.12073>
- Bignell, D.E., Eggleton, P., 2000. Termites in ecosystems, in: Abe, Y., Edward Bignell, D., Higashi, T. (Eds.), *Termites: evolution, sociality, symbioses, ecology*. Kluwer Academic, Massachusetts, pp. 363-387.
- Blackwell, M., 2017. Yeasts in insects and other invertebrates, in: Buzzini, P., Lachance, M.A., Yurkov, A. (Eds.), *Yeasts in natural ecosystems: diversity*. Springer, pp. 397–433. Cham. <https://doi.org/10.1007/978-3-319-62683-3>
- Bourguignon, T., Lo, N., Dietrich, C., Šobotník, J., Sidek, S., Roisin, Y., Brune, A., Evans, T.A., 2018. Rampant host switching shaped the termite gut microbiome. *Curr. Biol.* 28, 649-654.e642. <https://doi.org/10.1016/j.cub.2018.01.035>
- Bozic, J., Capone, A., Pediconi, D., Mensah, P., Cappelli, A., Valzano, M., Mancini, M.V., Scuppa, P., Martin, E., Epis, S., 2017. Mosquitoes can harbour yeasts of clinical significance and contribute to their environmental dissemination. *Environ. Microbiol. Rep.* 9, 642-648. <https://doi.org/10.1111/1758-2229.12569>

- Brune, A., 2014. Symbiotic digestion of lignocellulose in termite guts. *Nat. Rev. Microbiol.* 12, 168-180. <https://doi.org/10.1038/nrmicro3182>
- Brune, A., Dietrich, C., 2015. The gut microbiota of termites: digesting the diversity in the light of ecology and evolution. *Annu. Rev. Microbiol.* 69, 145-166. <https://doi.org/10.1146/annurev-micro-092412-155715>
- Brune, A., Ohkuma, M., 2011. Role of the termite gut microbiota in symbiotic digestion, in: Bignell, D. E., Roisin, Y., Lo, N. (Eds.), *Biology of termites: a modern synthesis*. Springer, Dordrecht, the Netherlands. pp. 439-475.
- Chouvenc, T., Elliott, M.L., Sobotnik, J., Efstathion, C.A., Su, N.Y., 2018. The termite fecal nest: a framework for the opportunistic acquisition of beneficial soil *Streptomyces* (Actinomycetales: Streptomycetaceae). *Environ. Entomol.* 47, 1431-1439. <https://doi.org/10.1093/ee/nvy152>
- Colman, D.R., Toolson, E.C., Takacs-Vesbach, C., 2012. Do diet and taxonomy influence insect gut bacterial communities? *Mol. Ecol.* 21, 5124-5137. <https://doi.org/10.1111/j.1365-294X.2012.05752.x>
- de França Passos, D., Pereira, N., de Castro, A.M., 2018. A comparative review of recent advances in cellulases production by *Aspergillus*, *Penicillium* and *Trichoderma* strains and their use for lignocellulose deconstruction. *Curr. Opin. Green Sustain. Chem.* 14: 60-66 <https://doi.org/10.1016/j.cogsc.2018.06.003>
- Dedeine, F., Weinert, L.A., Bigot, D., Josse, T., Ballenghien, M., Cahais, V., Galtier, N., Gayral, P., 2015. Comparative analysis of transcriptomes from secondary reproductives of three *Reticulitermes* termite species. *Plos One* 10(12): e0145596. <https://doi.org/10.1371/journal.pone.0145596>
- Dillon, R., Dillon, V., 2004. The gut bacteria of insects: nonpathogenic interactions. *Ann. Rev. Entomol.* 49, 71-92. <https://doi.org/10.1146/annurev.ento.49.061802.123416>
- Edgar, R.C., 2013. UPARSE: highly accurate OTU sequences from microbial amplicon reads. *Nat. Methods* 10, 996-998. <https://doi.org/10.1038/nmeth.2604>
- Eggleton P, 2006. The termite gut habitat: its evolution and co-evolution, in: König, H., Varma, A. (Eds.), *Intestinal microorganisms of termites and other invertebrates*. Springer-Verlag Berlin Heidelberg, pp. 373-404. [https://doi.org/10.1007/3-540-28185-1\\_16](https://doi.org/10.1007/3-540-28185-1_16)
- Eutick, M., Veivers, P., O'brien, R., Slaytor, M., 1978. Dependence of the higher termite, *Nasutitermes exitiosus* and the lower termite, *Coptotermes lacteus* on their gut flora. *J. Insect Physiol.* 24, 363-368. [https://doi.org/10.1016/0022-1910\(78\)90076-8](https://doi.org/10.1016/0022-1910(78)90076-8)
- Fredensborg, B.L., Fossdal í Kálvalíð, I., Johannesen, T.B., Stensvold, C.R., Nielsen, H.V., Kapel, C.M., 2020. Parasites modulate the gut-microbiome in insects: A proof-of-concept study. *Plos one* 15, e0227561. <https://doi.org/10.1371/journal.pone.0227561>

- Fuentefria, A.M., Suh, S.-O., Landell, M.F., Faganello, J., Schrank, A., Vainstein, M.H., Blackwell, M., Valente, P., 2008. *Trichosporon insectorum* sp. nov., a new anamorphic basidiomycetous killer yeast. *Mycol. Res.* 112, 93-99. <https://doi.org/10.1016/j.mycres.2007.05.001>
- Griffiths, H. M., Ashton, L. A., Evans, T. A., Parr, C. L., & Eggleton, P., 2019. Termites can decompose more than half of deadwood in tropical rainforest. *Curr. Biol.* 29, R118-R119. <https://doi.org/10.1016/j.cub.2019.01.012>
- Gurung, K., Wertheim, B., Falcao Salles, J., 2019. The microbiome of pest insects: it is not just bacteria. *Entomol. Experimental. Appl.* 167, 156-170. <https://doi.org/10.1111/eea.12768>
- Hendee, E.C., 1935. The role of fungi in the diet of the common damp-wood termite, *Zootermopsis angusticollis*. *Hilgardia* 9, 429-525. <https://doi.org/10.3733/hilg.v09n10p499>
- Hongoh, Y., 2011. Toward the functional analysis of uncultivable, symbiotic microorganisms in the termite gut. *Cell. Mol. Life Sci.* 68, 1311-25. doi:<http://dx.doi.org/10.1007/s00018-011-0648-z>
- Idowu, A., Edema, M., Oyedepo, M., 2009. Extracellular enzyme production by microflora from the gut region of the variegated grasshopper *Zonocerus variegatus* (Orthoptera: Pyrgomorphidae). *Int. J. Trop. Insect Sc.* 29, 229-235. <https://doi.org/10.1017/S1742758409990312>
- Ihrmark, K., Bodeker, I.T., Cruz-Martinez, K., Friberg, H., Kubartova, A., Schenck, J., Strid, Y., Stenlid, J., Brandstrom-Durling, M., Clemmensen, K.E., Lindahl, B.D., 2012. New primers to amplify the fungal ITS2 region--evaluation by 454-sequencing of artificial and natural communities. *FEMS Microbiol. Ecol.* 82, 666-677. <https://doi.org/10.1111/j.1574-6941.2012.01437.x>
- Jayasimha, P., Henderson, G., 2007. Fungi isolated from integument and guts of *Coptotermes fonnosanus* and their antagonistic effect on *Gleophyllum trabeum*. *Ann. Entomol. Soc. Am.* 100, 703-710. [https://doi.org/10.1603/0013-8746\(2007\)100\[703:FIFIAG\]2.0.CO;2](https://doi.org/10.1603/0013-8746(2007)100[703:FIFIAG]2.0.CO;2)
- Jouquet, P., Dauber, J., Lagerlof, J., Lavelle, P., Lepage, M., 2006. Soil invertebrates as ecosystem engineers: Intended and accidental effects on soil and feedback loops. *Appl. Soil Ecol.* 32, 153-164. <https://doi.org/10.1016/j.apsoil.2005.07.004>
- Kayang, H., Sharma, G.D, Mishra, R.R., 1996. The influence of isopod grazing on microbial dynamics in decomposing leaf litter of *Alnus nepalensis* D. Don. *Europ. J. Soil Biol.* 32, 35-39.
- Koljalg, U., Nilsson, R.H., Abarenkov, K., Tedersoo, L., Taylor, A.F., Bahram, M., Bates, S.T., Bruns, T.D., Bengtsson-Palme, J., Callaghan, T.M., Douglas, B., Drenkhan, T., Eberhardt, U., Duenas, M., Grebenc, T., Griffith, G.W., Hartmann, M., Kirk, P.M., Kohout, P., Larsson, E., Lindahl, B.D., Lucking, R., Martin, M.P., Matheny, P.B., Nguyen, N.H., Niskanen, T., Oja, J., Peay, K.G., Peintner, U., Peterson, M., Poldmaa, K., Saag, L., Saar, I., Schussler, A., Scott, J.A., Senes, C., Smith, M.E., Suija, A., Taylor, D.L., Telleria, M.T., Weiss, M., Larsson, K.H., 2013. Towards a unified paradigm for sequence-based identification of fungi. *Mol. Ecol.* 22, 5271-5277. <https://doi.org/10.1111/mec.12481>




- König, H., Fröhlich, J., Hertel, H., 2006. Diversity and lignocellulolytic activities of cultured microorganisms, in: König, H., Varma, A. (Eds.), *Intestinal microorganisms of termites and other invertebrates*. Springer. pp. 271-301. <https://doi.org/334>. 10.1007/3-540-28185-1
- Krishna, K., Grimaldi, D.A., Krishna, V., Engel, M.S., 2013. *Treatise on the Isoptera of the World Introduction*. B. Am. Mus. Nat. Hist. 377, 1-200. <https://doi.org/10.1206/377.1>.
- Lai, G.C., Tan, T.G., Pavelka, N., 2019. The mammalian mycobiome: a complex system in a dynamic relationship with the host. *Wiley Interdisciplinary Reviews: Systems Biol. Medicine* 11, e1438. <https://doi.org/10.1002/wsbm.1438>
- Legendre, P., Gallagher, E.D., 2001. Ecologically meaningful transformations for ordination of species data. *Oecologia* 129, 271-280. <https://doi.org/10.1007/s004420100716>
- Legendre, P., Legendre, L.F., 2012. *Numerical ecology*. Elsevier.
- Lenth, R.V., 2016. Least-squares means: the R package lsmeans. *J. Stat. Softw.* 69, 1–33. <https://doi.org/10.18637/jss.v069.i01>
- Lenz, M., Amburgey, T.L., Dai, Zi-Rong, Mauldin, J.K., Preston, A.F., Rudolph, D., Williams, E.R., 1991. Interlaboratory studies on termite wood decay fungi associations. 2. Response of termites to *Gloeophyllum trabeum* grown on different species of wood (Isoptera: Mastotermitidae, Termopsidae, Rhinotermitidae, Termitidae). *Sociobiol.* 18:, 203-254. <http://hdl.handle.net/102.100.100/253965?index=1>
- Letunic, I., Bork, P., 2019. Interactive Tree Of Life (iTOL) v4: recent updates and new developments. *Nucl. Acid. Res.* 47, W256–W259. <https://doi.org/10.1093/nar/gkz239>
- Lo, N., Eggleton, P., 2011. Termite phylogenetics and co-cladogenesis with symbionts. in: Bignell, D.E., Roisin, Y., Lo, N. (Eds.), *Biology of termites: a modern synthesis*. Springer, Dordrecht. pp. 27-50. <https://doi.org/10.1007/978-90-481-3977-4>
- Lo, N., Watanabe, H., Sugimura, M., 2003. Evidence for the presence of a cellulase gene in the last common ancestor of bilaterian animals. *P. Roy. Soc. B-Biol. Sci.* 270, S69-S72. <https://doi.org/10.1098/rsbl.2003.0016>
- Menezes, L., Alvarez, T.M., Persinoti, G.F., Franco, J.P., Squina, F., Moreira, E.A., Paixão, D.A.A., Costa-Leonardo, A.M., da Silva, V.X., Clerici, M.T.P.S., 2018. Food storage by the savanna termite *Cornitermes cumulans* (Syntermitinae): A strategy to improve hemicellulose digestibility? *Microb. Ecol.* 76: 492-505. <https://doi.org/10.1007/s00248-017-1128-2>
- Mikaelyan, A., Meuser, K., Brune, A., 2017. Microenvironmental heterogeneity of gut compartments drives bacterial community structure in wood- and humus-feeding higher termites. *FEMS Microb. Ecol.* 93, fiw210 <https://doi.org/10.1093/femsec/fiw210>
- Mohammed, W.S., Ziganshina, E.E., Shagimardanova, E.I., Gogoleva, N.E., Ziganshin, A.M., 2018. Comparison of intestinal bacterial and fungal communities across various xylophagous beetle larvae (Coleoptera: Cerambycidae). *Sci. Rep.* 8, 10073. <https://10.1038/s41598-018-27342-z>

- Molnar, O., Schatzmayr, G., Fuchs, E., Prillinger, H., 2004. *Trichosporon mycotoxinivorans* sp. nov., a new yeast species useful in biological detoxification of various mycotoxins. *Syst. Appl. Microbiol.* 27, 661-671. <https://doi.org/10.1078/0723202042369947>
- Moraes, A., Junqueira, A., Costa, G., Celano, V., Oliveira, P., Coura, J., 2001. Fungal flora of the digestive tract of 5 species of triatomines vectors of *Trypanosoma cruzi*, Chagas 1909. *Mycopathologia* 151, 41-48. <https://doi.org/10.1023/A:1010905420001>
- Moreira, E.A., Alvarez, T.M., Persinoti, G.F., Paixão, D.A.A., Menezes, L.R., Cairo, J.P.F., Squina, F.M., Costa-Leonardo, A.M., Carrijo, T., Arab, A., 2018. Microbial communities of the gut and nest of the humus-and litter-feeding termite *Procornitermes araujoi* (Syntermitinae). *Curr. Microb.* 75, 1609-1618. <https://doi.org/10.1007/s00284-018-1567-0>
- Nguyen, N., Song, Z., Bates, S., Branco, S., Tedersoo, L., Menke, J., Schilling, J., Kennedy, P., 2016. FUNGuild: an open annotation tool for parsing fungal community datasets by ecological guild. *Fungal Ecol.* 20, 241–248. <https://doi.org/10.1016/j.funeco.2015.06.006>
- Oksanen, J., Blanchet, F., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P., O'Hara, R., Simpson, G., Solymos, P., 2018. *Vegan: Community Ecology Package*. R package version 2.5-2.
- Pérez, J., Infante, F., HOLGUÍN, F., MACÍAS, J., Valle, J., Nieto, G., Peterson, S.W., Kurtzman, C.P., O'donnell, K., 2003. Mycobiota associated with the coffee berry borer (*Hypothenemus hampei*) in Mexico. *Mycol. Res.* 107, 879-887. <https://doi.org/10.1017/S0953756203007986>
- Peterson, B.F., Scharf, M.E., 2016. Lower termite associations with microbes: synergy, protection, and interplay. *Front. Microb.* 7, 422. <https://doi.org/10.3389/fmicb.2016.00422>
- Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., R Core Team, 2018. nlme: Linear and nonlinear mixed effects models. R package version 3.1-137, URL:<https://CRAN.R-project.org/package=nlme>.
- Prillinger, H., König, H., 2006. The intestinal yeasts, in: König, H., Varma, A. (Eds.), *Intestinal microorganisms of termites and other invertebrates*. Springer-Verlag Berlin Heidelberg, pp. 319-334. <https://doi.org/10.1007/3-540-28185-1>
- Prillinger, H., Messner, R., König, H., Bauer, R., Lopandic, K., Molnar, O., Dangel, P., Weigang, F., Kirisits, T., Nakase, T., Sigler, L., 1996. Yeasts associated with termites: a phenotypic and genotypic characterization and use of coevolution for dating evolutionary radiations in asco- and basidiomycetes. *Syst. Appl. Microbiol.* 19, 265-283. [https://doi.org/10.1016/S0723-2020\(96\)80053-1](https://doi.org/10.1016/S0723-2020(96)80053-1)
- Rajagopal, S., Rao, D.R., Varma, A.K., 1979. Association of fungi in the termite gut. *Curr. Sci.* 48, 998–999.
- Rajagopal, S., Rao, D.R., Varma, A.K., 1981. Fungi from worker termite gut, *Odontotermes obesus* (Rambur) from northern India. *Nova Hedwigia* 34, 97–100.


- Ravenscraft, A., Berry, M., Hammer, T., Peay, K., Boggs, C., 2019. Structure and function of the bacterial and fungal gut microbiota of Neotropical butterflies. *Ecol. Monogr.* 89, e01346 <https://doi.org/10.1002/ecm.1346>
- Rojas-Jiménez, K., Hernández, M., 2015. Isolation of fungi and bacteria associated with the guts of tropical wood-feeding coleoptera and determination of their lignocellulolytic activities. *Int. J. of Microbiol.* 2015:285018. <https://doi.org/10.1155/2015/285018>
- Rouland-Lefèvre, C., 2000. Symbiosis with fungi, in: Abe, T., Bignell, D.E., Higashi, M. (Eds.) *Termites: evolution, sociality, symbioses, ecology.* Springer, Dordrecht, pp 289-306. [https://doi.org/10.1007/978-94-017-3223-9\\_14](https://doi.org/10.1007/978-94-017-3223-9_14)
- Salehzadeh, A., Tavacol, P., Mahjub, H., 2007. Bacterial, fungal and parasitic contamination of cockroaches in public hospitals of Hamadan, Iran. *J. Vector Borne Dis.* 44,105-10..
- Santana, R.H., Catao, E.C., Lopes, F.A., Constantino, R., Barreto, C.C., Kruger, R.H., 2015. The gut microbiota of workers of the litter-feeding termite *Syntermes wheeleri* (Termitidae: Syntermitinae): archaeal, bacterial, and fungal communities. *Microb. Ecol.* 70, 545-556. <https://doi.org/10.1007/s00248-015-0581-z>
- Schäfer, A., Konrad, R., Kuhnigk, T., Kämpfer, P., Hertel, H., König, H., 1996. Hemicellulose-degrading bacteria and yeasts from the termite gut. *J. Appl. Bacteriol.* 80, 471-478. <https://doi.org/10.1111/j.1365-2672.1996.tb03245.x>
- Slaytor, M., 1992. Cellulose digestion in termites and cockroaches: what role do symbionts play? *Comp. Biochem. Physiol. B., Comp. Biochem* 103, 775-784. [https://doi.org/10.1016/0305-0491\(92\)90194-V](https://doi.org/10.1016/0305-0491(92)90194-V)
- Stefanini, I., 2018. Yeast-insect associations: It takes guts. *Yeast* 35, 315-330. <https://doi.org/10.1002/yea.3309>
- Suh, S.-O., Blackwell, M., 2004. Three new beetle-associated yeast species in the *Pichia guilliermondii* clade. *FEMS Yeast Res.* 5, 87-95. <https://doi.org/10.1016/j.femsyr.2004.06.001>
- Suh, S.-O., Nguyen, N.H., Blackwell M, 2007. Yeasts isolated from plant-associated beetles and other insects: seven novel *Candida* species near *Candida albicans*. *FEMS Yeast Res.* 8, 88-102. <https://doi.org/10.1111/j.1567-1364.2007.00320.x>
- Tarayre, C., Bauwens, J., Brasseur, C., Mattéotti, C., Millet, C., Guiot, P.A., Destain, J., Vandenberg, M., Portetelle, D., De Pauw, E., Haubruge, E., Francis, F., Thonart, P., 2015. Isolation and cultivation of xylanolytic and cellulolytic *Sarocladium kiliense* and *Trichoderma virens* from the gut of the termite *Reticulitermes santonensis*. *Environ. Sci. Pollut. R.* 22, 4369-4382. <https://doi.org/10.1007/s11356-014-3681-2>
- Tartar, A., Wheeler, M.M., Zhou, X., Coy, M., Boucias, D., Scharf, M., 2009. Parallel metatranscriptome analyses of host and symbiont gene expression in the gut of the termite *Reticulitermes flavipes*. *Biotechnol. Biofuels* 2, 25. <https://doi.org/10.1186/1754-6834-2-25>

- R Core Team, 2018. R: A Language and Environment for Statistical Computing.
- Tedersoo, L., Anslan, S., Bahram, M., Põlme, S., Riit, T., Liiv, I., Kõljalg, U., Kisand, V., Nilsson, R.H., Hildebrand, F., 2015. Shotgun metagenomes and multiple primer pair-barcode combinations of amplicons reveal biases in metabarcoding analyses of fungi. *MycoKeys*, 1-43. <https://doi.org/10.3897/mycokeys.10.4852>
- Varma, A., Kollu, B.K., Paul, J., Saxena, S., König, H., 1994. Lignocellulose degradation by microorganisms from termite hills and termite guts: A survey on the present state of art. *FEMS Microbiol. Rev.* 15, 9-28. <https://doi.org/10.1111/j.1574-6976.1994.tb00120.x>
- Větrovský, T., Baldrian, P., Morais, D., 2018. SEED 2: a user-friendly platform for amplicon high-throughput sequencing data analyses. *Bioinformatics* 34, 2292-2294. <https://doi.org/10.1093/bioinformatics/bty071>
- Větrovský, T., Morais, D., Kohout, P., Lepinay, C., Algora Gallardo, C., Awokunle Hollá, S., Doreen Bahmann, B., Bílohnědá, K., Brabcová, V., D'Alò, F., Human, Z.R., Jomura, M., Kolařík, M., Kvasničková, J., Lladó, S., López-Mondéjar, R., Martinović, T., Mašinová, T., Meszárošová, L., Michalčíková, L., Michalová, T., Mundra, S., Navrátilová, D., Odriozola, I., Piché-Choquette, S., Štursová, M., Švec, K., Tláskal, V., Urbanová, M., Vlk L., Voříšková, J., Žifčáková, L., Baldrian, P. 2020. GlobalFungi, a global database of fungal occurrences from high-throughput-sequencing metabarcoding studies. *Scientific Data*, in press.
- Wakefield, W.S., Powell, M.J., Letcher, P.M., Barr, D.J.S., Churchill, P.F., Longcore, J.E., Chen, S.F., 2010. A molecular phylogenetic evaluation of the *Spizellomyces*. *Mycologia* 102, 596-604. <https://doi.org/10.2307/27811072>
- Watanabe, H., Tokuda, G., 2010. Cellulolytic systems in insects. *Annu. Rev. Entomol.* 55, 609-632. <https://doi.org/10.1146/annurev-ento-112408-085319>
- Woolfolk, S.W., Inglis, G.D., 2004. Microorganisms associated with field-collected *Chrysoperla rufilabris* (Neuroptera: Chrysopidae) adults with emphasis on yeast symbionts. *Biol. Control* 29, 155-168. [https://doi.org/10.1016/S1049-9644\(03\)00139-7](https://doi.org/10.1016/S1049-9644(03)00139-7)
- Zhang, D., Lax, A., Henrissat, B., Coutinho, P., Katiya, N., Nierman, W.C., Fedorova, N., 2012. Carbohydrate-active enzymes revealed in *Coptotermes formosanus* (Isoptera: Rhinotermitidae) transcriptome. *Insect Mol. Biol.* 21, 235-245. <https://doi.org/10.1111/j.1365-2583.2011.01130.x>
- Zhang, N., Suh, S-O., Blackwell, M., 2003. Microorganisms in the gut of beetles: evidence from molecular cloning. *J Invertebr. Pathol.* 84, 226-233. <https://doi.org/10.1016/j.jip.2003.10.002>
- Ziganshina, E.E., Mohammed, W.S., Shagimardanova, E.I., Vankov, P.Y., Gogoleva, N.E., Ziganshin, A.M., 2018. Fungal, bacterial, and archaeal diversity in the digestive tract of several beetle larvae (Coleoptera). *BioMed Res. Int.* 2018, 6765438. <https://doi.org/10.1155/2018/6765438>


Zoberi, M.H., Grace, J.K., 1990. Fungi associated with the subterranean termite *Reticulitermes flavipes* in Ontario. *Mycologia* 82, 289-294. <https://doi.org/10.2307/3759899>



**Fig. 1.** Sampling scheme. Termites were collected in foraging galleries, and workers and soldiers were used for voucher sample in 80% ethanol (\_A), while 10 workers for RNA later sample (\_R). Samples of foraging galleries (\_WF) and control wood (\_WC; roughly 10 cm from the closest termite gallery) were also stored in RNA later. Up to three sample sets were collected from the same log, with a distance at least one meter from each other. If more than a single focal termite (T1 and T2) was found in the same log, both were treated independently.




**Fig. 2.** Alpha diversity indices; Chao-1 index (A), Shannon-Wiener index (B) and evenness (C) calculated from the fungal OTUs found in termite bodies, their galleries and insect-free wood controls. Data from *Coptotermes testaceus*, *Heterotermes tenuis*, *Nasutitermes octopilis* are shown. Groups sharing a letter are not significantly different (Tukey HSD *post-hoc* tests,  $p < 0.05$ ).




**Fig. 3.** Relative abundance of sequence reads classified at the phylum (A) and order (B) level. Only the orders with higher relative abundance ( $\geq 1\%$  of reads in at least one sample type) are shown in Fig. 3B. Data from the wood control (W0), bodies (T) and galleries (W) of the termites *Coptotermes testaceus* (C), *Heterotermes tenuis* (H), and *Nasutitermes octopilis* (N) are shown. Artificial order Mucoromycotina GS23 was created for OTU285 (see Table 1) and OTUs with 95% similarity.






**Fig. 4.** Relative abundance of sequence reads classified at the order level. Data from the wood control (W0), bodies (T) and galleries (W) of the termites *Coptotermes testaceus* (C), *Heterotermes tenuis* (H), and *Nasutitermes octopilis* (N) are shown. The number of OTUs for each order are given in the parentheses. Abundant orders, which reached at least 1 % abundance in one fungal community were selected for presentation.



**Fig. 5.** Non-metric multidimensional scaling (NMDS) ordination of the sampling units by their fungal OTU ( $\geq 97\%$  identity) composition based on ITS2 rRNA gene metabarcoding. A) Ordination of raw fungal OTU composition (k = 3 dimensions, final stress = 0.24); B) ordination of residualized fungal OTU composition (i.e. the effect of log and triplet identity removed; k = 3 dimensions, final stress = 0.25). Sample type significantly affects fungal community composition (PERMANOVA: permutations = 999,  $P$ -value = 0.001,  $R^2 = 0.044$ ).





**Fig. 7.** Venn diagram of RDA variation partitioning of fungal OTU composition in A) whole termites and B) termite galleries. Numbers are adjusted R<sup>2</sup> values. \*,  $p < 0.05$ ; \*\*,  $p < 0.01$ ; \*\*\*,  $p < 0.001$ .

**Table 1.** Fungal OTUs specifically associated with termite bodies. First 30 OTUs with the highest fit to RDA1 axis (see Fig. 6) are presented for each termite species. Taxonomic identity of OTUs is based on the Blast similarity search against NCBI Genbank and Unite database. Abbreviations: A – Ascomycota, B – Basidiomycota, C – Chytridiomycota, M – Mucoromycotina.

<sup>1</sup> – 98% sequence similarity with the type of *Spizellomyces punctatus* NR\_111189

<sup>2</sup> – no reliable ITS2 sequence similarity to named taxa. The best hits ( $\leq 90\%$ ) are various *Mucoromycotina* spp. (e.g. HQ406814, LC189046)

<i>Coptotermes testaceus</i>					<i>Heterotermes tenuis</i>					<i>Nasutitermes octopilis</i>				
OTU #	RDA axis loadings		Best Hit description		OTU #	RDA axis loadings		Best Hit description		OTU #	RDA axis loadings		Best Hit description	
	RDA1	RDA2	Similarity / Coverage [%]	Taxon		RDA1	RDA2	Similarity / Coverage [%]	Taxon		RDA1	RDA2	Similarity / Coverage [%]	Taxon
14	-1.05	0.14	94.8/100 (KJ654590)	<i>Phlebia</i> sp. (B)	34	-1.02	-0.01		unidentified	4	-1.20	0.14	88.1/93.5 (HM162185)	<i>Podospora</i> sp. (A)
15	-0.95	0.12		unidentified	15	-0.60	-0.01		unidentified	104	-0.98	0.08	87.1/100 (AB846969)	GS23 sp. (M) <sup>2</sup>
82	-0.58	0.02	99.5/98.5 (HM771021)	<i>Hawksworthiomyces</i> sp. (A)	20	-0.58	0.06	91.2/89.5 (AY762623)	<i>Scytalidium lignicola</i> (A)	12	-0.96	0.01	100/100 (GU945354)	<i>Lasiodiplodia citricola</i> (A)
31	-0.54	0.09	100/100 (AY857228)	<i>Trichoderma harzianum</i> (A)	11	-0.57	-0.01	90.4/96.7 (KU214528)	<i>Arthrographis</i> sp. (A)	413	-0.56	0.17	97/100 (GU054100)	<i>Wrightoporia tropicalis</i> (B)
210	-0.47	0.08	98.3/100 (AY997092)	<i>Spizellomyces punctatus</i> (C) <sup>1</sup>	41	-0.54	0.04	90.4/76 (UDB014156)	<i>Auriculariales</i> sp. (B)	258	-0.48	0.12		unidentified
23	-0.45	0.06		unidentified	68	-0.53	-0.39		unidentified	11	-0.44	0.04	90.4/96.7 (KU214528)	<i>Arthrographis</i> sp. (A)
24	-0.44	0.06	91.3/44 (UDB028178)	<i>Hyphodontia pilaecystidiata</i> (B)	207	-0.50	0.07	99.6/100 (AY743636)	<i>Malassezia restricta</i> (B)	20	-0.37	0.02	91.2/89.5 (AY762623)	<i>Scytalidium lignicola</i> (A)
203	-0.40	0.08	100/100 (AY154939)	<i>Trichoderma spirale</i> (A)	286	-0.48	0.01	92/98 (GQ272617)	<i>Scytalidium ganodermophthorum</i> (A)	171	-0.31	-0.02	100/100 (EU401550)	<i>Trichoderma orientale</i> (A)
20	-0.35	-0.05	91.2/89.5 (AY762623)	<i>Scytalidium lignicola</i> (A)	24	-0.48	0.00	91.3/44 (UDB028178)	<i>Hyphodontia pilaecystidiata</i> (B)	885	-0.30	0.04	84.4/100 (AB846969)	GS23 sp. (M) <sup>2</sup>
285	-0.35	0.05	92.9/93.8 (AB846959)	GS23 sp. (M) <sup>2</sup>	214	-0.48	-0.03	87.8/100 (AB846969)	GS23 sp. (M) <sup>2</sup>	183	-0.30	-0.01	90.5/81 (AB846975)	GS23 sp. (M) <sup>2</sup>
286	-0.34	0.04	92/98 (GQ272617)	<i>Scytalidium ganodermophthorum</i> (A)	31	-0.45	-0.03	100/100 (AY857228)	<i>Trichoderma harzianum</i> (A)	121	-0.27	0.17		unidentified
22	-0.32	-0.17	87.6/72.9 (FJ231021)	<i>Penicillium curticaule</i> (A)	210	-0.44	0.12	98.3/100 (AY997092)	<i>Spizellomyces punctatus</i> (C) <sup>1</sup>	179	-0.27	0.04	94.8/100 (KT951335)	<i>Agaricus candidolutescens</i> (B)
197	-0.32	0.05	99.4/100 (KJ174211)	<i>Trichoderma koningiopsis</i> (A)	149	-0.42	-0.01	89.8/90.3 (AB846969)	GS23 sp. (M) <sup>2</sup>	256	-0.26	0.02	86.3/56.7 (KM103946)	Fungi sp. (unidentified)
41	-0.32	0.12	90.4/76 (UDB014156)	<i>Auriculariales</i> sp. (B)	145	-0.34	-0.03		unidentified	118	-0.26	-0.01	88.3/95.7 (UDB013022)	Sordariales sp. (A)
12	-0.32	0.06	100/100 (GU945354)	<i>Lasiodiplodia citricola</i> (A)	22	-0.32	0.03	87.6/72.9 (FJ231021)	<i>Penicillium curticaule</i> (A)	59	-0.25	0.04		unidentified
154	-0.31	0.02	90.7/100 (GQ272617)	<i>Scytalidium ganodermophthorum</i> (A)	629	-0.31	0.00	98.6/100 (KU164491)	<i>Malassezia restricta</i> (B)	156	-0.24	-0.07	100/19.5 (GQ280589)	<i>Calonectria leguminum</i> (A)
38	-0.30	0.13		unidentified	85	-0.30	-0.07	100/100 (HM770996)	<i>Helotiales</i> sp. (A)	1923	-0.24	-0.05	100/98.8 (EU280098)	<i>Trichoderma citrinoviride</i> (A)
28	-0.30	-0.02	86.9/100 (JX857794)	<i>Corticium</i> sp. (B)	203	-0.29	0.06	100/100 (AY154939)	<i>Trichoderma spirale</i> (A)	6	-0.24	0.01	86/47.6 (DQ826552)	<i>Resinicium monticola</i> (B)
34	-0.30	0.04		unidentified	248	-0.28	0.04	100/100 (GU237707)	<i>Boeremia exigua</i> (A)	32	-0.24	-0.01		unidentified

158	-0.28	0.00	89.2/97.4 (GQ272617)	<i>Scytalidium ganodermophthorum</i> (A)	202	-0.27	0.00		unidentified	229	-0.23	0.03	99.6/100 (UDB014090)	Trechisporales sp. (B)
257	-0.27	-0.02	88.6/100 (KU295549)	GS23 sp. (M) <sup>2</sup>	257	-0.26	-0.02	88.6/100 (KU295549)	GS23 sp. (M) <sup>2</sup>	34	-0.22	0.02		unidentified
152	-0.26	0.03		unidentified	259	-0.26	0.00	100/98.1 (JN626104)	<i>Penicillium mallochii</i> (A)	25	-0.22	-0.04	85.6/96.1 (HF677173)	<i>Cordana terrestris</i> (A)
94	-0.26	-0.22		unidentified	302	-0.26	-0.01	99.5/100 (KF472157)	<i>Verticillium leptobactrum</i> (A)	172	-0.21	0.03		unidentified
133	-0.25	0.03	92.9/90.4 (EF127890)	<i>Hawksworthiomyces lignivorus</i> (A)	223	-0.25	0.00		unidentified	199	-0.21	0.04	86/92.6 (KU975068)	<i>Pseudoproboscispora</i> sp. (A)
267	-0.25	0.00	94.3/98.1 (AF033470)	<i>Penicillium sclerotigenum</i> (A)	946	-0.25	-0.01	93.7/100 (AB846969)	GS23 sp. (M) <sup>2</sup>	82	-0.21	0.07	99.5/98.5 (HM771021)	<i>Hawksworthiomyces</i> sp. (A)
30	-0.24	0.03	90.9/43.4 (JX675137)	<i>Gymnoascus</i> sp. (A)	12	-0.25	-0.03	100/100 (GU945354)	<i>Lasiodiplodia citricola</i> (A)	7	-0.20	-0.74	99/100 (KT224922)	unidentified
509	-0.24	0.03	99.4/100 (DQ109528)	<i>Trichoderma lieckfeldiae</i> (A)	83	-0.25	-0.01	99.3/100 (GQ272617)	<i>Scytalidium ganodermophthorum</i> (A)	88	-0.20	0.39	100/100 (HM148090)	<i>Cladosporium exasperatum</i> (A)
402	-0.23	0.03	91.1/93.2 (AB846969)	GS23 sp. (M) <sup>2</sup>	122	-0.24	-0.06	99.3/100 (AY273308)	Ascomycota sp. (A)	299	-0.18	0.04	100/19.4 (GQ280589)	<i>Calonectria leguminum</i> (A)
399	-0.22	-0.04	90.8/62.5 (AB846970)	GS23 sp. (M) <sup>2</sup>	14	-0.23	0.00	94.8/100 (KJ654590)	<i>Phlebia</i> sp. (B)	425	-0.18	0.04	86.6/79.2 (KY687694)	GS23 sp. (M) <sup>2</sup>
207	-0.22	0.05	99.6/100 (AY743636)	<i>Malassezia restricta</i> (B)	234	-0.23	0.03	99.3/100 (FR682163)	<i>Malassezia</i> sp. (B)	190	-0.18	0.04	88.7/100 (JX545187)	GS23 sp. (M) <sup>2</sup>