



International Journal of Aging Research (ISSN:2637-3742)



Exergaming improves functional fitness in MCI patients. Does the APOE genotype moderate the outcome?

Anthoula C. Tsolaki^{1,2*}, Magda Tsolaki², Niki Pandria¹, Eftychia Lazarou², Olympia Gkatzima³, Vasiliki Zilidou¹, Maria Karagianni¹, Zafiroura Iakovidou- Kritsi⁴, Vasilios K. Kimiskidis², Panagiotis D. Bamidis¹

¹Medical Physics Laboratory, School of Medicine, Aristotle University of Thessaloniki, Greece., orcid: 0000-0002-5563-7776, orcid: 0000-0002-4859-6996, orcid: 0000-0002-4492-4382, orcid: 0000-0002-7130-5739, orcid: 0000-0002-9936-5805. ²1st Department of Neurology, AHEPA University Hospital, Aristotle University of Thessaloniki, Greece, orcid: 0000-0003-1272-9782, orcid: 0000-0002-2072-8010, orcid: 0000-0002-3335-3019. ³Panhellenic Institute of Neurodegenerative Diseases, Thessaloniki, Greece, orcid: 0000-0002-2415-0498. ⁴Laboratory of Medical Biology-Genetics Department, School of Medicine, Aristotle University of Thessaloniki, Greece, orcid: 0000-0001-6800-2369.

ABSTRACT

Objectives: Mild cognitive impairment could be defined as the condition between healthy aging and dementia. MCI patients seem to retain the neuroplasticity to benefit from Physical Exercise (PE) interventions delaying the progression to dementia.

The present study investigates the impact of PE via “Exergaming” on the functional fitness of MCI adults, depending on the presence of the APOE ϵ 4 allele.

Methods: 159 MCI participants were recruited. They were separated to two groups (performing PE or not). The Fullerton Functional Test was used as a primary outcome measure in two-time points (prior to and after PE).

Results: The Active group showed more considerable improvement compared to the Passive group in all Fullerton components despite the presence of APOE ϵ 4.

Discussion: PE via exergaming has a beneficial functional effect in MCI patients, whether carrying the APOE ϵ 4 allele or not.


Keywords: Mild Cognitive Impairment; APOE ϵ 4; Exergaming; Functional fitness; Physical Exercise; Fullerton

*Correspondence to Author:

Anthoula C. Tsolaki, MD, Ph.D
1st Department of Neurology, AHEPA University Hospital, Aristotle University of Thessaloniki. Medical Physics Laboratory, School of Medicine, Aristotle University of Thessaloniki. PO Box 376, 54124, Thessaloniki, Greece.
Tel. 0030 6947568776

How to cite this article:

Anthoula C. Tsolaki, M. Tsolaki, N. Pandria, E. Lazarou, O. Gkatzima, V. Zilidou, M. Karagianni, Z. Iakovidou- Kritsi, V. K. Kimiskidis, P. D. Bamidis. Exergaming improves functional fitness in MCI patients. Does the APOE genotype moderate the outcome?. International Journal of Aging Research, 2020; 3:74.

 eSciPub
eSciPub LLC, Houston, TX USA.
Website: <https://escipub.com/>

1. Introduction

The motto “Exercise is medicine” was started by the American College of Sports Medicine [1]. This notion is based on the fact that lifelong physical exercise (PE) is related to a longer lifespan and delays the onset of more than forty (40) chronic conditions/diseases [2].

Mild cognitive impairment (MCI) could be defined as the condition between normal aging and dementia [3]. MCI patients present a neuronal loss of about 36.5% at that stage, along with a synaptic dysfunction [4]. However, they seem to retain sufficient neuroplasticity to benefit from PE interventions, delaying the progression to dementia [5]

APOE ϵ 4 presence was found to have a considerable impact on brain metabolism and structure [6]. It is also related to the progression of healthy elderly to MCI and AD [6]. However, all APOE ϵ 4 carriers will not develop AD suggesting the interactive effects of APOE genotype with other genetic or environmental factors [7].

PE could improve cognitive function and play a protective role against neurodegeneration, the underlying mechanisms, though, for these protective effects, have not yet been fully elucidated [8]. There is evidence indicating that PE activates the release of neurotrophic factors [9] and promotes angiogenesis [10], facilitates neurogenesis and synaptogenesis, which in turn improve memory and cognitive functions [11]. Research has also shown that PE has a neuroprotective role [12].

Recent research though suggests that prolonged exercise habits are required for cognitive enhancement, while benefits on functioning can be observed when individuals take up an exercise routine later in life or even after the cognitive impairment onset [13].

Exergaming is an effective training or rehabilitation tool for elderly [14], comparable to conventional forms of PE, with a personalized approach adapting the level of challenge according to the subject's performance. Hence, it provides a safe and effective solution with constant monitoring in older adults living in

institutions or rural areas away from organized Day care centers.

This study is part of a more extensive study, the Long Lasting Memories (LLM) Project [15] originally funded by the ICT-CIP-PSP Program of the European Commission, assessing both cognitive and somatometric/ physical effects of non-pharmacological computer- based interventions in MCI patients based on the presence of APOE ϵ 4. Because of the amount of the factors under investigation, the results have been split in different manuscripts, the first already published [16].

The aim of this study is to investigate the impact of the computerized PE via “Exergaming” on the functional fitness of older adults with MCI in general and its likely dependence in the presence of APOE ϵ 4.

2. Materials and methods

A Quasi-Experimental Study Design took place in Thessaloniki from 2009 to 2017, in order to explore the efficacy of different computer-based interventions in elderly.

The randomization of the participants was not feasible mainly for practical issues, time and financial limitation of the study. Allocation to groups was driven by non- systematic practical and logistic reasons (National holiday time, number of successfully screened participants at a given time point etc.). Moreover, the allocation was not influenced by the participant's choice, motivation or compliance. The type of the intervention applied each time in each place (Spiritual center, Open Care Centers) was determined before the initiation of the screening procedure and it was an open call to the elderly whether were cognitively intact or not. Thus, interventions took place both in the East and West area of Thessaloniki, minimizing the geographical socioeconomic differences of the participants. From all the participants who enrolled in the LLM project, we retrospectively collected and analyzed all of those who had an initial diagnosis of MCI, fulfilling the inclusion-exclusion criteria, to explore the impact of the computerized PE on the functional fitness of

older adults with MCI in general, and its likely dependence on the presence of APOEε4 allele. One hundred fifty-nine (159) MCI participants were recruited during a thorough screening procedure (see Figure 1). Males and females, aged ≥60 years old, fluent in Greek, were invited to participate. The call was made in Church Spiritual Centers, in Open Care Centers for the elderly in East and West area of Thessaloniki,

and the Day Care centers of Greek Alzheimer Association. Participants provided written informed consent and were compensated for their participation in the study. The study protocol was approved by the Ethics Committee of Aristotle University of Thessaloniki and Greek Alzheimer Association (Protocol No **38/5.6.2013** and **221/14-09-2013**). (see Table 1A)

Table 1: A Demographic characteristic of the study groups. Total sample and B. Neuropsychological assessment of the Total sample

A.					
Groups	Gender Male/Female	Age Median, IQR	Education years Median, IQR	CIRS Median, IQR	BMI cm/Kg Mean (SD)
Active N=82	12/70	68.75, [65.00,73.00]	6.0, [6.00, 9.25]	1.67, [1.44,2.00]	28.73 (3.57)
Passive N=77	17/60	68.00, [63.00,72.50]	6.0, [6.00,11.00]	1.50, [1.33,1.83]	29.77 (4.40)
Total	29/130	68.5, [64.00, 73.00]	6.0, [6.00, 10.00]	1.57, [1.40, 2.00]	29.29 (4.06)
B.					
Groups	MMSE Median, IQR	MoCA Median, IQR	Trail B Median, IQR	IADL Mean rank	GDS Median, IQR
Active	27.00, [26.00, 28.00]	23.00, [21.00, 25.00]	202.00, [145.00, 256.00]	81.86	2.00, [1.00, 4.00]
Passive	26.00, [24.00, 28.00]	23.00, [19.75, 25.00]	152.00, [115.00, 216.00]	68.79	1.00, [0.00, 4.00]
test statistic p-value	U=2345.50; p=0.046	U=2464.50; p=0.858	U=1496.50; p=0.012	U=2321.00; p=0.002	U=2509.50; p=0.165

MMSE: Mini Mental State Examination; MoCA: Montreal Cognitive Assessment; IADL: Instrumental Activities of Daily Living; GDS: Geriatric Depression Scale

Exclusion criteria included any severe physical illness, current psychiatric or other neurological disorder, history of drug or alcohol abuse and use of neuro-modifying drugs (other than cholinesterase inhibitors). All participants reported normal or corrected-to-normal vision and hearing. Prior to the PE intervention, all the participants underwent a neuropsychological evaluation (see Table 1B). The Senior Fitness Fullerton Test was used to assess the functional fitness of older adults. It consists of six tests and evaluates the flexibility of low back and hamstrings, the functional capacity of individuals through the strength of the lower limbs and the dynamic balance, the speed, agility and balance during movement [17,18]. Specifically, the tests used are Chair Stand Test (CST), Arm Curl Test

(ACT), 2- Minute Step Test (2-MST), Chair Sit-and-Reach Test (CSaRT), Back Scratch Test (BST) and Foot Up-and-Go Test (FUaG). They were also subjected to blood tests and neuroimaging via CT/MRI. The neuropsychological, medical and laboratory results were evaluated by a cognitive neurologist (MT). The diagnosis and categorization were based on clinical criteria [3,19] and supported by neuroimaging evidence of media-temporal atrophy. The diagnosis was performed by a dementia expert neurologist (MT). At first, during the LLM project, the participants were assigned to four different interventional groups and one passive group. The first group attended a combined cognitive and PE implemented in the framework of the LLM project

[20]. All the PE programs were implemented via the exergaming platform Fit For All (FFA) [21]. The second group was the Cognitive training control (CTC) group which performed only cognitive sessions using the Brain HQ software, the third group was the Physical Training Control (PTC) group which was exposed to PE by using the Fit For All platform, and the fourth group was the Active control (AC) group which was exposed to a computerized cognitive training via a short documentary video about nature, art, history, and culture. All the interventional groups had an identical number of sessions. The fifth group was the Passive control group, which did not receive any training and it was on the waiting list (see Figure 1).

The program lasted 8-12 weeks and participants completed at least 24 sessions of cognitive training and/or 16 sessions of PE. The screening neuropsychological evaluation was conducted 1-2 weeks prior to the intervention, the post-test evaluation 1-2 weeks after the end of the training, and the follow-up was six months later.

Neuropsychologists and physical trainers who performed the neuropsychological and somatometric assessment both before and after the training were different from those who administered the program.

For the purposes of the present study, all the above participants were divided into two larger groups depending on whether they did PE or not, namely, Active (LLM and PTC participants) and Passive physical training group (AC, CTC and Passive participants) (see Figure 1), to investigate the impact of the computerized PE via “Exergaming” on the functional fitness of older adults with MCI in general and its likely dependence on the presence of APOEε4 allele. A sample of some 159 participants, males and females (29/130), with a median age of 68.5 years, with a median education level of 6 years and a median Cumulative Illness Rating Scale (CIRS) of 1.57 has been included in this study. Eighty-two (82) of these were included in the Active group, while seventy-seven (77) in the Passive group (see Table 1A).




Figure 1: Study Flowchart.

Blood samples used for genotyping were (EDTA)-containing receptacles. DNA was collected in ethylenediaminetetraacetic acid extracted from peripheral blood using the

QIAamp Blood DNA purification kit (Qiagen Inc, USA). To determine the APOE genotype, part of the APOE gene (228 bp) containing both polymorphic sites (amino acid positions 112 and 158) was amplified by PCR analysis, using the following primers: forward:

5"-GGCACGGCTGTCCAAGGAGCTGCA-3"

and reverse:

5"-GCCCCGGCCTGGTACTACTGCCAG-3",

according to the method described in [22].

Statistical analysis was performed by means of IBM SPSS (version 23) and the significance level was set at 5%. The alpha inflation problem due to multiple comparisons was counteracted by applying Bonferroni correction.

Demographics characteristics were explored for normality assumption in the pool of participants as well as in both groups (Active, Passive). Depending on the normality assumption, appropriate descriptive statistics were calculated, and between-groups comparisons were performed. Comparisons of demographic data between groups were run using either Mann-Whitney U test or independent samples t-test.

Fullerton data

Fullerton components were measured both before and after the PE. Initially, the baseline performance in Fullerton test was compared between groups by means of Mann-Whitney U test. Subsequently, data collected were analyzed having as within-factor the time (before and after PE) and as between-factor the group (Active, Passive). Although Mixed Model Analysis of Variance should be the analysis of choice for the data collected, the assumptions of the corresponding test were not satisfied in each cell of the design. Thus, score differences (post-pre) were calculated and then tested for normality assumption. Within-group changes were investigated using Wilcoxon Signed Ranks test after grouping data by group (Active, Passive). Score change was compared between groups performing Mann-Whitney U test.

Moreover, the genotype of participants was also available. Therefore, we planned comparisons

taking into account the presence or the absence of APOEε4 allele. We determined as within-factor the time (before and after PE) and as between-factors the group (Active, Passive) and the genotype (non-ε4-carriers, ε4-carriers). Even though, we investigated the assumptions of the Mixed Model Analysis of Variance, they were not fulfilled. Therefore, score changes within each group, Active/Passive or Carriers/Non-Carriers respectively, were investigated using Wilcoxon Signed Ranks test after grouping data by group and genotype. Additionally, score changes between groups were explored using Mann-Whitney U test grouping data by group type (Active, Passive) and genotype (Carrier, Non-carrier).

3. Results

Active and Passive participants were not significantly different in their demographic data such as age, education years, CIRS and BMI (*Age: U=2891.50; p=0.359; Education: U=3082.50; p=0.782; CIRS: U=2607.50; p=0.056; BMI: t (141) =-1.543; p=0.125*) (see Table 1A).

Table 1 about here

The baseline performance between the two groups revealed a statistically significant better performance of the passive group in two tasks ACT (*U=1816.50; p<0.001*) and Foot-up-and-go (*U=1655.00; p<0.001*).

Participants recruited in the Active group significantly improved their performance in all Fullerton components when comparing their scores at the two time points (*All p values <0.001*). On the other hand, Passive participants significantly improved their performance only in CST task (*W=-2.265; p=0.023*) (See Table 2A and Figure 2).

Planned comparisons in score changes between groups (Active, Passive) revealed group-differences in all Fullerton components. More precisely, the Active group showed greater improvement compared to the Passive group in all Fullerton components (*All p values ≤0.005*). (See Table 2B).




Figure 2: Within-Group Comparisons at different Fullerton tasks before and after the training period.
 * p<0.05, ** p<0.01 &*** p<0.001.

Table 2: Fullerton Assessment before and after training period.
 Within group (A) and between groups (B) comparisons.

A.					
Fullerton components	Group	N	Descriptive statistics		Test results
			Before training	After training	
<i>Within-group comparison</i>					
Chair stand	Active	82	13.00, [11.00, 15.00]	15.00, [12.00, 18.00]	W=-5.693; p<0.001
	Passive	77	13.00, [10.00, 15.00]	14.00, [10.50, 17.00]	W=-2.265; p=0.023
Arm curl	Active	82	16.00, [14.00, 20.00]	21.00, [17.00, 25.00]	W=-6.015; p<0.001
	Passive	77	21.00, [17.00, 24.00]	21.00, [18.00, 25.00]	W=-1.583; p=0.114
2- Minutes Step in Place	Active	82	70.00, [57.00, 92.25]	83.00, [66.50, 103.50]	W=-4.524; p<0.001
	Passive	77	80.00, [67.50, 91.50]	78.00, [66.50, 92.00]	W=-0.531; p=0.595
Back scratch	Active	82	-10.00, [-17.25, 1.25]	-6.00, [-14.00, 3.25]	W=-5.839; p<0.001
	Passive	77	-7.00, [-15.00, 2.00]	-7.00, [-14.50, 2.00]	W=-1.451; p=0.147
Sit and Reach	Active	82	1.50, [-6.25, 7.00]	5.50, [0.00, 13.00]	W=-6.173; p<0.001
	Passive	77	2.00, [-5.00, 7.00]	2.00, [-2.50, 6.50]	W=-0.028; p=0.978
Foot-up-and-go	Active	82	5.82, [5.20, 6.82]	5.26, [4.60, 6.00]	W=-5.636; p<0.001
	Passive	77	5.00, [4.68, 5.41]	5.03, [4.61, 5.60]	W=-0.401; p=0.688
B.					
<i>Between-group comparison</i>					
Chair stand	Active	82	2.00, [0.00, 4.00]		U=2341.00; p=0.005
	Passive	77	0.00, [-1.00, 3.00]		
Arm curl	Active	82	3.00, [0.00, 8.00]		U=1813.00; p<0.001
	Passive	77	0.00, [-2.00, 2.00]		
2- Minutes Step in Place	Active	82	10.00, [0.00, 21.00]		U=1926.50; p<0.001
	Passive	77	1.00, [-7.50, 5.50]		
Back scratch	Active	82	2.50, [0.00, 5.00]		U=1880.00; p<0.001
	Passive	77	0.00, [-1.00, 2.00]		
Sit and Reach	Active	82	4.00, [1.00, 8.00]		U=1585.00; p<0.001
	Passive	77	0.50, [-2.00, 2.00]		
Foot-up-and-go	Active	82	-0.60, [-1.16, -0.03]		U=1624.50; p<0.001
	Passive	77	0.02, [-0.26, 0.34]		

Non-ε4-carriers of the Active group showed significant changes in all Fullerton components (All *p* values <0.001). Active ε4-carriers significantly improved their performance at all Fullerton tasks apart from 2-Minutes Steps in Place test (CST: *W*=-3.224; *p*=0.001; ACT: *W*=3.063; *p*=0.002; CSaRT: *W*=-3.055; *p*=0.002;

BST: *W*=-2.568; *p*=0.010; *FUaG*: *W*=-2.273; *p*=0.023).

Non-ε4-carriers of the Passive group increased their scores only at CST (*W*=-2.165; *p*=0.030). On the other hand, Passive ε4-carriers did not change their performance at any test (see Table 3A and Figure 3).




Figure 3: Within-Group Comparisons at different Fullerton tasks, based on genotype, before and after the training period. * *p*<0.05, ** *p*<0.01 & *** *p*<0.001

The comparison of performance change between the APOEε4-carriers and non-ε4-carriers within each group (Active, Passive) did not show significant differences in their performance change at the Fullerton tasks in the two-time points either in Active or in the Passive group (see Table 4).

Additionally, Active non-ε4-carriers were found to show superior performance compared to the Passive non-ε4-carriers in all Fullerton tasks

(CST: *U*=1542.50; *p*=0.021; ACT: *U*=1194.00; *p*<0.001; CSaRT: *U*=904.50; *p*<0.001; *BST*: *U*=1311.50; *p*=0.001; *FUaG*: *U*=1009.50; *p*<0.001; 2-MST: *U*=1210.00; *p*<0.001). On the other hand, Active ε4-carriers showed greater performance at the ACT (*U*=68.00; *p*=0.024), the *BST* (*U*=48.50; *p*=0.003) and the *FUaG* (*U*=72.50; *p*=0.038) compared to the Passive ε4-carriers (see Table 3B).

Table 3: Fullerton Assessment before and after training period. Within group based on APOEε4 (A) and between groups based on APOEε4(B) comparisons.

A.					
Fullerton components	Group	N	Descriptive statistics		Test results
			Before training	After training	
<i>Within-group comparison in ε4-Carriers</i>					
Chair stand	Active	17	13.00, [10.50, 14.00]	15.00, [12.00, 19.00]	W=-3.224; p=0.001
	Passive	15	12.00, [11.00, 17.00]	12.00, [6.00, 18.00]	W=-0.669; p=0.504
Arm curl	Active	17	16.00, [15.50, 19.00]	21.00, [16.50, 25.00]	W=-3.063; p=0.002
	Passive	15	21.00, [18.00, 23.00]	21.00, [19.00, 25.00]	W=-1.303; p=0.192
2- Minutes Step in Place	Active	17	73.00, [59.00, 101.50]	83.00, [70.50, 93.00]	W=-1.553; p=0.120
	Passive	15	80.00, [63.00, 92.00]	75.0, [60.00, 87.00]	W=-0.881; p=0.378
Back scratch	Active	17	-15.00, [-22.00, 0.50]	-12.00, [-19.50, 2.25]	W=-2.568; p=0.010
	Passive	15	-7.00, [-12.00, 2.00]	-7.00, [-12.00, 1.00]	W=-0.931; p=0.352
Sit and Reach	Active	17	2.00, [0.50, 7.00]	5.00, [3.00, 12.00]	W=-3.055; p=0.002
	Passive	15	0.00, [-8.00, 4.00]	2.00, [-7.00, 7.00]	W=-1.282; p=0.200
Foot-up-and-go	Active	17	5.72, [4.99, 6.21]	5.02, [4.50, 5.83]	W=-2.273; p=0.023
	Passive	15	4.99, [4.40, 5.63]	5.15, [4.69, 5.67]	W=-0.227; p=0.820
<i>Within-group comparison in Non-ε4-carriers</i>					
Chair stand	Active	65	13.00, [11.00, 15.50]	15.00, [12.00, 18.00]	W=-5.693; p<0.001
	Passive	62	13.00, [9.75, 15.00]	14.00, [10.75, 17.00]	W=-2.165; p=0.030
Arm curl	Active	65	16.00, [13.50, 20.00]	21.00, [17.00, 25.00]	W=-6.015; p<0.001
	Passive	62	21.00, [17.00, 24.25]	21.00, [17.00, 25.00]	W=-1.097; p=0.272
2- Minutes Step in Place	Active	65	69.00, [57.00, 91.00]	82.00, [66.00, 105.00]	W=-4.524; p<0.001
	Passive	62	79.50, [69.50, 92.25]	79.50, [66.75, 93.00]	W=-0.083; p=0.934
Back scratch	Active	65	-9.00, [-15.50, 2.00]	-6.00, [-12.00, 5.00]	W=-5.839; p<0.001
	Passive	62	-7.00, [-15.75, 2.00]	6.00, [-15.00, 2.00]	W=-1.827; p=0.068
Sit and Reach	Active	65	1.00, [-7.00, 7.00]	6.00, [0.00, 13.00]	W=-6.173; p<0.001
	Passive	62	2.75, [-5.25, 8.00]	2.00, [-0.50, 6.25]	W=-0.806; p=0.420
Foot-up-and-go	Active	65	5.90, [5.30, 7.04]	5.30, [4.65, 6.00]	W=-5.636; p<0.001
	Passive	62	5.00, [4.78, 5.41]	5.03, [4.60, 5.60]	W=-0.276; p=0.782
B.					
<i>Between-group comparison in ε4-Carriers</i>					
Chair stand	Active	17	2.00, [1.50, 4.00]		U=90.00; p=0.154
	Passive	15	0.00, [1.00, 4.00]		
Arm curl	Active	17	5.00, [1.50, 8.00]		U=68.00; p=0.024
	Passive	15	1.00, [-1.00, 3.00]		
2- Minutes Step in Place	Active	17	5.00, [-3.50, 16.50]		U=81.00; p=0.079
	Passive	15	-4.00, [-7.00, 4.00]		
Back scratch	Active	17	3.00, [0.75, 5.00]		U=48.50; p=0.003
	Passive	15	-1.00, [-1.00, 0.00]		
Sit and Reach	Active	17	4.00, [1.00, 7.00]		U=95.00; p=0.218

	Passive	15	2.00, [-2.00, 5.00]	
Foot-up-and-go	Active	17	-0.41, [-1.10, 0.13]	U=72.50; p=0.038
	Passive	15	-0.06, [-0.2, 0.4]	
<i>Between-group comparison in Non-ε4-carriers</i>				
Chair stand	Active	65	1.00, [0.00, 3.50]	U=1542.50; p=0.021
	Passive	62	0.50, [-1.00, 2.25]	
Arm curl	Active	65	2.00, [0.00, 9.00]	U=1194.00; p<0.001
	Passive	62	0.00, [-2.00, 2.00]	
2- Minutes Step in Place	Active	65	0.00, [-2.25, 2.00]	U=1210.00; p<0.001
	Passive	62	1.50, [-8.00, 6.00]	
Back scratch	Active	65	2.00, [0.00, 5.50]	U=1311.50; p=0.001
	Passive	62	0.00, [-1.00, 2.25]	
Sit and Reach	Active	65	4.00, [0.50, 8.50]	U=904.50; p<0.001
	Passive	62	0.00, [-2.25, 2.00]	
Foot-up-and-go	Active	65	-0.60, [-1.25, -0.03]	U=1009.50; p<0.001
	Passive	62	0.04, [-0.29, 0.31]	

Table 4: Fullerton Assessment comparison between ε4-carriers and nonε4-carriers within Active and Passive group respectively, before and after training.

Fullerton components Change in time (post-pre)	Group	N	Descriptive statistics	Test results
<i>Between-group comparison in the Active group</i>				
Chair stand	Carriers	17	2.00, [1.50, 4.00]	U=461.50; p=0.294
	Non-carriers	65	1.00, [0.00, 3.50]	
Arm curl	Carriers	17	5.00, [1.50, 8.00]	U=486.50; p=0.448
	Non-carriers	65	2.00, [0.00, 9.00]	
2- Minutes Step in Place	Carriers	17	5.00, [-3.50, 16.50]	U=467.00; p=0.328
	Non-carriers	65	14.00, [0.00, 21.00]	
Back scratch	Carriers	17	3.00, [0.75, 5.00]	U=549.00; p=0.968
	Non-carriers	65	2.00, [0.00, 5.50]	
Sit and Reach	Carriers	17	4.00, [1.00, 7.00]	U=531.50; p=0.809
	Non-carriers	65	4.00, [0.50, 8.50]	
Foot-up-and-go	Carriers	17	-0.41, [-1.10, 0.13]	U=513.00; p=0.651
	Non-carriers	65	-0.60, [-1.25, -0.03]	
<i>Between-group comparison in the Passive group</i>				
Chair stand	Carriers	15	0.00, [-1.00, 4.00]	U=451.50; p=0.861
	Non-carriers	62	0.50, [-1.00, 2.25]	
Arm curl	Carriers	15	1.00, [-1.00, 3.00]	U=404.50; p=0.433
	Non-carriers	62	0.00, [-2.00, 2.00]	
2- Minutes Step in Place	Carriers	15	-4.00, [-7.00, 4.00]	U=404.00; p=0.432
	Non-carriers	62	1.50, [-8.00, 6.00]	
Back scratch	Carriers	15	-1.00, [-1.00, 0.00]	U=329.50; p=0.076
	Non-carriers	62	0.00, [-1.00, 2.25]	
Sit and Reach	Carriers	15	2.00, [-2.00, 5.00]	U=324.50; p=0.070
	Non-carriers	62	0.00, [-2.25, 2.00]	
Foot-up-and-go	Carriers	15	-0.06, [-0.20, 0.40]	U=460.00; p=0.949
	Non-carriers	62	0.04, [-0.29, 0.31]	

Discussion

To our knowledge, there is no similar study in MCI seniors to investigate the impact of the computerized PE on functional fitness test depending on the APOE ϵ 4 allele presence.

In our study, the analysis within the groups before and after the training showed that the active group improved in all the physical parameters (flexibility of low back and hamstrings, the functional capacity of individuals through the strength of the lower limbs and the dynamic balance, the speed, agility and balance during movement) under investigation, which is in line with previous studies [23]. On the contrary, the Passive group showed minor or no improvement at all. The between groups analysis underlined, once again, the multiple training benefits on the Active group, even though, the baseline assessment was in favor of the passive group.

The analysis which was carried out based on the presence of the APOE ϵ 4 allele, a well-known risk factor for AD, showed similar improvement of the functional fitness variables in ϵ 4-carriers and non- ϵ 4-carriers of the Active group. These findings deviate from what is known so far regarding the questionable benefit of cognitive or combined exercise on the MCI APOE ϵ 4-carriers cognitive performance has been shown [24]. On the other hand, Passive non- ϵ 4-carriers improved only in one task whereas Passive ϵ 4-carriers did not improve at all.

Prevailing evidence suggests that differential effects of APOE isoforms on A β aggregation and clearance play a major role in neurodegeneration [25]. Sedentary elderly APOE ϵ 4 individuals are at augmented risk for cerebral amyloid deposition, synaptic deterioration, hippocampal atrophy and decreased brain metabolism in distinct brain areas. Thus, APOE ϵ 4 has a direct pathological effect on the cerebrovascular system [26].

PE modifies the metabolic, structural, and functional brain state. The regular, moderate-intensity exercise may improve brain health by neuroprotective mechanisms. It also leads to cardiovascular disease risk reduction via

numerous different mechanisms at a molecular and cellular level and via its sustainability, it enables gross anatomical remodeling [27].

Exergaming show comparable outcomes in physical [28] and cognitive functions [29] as well as in the psychosocial well-being of the elderly compared to the traditional physical exercise. It is also notable that exergames may reduce the incidence of falls [30], which is a leading cause for loss of independence among older adults.

MCI patients is a population at risk with preserved functionality through compensatory mechanisms, responding well in physical and cognitive intervention training [20]. Even though there is a growing interest in the PE cognition effect, existing knowledge is limited regarding how the functional fitness aspects, such as strength, endurance, balance, flexibility, of the MCI patients alter due to training and whether those changes, if any, are differentiated depending on the presence of APOE ϵ 4 allele.

Our results indicate that both MCI APOE ϵ 4 carriers and non- ϵ 4-carriers experience beneficial effects induced by exergaming in multiple physical parameters. Even though APOE ϵ 4-carriers have been found to gain less cognitive benefits, we showed a prominent positive outcome on the physical fitness of MCI ϵ 4-carriers seniors. Moreover, PE in this high-risk population may be crucial for the improvement of other important factors such as the cardiovascular disease load, the functional impairment, the social isolation, the falls, and the subsequent hospitalization, institutionalization and finally the mortality. Accordingly, PE may not improve directly cognition in MCI APOE ϵ 4-carriers [16], but it may eliminate the additional risk of all the aforementioned co-factors strengthening the notion that physical exercise may facilitate the benefits of any cognitive training, but also that "Exercise is medicine" [1].

5. Conclusion

PE via exergaming has a beneficial functional effect in MCI patients, whether carrying the APOE ϵ 4 allele or not. A continued and personalized, PE program is needed to maintain the physical status of the elderly, by improving

the strength, the endurance and the balance of this high-risk elderly group. The “exergaming” may be one of the optimal choices for the elderly without access to organized social structures providing physical training.

Practical Implications

- PE via exergaming has a beneficial functional effect in MCI patients, whether carrying the APOEε4 allele or not
- Active group had a significant improvement at the post-training period compared to the baseline to all the physical parameters under investigation.
- A continued and personalized, PE program, is necessary to maintain an elderly his/her physical status.
- The “exergaming” may be one of the optimal choices for the elderly without access to organized social structures providing physical training.

Disclosure

PDB discloses potential (non-financial and beyond the context of the submitted work) conflicts of interest with PositScience: there is a co-marketing agreement between the company and the Aristotle University of Thessaloniki (AUTH) to exploit Brain HQ within the LLM Care self-funded initiative that emerged as the non-for-profit business exploitation of the LLM Project originally funded by the ICT-CIP-PSP Program of the European Commission. FitForAll (FFA) has been developed in the AUTH during the LLM Project. It now forms part of LLM Care. The rest of the authors report no conflicts of interest in this work.

Lossary of Abbreviations

2-MST: 2- Minute Step Test

AC: Active Control

ACT: Arm Curl Test

AD: Alzheimer’s disease

APOE: Apolipoprotein E

BDNF: Brain-derived neurotrophic factor

BMI: Body Mass Index

BST: Back Scratch Test

CIP: Competitiveness and Innovation framework Program

CIRS: Cumulative illness Rate Scale

CSaRT: Chair Sit-and-Reach Test

CST: Chair Stand Test

CT: Computed Tomography

CTC: Cognitive training Control

FFA: Fit For All

FRSSD: Functional Rating Scale of Symptoms of Dementia

FUaG: Foot Up-and-Go Test

FUCAS: Functional and Cognitive Assessment Test

GDS: Geriatric Depression Scale

IADL: Instrumental Activities of Daily Living scale

ICT: Information and Communication Technologies

LLM: Long Lasting Memories

MCI: Mild Cognitive Impairment

MMSE: Mini-Mental State Examination

MOCA: Montreal Cognitive Assessment

MRI: Magnetic Resonance Imaging

PSP: Policy Support Program

PTC: Physical training Control

TrailB: Trail Making Test part-B

References

- [1] MacAuley D, Bauman A, Frémont P. Exercise: not a miracle cure, just good medicine. *Br J Sports Med.* 2016;50(18):1107-1108. doi:10.1136/bmj.h1416
- [2] Ruegsegger GN, Booth FW. Health Benefits of Exercise. *Cold Spring Harb Perspect Med.* Published online May 15, 2017;a029694. doi: 10.1101/cshperspect.a029694
- [3] Albert MS, DeKosky ST, Dickson D, et al. The diagnosis of mild cognitive impairment due to Alzheimer’s disease: recommendations from the National Institute on Aging-Alzheimer’s Association workgroups on diagnostic guidelines for Alzheimer’s disease. *Alzheimers Dement.* 2011;7(3):270-279. doi:10.1016/j.jalz.2011.03.008
- [4] Counts SE, He B, Nadeem M, Wu J, Scheff SW, Mufson EJ. Hippocampal Drebrin Loss in Mild Cognitive Impairment. *Neurodegener Dis.* 2012;10(1-4):216-219. doi:10.1159/000333122
- [5] Krell-Roesch J, Feder NT, Roberts RO, et al. Leisure-Time Physical Activity and the Risk of Incident Dementia: The Mayo Clinic Study of Aging. *J Alzheimers Dis.* 2018;63(1):149-155.

- doi:10.3233/JAD-171141
- [6] O'Donoghue MC, Murphy SE, Zamboni G, Nobre AC, Mackay CE. APOE genotype and cognition in healthy individuals at risk of Alzheimer's disease: A review. *Cortex*. 2018;104(9):103-123. doi:10.1016/j.cortex.2018.03.025
- [7] Wolf AB, Caselli RJ, Reiman EM, Valla J. APOE and neuroenergetics: an emerging paradigm in Alzheimer's disease. *Neurobiol Aging*. 2013;34(4):1007-1017. doi:10.1016/j.neurobiolaging.2012.10.011
- [8] Jensen CS, Simonsen AH, Siersma V, et al. Patients with Alzheimer's disease who carry the APOE ϵ 4 allele benefit more from physical exercise. *Alzheimer's Dement Transl Res Clin Interv*. 2019;5:99-106. doi:10.1016/j.trci.2019.02.007
- [9] Anderson-Hanley CP, Barcelos NPM, ZIMMERMAN EAM, et al. The Aerobic and Cognitive Exercise Study (ACES) for Community-Dwelling Older Adults With or At-Risk for Mild Cognitive Impairment (MCI): Neuropsychological, Neurobiological and Neuroimaging Outcomes of a Randomized Clinical Trial. *Front Aging Neurosci*. 2018;10(MAY):76. Accessed April 12, 2018. www.frontiersin.org
- [10] Maffei L, Picano E, Andreassi MG, et al. Randomized trial on the effects of a combined physical/cognitive training in aged MCI subjects: the Train the Brain study. *Sci Rep*. 2017;7:39471. doi:10.1038/srep39471
- [11] Krell-Roesch J, Syrjanen JA, Vassilaki M, et al. Association of non-exercise physical activity in mid- and late-life with cognitive trajectories and the impact of APOE ϵ 4 genotype status: the Mayo Clinic Study of Aging. *Eur J Ageing*. 2019;16(4):491-502. doi:10.1007/s10433-019-00513-1
- [12] Jeon SY, Byun MS, Yi D, et al. Midlife Lifestyle Activities Moderate APOE ϵ 4 Effect on in vivo Alzheimer's Disease Pathologies. *Front Aging Neurosci*. 2020;12. doi:10.3389/fnagi.2020.00042
- [13] Ma DY, Wong CHY, Leung GTY, Fung AWT, Chan WC, Lam LCW. Physical Exercise Helped to Maintain and Restore Functioning in Chinese Older Adults With Mild Cognitive Impairment: A 5-Year Prospective Study of the Hong Kong Memory and Ageing Prospective Study (HK-MAPS). *J Am Med Dir Assoc*. 2017;18(4):306-311. doi:10.1016/j.jamda.2016.10.003
- [14] Karssemeijer EGA, Aaronson JA, Bossers WJR, Donders R, Olde Rikkert MGM, Kessels RPC. The quest for synergy between physical exercise and cognitive stimulation via exergaming in people with dementia: A randomized controlled trial 17 *Psychology and Cognitive Sciences* 1701 *Psychology* 11 *Medical and Health Sciences* 1103 *Clinical Sciences*. *Alzheimer's Res Ther*. 2019;11(1):1-13. doi:10.1186/s13195-018-0454-z
- [15] Long Lasting Memories. <http://www.longlastingmemories.eu/>
- [16] Tsolaki AC, Tsolaki M, Pandria N, et al. Web-Based Intervention Effects on Mild Cognitive Impairment Based on Apolipoprotein E Genotype: Quasi-Experimental Study. *J Med Internet Res*. 2020;22(5):e14617. doi:10.2196/14617
- [17] Rikli RE, Jones CJ. Functional Fitness Normative Scores for Community-Residing Older Adults, Age 60-94. *J Aging Phys Act*. 1999;7:162-181.
- [18] Jones CJ, Rikli RE. Measuring functional fitness of older adults. *J Act Aging*. 2002;1:25-30. Accessed February 18, 2019. <http://www.dnbn.univr.it/documenti/Occorrenzan/matdid/matdid182478.pdf>
- [19] Petersen RC. Mild cognitive impairment as a diagnostic entity. *J Intern Med*. 2004;256(3):183-194. doi:10.1111/j.1365-2796.2004.01388.x
- [20] Bamidis PD, Fissler P, Papageorgiou SG, et al. Gains in cognition through combined cognitive and physical training: the role of training dosage and severity of neurocognitive disorder. *Front Aging Neurosci*. 2015;7(JUL):152. doi:10.3389/fnagi.2015.00152
- [21] Konstantinidis EI, Billis AS, Mouzakidis CA, Zilidou VI, Antoniou PE, Bamidis PD. Design, Implementation, and Wide Pilot Deployment of FitForAll: An Easy to use Exergaming Platform Improving Physical Fitness and Life Quality of Senior Citizens. *IEEE J Biomed Heal Informatics*. 2016;20(1):189-200. doi:10.1109/JBHI.2014.2378814
- [22] Koutroumani M, Daniilidou M, Giannakouros T, et al. The deletion variant of α 2b-adrenergic receptor is associated with decreased risk in Alzheimer's disease and mild cognitive impairment. *J Neurol Sci*. 2013;328(1-2):19-23. doi:10.1016/j.jns.2013.02.003
- [23] Douka S, Zilidou VI, Lilou O, Tsolaki M. Greek Traditional Dances: A Way to Support Intellectual, Psychological, and Motor Functions in Senior Citizens at Risk of Neurodegeneration. *Front Aging Neurosci*. 2019;11:6. doi:10.3389/fnagi.2019.00006
- [24] Poptsi E, Moraitou D, Eleftheriou M, et al. Normative Data for the Montreal Cognitive Assessment in Greek Older Adults With

- Subjective Cognitive Decline, Mild Cognitive Impairment and Dementia. *J Geriatr Psychiatry Neurol.* 2019;32(5):265-274. doi:10.1177/0891988719853046
- [25] Brown BM, Peiffer J, Rainey-Smith SR. Exploring the relationship between physical activity, beta-amyloid and tau: A narrative review. *Ageing Res Rev.* 2019;50:9-18. doi:10.1016/j.arr.2019.01.003
- [26] Brandon JA, Farmer BC, Williams HC, Johnson LA. APOE and alzheimer's disease: Neuroimaging of metabolic and cerebrovascular dysfunction. *Front Aging Neurosci.* 2018;10(JUN). doi:10.3389/fnagi.2018.00180
- [27] Wilson MG, Ellison GM, Cable NT. Basic science behind the cardiovascular benefits of exercise. *Br J Sports Med.* 2016;50(2):93-99. doi:10.1136/bjsports-2014-306596rep
- [28] Skjæret-Maroni N, Vonstad EK, Ihlen EAF, Tan X-C, Helbostad JL, Vereijken B. Exergaming in Older Adults: Movement Characteristics While Playing Stepping Games. *Front Psychol.* 2016; 7:964. doi:10.3389/fpsyg.2016.00964
- [29] Puleio A. Neuro-exergaming for older adults with mild cognitive impairment (MCI): A single bout of interactive Physical and Cognitive Exercise (iPACES v2.5). Honor Theses. Published online March 1, 2019. Accessed September 8, 2020. <https://digitalworks.union.edu/theses/2341>
- [30] Fu AS, Gao KL, Tung AK, Tsang WW, Kwan MM. Effectiveness of Exergaming Training in Reducing Risk and Incidence of Falls in Frail Older Adults With a History of Falls. *Arch Phys Med Rehabil.* 2015;96(12):2096-2102. doi:10.1016/j.apmr.2015.08.427

