

- Forest of Minnesota. U.S. Department of Agriculture Forest Service Research Paper NC-97.
- Mech, L. D.** 1986. Wolf numbers and population trend in the Superior National Forest, 1967-1985. Research Paper NC-270. U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station, St. Paul, Minnesota. 6 pages.
- Mech, L. D., and L. Boitani.** 2003. Social ecology. Pages 1-34 in *Wolves: Ecology, Behavior, and Conservation*. Edited by L. D. Mech and L. Boitani. University of Chicago Press, Chicago, Illinois.
- Mech, L. D., and M. E. Nelson.** 2000. Do Wolves affect white-tailed buck harvest in Minnesota? *Journal of Wildlife Management* 64: 129-136.
- Nelson, M. E., and L. D. Mech.** 1981. Deer social organization and Wolf predation in northeastern Minnesota. *Wildlife Monographs* 77: 1-53.
- Odum, E. P., and E. J. Kuenzler.** 1955. Measurement of territory and home range size in birds. *Auk* 72: 128-137.
- Scott, B. M. V., and D. M. Shackleton.** 1982. A preliminary study of the social organization of the Vancouver Island Wolf (*Canis lupus crassodon*). Pages 12-41 in *Wolves of the world*. Edited by F. H. Harrington and P. C. Paquet. Noyes Publication, Park Ridge, New Jersey.

Received 25 August 2002

Accepted 3 June 2004

Extraordinary Size and Survival of American Black Duck, *Anas rubripes*, Broods

JERRY R. LONGCORE and DANIEL G. MCAULEY

United States Geological Survey, Patuxent Wildlife Research Center-Orono, 5768 South Annex A, Orono, Maine 04469-5768 USA

Longcore, Jerry R., and Daniel G. McAuley. 2004. Extraordinary size and survival of American Black Duck, *Anas rubripes*, broods. *Canadian Field-Naturalist* 118(1): 129-131.

Two female American Black Ducks (*Anas rubripes*) were initially observed during June 1982 with 20 Class Ib or 18-22 Class Ia-b ducklings in two wetlands in Hancock County, Cherryfield, Maine. Fifteen of 20 ducklings (75%) in one brood and 16 of 18-22 ducklings (72-89%) in the other brood survived to fledge. These large broods probably resulted from post-hatch brood amalgamation.

Key Words: American Black Duck, *Anas rubripes*, brood size, duckling survival, post-hatch brood amalgamation, Maine.

Exceptionally large broods of North American dabbling ducks (*Anas* spp.) that exceed average clutch size (8-10 eggs, Zammuto 1986) occur when females (1) lay extraordinarily large clutches, (2) lay eggs in nests of conspecifics ("pre-hatch brood amalgamation" (Pre-HBA), or brood parasitism; Eadie et al. 1988), or (3) hatch their own clutches and acquire the brood of another female ("post-hatch brood amalgamation" (Post-HBA); Eadie et al. 1988). Pre-HBA, which can be either inter- or intra-specific, and post-HBA occur infrequently in the Anatini; only 3 of 9 species of Anatini were reported by Eadie et al. (1988) for either type. Tufts (1986) in Nova Scotia reported brood amalgamation for three American Black Duck broods when he released orphaned ducklings near females with broods. Herein, we report two records of probable post-HBA resulting in two extremely large broods of wild American Black Ducks in Maine.

Study Area and Methods

We observed the two broods in two Beaver (*Castor canadensis*) flowages (Snake Flowage, 44°37'N, 68°06'W; BFA Flowage, 44° 39'N, 68°07'W) that were 20 km northwest of Cherryfield, Maine, in township T10 SD, a forested area that has negligible acid-neutralizing

capacity and low nutrients in wetlands (Norton 1980). We obtained morphometric and water chemistry characteristics of wetlands by methods of McAuley and Longcore (1988). We mapped and classified (Cowardin et al. 1979) both wetlands, and we sampled invertebrates in one (Snake Flowage), as part of related fieldwork (J. R. Longcore, unpublished data). All observations of broods followed the survey protocol of Longcore and Ringelman (1980). Morning visits on wetlands began 0.5 hour before sunrise and lasted two hours; the 2-h evening visit ended ≥ 0.5 hour after sunset. Broods were always sought on both wetlands simultaneously, and observers scanned wetlands with binoculars (7x50) and spotting scopes (20-60x) from elevated (4-5 m high) platforms. We backdated brood age (Gollop and Marshall 1954*) to determine approximate dates that first eggs were laid.

Results

During 3 June – 12 July 1982, we observed two American Black Duck broods that were twice (20 and 18-22 ducklings) the size of average broods. Each brood was seen three times. Both broods were seen on the same wetland on the same day (8 June) by DGM. Both broods were observed on 12 July during the even-

Table 1. Characteristics of the two brood-rearing wetlands with the large American Black Duck broods in Cherryfield, Maine, 1982.

Variable	Snake Flowage	BFA Flowage
Wetland System, Class	Palustrine, Forested Wetland	Palustrine, Forested Wetland
Basin area (ha)	4.9	4.5
Surface water area (ha)	4.9	3.4
% Submergents	10	78
% Emergents	28	22
% Flooded timber	41	78
pH (in situ)	5.51	6.13
Alkalinity ($\mu\text{eq L}^{-1}$)	58.8	83.0
Conductivity ($\mu\text{S cm}^{-1}$)	24.0	23.0
Calcium ($\mu\text{eq L}^{-1}$)	59.4	77.5
Phosphorus ($\mu\text{g L}^{-1}$)	70	170
Color (Hazen units)	170	150

ing visit, each by a different observer on a different wetland, thereby corroborating that two different broods existed as first identified by plumage characteristics (Gollop and Marshall 1954*). The brood of 20 declined to 15 (75% survival) and the brood of 18-22 declined to 16 (73 or 89% survival) at Class III size near fledging. The features of the two wetlands are described by the variables in Table 1. We sampled invertebrates in Snake Flowage, which contained the highest ($P < 0.0001$) number of aquatic Insecta per sample (mean = 257) compared with nine other wetlands sampled in the area (J. R. Longcore, unpublished data).

Discussion

At Moosehorn National Wildlife Refuge, Maine, and at Lake Dalhousie, Nova Scotia (J. R. Longcore, unpublished data), we have observed Class III American Black Duck broods of 10 ducklings, which are near mean clutch size. The two large broods, however, were twice the clutch size (mean ± 1 SD) of American Black Ducks in Maine (10.4 ± 1.3), in Vermont (9.6 ± 1.8 ; Coulter and Miller 1968), in Maryland (9.1 ± 1.8 ; Stotts and Davis 1960), or in Quebec (9.2 ± 1.7 ; Reed 1970). Other studies of American Black Ducks have reported large clutches but rarely as large as the broods we observed. Reed (1970) reported sizes of mostly first clutches during 12-25 April, as 13, 14, 15, and 17 eggs in Quebec. Both Coulter and Miller (1968) and Stotts and Davis (1960) reported clutch sizes of 14 or 15 eggs, and Kremenz et al. (1991) reported that in 1982 American Black Ducks nesting on islands in Chesapeake Bay averaged 10.2 ± 3.0 eggs per clutch, ranging from 7-20 eggs, but it was unknown whether large clutches were from one female.

For our two wetlands adequate food seemed available for females to lay large clutches because the highest mean number of invertebrates per sweep net

sample was from Snake Flowage among 10 wetlands sampled (J. R. Longcore, unpublished data). By back-dating clutches we determined that these females, if they laid the entire clutch, would have initiated egg laying about 7 and 14 April, similar to early egg dates of 1-10 April for Maine and Vermont (Coulter and Miller 1968). Although ducks may adjust clutch size to environmental conditions (Skutch 1967), if these two females had laid 20 to 22 eggs per clutch, mass of the clutch would have been 1248 – 1373 g ($62.4 \text{ g/egg} \times 20$ or 22 eggs), which would have equaled or exceeded mass of a female and been >2 times the average mass of a usual clutch (mean = 580.3 g, Arnold 1988). Therefore, it is possible that each female produced the entire clutch for these large broods, but it seems improbable.

The possibility of intra-specific pre-HBA (hatch brood amalgamation) is supported by Stotts and Davis (1960) who reported two instances of American Black Duck females laying eggs in the nest of another female. They (Stotts and Davis 1960: 145) also commented that "Eleven others [clutches] may have been the result of similar parasitism (a total prevalence of 1.8 percent)." An example of inter-specific pre-HBA also was recorded by Stotts and Davis (1960) who documented that an American Black Duck female began laying in a Mallard (*Anas platyrhynchos*) nest that contained five eggs. The Mallard laid four more eggs before deserting, but the American Black Duck laid 11 eggs to equal a 20-egg mixed clutch. Also, they reported that two American Black Duck females nested within 46 cm of each other on an offshore duck blind in Chesapeake Bay, Maryland; one female gradually incorporated the other's clutch into her own and incubated all eggs (Stotts and Davis 1960: 142). Large clutches associated with ducks nesting on islands, however, may be related to high nest density (i.e., 25.2 – 35.7 nests/ha on Bodkin Island, Chesapeake Bay, Kremenz et al. 1991); but is uncommon elsewhere (e.g., 0.06 – 0.12 nest/ha in Maine bogs (Coulter and Miller 1968: 35). Because nest sites are not limiting in Maine or across the breeding range, nesting females are widely dispersed, except on islands (Coulter and Miller 1968), nests are well hidden (Bent 1923), and females are secretive when returning to nests in twilight hours (J. R. Longcore, unpublished data), it seems improbable that our large broods resulted from intra-specific pre-HBA (Beauchamp 1997). The third explanation for the large broods is intra-specific post-HBA. For this scenario, two additional clutches of 10-12 eggs must have hatched at the same time and ducklings in those broods were then acquired by the females that we observed. We know that two other American Black Duck broods (i.e., of five and 10 ducklings) used Snake Flowage at the same time as the large broods. To account for this scenario of post-HBA, six American Black Duck broods of similar age must have been associated with these two wetlands. Concentration of broods on high quality

wetlands is common (Longcore et al. 1998); indeed these two wetlands also supported one Green-winged Teal (*Anas crecca*), three Wood Duck (*Aix sponsa*), and four Hooded Merganser (*Lophodytes cucullatus*) broods. For the 20-duckling brood, JRL recorded on 24 June that some ducklings appeared to be in different age Classes (i.e. IIb and IIc), which suggests post-HBA; however, the duckling brood of 18-22 appeared as all the same age to DGM. The two brood-rearing females were large, extremely attentive, and especially adept in eliciting rapid responses from ducklings by uttering a few low calls. Although females are capable of laying large clutches and the remote possibility of nest parasitism (pre-HBA) exists, we conclude that the most plausible explanation for these two large broods was intra-specific post-HBA.

Acknowledgments

We thank P. E. Malicky for assistance during brood surveys and J. M. Eadie, T. W. Arnold, and A. J. Erskine for review and helpful suggestions on drafts of the manuscript.

Documents Cited (* marked in text)

Gollop, J. B., and W. H. Marshall. 1954. A guide for aging duck broods in the field. Mississippi Flyway Council Technical Report, Minneapolis, Minnesota. 14 pages.

Literature Cited

Arnold, T. W. 1988. Life histories of North American game birds: a reanalysis. *Canadian Journal of Zoology* 66: 1906-1912.

Beauchamp, G. 1997. Determinants of intraspecific brood amalgamation in waterfowl. *Auk* 114: 11-21.

Bent, A. C. 1923. Life histories of North American Wild Fowl. Order Anseres (Part). Smithsonian Institution, U.S. National Museum, Government Printing Office, Washington, D.C.

Coulter, M. W., and W. R. Miller. 1968. Nesting biology of Black Ducks and Mallards in northern New England. Vermont Fish and Game Department Bulletin 68-2. 74 pages.

Cowardin, L. M., V. Carter, F. C. Golet, and E. T. LaRoe. 1979. Classification of wetlands and deepwater habitats of the United States. U.S. Fish and Wildlife Service, FWS/OBS-79/31. 103 pages.

Eadie, J. McA., F. P. Kehoe, and T. D. Nudds. 1988. Pre-hatch and post-hatch brood amalgamation in North American Anatidae: a review of hypotheses. *Canadian Journal of Zoology* 6: 1709-1721.

Krementz, D. G., V. D. Stotts, D. B. Stotts, J. E. Hines, and S. L. Funderburk. 1991. Historical changes in laying date, clutch size, and nest success of American black ducks. *Journal of Wildlife Management* 55: 462-466.

Longcore, J. R., and J. R. Ringelman. 1980. Factors affecting waterfowl breeding density and productivity estimates in the northeast. *Transactions of the Northeast Section, The Wildlife Society* 37: 169-181.

Longcore, J. R., D. A. Clugston, and D. G. McAuley. 1998. Brood size of sympatric American black ducks and mallards in Maine. *Journal of Wildlife Management* 62: 142-151.

McAuley, D. G., and J. R. Longcore. 1988. Survival of juvenile Ring-necked Ducks on wetlands of different pH. *Journal of Wildlife Management* 52: 169-176.

Norton, S. A. 1980. Geological factors controlling the sensitivity of aquatic ecosystems to acid precipitation. Pages 521-531 in *Atmospheric sulfur deposition, environmental impact and health effects: Proceedings of the Second Life Sciences Symposium, Potential Environmental and Health Consequences of Atmospheric Sulfur Deposition, Gatlinburg, TN, October 14-18, 1979. Edited by D. S. Shriner, C. R. Richmond, and S. E. Lindberg.* Ann Arbor Science Publishing, Ann Arbor, Michigan.

Reed, A. 1970. The breeding ecology of the Black Duck in the St. Lawrence estuary. Ph.D. dissertation, Laval University, Quebec, Canada. 175 pages.

Skutch, A. F. 1967. Adaptive limitation of the reproductive rate of birds. *Ibis* 109: 579-599.

Stotts, V. D., and D. E. Davis. 1960. The Black Duck in the Chesapeake Bay of Maryland: breeding behavior and biology. *Chesapeake Science* 1: 127-154.

Tufts, R. W. 1986. *Birds of Nova Scotia*. Third Edition. Nimbus Publishing and Nova Scotia Museum, Halifax, Nova Scotia.

Zammuto, R. M. 1986. Life histories of birds: clutch size, longevity, and body mass among North American game birds. *Canadian Journal of Zoology* 64: 2739-2749.

Received 28 February 2002

Accepted 15 June 2004