

Conference or Workshop Item

Roy, Helen E.; Adrians, Tim; **Isaac, Nick J.B.**; Kenis, Marc; Onkelinx, Thierry; Martin, Gilles San; Brown, Peter M.J.; Hautier, Louis; Poland, Remy; **Roy, David B.**; **Comont, Richard**; Eschen, Rene; Frost, Robert; Zindel, Renate; Van Vlaenderen, Johan; Nedved, Oldrich; Ravn, Hans Peter; Gregoire, Jean-Claude; de Biseau, Jean-Christophe; Maes, Dirk. 2011 *Harmonia axyridis* implicated in native European ladybird declines. In: *2nd Meeting of the IOBC/WPRS Working Group "Benefits and Risks of Exotic Biological Control Agents", Hluboka, Czech Republic., 30 October - 3 November 2011.* (Unpublished)

This version available at <http://nora.nerc.ac.uk/16479>

NERC has developed NORA to enable users to access research outputs wholly or partially funded by NERC. Copyright and other rights for material on this site are retained by the authors and/or other rights owners. Users should read the terms and conditions of use of this material at <http://nora.nerc.ac.uk/policies.html#access>

Contact CEH NORA team at
noraceh@ceh.ac.uk

***Harmonia axyridis* implicated in native European ladybird declines**

Helen E. ROY^{1*}, Tim ADRIANS², Nick J.B. ISAAC¹, Marc KENIS³, Thierry ONKELINX³, Gilles San MARTIN⁴, Peter M.J. BROWN⁵, Louis HAUTIER⁶, Remy POLAND⁷, David B. ROY¹, Richard COMONT¹, René ESCHEN³, Robert FROST, Renate ZINDEL^{3,8}, Johan Van VLAENDEREN³, Oldřich NEDVĚD⁹, Hans Peter RAVN¹⁰, Jean-Claude GRÈGOIRE¹¹, Jean-Christophe de BISEAU¹², Dirk MAES²

¹*Centre for Ecology & Hydrology, Benson Lane, Crowmarsh Gifford, Oxfordshire, OX10 8BB, UK*

²*Research Institute for Nature and Forest (INBO), Kliniekstraat 25, B-1070 Brussels, Belgium*

³*CABI Europe-Switzerland, 1 Rue des Grillons, 2800 Delémont, Switzerland*

⁴*Universite catholique de Louvain, Earth and Life Institute, Biodiversity Research Centre, Behavioural Ecology and Conservation group Croix du Sud 4, B-1348 Louvain-la-Neuve, Belgium*

⁵*Animal & Environmental Research Group, Department of Life Sciences, Anglia Ruskin University, East Road, Cambridge, CB1 1PT, UK*

⁶*Unité Protection des plantes et écotoxicologie, Département Sciences du vivant, Centre wallon de Recherches agronomiques, Rue de Liroux, 2, B-5030 Gembloux, Belgium*

⁷*Clifton College, 32 College Rd, Clifton, Bristol, Avon, BS8 3JH, UK*

⁸*Department of Biology, University of Fribourg, Chemin du Musée 10, 1700 Fribourg, Switzerland*

⁹*University of South Bohemia, Faculty of Biological Sciences and Institute of Entomology, Academy of Sciences of the Czech Republic, Branišovská 31, CZ-37005 České Budějovice, Czech Republic*

¹⁰*University of Copenhagen, Forest & Landscape, Rolighedsvej 23, DK-1958 Frederiksberg C. Denmark*

¹¹*Lutte biologique et Ecologie spatiale (Biological Control and Spatial Ecology Lab), CP 160/12, Université Libre de Bruxelles, 50 av FD Roosevelt, 1050 Bruxelles, Belgium*

¹²*Evolution Biologique et Ecologie, Université Libre de Bruxelles, CP 160/12, Av. F.D. Roosevelt 50 - 1050 Bruxelles, Belgium*

Rates of global extinction are orders of magnitude higher than historical estimates and show no sign of slowing. The Convention on Biological Diversity and the 10th Conference of the Parties (Nagoya in 2010), identified invasive alien species (IAS) as one of five major pressures driving biodiversity loss, and ultimately extinction of species. However, there are few examples of causal relationships between IAS and species declines. IAS afford a unique opportunity to accurately assess threats to biodiversity because the time at which an IAS arrives within an ecosystem is often known, unlike other drivers of change. We examined trends in distribution of native ladybirds from large-scale and long-term annual citizen-science surveys before and after the arrival of the predatory harlequin (or Asian) ladybird

Harmonia axyridis, an IAS that is rapidly expanding across North America and Europe. We report rapid, dramatic and ongoing declines in the distribution of formerly common and widespread native ladybirds in direct response to the arrival of *H. axyridis* in Belgium and Britain. The dramatic decline of *A. bipunctata* over the five years following the arrival of *H. axyridis* is of particular note. Trends in ladybird abundance revealed similar patterns of declines in ladybirds across Belgium, Britain and Switzerland. Together, these parallel analyses show *H. axyridis* to be displacing native ladybirds with a high niche overlap, probably through predation and competition. Such rapid biotic homogenisation at the continental scale could impact on the resilience of ecosystems and severely diminish the services they deliver.