

## ARTICLE

## Open Access

# The placenta protects the fetal circulation from anxiety-driven elevations in maternal serum levels of brain-derived neurotrophic factor

Hayley Dingsdale<sup>1</sup>, Xinsheng Nan<sup>1</sup>, Samantha M. Garay<sup>1</sup>, Annett Mueller<sup>2</sup>, Lorna A. Sumption<sup>1</sup>, Pedro Chacón-Fernández<sup>1,3</sup>, Isabel Martinez-Garay<sup>1</sup>, Cedric Ghevaert<sup>2</sup>, Yves-Alain Barde<sup>1</sup> and Rosalind M. John<sup>1</sup>

## Abstract

Brain-derived neurotrophic factor (BDNF) plays crucial roles in brain function. Numerous studies report alterations in BDNF levels in human serum in various neurological conditions, including mood disorders such as depression. However, little is known about BDNF levels in the blood during pregnancy. We asked whether maternal depression and/or anxiety during pregnancy were associated with altered serum BDNF levels in mothers ( $n = 251$ ) and their newborn infants ( $n = 212$ ). As prenatal exposure to maternal mood disorders significantly increases the risk of neurological conditions in later life, we also examined the possibility of placental BDNF transfer by developing a new mouse model. We found no association between maternal symptoms of depression and either maternal or infant cord blood serum BDNF. However, maternal symptoms of anxiety correlated with significantly raised maternal serum BDNF exclusively in mothers of boys ( $r = 0.281$ ;  $P = 0.005$ ;  $n = 99$ ). Serum BDNF was significantly lower in male infants than female infants but neither correlated with maternal anxiety symptoms. Consistent with this observation, we found no evidence for BDNF transfer across the placenta. We conclude that the placenta protects the developing fetus from maternal changes in serum BDNF that could otherwise have adverse consequences for fetal development.

## Introduction

Perinatal mental health problems impact nearly one-quarter of all pregnancies<sup>1</sup>. In the UK alone, this costs society > £8 billion for each year of births<sup>1</sup> due to the risk of substantial morbidity to mother and child<sup>2</sup> and negative outcomes for the child<sup>3–5</sup>. Despite the considerable clinical and financial burden to society, the biological mechanisms linking maternal mood disorders to adverse outcomes are unknown.

One mechanism potentially accounting for the programming of disease risk is passage across the placenta of

maternal factors impacting fetal development during sensitive periods, as has been suggested for the stress hormone cortisol<sup>3,6</sup>. Brain-derived neurotrophic factor (BDNF) is a member of the nerve growth factor family expressed in the central and peripheral nervous system with a key role in the development and function of neurons<sup>7</sup>. In primates, but not mice, significant levels of BDNF are detected in circulating blood platelets<sup>8</sup>, originating from megakaryocytes (MKs)<sup>9</sup>. Decreased serum BDNF has been reported in numerous conditions, including depression<sup>10–15</sup>, Huntington's<sup>16</sup> and Alzheimer's disease<sup>17</sup>, whilst increased BDNF has been associated with autism spectrum disorder<sup>18</sup>. Circulating BDNF is also known to fluctuate with exercise<sup>19</sup>.

Pregnancy is a unique state where the blood systems of two individuals, the mother and the fetus, are intimately associated but not in direct contact. In humans, there is a

Correspondence: Rosalind M. John ([JohnRM@cardiff.ac.uk](mailto:JohnRM@cardiff.ac.uk))

<sup>1</sup>School of Biosciences, Cardiff University, Cardiff CF10 3AX, UK

<sup>2</sup>Department of Haematology, University of Cambridge, Cambridge CB2 0PT, UK

Full list of author information is available at the end of the article

These authors contributed equally: Hayley Dingsdale, Xinsheng Nan

© The Author(s) 2021



**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

barrier between the maternal and fetal circulations composed of a single continuous layer of syncytiotrophoblast over a layer of villous cytotrophoblast covering the endothelial cell-lined fetal blood vessels<sup>20</sup>. Despite significant morphological differences, rodents also possess the same hemochorial arrangement<sup>21</sup>, with the barrier consisting of a discontinuous layer of sinusoidal trophoblast giant cells and two continuous layers of syncytiotrophoblast covering fetal blood vessels<sup>22</sup>. Thus, in both cases, there are three cellular layers between maternal and fetal blood. Most proteins are too large to cross the placenta. One exception is maternal antibodies which cross the human syncytiotrophoblast by receptor-mediated pinocytosis, an active process requiring the neonatal Fc receptor<sup>23</sup>, and we note that a previous study in mice concluded that maternal BDNF may be another protein that can cross the placenta<sup>24</sup>.

Fluctuations in maternal serum BDNF may consequently adversely influence fetal development. A small number of studies have reported on maternal or fetal serum BDNF levels (see Supplementary Table 1<sup>25–46</sup>). Lower maternal BDNF has been associated with postnatal depression symptoms<sup>46</sup> and lower cord blood BDNF associated with major depression in pregnancy<sup>39</sup> but other studies report no association<sup>33,37</sup>. Two studies reported a relationship between cord blood serum BDNF and maternal anxiety symptoms, again with contrasting findings<sup>33,42</sup>. To date, no study has reported on the status of BDNF in blood from paired maternal and fetal samples in relation to maternal symptoms of both depression and anxiety.

In this study, we quantified BDNF levels in maternal serum and matched cord blood serum in relation to maternally reported symptoms of depression and anxiety, generated as part of the Grown in Wales study<sup>47</sup>. Mouse models have been extensively used to understand the function of brain-derived BDNF<sup>48</sup>; however, BDNF is not detectable in mouse MKs<sup>9</sup> or blood<sup>49</sup>. Consequently, we genetically engineered BDNF expression from MKs to determine the potential for MK-derived maternal BDNF transfer across the placenta near term.

## Methods

### Grown in Wales study

GiW study is a longitudinal pregnancy cohort study. Women aged 18–45 with a singleton, term pregnancy without fetal abnormalities or infectious diseases were recruited at pre-surgical appointments prior to an elective caesarean section (ELCS) at the University Hospital of Wales. In total, 355 women provided written consent; 7 women subsequently withdrew. Perceived depression symptoms were assessed at recruitment using the Edinburgh Postnatal Depression Scale (EPDS)<sup>50</sup> and trait anxiety by form Y-2 on the

Spielberger State-Trait Anxiety Inventory (STAI) test<sup>51</sup>. Maternal venous serum samples were obtained at recruitment and 1–4 days later cord venous blood taken within 2 h of delivery. Blood was collected into Vacutainer blood collection tubes (BD Biosciences, 367837), inverted several times and incubated at room temperature (RT) for 1–2 h. After centrifugation at  $3000 \times g$  for 10 min at 4 °C, serum aliquots were stored at –80 °C. Prior to data analysis, samples were excluded based on the following pre-established *sample exclusion criteria*: women who went into natural labor prior to their planned ELCS were included in the analyses of maternal serum but their cord blood serum samples were excluded; 36 samples from <36 weeks<sup>25</sup> and non-Caucasian<sup>52</sup> pairs were also excluded.

### Serum BDNF ELISA

ELISA measurements of human serum were performed blind to all participant information except participant ID and type (maternal versus fetal), and as described<sup>53</sup>, with minor changes. Briefly, NeutrAvidin-coated plates (ThermoFisher Scientific, 15509) were washed with Buffer A (0.1% TritonX-100 in 0.1 M phosphate buffer: 0.1 M  $\text{KH}_2\text{PO}_4$ , 0.1 M  $\text{Na}_2\text{HPO}_4$ , pH 7.6) before incubating with 13  $\mu\text{g}/\text{ml}$  biotin-BDNF-Ab#1<sup>54</sup> for 2 h at RT. Plates were then washed with Buffer B (Buffer A with 1% BSA), before the addition of 150  $\mu\text{l}$  Buffer A. Next, 50  $\mu\text{l}$  of either sample (1:20 dilution in Buffer B) or standard (in Buffer B) was added and incubated for 6 h at RT while shaking. Following Buffer A washes, 1.25  $\mu\text{g}/\text{ml}$  HRP-BDNF-Ab#9<sup>54</sup> (in Buffer B) was added for 3 h at RT, again shaking. After further Buffer A washes, the chemiluminescent substrate (Roche, 11582950001) was added and signal detected by a microplate reader (FLUOstar OMEGA, BMG Labtech). All plates contained two repeated samples, and samples were analyzed in triplicate. ELISA measurements of mouse serum were performed as above, with the exceptions that experimenters were not blinded to sample information, and the sample incubation step was reduced to 3 h.

### Western blot analysis

Western blots were performed as described<sup>53</sup>. Proteins from 1  $\mu\text{l}$  serum were separated on 4–12% Bis-Tris gradient gels (Invitrogen, NP0321BOX) and transferred to nitrocellulose membranes. After blocking for 1 h in 3% BSA (Sigma-Aldrich, A7906), 3% ECL Prime Blocking reagent (GE Healthcare) in TBS-T, membranes were incubated overnight with anti-BDNF 3C11 antibody (1:2000, Icosagen, 327-100). After washing, 1 h incubation with HRP-conjugated goat anti-mouse IgG1 (1:2000, Invitrogen, PA1-74421) followed, and membranes washed and developed.

### Generation of Rosa26-LSL-BDNF-myc-IG mice

The pZDonor-Mouse-Rosa26 plasmid (Sigma-Aldrich, D9196) was used as the backbone of a Rosa26 targeting vector. We inserted a 4185 bp PCR fragment, containing chimeric intron–exon-loxP-NeopA-loxP-MCS-IRES-eGFP-pA amplified from the pCAGfloxNeoIRESeGFP plasmid (gift from Professor Meng Li, Cardiff University) as a template, to the EcoRI/XmaI site to generate pZDR-LSL-IG. To generate pZDR-LSL-BDNF-myc-IG, a fragment containing the BDNF-myc coding sequence was isolated by double digestion with EcoRI/SacI from pCMV-BDNF-myc (constructed by adding one myc-tag copy at the C-terminus of WT mouse BDNF following the deletion of the last three amino acids<sup>55</sup>) and inserted into the PmeI site of pZDR-LSL-IG.

Mouse ES cells E14TG2a were cultured in gelatin-coated flasks or Petri dishes in GMEM medium (ThermoFisher Scientific, 11710035) with LIF (homemade) and 10% FCS (Biosera, FB-1001/500). To generate Rosa26-LSL-BDNF-myc-IG mouse ES cells, cells were transfected with pZDR-LSL-BDNF-myc-IG and pCMV-RosaL6 ELDmut and pCMV-RosaR4 KKRmut (gifts from Charles Gersbach, Addgene plasmid #37198 and #37199<sup>56</sup>) in a 10:1:1 ratio. Transfection was performed in Nucleofector solution P3 (Lonza, V4XP-3024) using 4D-Nucleofector X Unit (Lonza, AAF-1002X), program CG-104. 5'-homologous recombination was confirmed by generation of a 2.2 kb PCR fragment using primer pair (GTGGAGCCGTTCTGTGAGAC and GTCATAGCCG AATAGCCTCTCCAC), 3' homologous recombination by a 1.3 kb PCR fragment using primer pair (TCGAG CTGGACGGCGACGTAAAC and TTGGGGGAGGAG ACATCCACCTGG).

Rosa26-LSL-BDNF-myc-IG mice (BDNF-myc hereafter) were generated by injection of Rosa26-LSL-BDNF-myc-IG mouse ES cells into C57BL/6 blastocysts. Germ-line transmission was genotyped by PCR as above. Subsequent genotyping was conducted by PCR using two primer pairs (Bdnf pair: GAACTACCCAATCG-TATGTTTCG and CTACAAGTCTTCTTCAGAAATAA GCTT; Rosa26 pair: AAGGGAGCTGCAGTGGAGTA and GTCCCTCCAATTTTACACC) to identify heterozygotes (140 bp and 275 bp products), homozygotes (140 bp), and wild-type animals (275 bp).

### Generation of animals expressing transgene BDNF-myc in platelets

Pf4-Cre mice<sup>57</sup> (The Jackson Laboratory, 008535, C57BL/6 background) were crossed with BDNF-myc animals to generate het BDNF-myc::Pf4-Cre mice. Homozygous BDNF-myc::Pf4-Cre mice were generated by crossing homozygous BDNF-myc with homozygous BDNF-myc::Pf4-Cre mice. Pf4-Cre genotyping was by the production of a 420 bp Pf4-Cre product (primer pair

CCAAGTCTACTGTTTCTCACTC and TGCACAGT CAGCAGGTT) and an internal 302 bp (Y chromosome) and/or 331 bp (X chromosome) (primer pair CTGAA GCTTTTGGCTTTGAG and CCACTGCCAAATTCT TTGG).

### Mouse megakaryocyte culture

Mouse megakaryocytes (Mks) were differentiated from bone marrow as described<sup>9</sup>. Briefly, bone marrow was flushed from femurs and tibias with Hank's balanced salt solution containing 0.38% sodium citrate, 1 mM adenosine, 2 mM theophylline, and 5% FBS. Dissociated cells were cultured in Iscove's modified Dulbecco's medium with L-glutamine, 25 mM HEPES, 5% FBS, 100 units/ml penicillin, 100 µg/ml streptomycin, 25 ng/ml stem cell factor and TPO. Mature Mks were purified on a 1.5–3% BSA gradient and cultured in Cellgro stem cell growth media (CellGenix) for up to 16 h. For immunocytochemistry, Mks were cultured on 200 µg/ml human fibrinogen-coated coverslips for 6 h, fixed for 30 min in 4% paraformaldehyde in PBS, and permeabilized for 15 min with 0.5% TritonX-100 in PBS. Two male animals at the age of nine weeks were used for each genotype group, experimental replicates were biological.

### Whole blood cell and platelet counts

Following dilution of 20 µl of blood with 30 µl acid-citrate-dextrose (ACD) buffer (85 mM trisodium citrate, 65 mM citric acid, 100 mM glucose, pH 5.0) and 50 µl PBS, 1 µl PE Rat Anti-Mouse CD41 (BD Pharmingen, #561850) antibody was added to stain the platelets, and incubated at RT for 30 min. To this, 900 µl of fixative (0.2% formyl saline) was added and incubated for a further 10 min at RT. Finally, 10 µl of this fixed cell suspension was then added to 990 µl PBS. A hemocytometer was used to count whole blood cells using transmitted light, and platelets counted using blue epifluorescence. Four animals at the age of four months and of mixed sex were used for each genotype group, experimental replicates were biological.

### FACS analysis of platelets and leukocytes

Blood was collected by cardiac puncture into 100 µl ACD buffer and mixed with 1 ml DMEM with 10% heparin. To this, 8 ml Erythrocyte-Lysis-Buffer (10 mM KHCO<sub>3</sub>, 155 mM NH<sub>4</sub>Cl, 0.1 mM EDTA, pH 7.5) was added before 15 min incubation on ice. Next, 5 ml PBS/0.5% BSA was added, before centrifugation for 10 min at 200 × g at 4 °C, with the resulting cell pellet resuspended in 1 ml PBS/0.5% BSA. In all, 200 µl of cell suspension was used for immunostaining (20 min on ice) to examine immune cell subpopulations (all antibodies from BD Pharmingen, 1:400 dilutions). The cell suspension was washed with 1 ml PBS/0.5% BSA and centrifuged at 600 ×

g for 6 min at 4 °C. To detect all immune cells as well as the myeloid fraction, APC-conjugated anti-CD11b (#561690) and APC-conjugated anti-CD45 (#561018) were used. T and B cells were identified by PE anti-CD4 (#557308), PerCP anti-CD8a (#561092), and APC anti-CD19 (#561738) antibodies. Prior to measurement, the cell pellet was resuspended in 300 µl PBS/0.5% BSA, and DAPI was used to exclude dead cells.

To stain platelets, 30 µl ACD/PBS (1:10) with PE rat anti-mouse CD41 (#561850) or PE rat IgG1 control (#551979) antibodies were added to 20 µl blood and incubated for 30 min at RT. 1 ml 0.2% formyl saline and DAPI was then added before FACS analysis. PE rat IgG1 isotype controls were used for gating. For analysis, doublets were gated out on DAPI-negative events. Two female animals at the age of 4 months were used for each genotype group, experimental replicates were biological.

#### Platelet isolation

Blood was collected by cardiac puncture into 100 µl ACD buffer. Blood cells were removed by centrifugation at 200 × g for 5 min at RT without brake and platelet-rich plasma (PRP) collected. After centrifugation at 200 × g for 5 min at RT without brake to remove residual blood cells, the further cleared PRP was collected and centrifuged at 900 × g for 10 min at RT without brake. Precipitated platelets were resuspended in ACD and Ca<sup>2+</sup>-free Tyrodes buffer (134 mM NaCl, 2.9 mM KCl, 0.34 mM Na<sub>2</sub>HPO<sub>4</sub>, 12 mM NaHCO<sub>3</sub>, 20 mM HEPES, 1 mM MgCl<sub>2</sub>, 5 mM glucose) and centrifuged at 200 × g for 5 min at RT without brake. The further cleared PRP was collected and centrifuged at 900 × g for 10 min at RT without brake. The platelet pellet was then resuspended in Ca<sup>2+</sup>-free Tyrode's buffer for platelet spreading assays.

#### Platelet spreading assay

To stimulate activation, 2 mM CaCl<sub>2</sub> and 0.01 U/ml thrombin (Sigma-Aldrich) were added to 200 µl of rested platelets, and platelets transferred immediately onto fibrinogen-coated coverslips. Following incubation at 37 °C for 45 min, attached platelets were fixed using 10% neutral buffered formalin solution for 20 min. After washing with PBS/0.1% BSA, Alexa Fluor 555 Phalloidin (1:1000 in PBS/0.1% BSA, ThermoFisher Scientific, A34055) was used for staining. Activated platelets were imaged by confocal microscopy (LSM 780, Zeiss), and platelet sizes analyzed using ImageJ<sup>58</sup>. Two female animals at the age of four months were used for each genotype group, experimental replicates were biological.

#### Mouse serum generation

BDNF-myc<sup>+/+</sup>;Cre<sup>+/-</sup> females were crossed with BDNF-myc<sup>+/+</sup>;Cre<sup>-/-</sup> male mice to generate progeny that were either BDNF-myc<sup>+/+</sup>;Cre<sup>+/-</sup> or BDNF-myc<sup>+/+</sup>;

Cre<sup>-/-</sup>. On 3 consecutive days, dams were sacrificed at E18.5 and blood collected by cardiac puncture. Fetuses were removed, before decapitation to collect blood. Blood was incubated for 1 h at RT, followed by 1 h at 4 °C and centrifugation at 2000 × g for 10 min at 4 °C; serum aliquots were stored at -80 °C. Three litters of fetuses were analyzed at E18.5 with a total of 13 BDNF-myc<sup>+/+</sup>;Cre<sup>+/-</sup> pups and 11 BDNF-myc<sup>+/+</sup>;Cre<sup>-/-</sup> fetuses, experimental replicates were primarily biological. For adult mice, blood was collected by cardiac puncture, and nine Cre<sup>-/-</sup>, six BDNF-myc<sup>+/-</sup>;Cre<sup>+/-</sup>, and five BDNF-myc<sup>+/+</sup>;Cre<sup>+/-</sup> animals, 2–3 months in age and of mixed sex were analyzed by western blot. Data shown is representative of four experiments, experimental replicates were biological.

#### Statistics

RStudio V1.2.5001<sup>59</sup> was used for graphs and statistical analysis (packages: “car”<sup>60</sup>, “dplyr”<sup>61</sup>, “ggplot2”<sup>62</sup>, “ggpubr”<sup>63</sup>, “pastecs”<sup>64</sup>, and “reshape2”<sup>65</sup>). Median (interquartile range (IQR)) are presented unless otherwise stated. Normality was analyzed by Shapiro–Wilk test, and variance by Levene's test where appropriate. Group differences were analyzed by Student's *t* test (two-sided, with unequal variance corrected for as a default parameter; normally distributed data), Wilcoxon signed-rank test (paired data), Wilcoxon rank test (non-paired data), or linear modeling, and associations by Spearman's rank correlation or linear modeling. Confounding variables controlled for in linear models of human serum values were maternal hypertension<sup>66</sup>, gestational diabetes<sup>35</sup>, maternal BMI at booking<sup>67</sup>, and smoking<sup>41</sup>, alcohol<sup>68</sup>, strenuous exercise<sup>19</sup>, or antidepressant prescription<sup>10</sup> at any point during pregnancy, determined by previous literature. Analysis of human serum samples used samples collected through the Grown in Wales study<sup>47</sup>. Although this study was not designed to study serum BDNF, it was considered to be sufficiently powered based on a previous study on BDNF levels in human serum which indicated that a cohort size of 60 was required to detect a 20% difference between populations, and 200 for a 10% difference<sup>53</sup>. For analysis of mouse serum samples, no power calculation was performed, as initial data fell into the normal range. Randomization and blinding of mouse samples were not performed, as groups were compared across genotypes.

#### Study approval

Full ethical approval for the GiW Study cohort was obtained from the Wales Research Ethics Committee REC reference 15/WA/0004. The research was performed in line with the principles of the Declaration of Helsinki as revised in 2008. Mouse studies were approved by the Cardiff University Ethical Review Board, and all

experiments performed within the guidelines of the Home Office Animals (Scientific Procedures) Act, 1986.

## Results

### Higher levels of maternal serum BDNF compared to fetal BDNF

BDNF levels were measured in maternal and fetal serum samples from the GiW cohort<sup>47</sup>. After filtering by the *sample exclusion criteria* (“Methods”), BDNF levels were measured by ELISA in 251 maternal and 212 fetal serum samples (cohort demographics: Table 1).

BDNF values were not normally distributed in maternal (Fig. 1A;  $P = 5.52e-05$ ) or fetal (Fig. 1B;  $P = 1.77e-04$ ) serum. Values were significantly lower in fetal compared to maternal circulation (Fig. 1C; median (IQR)—maternal: 15.15 (5.28); fetal: 9.6 (4.46) ng/ml;  $P < 2.20e-16$  from 173 pairs,  $r = -0.575$ ) and lower in male compared to female cord blood (Fig. 1D; Supplementary Table 2), similar to previous studies<sup>26,35,41,45</sup>. Fetal sex had no overall effect on maternal serum BDNF values. Mothers with female infants had serum BDNF levels of 15.47 ng/ml (4.88); male infants: 15.2 ng/ml (6.12). Analysis of the relationship between individual maternal and fetal serum samples (Fig. 1E) showed a moderate correlation (Spearman’s rank correlation;  $r = 0.290$ ,  $P = 1.09e-04$ ,  $n = 173$  pairs).

### No association between BDNF and maternal EPDS scores

Maternal BDNF levels were not correlated with EPDS scores reported at term when all samples were analyzed (Fig. 2A;  $r = -0.022$ ,  $P = 0.733$ ,  $n = 244$ ) or when samples were analyzed by fetal sex (Fig. 2B; male infants:  $r = 0.06$ ;  $P = 0.522$ ;  $n = 113$ . Female infants:  $r = -0.1$ ;  $P = 0.258$ ;  $n = 131$ ). There was no relationship between maternal BDNF and categorically defined depression using the EPDS  $\geq 13$  cutoff<sup>50</sup> (Fig. 2C). There was also no association between fetal BDNF and EPDS scores (Fig. 2D–F).

### Significant association between maternal BDNF and maternal STAI scores

Maternal BDNF levels were not correlated with STAI scores when all samples were analyzed (Fig. 3A;  $r = 0.119$ ;  $P = 0.078$ ;  $n = 220$ ). However, when analyzed by fetal sex STAI scores were significantly associated with maternal serum BDNF in mothers of male (Fig. 3B;  $r = 0.281$ ;  $P = 0.005$ ;  $n = 99$ ; Supplementary Table 3) but not female infants (Fig. 3B;  $r = -0.014$ ;  $P = 0.882$ ;  $n = 121$ ). When analyzed categorically using STAI  $\geq 40$  suggesting anxiety<sup>51</sup>, BDNF levels between groups were not significantly different (Fig. 3C).

Fetal serum BDNF levels were not correlated with maternal STAI scores when all data were analyzed (Fig. 3D;  $r = 0.056$ ;  $P = 0.437$ ;  $n = 192$ ), or by fetal sex (Fig. 3E; male:  $r = 0.131$ ;  $P = 0.226$ ;  $n = 87$ . Female:  $r = 0.073$ ;  $P = 0.458$ ;  $n = 105$ ), or by group (Fig. 3F).

**Table 1 Cohort demographic data.**

Demographics	Maternal: % (n) or median (IQR)	P	Fetal: % (n) or median (IQR)	P
Maternal age at booking	33 (6)	0.223	33.5 (7)	<b>0.020</b>
Maternal BMI at booking	26.32 (7.21)	0.645	26.14 (6.84)	0.344
Maternal ethnicity, % (n)		–		–
Caucasian	100 (251)		100 (212)	
Fetal sex, % (n)		0.498		<b>2.02e-05</b>
Female	54 (136)		54 (115)	
Male	46 (115)		46 (97)	
Parity, % (n)		0.218		0.374
Multiparous	81 (204)		79 (168)	
Nulliparous	19 (47)		21 (44)	
Gestational age (weeks)	39 (0)	0.159	39 (0)	<b>0.026</b>
Smoking at any point in pregnancy, % (n)		0.332		0.633
No	89 (224)		89 (189)	
Yes	10 (24)		9 (20)	
Missing	1 (3)		1 (3)	
Alcohol at any point in pregnancy, % (n)		0.819		0.330
No	58 (146)		60 (127)	
Yes	40 (100)		38 (81)	
Missing	2 (5)		2 (4)	
Exercise, % (n)		0.48		0.502
No	81 (203)		82 (173)	
Yes	18 (46)		17 (36)	
Missing	1 (2)		1 (3)	
Hypertension, % (n)		0.735		<b>0.019</b>
No	95 (239)		96 (203)	
Yes	4 (9)		3 (7)	
Missing	1 (3)		1 (2)	
Gestational diabetes, % (n)		0.073		0.432
No	92 (232)		93 (198)	
Yes	6 (14)		5 (10)	
Missing	2 (5)		2 (4)	
Highest education level, % (n)		0.560		0.405
Left before GCSE	5 (13)		7 (14)	
GCSE & vocational	22 (55)		20 (42)	
A-level	12 (31)		11 (24)	
University	29 (74)		30 (64)	
Postgraduate	26 (66)		28 (60)	
Missing	5 (12)		4 (8)	
Family income, % (n)		0.064		0.154
<18,000	7 (18)		7 (14)	
18–25,000	10 (24)		8 (16)	
25–43,000	20 (49)		19 (41)	
>43,000	51 (127)		51 (108)	
Do not wish to say	10 (25)		12 (26)	
Missing	3 (8)		3 (7)	
WIMD score	1270 (1183)	0.408	1368 (1158)	0.693
Antidepressants prescribed in pregnancy, % (n)		0.198		0.535
No	91 (228)		93 (197)	
Yes	9 (23)		7 (15)	

**Table 1** continued

Demographics	Maternal: % (n) or median (IQR)	P	Fetal: % (n) or median (IQR)	P
EPDS score	7 (6)	0.733	7 (6)	0.394
Missing, n	7		6	
STAI score	33.5 (12)	0.078	33.5 (13)	0.437
Missing, n	31		20	

EPDS Edinburgh Postnatal Depression Scale, STAI State-Trait Anxiety Inventory, WIMD Welsh Index of Multiple Deprivation. For WIMD scores, please see [gov.wales/welsh-index-multiple-deprivation](http://gov.wales/welsh-index-multiple-deprivation). Low scores indicate high deprivation, whilst high scores indicate low levels of deprivation. P values indicate the significance of the relationship between serum BDNF levels and the corresponding variable. Bold values indicate  $P < 0.05$ .

### Maternal BDNF does not cross the placenta

To assess the potential for MK-derived BDNF to cross the placenta, we exploited the fact that wild-type mice do not have BDNF present in their blood<sup>69</sup> and engineered a novel mouse model in which BDNF was expressed from MKs. We inserted a myc-tagged *BDNF-IRES-eGFP* into the *Rosa26* locus (Fig. 4A). When this line is crossed to mice expressing PF4-Cre, driving Cre expression specifically within MKs<sup>57</sup>, mice should express BDNF from MKs (Fig. 4A) as in humans<sup>9</sup>. Through breeding, we generated animals of the following genotypes: *BDNF-myc<sup>+/+</sup>;Cre<sup>+/-</sup>* and *BDNF-myc<sup>+/+</sup>;Cre<sup>-/-</sup>* (homozygous for the *BDNF-myc* allele, positive or negative for expression of PF4-Cre), *BDNF-myc<sup>+/-</sup>;Cre<sup>+/-</sup>* and *BDNF-myc<sup>+/-</sup>;Cre<sup>-/-</sup>* (heterozygous for *BDNF-myc*, positive or negative for PF4-Cre) and lastly *BDNF-myc<sup>-/-</sup>;Cre<sup>+/-</sup>* and *BDNF-myc<sup>-/-</sup>;Cre<sup>-/-</sup>* (wild-type in the *BDNF-myc* allele, positive or negative for PF4-Cre). All genotypes were viable and fertile, with no overt phenotype. The hematological profile of mice carrying one or two alleles of *BDNF-myc* and the Cre transgene (BCre; recombined scenario) was similar to mice carrying a *BDNF-myc* allele in the absence of the Cre transgene (unrecombined) (B) (Fig. 4B, C). In animals carrying the Cre transgene and one or two alleles of the *BDNF-myc* (BCre: *BDNF-myc<sup>+/+</sup>;Cre<sup>+/-</sup>* or *BDNF-myc<sup>+/-</sup>;Cre<sup>+/-</sup>*), BDNF could be detected in MKs (Fig. 4D). To reach the bloodstream, MK-expressed BDNF must be packaged into platelets, from which it can be released during platelet activation. Activation-induced platelet spreading (Fig. 4E) and platelet counts (Fig. 4F) were similar between BCre and B mice. In animals without the Cre transgene (B), ~0.5% of white blood cells ( $0.52 \pm 0.1\%$  (SD)) were positive for eGFP whereas over 85% of platelets from BCre mice expressed eGFP (Fig. 4G). BDNF-myc was detectable by western blot in the serum of BCre mice and not detectable in mice without the Cre transgene (Fig. 4H and Supplementary Fig. 1). By ELISA, the mean ( $\pm$  standard error of the mean) serum BDNF values in young adult mice (12 weeks old) were  $10.6 \pm 1.45$  ng/ml in *BDNF-myc<sup>+/+</sup>;Cre<sup>+/-</sup>* animals ( $n = 4$ ; three female, one male), and

$32.3 \pm 3.70$  ng/ml in *BDNF-myc<sup>+/+</sup>;Cre<sup>+/-</sup>* animals ( $n = 3$ ; three female). Thus, in this model MK-derived BDNF is present in serum at levels comparable to those determined in adult human serum<sup>53</sup>.


Having confirmed MK expression of BDNF, this model was used to test if MK-derived BDNF in maternal blood could cross the placenta. *BDNF-myc<sup>+/+</sup>;Cre<sup>+/-</sup>* females expressing MK-derived BDNF were crossed with *BDNF-myc<sup>+/+</sup>;Cre<sup>-/-</sup>* male mice to generate fetuses homozygous for the *BDNF-myc* insertion, with approximately half lacking the Cre transgene and thus intrinsic serum BDNF. Cre-negative fetuses (*BDNF-myc<sup>+/+</sup>;Cre<sup>-/-</sup>*) could only have serum BDNF if this were maternally-derived. Maternal and fetal blood were collected at E18.5 from 3 litters, serum generated and BDNF expression analyzed by western blot. BDNF-myc was detectable in maternal serum and serum from *BDNF-myc<sup>+/+</sup>;Cre<sup>+/-</sup>* fetuses but absent in serum from the *BDNF-myc<sup>+/+</sup>;Cre<sup>-/-</sup>* fetuses (Fig. 4I).

### Discussion

This is the first study to interrogate the relationship between maternal and fetal serum BDNF at term in the context of maternal symptoms of depression and anxiety. We discovered that maternal symptoms of anxiety correlated with significantly raised maternal serum BDNF exclusively in mothers of boys. We did not find similarly raised BDNF levels in the male infants, in line with the results obtained with our new animal model in which we found no evidence that MK-derived BDNF, the main source of BDNF in humans, crosses the placenta.


Elevated maternal BDNF was specifically associated with maternally reported anxiety symptoms and not symptoms of depression. Symptoms of depression have been associated with lower serum BDNF in non-pregnant adults<sup>10,11</sup>. Reports of associations between anxiety symptoms and serum BDNF have been mixed, with one systematic review concluding lower BDNF only in association with obsessive-compulsive disorder<sup>70</sup>. Animal studies suggest that overexpression of BDNF in the brain can simultaneously improve performance in tests related to depression while facilitating anxiety-like behavior<sup>71</sup>. This would be essentially consistent with our findings of a contrasting relationship of serum BDNF with anxiety and depressive symptoms. Testing the relationship between MK-derived BDNF and behavioral outcomes will now be possible using the animal model described here.

Our finding of elevated maternal BDNF was exclusive to the mothers of boys reporting higher anxiety symptoms. We currently do not know whether this is a causal relationship nor, if causal, the direction of causality. Anxiety symptoms, or factors associated with anxiety, may drive changes in maternal BDNF that are only permissible in those mothers with a male fetus. Alternatively,



under conditions of maternal anxiety, the male fetus or placenta may produce signals driving elevations in maternal BDNF. A third possibility is that elevated BDNF is causing anxiety only in mothers with boys. Subsequent studies with our new mouse model should help tease apart these complex relationships. In addition to

determining causality, it will also be important to establish whether changes in maternal BDNF occur as a result of alterations in MK proliferation, MK differentiation, BDNF expression within MKs, and/or platelet release of BDNF into the circulation. Another important question to address is the function of MK-derived BDNF in




development or disease. While further work is required, our identification of this sex-specific association between maternal anxiety and maternal serum BDNF provides a highly novel example of the influence of fetal sex on the maternal state.

Whether these perturbations in maternal serum BDNF are transmitted to the fetus is an important question, given the known critical function of BDNF in the central nervous system and our increasing understanding of its function in the periphery, including in energy homeostasis and regulation of glucose levels<sup>72,73</sup>. We found no evidence that BDNF can cross the placenta, at least in mice. Maternal blood BDNF is contained within platelets and is not found as free serum protein. Furthermore, the hemochorial placenta provides a physical barrier with three continuous cellular layers between the maternal and

fetal blood systems in both mice and humans<sup>22</sup>. Some proteins can cross the placenta by an active process termed receptor-mediated pinocytosis<sup>23</sup>. We were unable to detect TrkB or p75 by western blot analysis of placental extracts or by immunostaining in the labyrinth of the mouse placenta where transport occurs (data not shown) consistent with the lack of BDNF transplacental transfer in our model. While we cannot formally exclude the possibility that there are species differences in the placental transfer of BDNF as observed with the transfer of maternal antibodies<sup>74</sup>, BDNF receptors do not appear to be expressed in the mature human placenta<sup>75</sup>.

An important feature of our study is the number of paired maternal:fetal BDNF samples measured at term. However, we only measured mood symptoms once during pregnancy. We do not know when anxiety symptoms first






manifested but the persistence of symptoms after birth<sup>47</sup> and the nature of the questionnaire, measuring trait not state anxiety, suggests prolonged symptomology. Our cohort is relatively homogeneous and it will be important to replicate these findings in more diverse cohorts, and across other modes of delivery.

In summary, we have discovered a sex-specific association between maternally reported symptoms of anxiety and elevated maternal serum BDNF at term, specifically in pregnancies with male infants. We find no evidence that maternal serum BDNF crosses the placenta either from our human study or in our new humanized mouse model. We conclude that the placenta provides a barrier

protecting the fetus against potentially damaging fluctuations in maternal BDNF that occur under conditions of stress.

#### Acknowledgements

We gratefully acknowledge the work of Bill Mansfield (University of Cambridge) in the generation of the *stop-flox Bdnf* line used in this study. H.D. was supported by the European Union's Horizon 2020 Research and Innovation Programme under the Marie Skłodowska-Curie Grant Agreement 663830. X.N., P.C.-F., and Y.-A.B. were supported by the Welsh Government's Sêr Cymru programme. S.M.G. was supported by a MRC GW4 BioMed PhD Studentship (MR/N013794/1), L.A.S. was supported by a BBSRC SWBio PhD studentship (BB/M009122/1). A.M. was supported by the German Research Foundation (DFG, MU 4099/1-1), I.M.-G. was supported by BBSRC grant BB/S002359/1, and C.G. was supported by NHS Blood and Transplant. The Grown in Wales study was funded by MRC grant (MR/M013960/1).



#### Author details

<sup>1</sup>School of Biosciences, Cardiff University, Cardiff CF10 3AX, UK. <sup>2</sup>Department of Haematology, University of Cambridge, Cambridge CB2 0PT, UK. <sup>3</sup>Present address: Hospital Universitario Virgen Macarena-FISEVI, University of Seville, E41009 Seville, Spain

#### Author contributions

Grown in Wales Study: H.D. and R.M.J. designed research. S.M.G. and L.A.S. performed data curation. H.D. performed research. S.M.G. and H.D. performed the statistical analysis, R.M.J. initiated the cohort. Mouse Model: X.N., H.D., P.C.-F., C.G., R.M.J., I.M.-G., and Y.-A.B. designed research. X.N., H.D., A.M., P.C.F., and R.M.J. performed research. H.D. and X.N. performed the statistical analysis, H.D., X.N., Y.-A.B., & R.M.J. wrote the paper.

#### Conflict of interest

The authors declare that they have no conflict of interest.

#### Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Supplementary Information** accompanies this paper at (<https://doi.org/10.1038/s41398-020-01176-8>).

Received: 9 June 2020 Revised: 16 November 2020 Accepted: 30 November 2020

Published online: 18 January 2021

#### References

- Bauer, A., Parsonage, M., Knapp, M., Lemmi, V. & Adelaja, B. The costs of perinatal mental health problems. *LSE. Cent. Ment. Heal.* <https://doi.org/10.13140/2.1.4731.6169> (2014).

2. Stewart, D. E. Depression during pregnancy. *N. Engl. J. Med.* **365**, 1605–1611 (2011).
3. Van den Bergh, B. R. H. et al. Prenatal developmental origins of behavior and mental health: the influence of maternal stress in pregnancy. *Neurosci. Biobehav. Rev.* <https://doi.org/10.1016/j.neubiorev.2017.07.003> (2017).
4. Kinsella, M. T. & Monk, C. Impact of maternal stress, depression and anxiety on fetal neurobehavioral development. *Clin. Obstet. Gynecol.* **52**, 425–440 (2009).
5. Brand, S. R. & Brennan, P. A. Impact of antenatal and postpartum maternal mental illness: how are the children? *Clin. Obstet. Gynecol.* **52**, 441–455 (2009).
6. O'Donnell, K., O'Connor, T. G. & Glover, V. Prenatal stress and neurodevelopment of the child: focus on the HPA axis and role of the placenta. *Dev. Neurosci.* **31**, 285–292 (2009).
7. Park, H. & Poo, M. M. Neurotrophin regulation of neural circuit development and function. *Nat. Rev. Neurosci.* **14**, 7–23 (2013).
8. Rosenfeld, R. D. et al. Purification and identification of brain-derived neurotrophic factor from human serum. *Protein Expr. Purif.* **6**, 465–471 (1995).
9. Chacón-Fernández, P. et al. Brain-derived neurotrophic factor in megakaryocytes. *J. Biol. Chem.* **291**, 9872–9881 (2016).
10. Molendijk, M. L. et al. Serum BDNF concentrations as peripheral manifestations of depression: evidence from a systematic review and meta-analyses on 179 associations (N=9484). *Mol. Psychiatry* **19**, 791–800 (2014).
11. Karege, F. et al. Decreased serum brain-derived neurotrophic factor levels in major depressed patients. *Psychiatry Res.* **109**, 143–148 (2002).
12. Sen, S., Duman, R. & Sanacora, G. Serum brain-derived neurotrophic factor, depression, and antidepressant medications: meta-analyses and implications. *Biol. Psychiatry* **64**, 527–532 (2008).
13. Kishi, T., Yoshimura, R., Ikuta, T. & Iwata, N. Brain-derived neurotrophic factor and major depressive disorder: evidence from meta-analyses. *Front. Psychiatry* <https://doi.org/10.3389/fpsy.2017.00308> (2018).
14. Brunoni, A. R., Lopes, M. & Fregni, F. A systematic review and meta-analysis of clinical studies on major depression and BDNF levels: implications for the role of neuroplasticity in depression. *Int. J. Neuropsychopharmacol.* **11**, 1169–1180 (2008).
15. Polyakova, M. et al. BDNF as a biomarker for successful treatment of mood disorders: a systematic & quantitative meta-analysis. *J. Affect. Disord.* **174**, 432–440 (2015).
16. Ciammola, A. et al. Low brain-derived neurotrophic factor (BDNF) levels in serum of Huntington's disease patients. *Am. J. Med. Genet. B Neuropsychiatr. Genet.* **144B**, 574–577 (2007).
17. Ng, T. K. S., Ho, C. S. H., Tam, W. W. S., Kua, E. H. & Ho, R. C. M. Decreased serum brain-derived neurotrophic factor (BDNF) levels in patients with Alzheimer's disease (AD): a systematic review and meta-analysis. *Int. J. Mol. Sci.* **20**, 1–26 (2019).
18. Qin, X. Y. et al. Association of peripheral blood levels of brain-derived neurotrophic factor with autism spectrum disorder in children: a systematic review and meta-analysis. *JAMA Pediatr.* **170**, 1079–1086 (2016).
19. Szuhany, K. L., Bugatti, M. & Otto, M. W. A meta-analytic review of the effects of exercise on brain-derived neurotrophic factor. *J. Psychiatr. Res.* **60**, 56–64 (2015).
20. Burton, G. J. & Fowden, A. L. The placenta: a multifaceted, transient organ. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* <https://doi.org/10.1098/rstb.2014.0066> (2015).
21. Soares, M. J., Varberg, K. M. & Iqbal, K. Hemochorial placentation: development, function, and adaptations. *Biol. Reprod.* **99**, 196–211 (2018).
22. Woods, L., Perez-Garcia, V. & Hemberger, M. Regulation of placental development and its impact on fetal growth—new insights from mouse models. *Front. Endocrinol.* <https://doi.org/10.3389/fendo.2018.00570> (2018).
23. Palmeira, P., Quinello, C., Silveira-Lessa, A. L., Zago, C. A. & Carneiro-Sampaio, M. IgG placental transfer in healthy and pathological pregnancies. *Clin. Dev. Immunol.* <https://doi.org/10.1155/2012/985646> (2012).
24. Kodomari, I., Wada, E., Nakamura, S. & Wada, K. Maternal supply of BDNF to mouse fetal brain through the placenta. *Neurochem. Int.* **54**, 95–98 (2009).
25. Chouthai, N. S., Sampers, J., Desai, N. & Smith, G. M. Changes in neurotrophin levels in umbilical cord blood from infants with different gestational ages and clinical conditions. *Pediatr. Res.* **53**, 965–969 (2003).
26. Malamitsi-Puchner, A., Economou, E., Rigopoulou, O. & Boutsikou, T. Perinatal changes of brain-derived neurotrophic factor in pre- and fullterm neonates. *Early Hum. Dev.* **76**, 17–22 (2004).
27. Hodyl, N. A. et al. Antenatal steroid exposure in the late preterm period is associated with reduced cord blood neurotrophin-3. *Early Hum. Dev.* **101**, 57–62 (2016).
28. Wang, Y. et al. Effects of prenatal exposure to cadmium on neurodevelopment of infants in Shandong, China. *Environ. Pollut.* **211**, 67–73 (2016).
29. Yu, X. et al. The role of cord blood BDNF in infant cognitive impairment induced by low-level prenatal manganese exposure: LW birth cohort, China. *Chemosphere* **163**, 446–451 (2016).
30. Cai, Q. Y. et al. Placental and cord blood brain derived neurotrophic factor levels are decreased in nondiabetic macrosomia. *Arch. Gynecol. Obstet.* **296**, 205–213 (2017).
31. Ferrari, N. et al. Exercise during pregnancy and its impact on mothers and offspring in humans and mice. *J. Dev. Orig. Health Dis.* **9**, 63–76 (2018).
32. Mardini, V. et al. TBARS and BDNF levels in newborns exposed to crack/cocaine during pregnancy: a comparative study. *Braz. J. Psychiatry* **39**, 263–266 (2017).
33. Akbaba, N. et al. Neurotrophins and neuroinflammation in fetuses exposed to maternal depression and anxiety disorders during pregnancy: a comparative study on cord blood. *Arch. Women's Ment. Health* **21**, 105–111 (2018).
34. Basu, S., Kumar, D., Anupurba, S., Verma, A. & Kumar, A. Effect of maternal iron deficiency anemia on fetal neural development. *J. Perinatol.* **38**, 233–239 (2018).
35. Briana, D. D. et al. Differential expression of cord blood neurotrophins in gestational diabetes: the impact of fetal growth abnormalities. *J. Matern. Neonatal Med.* **31**, 278–283 (2018).
36. Yusrwati Rina, G., Indrawati, L. N. & Machmud, R. Differences in brain-derived neurotrophic factor between neonates born to mothers with normal and low ferritin. *Asia Pac. J. Clin. Nutr.* **27**, 389–392 (2018).
37. Lommatzsch, M. et al. Maternal serum concentrations of BDNF and depression in the perinatal period. *Psychoneuroendocrinology* **31**, 388–394 (2006).
38. Pawluski, J. L., Brain, U., Hammond, G. L. & Oberlander, T. F. Selective serotonin reuptake inhibitor effects on neural biomarkers of perinatal depression. *Arch. Women's Ment. Health* **22**, 431–435 (2019).
39. Sonmez, E. O. et al. Effect of maternal depression on brain-derived neurotrophic factor levels in fetal cord blood. *Clin. Psychopharmacol. Neurosci.* **17**, 308–313 (2019).
40. Cannon, T. D., Yolken, R., Buka, S. & Torrey, E. F. Collaborative study group on the perinatal origins of severe psychiatric disorders. decreased neurotrophic response to birth hypoxia in the etiology of schizophrenia. *Biol. Psychiatry* **64**, 797–802 (2008).
41. Spulber, S. et al. Effects of maternal smoking and exposure to methylmercury on brain-derived neurotrophic factor concentrations in umbilical cord serum. *Toxicol. Sci.* **117**, 263–269 (2010).
42. Uguz, F. et al. Maternal generalized anxiety disorder during pregnancy and fetal brain development: a comparative study on cord blood brain-derived neurotrophic factor levels. *J. Psychosom. Res.* **75**, 346–350 (2013).
43. Dhiman, P., Say, A., Rajendiren, S., Kattimani, S. & Sagili, H. Association of foetal APGAR and maternal brain derived neurotrophic factor levels in postpartum depression. *Asian J. Psychiatr.* **11**, 82–83 (2014).
44. Garcés, M. F. et al. Brain-derived neurotrophic factor is expressed in rat and human placenta and its serum levels are similarly regulated throughout pregnancy in both species. *Clin. Endocrinol.* **81**, 141–151 (2014).
45. Flöck, A. et al. Determinants of brain-derived neurotrophic factor (BDNF) in umbilical cord and maternal serum. *Psychoneuroendocrinology* **63**, 191–197 (2016).
46. Gao, X., Wang, J., Yao, H., Cai, Y. & Cheng, R. Serum BDNF concentration after delivery is associated with development of postpartum depression: a 3-month follow up study. *J. Affect. Disord.* **200**, 25–30 (2016).
47. Janssen, A. B. et al. Persistence of anxiety symptoms after elective caesarean delivery. *BJPsych. Open* **4**, 354–360 (2018).
48. Rauskolb, S. et al. Global deprivation of brain-derived neurotrophic factor in the CNS reveals an area-specific requirement for dendritic growth. *J. Neurosci.* **30**, 1739–1749 (2010).
49. Klein, A. B. et al. Blood BDNF concentrations reflect brain-tissue BDNF levels across species. *Int. J. Neuropsychopharmacol.* **14**, 347–353 (2011).
50. Cox, J. L., Chapman, G., Murray, D. & Jones, P. Validation of the Edinburgh postnatal depression scale (EPDS) in non-postnatal women. *J. Affect. Disord.* **39**, 185–189 (1996).
51. Meades, R. & Ayers, S. Anxiety measures validated in perinatal populations: a systematic review. *J. Affect. Disord.* **133**, 1–15 (2011).
52. Christian, L. M., Mitchell, A. M., Gillespie, S. L. & Palettas, M. Serum brain-derived neurotrophic factor (BDNF) across pregnancy and postpartum: Associations

- with race, depressive symptoms, and low birth weight. *Psychoneuroendocrinology* **74**, 69–76 (2016).
53. Naegelin, Y. et al. Measuring and validating the levels of brain-derived neurotrophic factor in human serum. *eNeuro* <https://doi.org/10.1523/ENEURO.0419-17.2018> (2018).
  54. Kolbeck, R., Bartke, I., Eberle, W. & Barde, Y. A. Brain-derived neurotrophic factor levels in the nervous system of wild-type and neurotrophin gene mutant mice. *J. Neurochem.* **72**, 1930–1938 (1999).
  55. Matsumoto, T. et al. Biosynthesis and processing of endogenous BDNF: CNS neurons store and secrete BDNF, not pro-BDNF. *Nat. Neurosci.* **11**, 131–133 (2008).
  56. Perez-Pinera, P., Ousterout, D. G., Brown, M. T. & Gersbach, C. A. Gene targeting to the ROSA26 locus directed by engineered zinc finger nucleases. *Nucleic Acids Res.* **40**, 3741–3752 (2012).
  57. Tiedt, R., Schomber, T., Hao-Shen, H. & Skoda, R. C. Pf4-Cre transgenic mice allow the generation of lineage-restricted gene knockouts for studying megakaryocyte and platelet function in vivo. *Blood* **109**, 1503–1506 (2007).
  58. Schneider, C. A., Rasband, W. S. & Eliceiri, K. W. NIH Image to ImageJ: 25 years of image analysis. *Nat. Methods* **9**, 671–675 (2012).
  59. RStudio Team. RStudio: Integrated Development for R. <http://www.rstudio.com> (2019).
  60. Fox, J. & Weisberg, S. *An R Companion to Applied Regression*, Third Edition. Third edn. (Sage, Thousand Oaks, CA, 2019).
  61. Wickham, H., Francois, R., Henry, L. & Müller, K. Package 'dplyr'. A Grammar of Data Manipulation. R Package version 08.01 <https://CRAN.R-project.org/package=dplyr> (2019).
  62. Wickham, H. *ggplot2: Elegant Graphics for Data Analysis* (Springer-Verlag, New York, 2016).
  63. Kassambara, A. *ggpubr: 'ggplot2' Based Publication Ready Plots*. R package version 0.2.3. <https://CRAN.R-project.org/package=ggpubr> (2019).
  64. Grosjean, P., Ibanez F. *pastecs: Package for Analysis of Space-Time Ecological Series*. R package version 1.3.21 <https://CRAN.R-project.org/package=pastecs> (2018).
  65. Wickham, H. Reshaping data with the reshape package. *J. Stat. Softw.* **12**, 1–20 (2007).
  66. Bienertova-Vasku, J. et al. Brain-derived neurotrophic factor and ciliary neurotrophic factor in maternal plasma and umbilical cord blood from pre-eclamptic and physiological pregnancies. *J. Obstet. Gynaecol.* **33**, 359–363 (2013).
  67. Monteleone, P. et al. Opposite changes in the serum brain-derived neurotrophic factor in anorexia nervosa and obesity. *Psychosom. Med.* **66**, 744–748 (2004).
  68. Nanri, A., Kochi, T., Eguchi, M., Kabe, I. & Mizoue, T. Demographic and lifestyle correlates of brain-derived neurotrophic factor in a working population: The Furukawa Nutrition and Health Study. *Psychiatry Res.* **272**, 581–586 (2019).
  69. Radka, S. F., Holst, P. A., Fritsche, M. & Altar, C. A. Presence of brain-derived neurotrophic factor in brain and human and rat but not mouse serum detected by a sensitive and specific immunoassay. *Brain Res.* **709**, 122–130 (1996).
  70. Suliman, S., Hemmings, S. M. J. & Seedat, S. Brain-derived neurotrophic factor (BDNF) protein levels in anxiety disorders: systematic review and meta-regression analysis. *Front. Integr. Neurosci.* <https://doi.org/10.3389/fnint.2013.00055> (2013).
  71. Govindarajan, A. et al. Transgenic brain-derived neurotrophic factor expression causes both angiogenic and antidepressant effects. *Proc. Natl Acad. Sci. USA* **103**, 13208–13213 (2006).
  72. Marosi, K. & Mattson, M. P. BDNF mediates adaptive brain and body responses to energetic challenges. *Trends Endocrinol. Metab.* **25**, 89–98 (2014).
  73. Fulgenzi, G. et al. Novel metabolic role for BDNF in pancreatic  $\beta$ -cell insulin secretion. *Nat. Commun.* <https://doi.org/10.1038/s41467-020-15833-5> (2020).
  74. Bowman, C. J. et al. Placental transfer of Fc-containing biopharmaceuticals across species, an industry survey analysis. *Birth Defects Res. B Dev. Reprod. Toxicol.* **98**, 459–485 (2013).
  75. Uhlen, M. et al. Tissue-based map of the human proteome. *Science* <https://doi.org/10.1126/science.1260419> (2015).