

# Could ratio of hemoglobin to red cell distribution width and ratio of absolute lymphocyte count to absolute monocyte count be a prognostic tool in newly diagnosed multiple myeloma patients?

## Article history:

Received: 30.08.2019  
Accepted: 18.01.2020

Mehmet Baysal<sup>1\*</sup>,  
Ufuk Demirci<sup>1</sup>,  
Volkan Bas<sup>1</sup>,  
Sedanur Karaman Gulsaran<sup>1</sup>,  
Elif Umit<sup>1</sup>,  
Hakki Onur Kirkizlar<sup>1</sup>,  
Serap Baysal<sup>2</sup>,  
Ahmet Muzaffer Demir<sup>1</sup>

<sup>1</sup>Department of Hematology, Trakya University Faculty of Medicine, Edirne, Turkey

<sup>2</sup>Provincial Health Directorate, Edirne, Turkey

## Abstract

**Introduction:** Hemoglobin/red cell distribution width (RDW) ratio (HRR) and lymphocyte-to-monocyte ratio (LMR) are two novel biomarkers associated with overall survival (OS) and prognosis in several types of cancers. The aim of this study is to investigate the value of HRR and LMR in newly diagnosed multiple myeloma (MM) patients. **Methods:** A total of 180 patients were included in this study. Patients diagnosed with MM between May 2013 and May 2019 at a single center were evaluated. HRR was calculated by dividing hemoglobin to RDW, both measured from the same sample. LMR was calculated by dividing absolute lymphocyte count (ALC) to absolute monocyte count (AMC). **Results:** The cutoff value for HRR was taken as 0.61, and the cutoff value for LMR was taken as 3.28. Patients were divided into low HRR, high HRR, low LMR, and high LMR groups. OS of the patients with low HRR was found lower compared with high HRR (36.7 months for low HRR and 53.2 months for high HRR,  $p < 0.001$ ). Also, OS was found lower in the low LMR group (39.4 months for low LMR and 51.7 months for high LMR,  $p = 0.016$ ). On multivariate analysis, low HRR and low LMR were predictive factors of OS (hazard ratio (HR) 2.08, 95% confidence intervals (CI) 1.31–3.03, and  $p = 0.002$  for low HRR; HR 1.47, 95% CI 0.92–2.29, and  $p = 0.010$  for low LMR). **Conclusion:** Combining both HRR and LMR could be a prognostic biomarker and it reflects the status of the immune system in newly diagnosed MM patients.

© 2020 Polish Society of Hematology and Transfusion Medicine, Insitute of Hematology and Transfusion Medicine. Published by Sciendo. All rights reserved.

## Keywords:

multiple myeloma, hemoglobin, red cell distribution width, lymphocyte–monocyte ratio, prognostic

## Introduction

Multiple myeloma (MM) is described as a clonal proliferation of malignant plasma cells secreting monoclonal immunoglobulin with heavy or light chains. This incurable disease is accounted for 1–2% of all cancers and 15% of hematological malignancies [1, 2]. Usual presentation and end-organ diseases of MM include anemia, hypercalcemia, lytic bone lesions, and renal disease.

Anemia itself is accepted as myeloma defining event and is associated with involvement of bone marrow of malignant plasma cells [3]. As a surrogate of red blood cell size and variability, red cell distribution width (RDW) is investigated in several cancer types and associated with stage, grade, activity, and prognosis in certain types of cancers [4, 5]. Likewise, RDW is demonstrated to be correlated with inflammatory states and chronic conditions [6, 7]. Although erythropoiesis can be affected by any chronic condition, the variability of RDW, the inclusion of hemoglobin levels, and inflammatory markers to the equation might bring insight to the pathogenesis of how erythropoiesis is affected and the role of bone marrow microenvironment (BME) in MM [8]. Besides, RDW itself was proposed as a prognostic factor in symptomatic MM patients [9].

As mentioned earlier, BME holds an essential role in the pathogenesis of MM. The interactions between BME and malign plasma cells are demonstrated to be related to prognosis [10, 11]. These interactions cause an immune system to escape due to tumor-associated macrophages [12]. The immunosuppressive microenvironment and the effect of stromal cell–myeloma cell interferences stimulate the expansion of malign plasma cell clone. These immune escape and tumor growth are generated by myeloid-derived suppressor cells [13]. Peripheral absolute lymphocyte count (ALC) and absolute monocyte count (AMC) could preview the balance between the immune system and malignancy-associated immune escape. Monocyte count could be regarded as myeloid-derived suppressor cells, and therefore the ratio of ALC/AMC (LMR) may implicate the effect of the impaired immune system on MM [14]. There are studies regarding the effect of LMR on overall survival (OS), but combining hemoglobin/RDW ratio (HRR) with LMR may reveal the effects of immune dysregulation on plasma cell disease [15, 16, 17]. Besides, there is insufficient data on the effect of HRR in MM, especially in the era of novel agents. Therefore, in this study, we aimed to evaluate the possible impact of HRR and LMR on newly diagnosed MM patients treated with novel agents as the first line.

\* Corresponding author: Mehmet Baysal, Department of Hematology, Trakya University Faculty of Medicine, Balkan Campus 22030, Edirne, Turkey, phone: +902842357641-2687, +905359664188, e-mail: [drmehmetbaysal@gmail.com](mailto:drmehmetbaysal@gmail.com)

## Patients and methods

This retrospective study included 180 patients diagnosed with MM according to International Myeloma Working Group (IMWG) definitions between May 2013 and May 2019 in Trakya University Hematology Department. Baseline demographic features, including age and sex, and disease-related data, including heavy and light chain types of immunoglobulin, International staging system (ISS), whole blood count (WBC), results of the bone marrow sample analysis and treatments, were recorded from patient files. WBC was performed with Beckman Coulter DXH 800 device, and fresh blood samples were obtained after 8 h of fasting and before treatment.

Cytogenetic analysis was performed from bone marrow samples obtained at the time of diagnosis and included high-risk determinants of MM, such as ISS and revised ISS (R-ISS), according to the definitions of IMWG [18, 19]. The ratio of hemoglobin (g/dl) to RDW (%) was calculated by the formula hemoglobin/RDW, and LMR was calculated by the formula ALC/AMC. Overall survival (OS) was accepted as time from date of treatment to date of last follow-up or death from any cause.

This study was conducted according to the principles of the Helsinki declaration and was approved by the local ethical committee (TUTF-BAEK 2019-248).

### Statistical analysis

IBM Statistical Package for Social Sciences (SPSS) version 20 was used for statistical analysis. Kolmogorov-Smirnov test was performed to assess the distribution of the parameters. Descriptive statistics were used to evaluate the characteristics of patients. The correlation of HRR and LMR with different variables was assessed with Pearson's chi-squared test or Fisher's exact test for categorical variables and with Mann-Whitney U test for continuous variables. The OS is defined as the time from the diagnosis of MM to death from any cause. Kaplan-Meier OS estimates were conducted for survival analysis. Log-rank test and Cox regression analysis were performed to evaluate the estimate hazard ratios (HRs) and 95% confidence intervals (CI). A  $p$ -value  $<0.05$  was considered to be statistically significant. Receiver operating characteristic (ROC) curves and area under the curve (AUC) are constructed to define best cutoff values for HRR and LMR.

## Results

A total of 180 newly diagnosed MM patients were included in this study. The mean age of the patients was 66.77 years. Eighty-seven patients were male and 93 patients were female. Sixty-eight patients were classified as ISS-1, 51 patients were classified as ISS-2, and 61 patients were classified as ISS-3. Fifty-seven patients (31.6%) evaluated were R-ISS-1, 41 (22.8%) patients evaluated were R-ISS-2, and 82 (45.6%) patients evaluated were R-ISS-3. Thirty-six (20%) patients were regarded as high risk according to the analysis. Moreover, in autologous hematopoietic stem cell transplant (ASCT) eligible patients, ASCT was performed after 4–6 cycles of induction chemotherapy containing bortezomib. Ninety-seven (53.9%) patients

received bortezomib cyclophosphamide dexamethasone, 8 patients (4.4%) received bortezomib thalidomide dexamethasone, 33 patients (18.3%) received bortezomib dexamethasone, and 15 patients received VTD-PACE (bortezomib, thalidomide, dexamethasone, cyclophosphamide, doxorubicin, cisplatin, and etoposide). The demographic characteristics of the patients were summarized in table I.

The cutoff point for HRR was selected as 0.61 according to ROC analysis with an AUC value of 0.64 (95% CI 0.561–0.724). The most discriminative value of LMR was also calculated with ROC analysis and found to be 3.28 with an AUC value of 0.62 (95% CI 0.538–0.703). Patients were categorized as low LMR  $< 3.28$ , high LMR  $> 3.28$ , low HRR  $< 0.61$ , and high HRR  $> 0.61$ . Patients with low LMR had a lower OS compared with patients with high LMR ( $p = 0.018$ , HR 1.67, 95% CI 1.09–2.57), and patients with low HRR had a lower OS compared with high HRR ( $p = 0.001$ , HR 2.046, 95% CI 1.33–3.13) (Kaplan-Meier OS analysis for HRR and LMR was shown in figures 1 and 2, respectively). Univariate analysis and characteristics of the patients according to LMR and HRR were given in tables II and III.

**Table I. Demographic and clinical characteristics data of the patients**

Characteristics	
Age	66.77 ± 11.16 (28–93)
Gender	
Male	87 (48.3%)
Female	93 (51.7%)
Laboratory parameters	
Hemoglobin (g/dl)	11.01 (6.1–15.7)
Creatinine (mg/dl)	1.36 (0.42–6.90)
Calcium (mg/dl)	9.6 (7.1–18.4)
Albumin (g/dl)	3.6 (1.8–4.7)
LDH (U/L)	229.3 (92–662)
Beta-2 microglobulin (ng/ml)	5,687.3 (1,301–20,000)
CRP (mg/dl)	1.62 (0.1–17.9)
ISS	
1	68 (37.8%)
2	51 (28.3%)
3	61 (31.9%)
Revised ISS	
1	57 (31.7%)
2	41 (22.8%)
3	82 (45.6%)
Cytogenetic risk	
High risk	36 (20%)
Standard risk	144 (80%)
Frontline treatment	
Bortezomib cyclophosphamide dexametasone	97 (53.9%)
Bortezomib thalidomide dexametasone	8 (4.4%)
Bortezomib dexametasone	33 (18.3%)
VTD-PACE	15 (8.3%)
Others (melphalan-prednisolone, etc)	27 (15.1%)
Upfront ASCT	
Yes	61 (33.9%)
No	119 (66.1%)
Immunoglobulin type	
IgG	96 (53.3%)
IgA	43 (23.9%)
Light chain	38 (21.1%)
Non-secretory	2 (1.1%)
IgM	1 (0.6%)
Lytic lesion on presentation	
Yes	156 (86.7%)
No	24 (13.3%)




Fig. 1. Kaplan-Meier curve for overall survival of low HRR and high HRR




Fig. 2. Kaplan-Meier curve for overall survival of low LMR and high LMR

**Table II. Characteristics of the multiple myeloma patients according to LMR**

	Low LMR <3.28 (n:87)	High LMR >3.28 (n:93)	p-value
Age			
>65 years old	55 (63%)	45 (48%)	0.072
<65 years old	32 (36%)	48 (52%)	
Gender			
Male	45 (52%)	42 (45%)	0.456
Female	42 (48%)	51 (55%)	
Cytogenetic risk			
High risk	19 (22%)	17 (18%)	0.580
Standard risk	68 (78%)	76 (82%)	
ISS			
1	26 (29%)	42 (45%)	0.069
2	25 (28%)	26 (28%)	
3	36 (43%)	25 (27%)	
Revised ISS			
1	21 (25%)	36 (39%)	0.082
2	20 (22%)	21 (22%)	
3	46 (53%)	36 (29%)	
Immunoglobulin type			
IgG	41 (47%)	55 (59%)	0.125
IgA	19 (22%)	24 (26%)	
Light chain	25 (29%)	13 (14%)	
Non-secretory	1 (1%)	1 (1%)	
IgM	1 (1%)	0	
Laboratory parameters			
Hemoglobin (g/dl)	10.8	11.1	0.27
Creatinine (mg/dl)	1.51	1.19	<b>0.014</b>
Calcium (mg/dl)	9.54	9.87	0.176
Albumin (g/dl)	3.61	3.76	0.087
LDH (U/L)	234	224	0.527
Beta-2 microglobulin (ng/ml)	6,702	4,737	<b>0.004</b>
CRP (mg/dl)	1.90	1.35	0.18
Frontline treatment			
Bortezomib cyclophosphamide dexametasone	46 (53%)	51 (55%)	0.833
Bortezomib thalidomide dexametasone	4 (4%)	4 (4%)	
Bortezomib dexametasone	17 (19%)	16 (17%)	
VTD-PACE	6 (7%)	9 (10%)	
Others (melphalan-prednisolone, etc)	14 (16%)	13 (14%)	
Upfront ASCT			
Yes	22 (22%)	39 (41%)	<b>0.027</b>
No	65 (78%)	54 (58%)	
Overall survival months	39.44 ± 3.72	51.79 ± 3.68	<b>0.016</b>

Univariate analysis for low LMR, HR: 1.67 (95% CI 1.09–2.57)

On multivariate analysis, low HRR and low LMR are associated with lower OS (HR for HRR 2.08, 95% CI 1.31–3.03,  $p = 0.002$ ; and HR for low LMR 1.47, 95% CI 0.92–2.29,  $p = 0.010$ ). Also ASCT, high-risk MM, and ISS-3 were associated with lower OS on multivariate analysis (details of the multivariate cox-regression analysis was given in table. IV). Furthermore, we have stratified our patients into four groups: low LMR and low HRR, low LMR and high HRR, high LMR and low HRR, and high LMR and high HRR. The OS of these four groups was 34.6, 44.8, 40.5, and 56.1 months, respectively. OS was lower in the low LMR and low HRR groups, which was higher in high LMR and high HRR groups.

## Discussion

MM is an incurable disease with complex biological heterogeneity. The underlying complex biology plays a unique role in the clinical

course and prognosis of the disease. In this study, baseline WBC parameters were analyzed for their probable role in the OS of MM patients. These findings underline that no matter how complicated diseases and treatments become in the real world, simple laboratory tests such as complete blood count will always be valuable. In this regard, both HRR and LMR have been associated with prognosis in several types of cancers [4, 20]. In two recent analyses, HRR is suggested as a prognostic factor for survival in both non-small cell lung cancer and head and neck cancer [21, 22].

Anemia itself is a common phenomenon in cancer patients and is caused by several mechanisms. An association with anemia and prognosis is demonstrated in several types of solid and hematological malignancies [23, 24]. Development of anemia in MM is caused by the infiltration of malign plasma cells in the bone marrow, which results in suppressing erythropoiesis and dysregulated apoptosis of plasma cells [3, 25]. However, stromal and endothelial cells in

**Table III. Characteristics of the multiple myeloma patients according to HRR**

	Low HRR <0.61 (n:82)	High HRR >0.61 (n:98)	p-value
Age			
>65 years old	31 (38%)	49 (50%)	0.132
<65 years old	51 (62%)	49 (50%)	
Gender			
Male	40 (49%)	47 (48%)	0.516
Female	42 (51%)	51 (52%)	
Cytogenetic risk			
High risk	20 (24%)	16 (16%)	0.195
Standard risk	62 (76%)	82 (84%)	
ISS			
1	18 (22%)	50 (51%)	<b>0.001</b>
2	30 (36%)	21 (21%)	
3	34 (42%)	27 (28%)	
Revised ISS			
1	12 (14.6%)	45 (45.9%)	<b>0.001</b>
2	17 (20.7%)	24 (24.5%)	
3	53 (64.6%)	29 (29.6%)	
Immunoglobulin type			
IgG	41 (50%)	55 (56.1%)	0.118
IgA	26 (31.7%)	17 (17.3%)	
Light chain	13 (15.9%)	25 (25.5%)	
Non-secretory	1 (1.2%)	1 (1.0%)	
IgM	1 (1.2%)	0	
Laboratory parameters			
Hemoglobin (g/dl)	9.7	12.0	<b>0.000</b>
Creatinine (mg/dl)	1.60	1.16	<b>0.014</b>
Calcium (mg/dl)	9.6	9.7	0.883
Albumin (g/dl)	3.4	3.9	0.000
LDH (U/L)	237	222	0.338
Beta-2 microglobulin (ng/ml)	7,429	4,229	<b>0.000</b>
CRP (mg/dl)	2.19	1.11	<b>0.011</b>
Frontline treatment			
Bortezomib cyclophosphamide dexametasone	38 (46.3%)	59 (60.2%)	0.168
Bortezomib thalidomide dexametasone	5 (6.1%)	3 (3.1%)	
Bortezomib dexametasone	17 (20.7%)	16 (16.3%)	
VTD-PACE	9 (11.0%)	6 (6.1%)	
Others (melphalan-prednisolone, etc)	13 (15.7%)	14 (14.2%)	
Upfront ASCT			
Yes	23 (28%)	38 (39%)	0.155
No	59 (72%)	60 (61%)	
Overall survival months	36.70 ± 3.77	53.32 ± 3.54	<b>0.001</b>

Univariate analysis for low HRR, HR; 2.046 (95% CI 1.33–3.13)

**Table IV. Multivariate analysis for overall survival**

Multivariate analysis			
	Hazard ratio (HR)	95% CI	p-value
LMR < 3.28	1.47	0.92–2.29	<b>0.010</b>
HRR < 0.61	2.08	1.31–3.03	<b>0.002</b>
Gender	0.74	0.47–1.14	0.178
Age	0.99	0.97–1.02	0.782
No ASCT versus ASCT	0.14	0.07–0.27	<b>0.000</b>
High risk MM	0.35	0.21–0.59	<b>0.000</b>
ISS	1.89	1.09–3.30	<b>0.023</b>
Revised ISS	0.79	0.35–1.19	<b>0.008</b>

bone marrow contribute to the development of anemia by producing specific cytokines, particularly interleukin 6 (IL-6) [8]. Likewise, renal insufficiency also contributes to the development of anemia in MM patients via erythropoietin deficiency [26]. Furthermore, the interaction of plasma cells with the tumor microenvironment might play a role in the progression of the disease and the worsening of anemia [11].

Indeed, RDW itself is a reliable marker of red cell size variability. Several relations with RDW and age-related clonal hematopoiesis were reported. Even high RDW was associated with the risk of acute myeloid leukemia development in healthy individuals [27]. Besides, there was an association of increased RDW with worse OS in myelodysplastic syndrome (MDS) patients, especially in patients with refractory anemia [28]. This observation might be regarded as the effects of dysregulated erythropoiesis on RDW in MDS [28]. RDW is also associated with several types of solid and hematological malignancies [4, 29]. For example, in a recent analysis, high RDW was associated with lower progression-free survival and OS in MM [30]. A retrospective analysis including 161 patients with diffuse large B cell lymphoma evaluated the relationship of RDW with OS [31]. The authors reported that high RDW was associated with lower progression-free survival and OS [31]. Considering the high incidence of anemia in solid and hematological cancers, it is noteworthy that high RDW was associated with lower survival in several hematological malignancies. Therefore, HRR is thought to be a logical and surrogate prognostic marker. However, data including HRR in hematological malignancies are scarce. Several studies including data of real-world patients are needed in hematological cancers, especially in MM. Although hemoglobin or RDW alone has been shown to have an impact on prognosis in MM patients, we have shown that HRR may affect OS and prognosis alone. Considering that both low hemoglobin and high RDW have an impact on prognosis, it will not be difficult and surprising to predict that HRR may also affect prognosis. However, multivariate analysis showed that HRR is associated with poor prognosis compared with high HRR. A second important issue is that both hemoglobin and RDW might be affected by nutritional deficiencies besides the effect of MM infiltration and cytokines. In this case, HRR can be suggested as a more powerful and reliable marker.

The BME holds a critical role, especially in the progress of MM [10]. The interactions in malign plasma cells with BME cause an immunosuppressive state and may result in an immune escape causing myeloma cell growth [11, 13]. These interplays stimulate several regulatory or inhibitory cells. Various suggestions have been made to measure and show the contribution of immunosuppression induced by tumor microenvironment [13, 32]. It has been shown that both ALC and AMC could be an indirect indicator of immune surveillance. Thus, LMR, which is obtained by dividing ALC by AMC, could also be an additional determiner [12, 16]. Several studies investigated the effect of LMR in MM. The prognostic value of ALC, AMC, and LMR was analyzed in 189 MM patients retrospectively [17]. In multivariate analysis, low LMR was an independent poor prognostic factor. Low LMR is reported to be

<2.9 in this study. In another retrospective analysis, the authors stated that peripheral blood monocyte count could be correlated with tumor-associated macrophages, thus reflecting immune status [16]. This study included 372 newly diagnosed MM patients and cutoff value for LMR was determined as 3.6. Low LMR is associated with lower progression-free survival and OS on multivariate analysis [16]. Another study questioned the value of LMR evaluated in 285 MM patients [33]. In this study, the cutoff for LMR was 4.2, and low LMR is found to be an independent factor for OS [33]. These studies indicated the value of LMR in MM patients. In our study, fewer patients in the low LMR group proceeded to the ASCT, which may be the explanation for lower OS in the low LMR group. As we mentioned earlier, the search for a basic and reliable marker for immune regulation and prognosis is mandatory. The main limitation of the studies involving LMR is that lymphocyte and monocyte counts can be affected by a variety of reasons that do not affect the disease, mainly infections. In addition, different cutoff values are taken for LMR in different studies. This situation is also a confounding factor. Therefore, we consider that the combination of LMR and HRR might partially overcome these limitations.

Although medicine is becoming more and more complicated every day, simple, easily accessible, and useful markers are required to predict the prognosis of malignant diseases, especially in developing countries. In this study, we found that two complete blood count parameters that can be easily obtained, such as HRR and LMR, might be independent factors in predicting prognosis in MM patients.

#### **Authors' contributions**

EU – reviewed the data and the manuscript. HOK – data acquisition, data interpretation, reviewed the manuscript. SB – statistical analysis, reviewed the manuscript, data analysis. AMD – edited the data, reviewed the statistical analysis, reviewed the manuscript and a critical review. MB – collected the data, wrote the manuscript, statistical analysis. UD – collected the data, literature search. VB – collected and edited the data. SKG – data acquisition and interpretation, reviewed the manuscript.

#### **Conflict of interest**

None.

#### **Financial support**

None.

#### **Ethics**

The work described in this article has been carried out in accordance with The Code of Ethics of the World Medical Association (Declaration of Helsinki) for experiments involving humans; EU Directive 2010/63/EU for animal experiments; Uniform requirements for manuscripts submitted to biomedical journals.

## References

- [1] Siegel RL, Miller KD, Jemal A. Cancer statistics, 2019. *CA Cancer J Clin* 2019;69:7–34.
- [2] Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 2018;68:394–424.
- [3] Kyle RA, Gertz MA, Witzig TE, et al. Review of 1027 patients with newly diagnosed multiple myeloma. *Mayo Clin Proc* 2003;78:21–33.
- [4] Hu L, Li M, Ding Y, et al. Prognostic value of RDW in cancers: a systematic review and meta-analysis. *Oncotarget* 2017;8:16027–35.
- [5] Sun P, Zhang F, Chen C, et al. The ratio of hemoglobin to red cell distribution width as a novel prognostic parameter in esophageal squamous cell carcinoma: a retrospective study from southern China. *Oncotarget* 2016;7:42650–60.
- [6] Xu WS, Qiu XM, Ou QS, et al. Red blood cell distribution width levels correlate with liver fibrosis and inflammation: a noninvasive serum marker panel to predict the severity of fibrosis and inflammation in patients with hepatitis B. *Medicine (Baltimore)* 2015;94:e612.
- [7] Cetinkaya E, Senol K, Saylam B, Tez M. Red cell distribution width to platelet ratio: new and promising prognostic marker in acute pancreatitis. *World J Gastroenterol* 2014;20:14450–4.
- [8] Sharma S, Nemeth E, Chen YH, et al. Involvement of hepcidin in the anemia of multiple myeloma. *Clin Cancer Res* 2008;14:3262–7.
- [9] Lee H, Kong SY, Sohn JY, et al. Elevated red blood cell distribution width as a simple prognostic factor in patients with symptomatic multiple myeloma. *Biomed Res Int* 2014;2014:145619.
- [10] Stakiw J, Bosch M, Goubran H. A closer look at the bone marrow microenvironment in multiple myeloma. *Tumor and Microenvironment* 2018;1:1–8.
- [11] Manier S, Sacco A, Leleu X, Ghobrial IM, Roccaro AM. Bone marrow microenvironment in multiple myeloma progression. *J Biomed Biotechnol* 2012;2012:157496.
- [12] Suyani E, Sucak GT, Akyürek N, et al. Tumor-associated macrophages as a prognostic parameter in multiple myeloma. *Ann Hematol* 2013;92:669–77.
- [13] Green CE, Liu T, Montel V, et al. Chemoattractant signaling between tumor cells and macrophages regulates cancer cell migration, metastasis and neovascularization. *PLoS One* 2009;4:e6713.
- [14] Koh YW, Kang HJ, Park C, et al. The ratio of the absolute lymphocyte count to the absolute monocyte count is associated with prognosis in Hodgkin's lymphoma: correlation with tumor-associated macrophages. *Oncologist* 2012;17:871–80.
- [15] Binder M, Rajkumar SV, Lacy MQ, et al. Peripheral blood biomarkers of early immune reconstitution in newly diagnosed multiple myeloma. *Am J Hematol* 2019;94:306–11.
- [16] Dosani T, Covut F, Beck R, Driscoll JJ, de Lima M, Malek E. Significance of the absolute lymphocyte/monocyte ratio as a prognostic immune biomarker in newly diagnosed multiple myeloma. *Blood Cancer J* 2017;7:e579.
- [17] Shin SJ, Roh J, Kim M, et al. Prognostic significance of absolute lymphocyte count/absolute monocyte count ratio at diagnosis in patients with multiple myeloma. *Korean J Pathol* 2013;47:526–33.
- [18] Palumbo A, Avet-Loiseau H, Oliva S, et al. Revised International Staging System for multiple myeloma: a report from International Myeloma Working Group. *J Clin Oncol* 2015;33:2863–9.
- [19] Sonneveld P, Avet-Loiseau H, Lonial S, et al. Treatment of multiple myeloma with high-risk cytogenetics: a consensus of the International Myeloma Working Group. *Blood* 2016;127:2955–62.
- [20] Gu L, Li H, Chen L, et al. Prognostic role of lymphocyte to monocyte ratio for patients with cancer: evidence from a systematic review and meta-analysis. *Oncotarget* 2016;7:31926–42.
- [21] Bozkaya Y, Kurt B, Gurler F. A prognostic parameter in advanced non-small cell lung cancer: the ratio of hemoglobin-to-red cell distribution width. *Int J Clin Oncol* 2019;24:798–806.
- [22] Tham T, Olson C, Wotman M, et al. Evaluation of the prognostic utility of the hemoglobin-to-red cell distribution width ratio in head and neck cancer. *Eur Arch Otorhinolaryngol* 2018;275:2869–78.
- [23] Tas F, Eralp Y, Basaran M, et al. Anemia in oncology practice: relation to diseases and their therapies. *Am J Clin Oncol* 2002;25:371–9.
- [24] Littlewood T, Mandelli F. The effects of anemia in hematologic malignancies: more than a symptom. *Semin Oncol* 2002;29(Suppl 8): 40–4.
- [25] Tucci M, Grinello D, Cafforio P, Silvestris F, Dammacco F. Anemia in multiple myeloma: role of deregulated plasma cell apoptosis. *Leuk Lymphoma* 2002;43:1527–33.
- [26] Katagiri D, Noiri E, Hinoshita F. Multiple myeloma and kidney disease. *Scientific World Journal* 2013;2013:487285.
- [27] Abelson S, Collord G, Ng SWK, et al. Prediction of acute myeloid leukaemia risk in healthy individuals. *Nature* 2018;559:400–4.
- [28] Baba Y, Saito B, Shimada S, et al. Association of red cell distribution width with clinical outcomes in myelodysplastic syndrome. *Leuk Res* 2018;67:56–9.
- [29] Montagnana M, Danese E. Red cell distribution width and cancer. *Ann Transl Med* 2016;4:399.
- [30] Ma Y, Jin Z, Zhou S, Ye H, Jiang S, Yu K. Prognostic significance of the red blood cell distribution width that maintain at high level following completion of first line therapy in multiple myeloma patients. *Oncotarget* 2018;9:10118–27.
- [31] Zhou S, Fang F, Chen H, et al. Prognostic significance of the red blood cell distribution width in diffuse large B-cell lymphoma patients. *Oncotarget* 2017;8:40724–31.
- [32] Rabinovich GA, Gabrilovich D, Sotomayor EM. Immunosuppressive strategies that are mediated by tumor cells. *Annu Rev Immunol* 2007;25:267–96.
- [33] Tian Y, Zhang Y, Zhu WQ, Chen XL, Zhou HB, Chen WM. Peripheral blood lymphocyte-to-monocyte ratio as a useful prognostic factor in newly diagnosed multiple myeloma. *Biomed Res Int* 2018;2018:9434637.