

**UNIVERSIDAD DE SAN CARLOS DE GUATEMALA**  
**FACULTAD DE MEDICINA VETERINARIA Y ZOOTECNIA**  
**ESCUELA DE MEDICINA VETERINARIA**



**ANTIBIOTIC-RESISTANT SALMONELLA, ISOLATED FROM  
CLOACAL SWAB SAMPLES FROM TURTLES IN  
GUATEMALA - SALMONELLA RESISTENTE A  
ANTIBIÓTICOS, AISLADA DE MUESTRAS DE HISOPADOS  
CLOACALES EN TORTUGAS DE GUATEMALA**

**JOANA GABRIELA BORRAYO GODINEZ**

**MÉDICA VETERINARIA**

**GUATEMALA, OCTUBRE DE 2020**

**UNIVERSIDAD DE SAN CARLOS DE GUATEMALA  
FACULTAD DE MEDICINA VETERINARIA Y ZOOTECNIA  
ESCUELA DE MEDICINA VETERINARIA**



**ANTIBIOTIC-RESISTANT SALMONELLA, ISOLATED FROM  
CLOACAL SWAB SAMPLES FROM TURTLES IN  
GUATEMALA**

**SALMONELLA RESISTENTE A ANTIBIÓTICOS, AISLADA DE  
MUESTRAS DE HISOPADOS CLOACALES EN TORTUGAS  
DE GUATEMALA**

**TRABAJO DE GRADUACIÓN**

**PRESENTADO A LA HONORABLE JUNTA DIRECTIVA DE LA FACULTAD DE  
MEDICINA VETERINARIA Y ZOOTECNIA**

**POR**

**JOANA GABRIELA BORRAYO GODINEZ**

**Al conferírsele el título profesional de**

**Médica Veterinaria**

**En el grado de Licenciado**

**GUATEMALA, OCTUBRE DE 2020**

**UNIVERSIDAD DE SAN CARLOS DE GUATEMALA**  
**FACULTAD DE MEDICINA VETERINARIA Y ZOOTECNIA**  
**JUNTA DIRECTIVA**

<b>DECANO:</b>	M.V. Gustavo Enrique Taracena Gil
<b>SECRETARIO:</b>	Dr. Hugo René Pérez Noriega
<b>VOCAL I:</b>	M.Sc. Juan José Prem González
<b>VOCAL II:</b>	Lic. Zoot. Miguel Ángel Rodenas Argueta
<b>VOCAL III:</b>	Lic. Zoot. Alex Rafael Salazar Melgar
<b>VOCAL IV:</b>	P. Agr. Luis Gerardo López Morales
<b>VOCAL V:</b>	Br. María Jose Solares Herrera

**ASESORES**

PHD. DENNIS SIGFRIED GUERRA CENTENO

DR. SC. CARLOS VALDEZ SANDOVAL

MSC. MANUEL LEPE LÓPEZ

En cumplimiento con lo establecido por los reglamentos y normas de la Universidad de San Carlos de Guatemala, presento a su consideración el artículo científico aceptado para publicación en una revista indexada, titulado:

**ANTIBIOTIC-RESISTANT SALMONELLA, ISOLATED FROM  
CLOACAL SWAB SAMPLES FROM TURTLES IN  
GUATEMALA**

**SALMONELLA RESISTENTE A ANTIBIÓTICOS, AISLADA DE  
MUESTRAS DE HISOPADOS CLOACALES EN TORTUGAS  
DE GUATEMALA**

Que fuera aprobado por la Honorable Junta Directiva de la Facultad de  
Medicina Veterinaria y Zootecnia

Como requisito previo a optar al título de:

**MÉDICA VETERINARIA**

## **ACTO QUE DEDICO**

- A DIOS:** Por ser el pilar más valioso para mí, por acompañarme y guiarme en cada momento de mi vida y ser una fuente inagotable de amor, valentía, fuerza y sabiduría. Por permitirme alcanzar ésta meta y cumplir uno de mis sueños.
- A MIS PADRES:** Anibal Borrayo y Alma de Borrayo, por darme la vida y ser un ejemplo de esfuerzo, superación y perseverancia. Porque nunca dejaron de creer en mí y siempre lo han dado todo para que yo pudiera llegar hasta esta meta. Por su apoyo y amor incondicional, éste triunfo también es de ustedes.
- A MI HERMANA:** Cynthia Borrayo, por ser mi ayuda idónea, por siempre estar en los momentos buenos y malos durante toda mi carrera, por acompañarme en las noches de desvelos y siempre que te necesito.
- A MI ESPOSO:** Andrés Pivaral, por estar a mi lado desde que este sueño comenzó a fluir, por animarme y motivarme a dar siempre lo mejor de mí, por todo tu amor y paciencia.
- A MIS ABUELOS:** Rubén Moncada y Ángela Veliz que el día de hoy ya no están conmigo pero dejaron su legado en mí, por llenar nuestra familia de valores, humildad y amor.

## **AGRADECIMIENTOS**

- A DIOS:** Por su amor, gracia, misericordia, bondad y por su guía.
- A MIS PADRES:** Gracias ayudarme a alcanzar esta meta, por todo su sacrificio, esfuerzo y todas las desveladas, porque nunca me hizo falta nada en la vida y durante mi estadía en la universidad.
- A MI HERMANA:** Cynthia, por tu amor, tus ánimos y apoyo incondicional y por todos los momentos que hemos compartido.
- A MI ESPOSO:** Andrés, por ser ese pilar que me ayuda a diario, que me edifica y me hace creer en mí y en mis habilidades, por ser el mejor compañero y por compartir esa pasión de amor por los animales. Te amo mi amor sos el regalo más grande que Dios me ha dado.
- A MI FAMILIA:** Los Forkel, los Borrayo, los Moncada, los Contreras, los Sierra, los Pivaral, los Santizo, por todo su amor y apoyo.
- A MIS MADRINAS:** Joss y Lu porque han llegado a ser más que mis mejores amigas mis hermanas, por estar en cada paso importante y no tan importante y porque nunca han dejado de creer en mí, las amo.
- A MIS AMIGOS:** A cada uno por nombre, por todos los momentos que hemos compartido, por sus muestras de cariño y por su valiosa amistad por la ayuda constante y porque nunca me han dejado sola en este camino.

**A LA FMVZ:**

Por ser mi casa de estudios y por la formación académica y profesional.

**A MI ASESOR:**

Por su tiempo, buena disposición, amabilidad y apoyo.

**A:**

Todas las personas que de una u otra forma ayudaron a la realización del estudio, muchas gracias.

## ÍNDICE

ABSTRACT .....	1
RESUMEN .....	2
INTRODUCTION.....	3
MATERIALS AND METHODS .....	4
Study site .....	4
Sample collection.....	4
Laboratory procedures for <i>Salmonella</i> isolation .....	5
Antibiotic resistance test .....	5
RESULTS.....	5
DISCUSSION.....	6
ACKNOWLEDGEMENTS .....	9
REFERENCES.....	10



**ANTIBIOTIC-RESISTANT *SALMONELLA*,  
ISOLATED FROM CLOACAL SWAB SAMPLES FROM TURTLES IN  
GUATEMALA**

***SALMONELLA* RESISTENTE A ANTIBIÓTICOS,  
AISLADA DE MUESTRAS DE HISOPADOS CLOACALES EN  
TORTUGAS DE GUATEMALA**

**ABSTRACT**

Salmonellosis is a relevant public health threat worldwide. Reptiles are commonly involved in human cases. A microbiological survey was conducted from August to October 2018 to isolate *Salmonella* bacteria and determine if they are resistant to regularly used antibiotics in five species of pet turtles (*Kinosternon* sp., *Rhinoclemmys* sp., *Staurotypus* sp., *Trachemys scripta* y *T. venusta*) in Guatemala City, Antigua Guatemala and San Lucas Sacatepéquez. Cloacal swabs were taken from 63 turtles and cultivated in the Microbiology Laboratory at the Veterinary Medicine and Animal Husbandry Faculty, University of San Carlos of Guatemala, in Guatemala City. Three samples were positive to the presence of *Salmonella* sp. One of these isolates (from *Trachemys scripta*) was resistant to gentamicin, penicillin and amikacin, other isolate (from *T. scripta*) was partially resistant to amoxicilin + clavulanic acid and penicillin, and other (from *T. venusta*) to penicillin. These findings highlight the need for better biosecurity practices and show the capacity of bacteria to develop survival strategies that involve resistance to harmful substances like antibiotics.

Key words: Salmonellosis, public health, epidemiology, zoonoses, multiresistant gram negative bacteria.

## RESUMEN

La salmonelosis es una importante enfermedad zoonótica considerada una amenaza a la salud pública a nivel mundial. Los reptiles están comúnmente involucrados en la transmisión animal-humano. Con el objetivo de determinar la presencia de *Salmonella* y determinar su resistencia a antibióticos de uso común, se realizó un estudio exploratorio en cinco especies de tortugas (*Kinosternon* sp., *Rhinoclemmys* sp., *Staurotypus* sp., *Trachemys scripta* y *T. venusta*) en Guatemala, Antigua Guatemala y en San Lucas Sacatepéquez. Se tomaron hisopados cloacales de 63 especímenes y se cultivaron en el Laboratorio de Microbiología de la Facultad de Medicina Veterinaria y Zootecnia de la Universidad de San Carlos de Guatemala. Tres muestras fueron positivas a la presencia de *Salmonella* sp. Uno de los aislados (de *Trachemys scripta*) fue resistente a gentamicina, penicilina y amikacina, otro aislado (de *T. scripta*) fue parcialmente resistente a amoxicilina + ácido clavulánico y a penicilina y un tercer aislado (de *T. venusta*) a penicilina. Estos hallazgos resaltan la necesidad de mejores prácticas de bioseguridad y muestran la capacidad de las bacterias para desarrollar estrategias de sobrevivencia que involucran la resistencia a sustancias que les son nocivas, como los antibióticos.

Palabras clave: Salmonelosis, salud pública, epidemiología, zoonosis, bacterias gram negativas multirresistentes.

## INTRODUCTION

Nontyphoidal *Salmonella* is an important public health threat, causing an estimated of 93.8 million cases of gastroenteritis globally, including 155,000 deaths (Majowicz et al., 2010). Over 2,500 *Salmonella* serotypes have been identified and more than half of them belong to *Salmonella enterica* subspecies *enterica* which causes most *Salmonella* infections in humans (Eng et al., 2015). Not only is salmonellosis one of the diseases affecting socio-economic development worldwide (World Health Organization, 2015), but is also a major contemporary concern due to its dynamic epidemic behavior (Ao et al., 2015; Besser, 2018) and to the growing ability of the causing bacteria serotypes to develop multiple resistance to antibiotics (Blair et al., 2015; Ventola, 2015; Wong et al., 2015).

Many animal species, both domestic and wild, are reservoirs of *Salmonella* and harbor these bacteria in their gastrointestinal tracts with no apparent signs of illness (MacDonald et al., 2019; Oludairo et al., 2013; Sanchez et al., 2002). Salmonellosis outbreaks are often associated with exotic pet ownership (Woodward et al., 1997). Turtles, specially inland species, were common sources of *Salmonella* in many of these outbreaks in United States (Bosch et al., 2016; Centers for Disease Control and Prevention [CDC], 2007; Harris et al., 2009; Harris et al., 2010; Warwick et al., 2001), Spain (Hernández et al., 2012; Hidalgo-Vila et al., 2007) and Japan (Nagano et al., 2006). By 1971, 280,000 turtle-associated cases were estimated to occur only in the United States (Lamm et al., 1972) and although some public health measures have been taken since then, turtles are still common sources of *Salmonella* infections (Bosch et al., 2016; Cummings et al., 2012; Hale et al., 2012). This, because turtles are the second most popular pets after birds (Bush et al., 2014) and sometimes live in frequent physical contact with their owners (Siegmund & Biermann, 1988).

Although salmonellosis is an important zoonosis in Latin America (Gil & Samartino, 2001), very few studies have addressed the role of domestic turtles in the cases or outbreaks in this region (Braun et al., 2015; Sacristán-Rodríguez et al., 2014; Tauxe et al., 1985). Considering the role of pet turtles in the epidemiology of salmonellosis in other parts of the world (Hoelzer et al., 2011; Sodgari et al., 2020), the objectives in this exploratory survey were to look for *Salmonella* organisms in turtle cloacal swab samples and to test the isolates for antibiotic resistance. The findings provide the first report on the circulation of *Salmonella*, including antibiotic-resistant isolates, in pet turtles in Guatemala.

## **MATERIALS AND METHODS**

### **Study site**

Samples were taken from households in Guatemala City and San Lucas Sacatepéquez, and from a rescue center in Antigua Guatemala. Turtle owners were contacted through a call placed in social networks and in the Veterinary Faculty of University of San Carlos of Guatemala website. A maximum of three specimens were collected from each sampling sites except in the rescue center where nine samples were taken from turtles that had recently been brought from a domestic environment. Figure 1 shows the sampling sites distribution.

### **Sample collection**

In a period of three months (August to October 2018) 63 cloacal swabs were taken from turtles of eight species (1 *Kinosternon acutum*, 3 *K. leucostomum*, 4 *K. scorpioides*, 14 *Rinoclemmys pulcherrima*, 3 *R. areolata*, 3 *Staurotypus salvinii*, 22 *Trachemys scripta* and 13 *Trachemys venusta*). The sampled turtles were kept either in backyards or in

aquaria and all were being fed commercial food for turtles and, occasionally, vegetables.

Samples were collected from the cloaca, using a sterile peptone water-soaked swab, and transported in test tubes with peptone water to the Microbiology Laboratory of the Veterinary Faculty of University of San Carlos of Guatemala. Samples were received for processing when they had less than 4 hours of being taken.

### **Laboratory procedures for *Salmonella* isolation**

*Salmonella* isolation was performed according to the ISO 6579 International Standard for *Salmonella* spp. detection (International Organization for Standardization, 2002). The samples were processed in duplicate to improve the chances of *Salmonella* isolation. Since this was an exploratory investigation, *Salmonella* serovars were not identified.

### **Antibiotic resistance test**

The *Salmonella* isolates were tested for resistance against amoxiciline + clavulanic acid, cefotetan, oxytetracycline, gentamicine, penicillin, amikacin, nalidixic acid, doxycycline, and ceftriaxone, following the disk sensitivity test procedures presented by Sacsquispe and Velásquez (2002).

## **RESULTS**

*Salmonella* bacteria was isolated from three of the 63 cloacal swabs. Even though specimens were sampled from eight turtle species reported for Guatemala (Acevedo et al., 2010) the three isolates came only from turtles of the genus *Trachemys* (Table 1). Although not an objective of this study, *Escherichia coli*, *Proteus* sp., and *Enterobacter* sp. were common findings in all cloacal swabs.

The *Salmonella* isolates were resistant or partially resistant to at least one antibiotic. The first isolate (from *Trachemys scripta*) was resistant to amikacin, gentamicin and penicillin. The second isolate (from *T. venusta*) was partially resistant to amoxicillin/clavulanic acid and penicillin. The third isolate (from *T. scripta*) was partially resistant to penicillin.

## DISCUSSION

The prevalence of *Salmonella* isolates in our sample was rather low (4.7%). This level of detection is consistent with other investigations in turtle populations where the prevalence was close to zero (Brenner et al., 2002; Mitchell & McAvoy, 1990; Richards et al., 2004; Saelinger et al., , 2006; Way & Allender, 2011). However, other studies have detected prevalences ranging from 8 to 18.75% (Charles-Smith et al., 2009; Readell et al., 2010; Sharma et al., 2020; Tomastikova et al., ; Zhang et al., 2016). It is known that *Salmonella* shedding could be intermittent (Kaufmann et al., 1967) and stress-related in reptiles (Burnham et al., 1998; Chiodini & Sundberg, 1981; DuPonte et al., 1978), so, this apparent prevalence should be taken with caution. Cloacal swab sampling could as well not be the best technique for *Salmonella* detection in comparison to better –but more invasive– methods like mucosa scrapings. On the other hand, *Salmonella* has been a common finding in imported turtles from captive breeding facilities, mostly from the USA (D’Aoust et al., 1990; Lukac et al., 2015; Shane et al., 1990).

It should be noted that *Trachemys* turtles were the only source of the *Salmonella* isolates in this study. *Trachemys scripta* is a common pet turtle worldwide. In Guatemala, it is sold at a very affordable price (around 2.5 US \$ each) in most pet stores around the country. Over 36,000 *T. scripta* turtles were imported to Guatemala in 2018 to supply the pet market (R. De León, personal communication, April 20, 2019).


This turtle species has been considered one of the most common sources of *Salmonella* to humans (Hidalgo-Vila et al., 2008; Lamm et al., 1972; Nagano et al., 2006; Pasmans et al., 2002; Readell et al., 2010; Shen et al., 2011) and also an alien invasive species (Kuzmanova et al., 2018; Ma & Shi, 2017; Zhang et al., 2020).

Based on the premises that some of the sampled turtles in the present investigation were shedding *Salmonella* bacteria with some degree of antibiotic resistance and that thousands of turtles are being imported annually to Guatemala for the pet market, we can assume that there is some risk of acquiring salmonellosis from the contact with these animals or their environment. It has been stressed out that the risk of turtles as a source of human salmonellosis, especially for young children, should not be underestimated (Sodagari et al., 2020). Furthermore, not only could *Salmonella* cause highly pathogenic cases in humans (Nagano et al., 2006), but also hard to treat cases due to antibiotic resistance (Eng et al., 2015; Liang et al., 2015).

Resistance to commonly use antibiotics –such as erythromycin, sulfisoxazole, gentamycin, amoxicillin and ampicillin– by *Salmonella* isolates have been previously reported for various species of reptiles (Ebani et al., 2005). *Salmonella* strains from turtles imported from the USA have shown to be resistant to gentamicin and other antibiotics commonly used for treating turtles after the 1974 small turtle trade ban (Shane et al., 1990). Antibiotic use as supplement in livestock and other captive bred animals like turtles has been identified as one of the main causes of the antibiotic resistance crisis (Angulo et al., 2014; Eng et al., 2015; Ventola, 2015). As the household size become smaller in modern societies, people look for more suitable pets like turtles to fit small apartments. Children could then be the more exposed age group to become infected with *Salmonella* from turtles because they use to play with their pets (Kaibu et al., 2006; Pees et al., 2013; Siegmund & Biermann, 1988) and because turtles are one of the most commercialized pets worldwide (Bush et al., 2014).

Our preliminary data suggests that only *Trachemys* turtles harbor *Salmonella* bacteria amongst inland turtle species in Guatemala. Interesting also, is the fact that we found the bacteria in imported (*T. scripta*) and native (*T. venusta*) specimens of this genus of turtle. This could be the result of inter-species transmission between imported *T. scripta* and native *T. venusta* housed together or coexisting in free ranging conditions. *Salmonella* was already found in wild populations of naturalized *T. scripta* in China (Gong et al., 2013). *Salmonella* shedding and transmission to naturalized turtles could even threaten native turtle populations. More research needs to be done to understand the ecology of *Salmonella* bacteria across the inland and marine species of turtles in Guatemala in domestic or wild environments.

The results of this investigation reinforce the idea that captive raising of farm or wild animals could provide the necessary conditions for the establishment and dissemination of *Salmonella*. Furthermore, the practice of administering preventive or therapeutic antibiotics to captive populations favors the evolution of multi-resistant bacteria strains which represent a contemporary public health issue (Kariuki et al, 2015; Medalla et al., 2017; Ventola, 2015).



**Figure 1.**

**Turtle sampling sites in Guatemala City, San Lucas Sacatepéquez and Antigua Guatemala**



**Table 1.**  
**Distribution of Salmonella isolates in the sampled turtle specimens**

<b>Species</b>	<b><i>n</i></b>	<b>Number of Isolates</b>
<i>Kinosternon acutum</i>	1	0
<i>Kinosternon leucostomum</i>	3	0
<i>Kinosternon scorpioides</i>	4	0
<i>Rhinoclemmys areolata</i>	3	0
<i>Rhinoclemmys pulcherrima</i>	14	0
<i>Staurotypus salvinii</i>	3	0
<i>Trachemys scripta</i>	22	2
<i>Trachemys venusta</i>	13	1
<b>Total</b>	<b>63</b>	<b>3</b>

## **ACKNOWLEDGEMENTS**

We thank Martin Cutzán of the Microbiology Laboratory, Veterinary and Animal Husbandry Faculty, University of San Carlos of Guatemala for processing the cloacal samples for *Salmonella* isolation and antibiotic resistance.

## REFERENCES

Acevedo, M., Wilson, L. D., Cano, E. B., & Vasquez-Almazán, C. (2010) Diversity and conservation status of the Guatemalan herpetofauna. In L. Wilson, J. Townsend & J. Johnson (Eds), *Conservation of Mesoamerican amphibians and reptiles* (pp. 407-434). Eagle Mountain, Utah: Eagle Mountain Publishing, LC.

Angulo, F. J., Baker, N. L., Olsen, S. J., Anderson, A., & Barrett, T. J. (2004, April). Antimicrobial use in agriculture: controlling the transfer of antimicrobial resistance to humans. *Seminars in Pediatric Infectious diseases*, 15(2), 78-85. <https://doi.org/10.1053/j.spid.2004.01.010>

Ao, T. T., Feasey, N. A., Gordon, M. A., Keddy, K. H., Angulo, F. J., & Crump, J. A. (2015). Global burden of invasive nontyphoidal Salmonella disease, 2010. *Emerging Infectious Diseases*, 21(6), 941. <https://doi.org/10.3201/eid2106.140999>

Besser, J. M. (2018). Salmonella epidemiology: A whirlwind of change. *Food Microbiology* 71, 55-59. <https://doi.org/10.1016/j.fm.2017.08.018>

Blair, J. M., Webber, M. A., Baylay, A. J., Ogbolu, D. O., & Piddock, L. J. (2015). Molecular mechanisms of antibiotic resistance. *Nature Reviews Microbiology*, 13(1), 42-51. <https://doi.org/10.1038/nrmicro3380>

Bosch, S., Tauxe, R. V., & Behravesh, C. B. (2016). Turtle-associated salmonellosis, United States, 2006–2014. *Emerging Infectious Diseases*, 22(7), 1149-1155. <https://doi.org/10.3201/eid2207.150685>

Braun, S., Spalloni, W., Ferreccio, F., Postigo, J., Fernández, A., Porte, L., ... Triantafilo, V. (2015). Gastroenteritis por *Salmonella* spp. en tres lactantes asociada a contacto con tortugas acuáticas. *Revista Chilena de Infectología*, 32, 334-338. <http://dx.doi.org/10.4067/S0716-10182015000400013>

Brenner, D., Lewbart, G., Stebbins, M., & Herman, D. W. (2002). Health survey of wild and captive bog turtles (*Clemmys muhlenbergii*) in North Carolina and Virginia. *Journal of Zoo and Wildlife Medicine*, 33(4), 311-317. [https://doi.org/10.1638/1042-7260\(2002\)033\[0311:HSOWAC\]2.0.CO;2](https://doi.org/10.1638/1042-7260(2002)033[0311:HSOWAC]2.0.CO;2)

Burnham, B. R., Atchley, D. H., DeFusco, R. P., Ferris, K. E., Zicarelli, J. C., Lee, J. H., & Angulo, F. J. (1998). Prevalence of fecal shedding of *Salmonella* organisms among captive green iguanas and potential public health implications. *Journal of the American Veterinary Medical Association*, 213(1), 48-50.

Bush, E. R., Baker, S. E., & Macdonald, D. W. (2014). Global trade in exotic pets 2006–2012. *Conservation Biology*, 28(3), 663-676. <https://doi.org/10.1111/cobi.12240>

Centers for Disease Control and Prevention. (2007). Turtle-associated salmonellosis in humans-United States, 2006-2007. *Morbidity and Mortality Weekly Report*, 56(26),649.

Charles-Smith, L. E., Lewbart, G. A., Aresco, M. J., & Cowen, P. (2009). Detection of *Salmonella* in Gopher Tortoises (*Gopherus polyphemus*) during two relocation efforts in Florida. *Chelonian Conservation and Biology*, 8(2), 213-216. <https://doi.org/10.2744/CCB-0771.1>

Chiodini, R. J., & Sundberg, J. P. (1981). Salmonellosis in reptiles: a review. *American Journal of Epidemiology*, 113(5), 494-499. <https://doi.org/10.1093/oxforjournals.aje.a113124>

Cummings, P. L., Sorvillo, F., & Kuo, T. (2012). The burden of salmonellosis in the United States. En B. S. Mahmoud. (Ed.), *Salmonella a dangerous foodborne pathogen* (pp. 1-20). Rijeka, Croatia: In Tech.

D'Aoust, J. Y., Daley, E., Crozier, M., & Sewell, A. M. (1990). Pet turtles: A continuing international threat to public health. *American Journal of Epidemiology*, 132(2), 233-238. <https://doi.org/10.1093/oxfordjournals.aje.a115652>

DuPonte, M. W., Nakamura, R. M., & Chang, E. M. (1978). Activation of latent Salmonella and Arizona organisms by dehydration of red-eared turtles, *Pseudemys scripta-elegans*. *American Journal of Veterinary Research*, 39(3), 529-530.

Ebani, V. V., Cerri, D., Fratini, F., Meille, N., Valentini, P., & Andreani, E. (2005). Salmonella enterica isolates from faeces of domestic reptiles and a study of their antimicrobial in vitro sensitivity. *Research in Veterinary Science*, 78(2), 117-121. <https://doi.org/10.1016/j.rvsc.2004.08.002>

Eng, S. K. Pusparajah, P., Ab Mutalib, N. S., Ser, H. L., Chan, K. G., & Lee, L. H. (2015). Salmonella: A review on pathogenesis, epidemiology and antibiotic resistance. *Frontiers in Life Science*, 8(3), 284-293. <https://doi.org/10.1080/21554769.2015.1051243>

Gil, A., & Samartino, L. (2001). Zoonosis en los sistemas de producción animal de las áreas urbanas y periurbanas de América Latina. Roma: FAO.

Gong, S., Wang, F., Shi, H., Zhou, P., Ge, Y., Hua, L., & Liu, W. (2013). Highly pathogenic Salmonella Pomona was first isolated from the exotic red-eared slider (*Trachemys scripta elegans*) in the wild in China: Implications for public health. *Science of the Total Environment*, 468, 28-30. <https://doi.org/10.1016/j.scitotenv.2013.08.025>

Hale, C. R., Scallan, E., Cronquist, A. B., Dunn, J., Smith, K., Robinson, T., ... & Clogher, P. (2012). Estimates of enteric illness attributable to contact with animals and their environments in the United States. *Clinical Infectious Diseases*, 54(suppl\_5), S472-S479. <https://doi.org/10.1093/cid/cis051>

Harris, J. R., Bergmire-Sweat, D., Schlegel, J. H. Winpisinger, K. A., Klos, R. F., Perry, C., ... Sotir, M. J. (2009). Multistate outbreak of *Salmonella* infections associated with small turtle exposure, 2007-2008. *Pediatrics*, *124*(5), 1388-1394. <https://doi.org/10.1542/peds.2009-0272>

Harris, J. R., Neil, K. P., Behravesh, C. B., Sotir, M. J., & Angulo, F. J. (2010). Recent multistate outbreaks of human *Salmonella* infections acquired from turtles: A continuing public health Challenge. *Clinical Infectious Diseases*, *50*(4), 554-559. <https://doi.org/10.1086/649932>

Hernández, E., Rodriguez, J. L., Herrera-León, S., García, I., De Castro, V., & Muniozguren, N. (2012). *Salmonella* Paratyphi B var Java infections associated with exposure to turtles in Bizkaia, Spain, September 2010 to October 2011. *Eurosurveillance*, *17*(25), 20201. <https://doi.org/10.2807/ese.17.25.20201-en>

Hidalgo-Vila, J., Díaz-Paniagua, C., de Frutos-Escobar, C., Jiménez-Martínez, C., & Pérez-Santigosa, N. (2007). *Salmonella* in free living terrestrial and aquatic turtles. *Veterinary Microbiology*, *119*(2-4), 311-315. <https://doi.org/10.1016/j.vetmic.2006.08.012>

Hidalgo-Vila, J., Díaz-Paniagua, C., Pérez-Santigosa, N., de Frutos-Escobar, C., & Herrero-Herrero, A. (2008). *Salmonella* in free-living exotic and native turtles and in pet exotic turtles from SW Spain. *Research in Veterinary Science*, *85*, 449-452. <https://doi.org/10.1016/j.rvsc.2008.01.011>

Hoelzer, K., Switt, A. I. M., & Wiedmann, M. (2011). Animal contact as a source of human non-typhoidal salmonellosis. *Veterinary Research*, *42*(1), 34. <https://doi.org/10.1186/1297-9716-42-34>

Kaibu, H., Iida, K., Ueki, S., Ehara, H., Simasaki, Y., Anzai, H., ... Shirono, S. (2006). Salmonellosis of infants presumably originating from an infected turtle in Nagasaki, Japan. *Japanese Journal of Infectious Diseases*, 59(4), 281.

Kariuki, S., Gordon, M. A., Feasey, N., & Parry, C. M. (2015). Antimicrobial resistance and management of invasive *Salmonella* disease. *Vaccine*, 33, C21-C29. <https://doi.org/10.1016/j.vaccine.2015.03.102>

Kaufmann, A. F., Feeley, J. C., & DeWitt, W. E. (1967). *Salmonella* excretion by turtles. *Public Health Reports*, 82(9), 840. <https://doi.org/10.2307/4593144>

Kuzmanova, Y., Natcheva, I., Koleva, V., Popgeorgiev, G., Slavchev, M., & Natchev, N. (2018). Public awareness of risks and recent marketing dynamics of Pond sliders (*Trachemys scripta*, Schoepff, 1792) in NE Bulgaria. *Zoology and Ecology*, 28(4), 329-336. <https://doi.org/10.1080/21658005.2018.1521675>

International Organization for Standardization. (2002). ISO 6579: Microbiology of food and animal feeding stuffs – Horizontal method for the detection of *Salmonella* spp. Geneva: Author.

Lamm, S. H., Taylor, A., Gangarosa, E. J., Anderson, H. W., Young, W., Clark, M. H., & Bruce, A. R. (1972). Turtle-associated salmonellosis. I. An estimation of the magnitude of the problem in the United States, 1970-1971. *American Journal of Epidemiology*, 95(6), 511-517. <https://doi.org/10.1093/oxfordjournals.aje.a121418>

Liang, Z., Ke, B., Deng, X., Liang, J., Ran, L., Lu, L., ... & Yu, H. (2015). Serotypes, seasonal trends, and antibiotic resistance of non-typhoidal *Salmonella* from human patients in Guangdong Province, China, 2009-2012. *BMC Infectious Diseases*, 15(1), 1-9. <https://doi.org/10.1186/s12879-015-0784-4>.

Lukac, M., Pedersen, K., & Prukner-Radovic, E. (2015). Prevalence of Salmonella in captive reptiles from Croatia. *Journal of Zoo and Wildlife Medicine*, 46(2), 234-240. <https://doi.org/10.1638/2014-0098R1.1>

Ma, K., & Shi, H. (2017). Red-eared slider *Trachemys scripta elegans* (Wied-Neuwied). In F. Wan, M. Jiang & A. Zhan (Eds.), *Biological invasions and its management in China* (pp. 49-76). Singapore: Springer.

MacDonald, E, White, R., Meixa, R., Bruun, T., Kapperud, G., Brandal., ... Vold, L. (2019). The role of domestic reservoirs in domestically acquired Salmonella infections in Norway: Epidemiology of salmonellosis, 2000–2015, and results of a national prospective case–control study, 2010–2012. *Epidemiology & Infection* 147, 1-8. <https://doi.org/10.1017/S0950268818002911>

Majowicz, S. E., Musto, J., Scallan, E., Angulo, F. J., Kirk, M., O'brien, S. J., ... & International Collaboration of Enteric Disease “Burden of Illness” Studies. (2010). The global burden of nontyphoidal Salmonella gastroenteritis. *Clinical Infectious Diseases*, 50(6), 882-889. <https://doi.org/10.1086/650733>

Medalla, F., Gu, W., Mahon, B. E., Judd, M., Folster, J., Griffin, P. M., & Hoekstra, R. M. (2017). Estimated incidence of antimicrobial drug–resistant nontyphoidal Salmonella infections, United States, 2004–2012. *Emerging Infectious Diseases*, 23(1), 29. <https://doi.org/10.3201/eid2301.160771>

Mitchell, J. C., & McAvoy, B. V. (1990). Enteric bacteria in natural populations of freshwater turtles in Virginia. *Virginia Journal of Science*, 41(3), 233-242.

Nagano, N., Oana, S., Nagano, Y., & Arakawa, Y. (2006). A severe *Salmonella enterica* serotype Paratyphi B infection in a child related to a pet turtle, *Trachemys scripta elegans*. *Japanese Journal of Infectious Diseases*, 59(2), 132.

Oludairo, O. O., Kwaga, J. K. P., Dzikwi, A. A., & Kabir, J. (2013). The genus *Salmonella*, isolation and occurrence in wildlife. *International Journal of Microbiology and Immunology Research*, 1(5), 47-52.

Pasmans, F., De Herdt, P., Dewulf, J., & Haesebrouck, F. (2002). Pathogenesis of infections with *Salmonella enterica* subsp. *enterica* serovar Muenchen in the turtle *Trachemys scripta scripta*. *Veterinary Microbiology*, 87(4), 315-325. [https://doi.org/10.1016/S0378-1135\(02\)00081-0](https://doi.org/10.1016/S0378-1135(02)00081-0)

Pees, M., Rabsch, W., Plenz, B., Fruth, A., Prager, R., Simon, S., ... Braun, P. (2013). Evidence for the transmission of *Salmonella* from reptiles to children in Germany, July 2010 to October 2011. *Euro Surveill*, 18(46), 1-10. <https://doi.org/10.2807/1560-7917.es2013.18.46.20634>

Readel, A. M., Phillips, C. A., & Goldberg, T. L. (2010). Prevalence of *Salmonella* in intestinal mucosal samples from free-ranging red-eared sliders (*Trachemys scripta*) in Illinois. *Herpetological Conservation and Biology*, 5(2), 207-213.

Richards, J. M., Brown, J. D., Kelly, T. R., Fountain, A. L., & Sleeman, J. M. (2004). Absence of detectable *Salmonella* cloacal shedding in free-living reptiles on admission to the wildlife center of Virginia. *Journal of Zoo and Wildlife Medicine*, 35(4), 562-564. <https://doi.org/10.1638/03-070>

Sacristán-Rodríguez, P. A., Pérez-Osorio, L. A. & Méndez-Rodríguez, I. (2014). Prevalencia y susceptibilidad antimicrobiana de *Salmonella* spp. aislada de animales exóticos que conviven con niños. *Medicina & Laboratorio*, 20(3-4), 169-184.

Sacsaquispe, R. & Velásquez, J. (2002). *Manual de procedimientos para la prueba de sensibilidad antimicrobiana por el método de Disco Difusión*. Lima: Ministerio de Salud, Instituto de Salud.



Saelinger, C. A., Lewbart, G. A., Christian, L. S., & Lemons, C. L. (2006). Prevalence of *Salmonella* spp in cloacal, fecal, and gastrointestinal mucosal samples from wild North American turtles. *Journal of the American Veterinary Medical Association*, 229(2), 266-268. <https://doi.org/10.2460/javma.229.2.266>

Sanchez, S., Hofacre, C. L., Lee, M. D., Maurer, J. J., & Doyle, M. P. (2002). Animal sources of salmonellosis in humans. *Journal of the American Veterinary Medical Association*, 221(4), 492-497. <https://doi.org/10.2460/javma.2002.221.492>

Shane, S. M., Gilbert, R., & Harrington, K. S. (1990). *Salmonella* colonization in commercial pet turtles (*Pseudemys scripta elegans*). *Epidemiology and Infection*, 105(2), 307-316. <https://doi.org/10.1017/S0950268800047907>

Sharma, R. N., Matthew-Belmar, V., Nicholas-Thomas, R., Arya, G., Holtlander, B., Hariharan, H., & Amadi, V. A. (2020). Prevalence of *Salmonella* spp. in red-footed tortoises (*Chelonoidis carbonaria*) from Grenada, West Indies. *International Journal of One Health*, 6(1), 12-16. <https://doi.org/10.1292/jvms.67.97>

Shen, L., Shi, H., Wang, R., Liu, D., & Pang, X. (2011). An invasive species red-eared slider (*Trachemys scripta elegans*) carrying *Salmonella* pathogens in Hainan Island. *Molecular Pathogens*, 2(4). <https://doi.org/10.5376/mp.2011.02.004>

Siegmund, R., & Biermann, K. (1988). Common leisure activities of pets and children. *Anthrozoös*, 2(1), 53-57. <https://doi.org/10.2752/089279389787058244>

Sodagari, H. R., Habib, I., Shahabi, M. P., Dybing, N. A., Wang, P., & Bruce, M. (2020). A review of the public health challenges of *Salmonella* and turtles. *Veterinary Sciences*, 7(2), 56. <https://doi.org/10.3390/vetsci7020056>

Tauxe, R. V., Rigau-Pérez, J. G., Wells, J. G., & Blake, P. A. (1985). Turtle-associated salmonellosis in Puerto Rico: Hazards of the global turtle trade. *Journal of the American Medical Association*, *254*(2), 237-239. <https://doi.org/10.1001/jama.1985.03360020069026>

Tomastikova, Z., Barazorda, S. B., Knotek, Z., & Karpiskova, R. (2017). Prevalence and characteristics of *Salmonella* species isolated from captive reptiles in the Czech Republic. *Veterinární Medicína*, *62*(8), 456-469. <https://doi.org/10.17221/44/2017-VETMED>

Ventola, C. L. (2015). The antibiotic resistance crisis: part 1: causes and threats. *Pharmacy and Therapeutics*, *40*(4), 277.

Warwick, C., Lambiris, A. J., Westwood, D., & Steedman, C. (2001). Reptile-related salmonellosis. *Journal of the Royal Society of Medicine*, *94*(3), 124-126. <https://doi.org/10.1177/014107680109400306>

Way, B. M., & Allender, M. C. (2011). Health assessment of wild eastern box turtles (*Terrapene carolina carolina*) in east Tennessee. *Journal of Herpetological Medicine and Surgery*, *21*(4), 107-112. <https://doi.org/10.5818/1529-9651-21.4.107>

Woodward, D. L., Kharkhria, R., & Johnson, W. M. (1997). Human salmonellosis associated with exotic pets. *Journal of Clinical Microbiology*, *35*(11), 1786-2790.

Wong, V. K., Baker, S., Pickard, D. J., Parkhill, J., Page, A. J., Feasey, N. A., ... & Edwards, D. J. (2015). Phylogeographical analysis of the dominant multidrug-resistant H58 clade of *Salmonella* Typhi identifies inter-and intracontinental transmission events. *Nature Genetics*, *47*(6), 632. <https://doi.org/10.1038/ng.3281>

World Health Organization. (2015). WHO estimates of the global burden of foodborne diseases: foodborne disease burden epidemiology reference group 2007-2015. No. 9789241565165. Geneva: Author.

Zhang, J., Kuang, D., Wang, F., Meng, J., Jin, H., Yang, X., ... & Xu, X. (2016). Turtles as a possible reservoir of nontyphoidal Salmonella in Shanghai, China. *Foodborne Pathogens and Disease*, 13(8), 428-433. <https://doi.org/10.1089/fpd.2015.2107>

Zhang, Y., Song, T., Jin, Q., Huang, Y., Tang, X., Sun, X., ... & Bao, W. (2020). Status of an alien turtle in city park waters and its potential threats to local biodiversity: the red-eared slider in Beijing. *Urban Ecosystems*, 23(1), 147-157. <https://doi.org/10.1007/s11252-019-00897-z>

**UNIVERSIDAD DE SAN CARLOS DE GUATEMALA**  
**FACULTAD DE MEDICINA VETERINARIA Y ZOOTECNIA**  
**ESCUELA DE MEDICINA VETERINARIA**

**ANTIBIOTIC-RESISTANT SALMONELLA, ISOLATED FROM CLOACAL  
SWAB SAMPLES FROM TURTLES IN GUATEMALA**

**SALMONELLA RESISTENTE A ANTIBIÓTICOS, AISLADA DE  
MUESTRAS DE HISOPADOS CLOACALES EN TORTUGAS DE  
GUATEMALA**



Br. Joana Gabriela Borrayo Godinez



Phd. Dennis Sigfried Guerra Centeno



Dr. Sc. Carlos Valdez Sandoval



MSc. Manuel Lepe López

IMPRÍMASE

M.A. Gustavo Enrique Taracena Gil

DECANO