

## Original Article

J Prev Med Public Health 2020;53:275-284 • <https://doi.org/10.3961/jpmph.20.095>

pISSN 1975-8375 eISSN 2233-4521



# Educational Intervention Based on the Health Belief Model to Modify Risk Factors of Cardiovascular Disease in Police Officers in Iran: A Quasi-experimental Study

Mohsen Saffari<sup>1,2</sup>, Hormoz Sanaeinasab<sup>1,2</sup>, Hassan Jafarzadeh<sup>2</sup>, Mojtaba Sepandi<sup>1,3</sup>, Keisha-Gaye N. O'Garro<sup>4</sup>, Harold G. Koenig<sup>4,5,6</sup>, Amir H. Pakpour<sup>7,8</sup>

<sup>1</sup>Health Research Center, Life Style Institute, Baqiyatallah University of Medical Sciences Tehran, Iran; <sup>2</sup>Health Education Department, Faculty of Health, Baqiyatallah University of Medical Sciences, Tehran, Iran; <sup>3</sup>Department of Epidemiology and Biostatistics, Baqiyatallah University of Medical Sciences, Tehran, Iran; <sup>4</sup>Department of Psychiatry and Behavioral Sciences, Duke University Medical Center, Durham, NC, USA; <sup>5</sup>Department of Medicine, King Abdulaziz University, Jeddah, Saudi Arabia; <sup>6</sup>School of Public Health, Ningxia Medical University, Yinchuan, China; <sup>7</sup>Social Determinants of Health Research Center, Qazvin University of Medical Sciences, Qazvin, Iran; <sup>8</sup>Department of Nursing, School of Health and Welfare, Jönköping University, Jönköping, Sweden

**Objectives:** Police officers may be at a greater risk for cardiovascular disease (CVD) than the general population due to their high-stress occupation. This study evaluated how an educational program based on the health belief model (HBM) may protect police officers from developing CVD.

**Methods:** In this single-group experimental study, 58 police officers in Iran participated in a 5-week intervention based on HBM principles. Outcomes included changes in scores on an HBM scale, time spent on moderate to vigorous physical activity (International Physical Activity Questionnaire), body mass index (BMI), blood lipid profile, blood glucose, and blood pressure. The intervention consisted of 5 HBM-based educational sessions. Follow-up was conducted at 3 months post-intervention. The paired *t*-test was used to examine differences between baseline and follow-up scores.

**Results:** All aspects of the HBM scale improved between baseline and follow-up ( $p < 0.05$ ), except the cues to action subscale. Self-efficacy and preventive behaviors improved the most. BMI decreased from  $26.7 \pm 2.9$  kg/m<sup>2</sup> at baseline to  $25.8 \pm 2.4$  kg/m<sup>2</sup> at follow-up. All components of the lipid profile, including triglycerides, cholesterol, high-density lipoprotein, and low-density lipoprotein, showed significant improvements post-intervention. Blood glucose and blood pressure also decreased, but not significantly. Nearly 25% of participants who were not physically active at baseline increased their physical activity above or beyond the healthy threshold.

**Conclusions:** A relatively brief educational intervention based on HBM principles led to a significant improvement in CVD risk factors among police officers. Further research is needed to corroborate the effectiveness of this intervention.

**Key words:** Health education, Lifestyle, Cardiovascular disease, Behavior change, Risk factors, Iran

Received: March 28, 2020 Accepted: May 20, 2020

**Corresponding author:** Mohsen Saffari, PhD  
Health Research Center, Life Style Institute, Baqiyatallah University of Medical Sciences, Sheykh-Bahaei Street, Tehran 1435916471, Iran  
**E-mail:** m.saffari@bmsu.ac.ir

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

## INTRODUCTION

Cardiovascular disease (CVD), which has been recognized as a major cause of mortality and morbidity around the world [1], includes disorders such as hypertension, coronary heart disease, atherosclerosis, and stroke that involve pathologies of the heart and blood vessels [2]. Nearly 18 million people died from CVD in 2016, accounting for one-third of all deaths at the