

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

5,800

Open access books available

142,000

International authors and editors

180M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Secretory Defense Response in the Bird's Gastro-Intestinal Tract and Nutritional Strategies to Modulate It

Luis-Miguel Gomez-Osorio, Zhengyu Jiang, Qian Zhang, Hui Yan, Ana-Maria Villegas and Todd Applegate

Abstract

The Gastrointestinal tract (GIT) is a very complex environment which converges a lot of players including nutrients, microorganisms, pathogens, cells, and peptides which determine the type of outcome against threats affecting feed efficiency and body weight gain. Traditionally, GIT is examined as a selective barrier which permit or deny the entrance of molecules, but beyond of that, it is a crucial place to produce important proteins for the host which, at least in part, determine the outcome for a threat such as microorganisms, toxins, anti-nutritional factors, among others. During the non-antibiotic promotants time, there is a necessity to understand how this system works and how we can modulate through nutrition, in part to take advantage of this, and support a better immune response and nutrient absorption in challenged poultry environments. The goal of this chapter is to review the different mechanism of immunity in the GIT emphasizing on secretory defense response and the nutritional strategies including fiber and fatty acids to improve it.

Keywords: antibiotic growth promotants, mucins, M cells, paneth cells, goblet cells, host defense peptides, trefoil factors, IgA, dietary fiber short chain fatty acids, medium chain fatty acids

1. Introduction

There is a tremendous interest in the understanding of immune response against pathogens and toxins on the gastrointestinal tract (GIT) of the birds due to in this specialized system, as it harbors 70 to 80 percent of the avian immune cells and molecules [1]. Additionally, there is an overwhelming interest in finding new alternatives to antibiotic growth promoters (AGP) because of regulations and consumer preference in many countries which these strategies are banned or regulated. The mucosal surface of the GIT is covered by a monolayer of columnar epithelial cells. This epithelium represents a vast surface that is vulnerable to foreign immunogens (i.e. food-borne antigens), microbial pathogens and toxins. By being in contact with a large number of potentially harmful substances and infectious organisms, the mucosal surface must provide a means to not only regulate active and passive absorption of macromolecules but also provide as a general and selective defenses

in part through secretory antibodies and other mucosal defense mechanisms. Consistent with these functions, the epithelial surface of the GIT is lubricated and protected by mucus secretion and by a highly specialized immune system underlying the epithelium which exports immunoglobulins into the intestinal mucosa. Secretory defenses are some of the most important means to protect the intestinal epithelium from enteric pathogens and toxins. Secretory IgA (sIgA) production, Goblet, Paneth, M cells and GALT tissues are the key cells in this defense. The objective of this review is to describe a variety of secretory immune responses against pathogens in GIT and the role of nutrients in immunomodulation.

2. Histology and physiology of the gastro-intestinal tract

Gallus species have villi which decrease in length from 1.5 mm in the duodenum to 0.4–0.6 mm in the ileum and rectum. The number of villi decreases from 1 to 10 days of age, but thereafter remains constant. Genetic selection for growth has altered villi morphology [2]. The villi of broilers are larger than White Leghorns, and show more epithelial cell protrusions from the apical surface of the duodenal villi. However, the villi from both types of chickens consist of a zig-zag arrangement which is thought to slow the passage rate. The intestinal wall contains four layers as including the mucosal, submucosal, muscle tunic, and the serosal layer. The mucosal layer consists of the muscularis mucosa, lamina propria, and epithelium. However, the muscularis mucosa and lamina propria are poorly developed in chickens, possibly because of the absence of a central lacteal. Although Brunner's glands, common to mammals, are absent [3] tubular glands possibly homologous to Brunner's glands, are present in some birds [4]. The epithelium has chief cells, goblet cells, and endocrine cells. The crypts of Lieberkühn are the source of epithelial cells lining the villi. The crypts contain undifferentiated cells, goblet cells, endocrine cells, and lymphocytes. Globular leukocytes and Paneth cells appear near the base of the crypts. The intestine contains extensive innervation from both the sympathetic and parasympathetic nervous system. As described [5], innervation is both cholinergic and adrenergic. Contraction of the rectum appears to be mediated by noncholinergic, non-adrenergic nerves [6, 7].

The mucosa of the GIT is a functional interface between the environment and the internal physiological compartments of the organism. As such, the mucosal and associated cells constitute a dynamic and metabolically active barrier possessing selective permeability [8]. This barrier has multiple functions that involve the digestion, transport and uptake of specific substances and nutrients and exclusion of microorganisms and toxins. The processes of digestion and absorption occur in a micro-environment modified by the intestinal mucosa, its secretions, and the ancillary organs (pancreas, liver). The importance of 'the intestinal barrier' as it relates to gut function and gut health in poultry has been reviewed [9, 10]. Optimal digestive and absorptive functions are essential for growth, development and health of the animal. In addition, the intestine must act as a physical barrier to pathogenic organisms and toxins and play a role in both innate and acquired immunity. The integration of the digestive, absorptive and immune function of the GIT and the genetic regulation of these processes are central to animal production and health.

3. Innate immunity of the GIT

The epithelial cell physical barrier in the GIT represents a vast surface area that is very vulnerable to intraluminal impacts. Continual confrontation by direct

contact with foreign substances, the mucosal system is tightly regulated in order to allow selective entry of macromolecules necessary for mucosal defense [11]. The cells and molecules that comprise the innate immune responses encompass both physical and chemical barrier mechanisms. For example, epithelial cells are tightly connected by multi-protein junctional complexes which regulate passage of solutes while providing an obstacle to luminal microbes and the lamina propria. Mucosal epithelial cells also produce non-specific macro-molecules (such as defensins) with antimicrobial action. Inflammatory and anti-viral responses are produced by specific mucosal cell types, which include: dendritic cells (DC), macrophages, and innate lymphoid cells (ILC). Pattern recognition receptors on these cells regulate many of these responses through interaction with microbial ligands [12].

3.1 Mucus and mucins

The intestine is protected by that substance, which forms a tightly adherent layer along the epithelial surface, followed by a more loosely adherent, partially hydrolyzed layer. It is also part of an integral process, and is secreted, forming and “unstirred” water gel layer covering the epithelial surface. This gelatinous molecular “coat” is subjected to continuous erosion by luminal fluid flow and rapid replenishment from epithelial secretion. The dynamics of mucus gel turnover contributes to a complex milieu where digestive events occur, nutrients approach epithelial cells, microbes build ecological niches, exfoliated enterocytes break down and immunological molecules (defensins, IgA, etc.) carry out surveillance. Consequently, the mucin layer, which encompasses all the of these components, constructs a gel-like biological barrier that shields the underlying tissue compartments, and eventually serves as an important component of the innate arm of the host system in the GIT [11]. In the small intestine the mucus layer is penetrable, but the bacteria are kept away from the epithelium by antibacterial mediators. In the large intestine, the inner mucus layer is impenetrable to bacteria whereas the outer mucus layer is expanded and serves as the habitat for bacteria (esp. mucolytic bacteria) [13]. Serving not only as a lubricant but also a protective barrier, the mucus gel layer(s) in the GIT is the largest area and of critical importance to the body both physiologically and nutritionally. Compromised mucin function is associated with many gastro-enteric disorders and nutritional insufficiencies. Particularly, many functional modulations of the GIT are closely related to expressional, structural, and physiological alterations of mucus and its major components [14].

The protective functions of mucus are attributable to mucus glycoproteins, the major macromolecules present in the mucus gel. Mucus glycoproteins, now widely known as mucins, are defined as a class of high-molecular-weight proteins that are heavily glycosylated with complex oligosaccharide chains [15]. The molecular weight of mucins has been estimated from early studies of ~1000 kDa with attached carbohydrates accounting for 80% of the mass [16].

According to cellular localization and distribution, mucins are broadly classified into secretory and membrane-associated proteins [17]. Structurally mucins are comprised of a linear protein backbone in the center and a large number of carbohydrate chains attached around it. The carbohydrate components, usually hetero-saccharides, are bound covalently to the peptide chains and terminated with sialic acid (sialoglycoproteins) or with both sialic acid and sulphate ester (sialosulphoglycoproteins) or with neutral ends (neutral glycoproteins). These ends determine the extent of negative charges on each mucin molecule [17].

Intestinal secretory mucins are synthesized and secreted by goblet cells, a specialized wine-goblet-shaped epithelial cell lineage dispersed along the intestinal lining. The dimerization and/or polymerization of mucin molecules and the

electrochemical properties of mucopolysaccharides are believed to determine the chemical and biophysical characteristics of mucus along the GIT [18].

Mucins have a key role in avoiding potential damage from microbes. The mechanism by which mucus controls microflora colonization is referred to as part of innate epithelial cells [19]. The role of mucin on microbe colonization is manifested in at least two distinct ways. First, some microbes are mucolytic, including Bacteroidetes, and use mucin glycoproteins and carbohydrates as an energy source and provide physical support for intestinal colonization. Moreover, these bacteria provide substrates for other bacteria in the outer mucus layer by degrading the mucins [20, 21]. Second, mucins are generally “toxic” to the proliferation of certain microbes. Mucus gel inhibits proliferation by entrapping microbes that are starved or killed by antimicrobial peptides, and/or expelled by the luminal flow. Mucus also provides a physicochemical barrier to prevent microbes from direct contact with epithelial cells.

Moreover, the mucus gel provides a matrix for antimicrobial molecules, which are mainly produced by Paneth cells. Direct interactions with mucins can facilitate the diffusion of these antimicrobial molecules [22]. Taken together, mucins have been proposed to play an important role in shaping microbial communities at the intestinal mucosa. Recent studies suggest the correlation between changes in mucin glycosylation profile and deviations of overall microbial community ecology as well as altered abundances of specific microbes [23, 24].

3.2 Trefoil factors

Co-expressed with mucin-secreting cells and in close relation with mucus, trefoil factors (TFF) demonstrate an interesting group of mucus molecules. Trefoil factors were initially discovered in the pig pancreas [25] and further characterization of this family has strikingly observed their abundant expression in the GIT and their efficacy as therapeutics especially for preventing and treating various GIT conditions [26, 27]. They are named as trefoil by their “three-leaf” structure and are a family of small (7-12 kDa in mammals) protease resistant peptides whose common unit is the trefoil motif [25].

It is now clear that TFF participate in the healing of mucosal injury in disease conditions by promoting cell migration over damaged areas (rather than promoting cell division), and inhibiting cell death, and are also believed to be involved in physiological repair of epithelia from daily apical sloughing against frequent luminal insults [25, 28, 29].

TFF have recently been found to participate in immune responses. It was showed that TFF2 deficiency or administration of recombinant TFF2 altered the expression of immune associated genes including defensin genes in Paneth cells [30]. The presence of TFF in immune organs, including spleen, thymus, lymph nodes and bone marrow [31], may suggest possible regulatory role(s) played there. TFF can be a potent mitogen by regulating chemotaxis, stimulating the migration of immune cells. The molecular basis of such may be supported by the recent *in vitro* evidence that recombinant TFF2 activates CXCR4 chemokine receptors and attenuates CXCR4 mediated chemotaxis [32]. This finding also highlights a molecular linkage between TFF and the immune system.

TFF are thought to cooperatively interact with mucins in the lumen to enhance the protective barrier properties of the adherent mucus layer against bacterial and toxic insults [25, 28]. Thim et al. [33] observed significant increase in the viscosity and elasticity of gastric mucin solutions because of TFF2 addition [33]. Increased viscosity could help prevent antigens from approaching the epithelium surface, especially in healing epithelia, which eventually benefits epithelium restitution and

alleviates immune system burden. In this scenario, TFF are predicted to be involved in mucus polymerization.

3.3 Goblet cells

Goblet cells together with absorptive enterocytes, Paneth cells (secreting anti-microbial peptides etc.) and enteroendocrine cells, represent the four principal cell types that are continuously renewed in the epithelium of the small intestine. During intestinal epithelial cell regeneration, pluripotent stem cells that reside at the bottom of the crypt divide to generate multiple cell lineages which migrate from the proliferative crypts to the villus tip [34]. While migrating along the crypt-to-villus axis, goblet cells are terminally differentiated from secretory cell lineage derived from a common Math1-expressing progenitor cell [35]. Goblet cell differentiation is controlled by winged helix transcription factors Foxa1/a2 which can also transactivate Muc2 promoters [36].

It is generally believed that goblet cells producing neutral mucins contain little sialic acid and represent an immature state; while goblet cells containing acidic mucins are more likely resistant to infections because they are normally “upregulated” in response to bacterial infection. In addition to mucins, several other molecules are co-expressed within the intestine such as ingobsin (localized in human and rat goblet cells) with endoproteolytic activity in the presence of both epidermal growth factor and cobalamin-binding protein haptocorrin [37]. TFFs are (specifically TFF3) along with mucins biomarkers of goblet cells.

3.4 M cells

M cells or Microfold cells (because of uneven microvilli) are classified as epithelial cells with large fenestrations in their membranes. These features enhancing the uptake of antigens from the gut lumen [38]. They have a capability for capturing luminal antigens and transporting them across the epithelium (“transcytosis”). They are placed in the gut epithelium called follicle associated epithelium overlying the domes of Peyer’s patches and other lymphoid organs. M cells are not professional antigen-presenting cells because they do not have the ability to process and present antigens to the major histocompatibility complex (MHC) molecules. Instead, they serve as antigen delivery cells, that is, as a functional equivalent to lymphoid nodes because they provide antigens to professional antigen-presenting cells, such as dendritic cells (DCs), macrophages as well as B lymphocytes. Indeed, many pathogens take advantage of their transport efficacy to invade the body [39–41]. M cells subsequently transfer these antigens to underlying DCs enabling the transfer of captured molecules through transcytosis mechanism (which remain to be elucidated) as well as intracellular material through microvesicles to underlying DCs [42]. In conclusion, M cells provide specialized full-service immune surveillance capabilities.

3.5 Paneth cells

Paneth cells are physiologically found at the distal small intestinal crypts of Lieberkühn and contain abundant secretory granules. Their unique histomorphological features implicate special functions in cellular homeostasis as well as in the establishment and configuration of the mucosal barrier as a physical and highly organized immune interface [43]. Previous studies suggesting the existence of Paneth cells in the chicken remained controversial. However, recent research has supported Paneth cells existence in the small intestine of the chicken by electron

microscopy confirming the presence of granulated secretory cells at the base of the crypts in the chicken small intestine. The researchers also confirmed by Western blot the expression of lysozyme protein, which is specifically secreted by the Paneth cells in the small intestine [44]. Paneth cells have the morphological characteristics of a professional secretory cells, including an extensive ER, a Golgi apparatus and an internal secretory granule. The first assumption that Paneth cells had a host-defense function emerged when lysozyme was identified as a product of these cells [45]. After that, it was discovered that Paneth cells secrete antimicrobial peptides (AMP) or host defense peptides (HDPs) which are important host-defense substances in the communication between host and microbiome. One of the most well characterized are β -defensins [46]. In addition to defensins, Paneth cells is able to secrete other AMPs including secretory phospholipase A2, Reg III, angiogenin 4 and cathelicidins [47–49].

3.6 Host defense peptides

HDPs are generally positively charged small peptides with amphipathic properties [50]. These peptides present in the GIT display an important, but often overlooked role in the first line of defense. With the first avian HDPs identified in 1990s [51], the information about avian HDPs has increased considerably in the subsequent decades. Currently, avian β -defensins and cathelicidins are the two major classes identified and extensively studied in chickens [52, 53].

HDPs were initially called antimicrobial peptides (AMPs), because they are characterized by the direct antimicrobial activities against a broad spectrum of numerous pathogens, including gram negative and positive bacteria, fungi, and even certain viruses [54–56]. Generally, the cytoplasmic membrane of pathogenic organisms is a frequent target for HDPs. The amphipathicity and cationic charge of HDPs allow the initial contact with membrane electrostatically, as most bacterial surfaces are hydrophobic and anionic. The peptides then insert into phospholipid bilayers and induce pore formation in membranes by toroidal pore formation, carpet formation and barrel-stave formation, resulting the cytoplasmic leakage and death of pathogens [54, 57–59]. Besides pore formation in membranes, some HDPs can directly penetrate into cells and interfere with intracellular molecules, interrupting cell wall formation, DNA and RNA synthesis, protein translation and post-translational modification [57, 60].

To be specific, chicken AvBD1, –2, and –7 exhibit high efficiency against a large variety of both gram-negative (*E. coli*, *S. enteritidis*, *S. typhimurium*, *C. jejuni*, and *K. pneumoniae*) and gram-positive (*S. aureus*, *B. cereus*, *L. monocytogenes*, *S. haemolyticus*, and *S. saprophytus*) bacteria [51, 61–64]. AvBD1 and –7 also efficiently kill *P. aeruginosa* and *E. cloaca*, while AvDB2 showed reduced efficacy [61, 64]. AvBD4, –5, and –11 protect host from invasion of *S. enteritidis* and *S. typhimurium*, however their antimicrobial activities on other bacteria species remain to be determined [63, 65, 66]. Although AvDB8, –9 and –13 are active against *E. coli*, respectively, they exhibit a minimal activity against several other bacteria [66–69]. Based on studies of different AvBD isoforms, it seems that both structure and catholicity are important for antimicrobial activity but disparity in the preference of gram-negative or positive bacteria.

All four chicken CATHs show antimicrobial capacities in the same order of magnitude against a wide range of gram-negative and positive bacteria, and fungi [70–73]. Similar to AvBDs, the structure and cationic charge are equally important for their antimicrobial activities. The presence of an alpha-helical region in N-terminal and hinge region around the center of the peptide are important for antimicrobial. Removal of N-terminal alpha-helix in CATH2 truncation or

disrupted helix formation in α -helical synthetic peptide leads to the loss of antimicrobial activity [72, 74, 75]. Although deletion of C-terminal α -helix in CATH2 reduces the activity against pathogens, the remaining truncation is still capable to kill bacteria [75]. The truncation of CATH2 with N-terminal α -helix alone shows increased antibacterial activity [76]. The hinge region plays a key role in the insertion of CATH into the bacterial membrane and pore formation [74, 77]. Disruption of the hinge region by point mutation or removal in the center of the CATHs largely decreases the antimicrobial activity [72, 74, 78]. The cationicity of CATH and AvBDs is important for the initial contact with the surface of bacteria. The higher cationic charge in CATH2 and the synthetic analogs results in the better antimicrobial outcomes [72, 75].

In addition to direct antimicrobial activity, the HDPs exhibit the immunomodulatory function, involving inflammation and chemotaxis. Chicken AvBD13 was reported as a direct TLR4 ligand [79], increases production of IFN- γ and IL-12 in mouse monocytes through activation of TLR4-NF κ B axis. Combined with the evidence that AvBD13 increases serum IgG and IgM levels in chicken and induces lymphocytes proliferation in spleen after the administration of the infectious bursal disease vaccine (IBDV) [80], activation of TLR signaling by AvBD13 indicates an immune enhancement rather than a merely pro-inflammatory effect. Moreover, chicken AvBD1 fusion protein expressed by IBDV enhances CD4⁺, CD8⁺, and CD3⁺ T-cell proliferation, increases antibody titers, improves survival rate in *in vivo* experiment [81]. Additionally, HDPs have been shown chemotactic effect. Investigations about immunomodulation by avian AvBDs and CATHs are mainly limited to NF- κ B activation, cytokine production, and direct immune activation. The similar findings in human and mouse studies suggest the conserved function of HDPs among species, providing the guideline for the application and future research in poultry area.

4. Adaptative immunity of the GIT

Unlike the innate immune system which attacks only general threats, adaptative mucosal immune system is triggered by exposure of potentially dangerous pathogens. However, sometimes it overlaps some of their functions. The three most key roles of that system are: the induction of an efficient and appropriate immune response to pathogenic invaders, the tolerance of the commensal microorganisms of the intestine as well as the induction of the tolerance of nutrients and other environmental immunogens. Responses of the systemic immune system can originate from or be modified by the mucosa; this is exemplified by the attenuation of systemic immune responses to a protein that has first been fed orally to the animal (oral tolerance). Thus, the mucosal immune system must maintain the delicate balance between responsiveness to pathogens and tolerance to a vast array of other harmless antigens encountered at mucosal sites. This balance is achieved through the interplay of innate and adaptative (B- and T-lymphocyte) mechanisms [82].

The adaptative immune system in the GIT has features that are distinct from adaptative immune systems in other organs. The major form of adaptative immunity in the gut is humoral immunity directed at microbes in the lumen. This function is mediated mostly by dimeric IgA antibodies that are secreted into the lumen of the gut. Cellular adaptative immunity is carried out by intraepithelial lymphocytes (IEL) in healthy adult bird includes major subsets of NK and T cells bearing the $\gamma\delta$ or $\alpha\beta$ form of the T cell receptor (TCR). In contrast to other tissues, B cells are almost entirely absent from the IEL and the T cells predominantly express the CD8 co-receptor with smaller populations of TCR $\alpha\beta$ ⁺ CD4⁺ and CD4⁺ CD8⁺ cells [83, 84].

Moreover, within the CD8⁺ IEL population the majority express CD8 $\alpha\alpha$ homodimers rather than the CD8 $\alpha\beta$ heterodimer commonly expressed on classical CD8⁺ T cells found at systemic sites [83–85]. The proportions of IEL belonging to each subpopulation differ according to age, genetics and environment (including infection). Numerically, B and T cells are the most common lymphocytes (~90%), the remainder being of the NK cell phenotype (CD3-Bu-1-). In contrast to the IEL population, the T cell population of the lamina propria contains a smaller proportion of $\gamma\delta$ -T cells (~10%); the much larger $\alpha\beta$ -T cell population is dominated by CD4⁺ T cells, with a less prominent CD8⁺ cell population [86].

4.1 Secretory IgA (sIgA) and its transporter, polymeric Ig receptor (pIgR)

The existence of sIgA in the bird has been established for quite some time, but studies are relatively limited compared with mammals. In humans, it is estimated that approximately 70% of the body's IgA-producing plasma cells (differentiated from activated B cells) reside in the lamina propria of intestinal mucosa [87–89]. Although sIgA belongs to adaptive immunity by definition, it plays an important role in the first lines of mucosal defense [87, 90]. There are three modes of defense modulated by sIgA on gut mucosal surfaces: (1) sIgA has been shown to interfere with the early steps of the infection process through directly blocking pathogens and toxins from attaching to the intestinal epithelium [91]; (2) sIgA exerts the protective immunity through immune exclusion, which is the prevention of pathogens and toxins from approaching to epithelium through the stepwise procedures involving antibody-mediated agglutination, entrapment in mucus, and clearance through intestine peristalsis [92, 93]; (3) sIgA exhibits the ability to neutralize intracellular pathogens, viruses, and toxins within intestinal epithelial cells, which requires binding of specific IgA and occupation of antigens by pIgR transportation vesicles, followed by the passage of antigens into the lumen. Notably, the intracellular neutralization of LPS by IgA indicates the potential role in anti-inflammation and deactivation of the proinflammatory pathways in epithelial cells [90, 94, 95].

T-cells generally produce high-affinity IgA antibodies. IgA has the specificity against previous exposure of the GIT by pathogens and more invasive commensal species [89, 96]. Conversely, low-affinity IgA antibodies can also be produced from T-cell-independent (TI) pathways. These low-affinity IgA antibodies act through coating commensal bacteria thereby augmenting the competitive inhibition of pathogens [88, 89, 96]. The production of both high and low-affinity production of IgA provides protective roles during an overt infection with a pathogen as well as during unchallenged/non-pathogenic bacterial exposure.

Presence of microflora in the the GIT may also regulate production of IgA. Studies with germ-free mice [99] and pigs [100, 101] have demonstrated that intestinal IgA and IgA-positive cells in the lamina propria are dramatically reduced versus conventionally reared animals. Further studies have shown that specific microflora (e.g. segmented filamentous bacteria and clostridia) when given to germ-free mice will stimulate the development of IgA-producing cells, while other microflora will have no effect or inhibit this development [97, 98]. Thus, other researchers have reported similar IgA production responses with poultry diets were supplemented with probiotics [99, 100]. Notably, IgA development in the hindgut of the bird early in life coincides with the rapidity of bacterial colonization [101].

Prior to development of IgA by the GIT, the chick is reliant upon maternal antibodies and physical defenses (such as mucins and intestinal turnover). In birds, a small amount of IgA (~ 0.3 mg) is transferred via the embryo imbibing amniotic

fluid prior to internal pipping [102, 103]. The endogenous IgA expression starts to increase after the second week post-hatch [101]. Bar-Shira et al. [103] suggested that the resistance of rapid depletion of maternal IgA may be due to unique uptake by goblet cells, which serves as a reservoir to slowly release maternal IgA.

Circulating IgA is predominantly in a monomeric form, whereas in intestinal secretions it is found in a dimeric configuration both in mammals and birds [96, 104]. IgA secreted by plasma cells accumulates in the lamina propria. To exert its protective effect, pIgR is constitutively expressed by epithelial cells to transport IgA through the epithelia from the lamina propria to intestinal lumen. During the transcytosis, IgA is bound by pIgR on the basolateral surface and transported to the apical surface. At this surface, cleavage of the extracellular portion of pIgR results in release of secretory component (SC) as part of the dimeric IgA, otherwise known as the sIgA complex [105]. In this complex, sIgA is thought to be protected against degradation by proteases and pH fluctuations in the gut [90]. Excess production of pIgR which is not utilized as an IgA chaperone is also secreted as “free SC”, which may have additional bacterial scavenger properties [106]. Once secreted, the N-glycans of SC can then bind to itself, and/or sIgA in the mucin layer thereby bridging these luminal defenses [105].

As one molecule of pIgR is required to bind and transport one dimeric IgA for secretion of sIgA into the intestinal lumen, pIgR's expression regulates sIgA capacity into the GIT [105, 107]. Expression regulation in mammals can be induced by numerous cytokines, including: interferon- γ (IFN- γ), tumor necrosis factor (TNF), interleukin-1 β (IL-1 β), and IL-4. These cytokines act through mediating a transcriptional response through activation of several transcription factor-binding sites in regulatory regions [105, 107–109]. In the chick, increases in IFN- γ , IL-1 β and IL-4 expression in the second week post-hatch [110, 111] may influence subsequent increases in expression of the chicken pIgR gene [111]. Additional bacterial binding to Toll-like receptors have also been shown to increase pIgR expression in epithelial cells [105, 107, 112].

5. Nutrition and secretory immune response

The GIT is an extremely expensive tissue in terms of energy and nutrient needs to maintain and facilitate the full range of barrier and energy/nutrient assimilation functions it displays. Cant et al. [113] estimated that the GIT consumes approximately 20% of dietary energy with a turn-over rate of 50 to 75% per day. However, the GIT is a dynamic organ system whose maintenance needs dramatically changes based on responsive demands. Applegate [114] elucidated some of these adaptive responses, including: changes to peristaltic rate, changes to enterocyte turnover, tight junction regulation, mucin production (quantity and composition), changes to differentiation direction of undifferentiated cells and changes to secretory defenses.

While we often think of presence of microbiota as an additional barrier cost, there is some symbiotic relationships that they convey to the host. For example, the presence of the ceca contributes approximately 3% to dry matter digestibility to the bird [115] in part through 8% of energy derived from microbial fermentation resulting in short-chain fatty acids [116, 117].

Due to limitations of space in this review, we were unable to address all nutrient impacts on the secretory immune defenses of the bird. Notably, recent reviews have published on roles of amino acids on physiological, immunological, and microbiological responses as well as quantification of changes to endogenous amino acid production in the bird [118, 119]; as well as implications of protein indigestibility in

the GIT and implications of microbial fermentation of protein in the hindgut of the animal [120]. Additional impact of microminerals (e.g. zinc, copper, and manganese) and plant bioactive compounds on intestinal functionality have been elucidated [121, 122]. Similarly, recent research has revealed modes of action of specific classes of feed additives that directly or indirectly influence the secretory immune responses of the GIT. For example, probiotic and phytogetic additives have had numerous reviews on these actions [123–125]. Further elucidation of contribution of specific fibrous and fatty acids to the intestinal secretory defenses are further elucidated.

5.1 Dietary fiber and intestinal health

Carbohydrates that are not hydrolyzed by endogenous enzymes in the upper GIT can be fermented by bacteria in the large intestine and ceca are designated as dietary fiber [126]. Dietary fiber (DF) resides in the indigestible portion of plant derived foods that include cell walls, non-starch polysaccharides (NSP), oligosaccharides and lignin [126, 127].

Polysaccharides of NSP include cellulose, pectins, β -glucans, pentosans, heteroxylans and xyloglucan [128]. There are two different types of NSP, soluble and insoluble. Such classification is based on their solubility in water. The ability of soluble NSP to mix with water, producing an increase in the viscosity of the digesta and decreasing the binding of digestive enzymes, negatively affects the digestibility of nutrients [129]. As a result of suboptimal digestion, there is an increase in GIT surface area and secretion of digestive enzymes, creating an increased endogenous energy cost of digestion and affecting bird productivity [130]. NSP from cereal-based diets are associated with low apparently metabolized energy, increased feed conversion rates and increased incidence of wet droppings.

Some previous studies have considered the effects of different cereal NSP based diets on the intestinal microbial immunity. Different types of cereal can modify specific members of the microbiota in the cecum of chickens in two different ways; by altering the viscosity and pH and/or by supplying nutrients to produce the selective growth of specific bacteria [131]. The increase in digesta viscosity with the subsequent reduction in feed passage rate leaves more undigestible feed in the intestine, which represents an ideal substrate for bacterial growth [131]. Chickens fed with a barley-based diet had a higher number of *Clostridium perfringens* in the ileum and ceca. Likewise, it has been reported that the use of wheat in poultry diets may favor the proliferation of pathogenic bacteria like *Escherichia coli*, *Salmonella* and *Campylobacter* [132].

In contrast, insoluble NSP is metabolized into short chain fatty acids (SCFA) including acetate, propionate, butyrate, valerate and isovalerate [116]. Those fermentable metabolites serve as sources of carbon and energy for the commensal microbiota in the lower intestine, specifically, for the bacterial population in the ceca of chickens [116] which provide up to 10% of the energy to the bird. In addition, cecal reverse peristalsis produces translocation of the cecal microbiota affecting energy metabolism and performance [133]. The fermentation metabolites produced by the intestinal bacteria depends on the availability of the substrate, fermentation mechanisms and bacteria specie involved in the process [117].

Dietary fiber has a direct, positive effect on the immune response in numerous species by increasing the abundance of some immune cells, specifically T cells, in the gut-associated lymphoid tissue [134], changing the cytokine secretion profile [135, 136] increasing mucosal immunoglobulins and by acting as a prebiotic substrate for beneficial bacteria [137]. For feed ingredients to be considered prebiotics,

they have to meet the following criteria: resistance to an acidic environment (indigestible), fermentation by intestinal microbiota and selective stimulation of beneficial bacterial populations [138]. Based on this concept, dietary fiber is classified as a prebiotic.

A number of studies have found that fiber-rich prebiotics can enhance immune function including direct production of SCFA [139, 140], augmentation of gut barrier function [141], influence on immune mediated inflammatory responses and restoration of the physiological function of bacterial populations.

In human nutrition, multiple benefits have been attributed to dietary fiber, including maintaining normal bowel structure and function, increasing water retention, blood flow, fluid, and electrolyte uptake in colonic intestinal mucosa [128, 142]. Moreover, fiber intake can reduce the risk of metabolic diseases such as obesity, coronary artery disease, diabetes, constipation, inflammatory bowel disease, colitis and colon cancer [128]. In diets rich in protein, the inclusion of dietary fiber such as arabinoxylan-oligosaccharides (AXOS) can potentially decrease the generation of toxic metabolites originated from proteolytic activity and increase the amount of health-promoting bacterial populations [143].

The addition of dietary fiber has also been widely adopted in swine nutrition in order to maximize the nutrient supply and intestinal health [144, 145]. Dietary fiber can change the physiological features of the digesta, most notably modifying the transit time, and the composition of digesta in terms of solubility, fermentability and water retention. Such changes have a direct impact on intestinal functions, bacteria population and fermentation. The inclusion of high to moderate levels of dietary fiber in pigs, remodel the gut microbiota since certain healthy bacteria species such as Lactobacilli and Bifidobacterium tend to increase. The proliferation of lactic acid producing bacteria decrease the pH of the intestinal lumen, resulting in decreased abundance of other pH sensitive enteropathogenic bacteria like *Escherichia coli*, Salmonella, Shigella and Clostridia [144, 145]. Other effects of dietary fiber have been demonstrated. Changes in the gut morphology, most remarkably inducing increases in crypt depth and altering cell division in growing pigs. This effect has been attributed to the trophic nature of SCFA, specifically butyrate [145]. In contrast, fiber is a feed ingredient poorly utilized in poultry nutrition due to antinutritional effects observed from soluble fiber sources that are mainly associated with increased viscosity of digesta and subsequent impair of nutrient absorption and performance parameters [129]. The effects of fiber are variable and depends on the fiber source, particle size, level of inclusion and chemical composition [146]. A number of studies have found that low levels of insoluble fiber can provide benefits from the point of view of gut health by improving nutrient digestibility [147], gizzard functionality, and resulting in modulation of digesta passage and higher nutrient retention [148, 149]. In the literature, a wide range of other effects of dietary fiber have been demonstrated in laying breeders and broilers chickens. In commercial layers supplemented with high fiber ingredients in the diets, environmental improvements have been demonstrated by reducing ammonia concentrations in manure [150], feather pecking [151], cannibalistic behavior and associated mortality [152].

A number of oligosaccharides including lactulose, inulin, galacto-oligosaccharide and mannan oligosaccharides have been proposed to use as prebiotics in chickens. Those non-digestible carbohydrates are metabolized by fermenting bacteria to produce SCFAs. SCFA are nutritional substrates required for an adequate function of the immune system [139]. When xylo-oligosaccharides were supplemented into a broiler chicken diet, the abundance of butyrate-producing bacteria in the colon and ceca, such as Bifidobacterium and Lactobacillus, significantly increased [153].

Similarly, Zhao et al. reported an increase in *Lactobacillus* counts in excreta when birds were fed with 0.15% inclusion of lactulose [154].

Production of butyrate is considered advantageous to maintain gut health. Butyrate is an important energy source for the enterocytes [140] and is characterized for having immunomodulatory properties. Butyrate can have an anti-inflammatory effect by modulating key inflammatory mediators including the reduction of IFN- γ and NF- κ B and the increase in the number of T reg cells and the expression of IL-10 which suppresses the activity of the immune system [155]. Likewise, inulin supplementation in broiler chickens (0.25-0.5%) induces an anti-inflammatory response by decreasing the gene expression of proinflammatory cytokines such as NF- κ B, LITAF, IL-6, iNOS and enhances the protective barrier function represented by increased expression of epithelial tightness components including MUC2 and claudin-1 [156].

Other major effects have been shown with the supplementation of oligosaccharides, such as the improvement of growth performance [153], the influence on the intestinal morphology reflected in an increase in crypt depth, villus height and villus area [157] and the reduction of pathogenic bacterial colonization. The increase in pathogen resistance due to prebiotic supplementation is associated with the simultaneous elevation of lactic acid producing bacteria and the decrease in the pH of the intestinal lumen, creating an unfavorable environment for pathogenic bacteria and thereby decreasing the colonization. In fact, a meta-analysis study showed a reduction of 0.61 log₁₀ cfu/g cecal *Salmonella* spp. in birds treated with lactose and its associated prebiotic products (lactulose, lacto-sucrose, whey and dried milk) [158].

5.2 Fatty acids and immune response

Short-chain and medium chain fatty acids play an important role on maintaining intestinal gut health and controlling enteric pathogens [159]. Endogenous metabolic pathways, including beta oxidation of fats, leads to the production of short chain fatty acids (SCFA) such as acetate, propionate and butyrate [160]. Long chain fatty acids can be converted into acetate via acetyl-coA or in propionate via propionyl-CoA [160]. SCFA can modulate multiple cellular metabolic activities by the interaction of nuclear cellular (G-protein couple receptors: GRPs), enzymatic receptors (histone deacetylases: HDACs), serving as a substrate for energy for enterocyte and Krebs's cycle and inducing apoptosis of cells [156]. Through these mechanisms, SCFA modulates gene transcription of cells involved in metabolic pathways, inflammation and immune response. In intestinal cells, butyrate and propionate has the ability to inhibit the HDAC activity which decrease the activation of NF κ B transcription factor and subsequently modulating the expression of inflammatory genes [161]. The anti-inflammatory effect of butyrate is produced by preventing the secretion of pro-inflammatory cytokines by macrophages through the NF κ B pathway [156].

Regarding the adaptive immune response, butyrate plays an important role in modifying various lymphocyte function including the inhibition of T-cell proliferation, and reduction of the secretion of pro-inflammatory cytokines such as IL-2, IFN- γ and promoting the production of the main anti-inflammatory cytokine, IL-10 [156, 161].

Due to its anti-inflammatory properties, SCFA has been used as a therapeutic alternative for intestinal diseases [162]. Direct delivery of SCFA by encapsulation allows the supplementation without the need for fermentation, increasing the release in the distal gastrointestinal section [163]. Multiple studies have shown that SCFAs are beneficial as a drinking water supplement and feed additive for the control of *Salmonella*, *Campylobacter* and *Clostridium* [164, 165].

In young chickens, *Salmonella enterica* spp. *enteritidis* cecal colonization significantly decreased when butyric acid was added to the feed [166, 167]. The

addition of SCFA in the drinking water has also been used as an efficient strategy for decreasing the recovery of *Salmonella enterica* spp. *typhimurium* in the crop and in pre-chilled carcasses at the processing plant [168]. The reduction in colonization of *Salmonella* by SCFA is related to the regulation of invasion genes (*hilA*, *invF* and *sipC*) located on the pathogenicity island (Sp1-1) [169]. In addition to the antimicrobial activity, SCFA can contribute to disease resistance by enhancing the expression of host defense peptides including Avian β -defensin 9 (*AvBD9*), *cathelicidin B1*, *AvBD3*, *AvBD4*, *AvBD8*, *AvBD10* and *AvBD14* which consequently reduce bacterial growth [170]. However, the ability of SCFA to control *Salmonella* is highly correlated with acid type and concentration. For example, the feed supplementation of butyric acid in the coated form is more effective in decreasing *Salmonella enterica* spp. *enteritidis* counts than when using the powder form [166]. Previous studies have also investigated the formic-propionic acid combination at 0.5 and 0.68% respectively, with a significant reduction of *Salmonella enterica* spp. *kedougou* [171]. Furthermore, the use of a combination of propionic and formic acid decreased the recovery of *Salmonella enterica* spp. *typhimurium* in the ceca by 3.61 log at 21 days [172]. Similarly, the combination of 1.5% of formic acid and 0.1% of sorbic acid were protective against *Campylobacter jejuni* colonization during infection trials in broiler chickens by reducing *C. jejuni* counts in the crop [173].

Among different classes of fatty acids, medium chain fatty acids (MCFA) have reported to be more inhibitory against *Salmonella* than short chain fatty acids [163]. MCFA are fatty acids composed by 6 to 12 carbons and include caproic, caprylic, capric and lauric fatty acids [174]. The greater antibacterial effect of MCFA is correlated with metabolic differences. Because of its smaller molecular size, MCFA can be absorbed more efficiently and therefore can be utilized more efficiently in the intestinal tract [175]. Indeed, the in-vitro antimicrobial activity of MCFA against *Salmonella* is observed at very low concentrations (between 10 nM- 50 nM) [176, 177]. Furthermore, in-vivo studies have shown reduction in *Salmonella* cecal counts with supplementation of caprylic acid [178, 179]. The supplementation with either 0.7 or 1% of caprylic acid significantly reduced the *Salmonella enterica* spp. *enteritidis* counts in cecal samples of birds fed caprylic acid 7 to 10 days post-challenge in 18 day-old chickens [179]. Another study, showed a reduction in cecal *Salmonella* *Salmonella enterica* spp. *enteritidis* counts in ceca, spleen and liver [178] in 3 and 6-week-old chickens. Similarly, the supplementation of caproic acid in broilers decrease the colonization of *Salmonella* through *hilA* gen suppression [177].

MCFA acid have also been used for controlling *Campylobacter jejuni*. Although, studies have shown inconsistent results, caprylic acid at 0.7 and 1.4% has shown to be effective in reducing *C. jejuni* counts by 3 to 5 log in infected chickens [180].

In conclusion, the application of fatty acids to reduce inflammation and intestinal pathogens is an alternative strategy for poultry nutritionists. Multiple studies support the important role of fatty acids as a modulation of intestinal health. Long chain fatty acids can modulate innate and adaptive immune responses and reduce inflammation produced by systemic diseases. On the other hand, SCFA and MCFA modulate the immune cell function to facilitate the elimination of pathogenic bacteria. Understanding the role of fatty acids in health and disease will increase the effectiveness of these compounds in a wide range of intestinal, metabolic and inflammatory diseases.

6. Conclusions

In summary, secretory defense host response and their players including host defense peptides, sIgA and pIgR among others, constitute the first line of intestinal

immune defense and bridge innate and adaptive immune responses at mucosal surfaces. Understanding the complex function and regulation of these immune components may offer new insights into the nutritional prevention and treatment of infectious and inflammatory diseases that originate at mucosal surfaces. Some studies have been addressed the role of key nutrients modulating this secretory defense system and aiding to the host to counteract the noxious effect of harmful microorganism. Based on that, nutrition would be considered as an important strategy in the reduction of antibiotic growth promotants. However, more studies are needed to understand the effects of nutrients on gut immune response against pathogens.

Author details

Luis-Miguel Gomez-Osorio^{1*}, Zhengyu Jiang², Qian Zhang³, Hui Yan⁴, Ana-Maria Villegas⁵ and Todd Applegate⁵

1 Alura Animal Health and Nutrition, Bogota, Colombia

2 Columbia University, Irving Medical Center, New York, NY, USA

3 DSM (China) Animal Nutrition Research Center Co., Ltd, Bazhou, P.R. China

4 Animal Nutrition Institute, Sichuan Agricultural University, Chengdu, P.R. China

5 Poultry Science Department, University of Georgia, Athens, USA

*Address all correspondence to: lgomez@alura-ahn.com

IntechOpen

© 2021 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

References

- [1] Kagnoff MF. Immunology of the Intestinal Tract. *Gastroenterology*. 1993;(105):1275-1280.
- [2] Yamauchi K-E, Isshiki Y. Scanning electron microscopic observations on the intestinal villi in growing white leghorn and broiler chickens from 1 to 30 days of age. 1991;(32):67-78.
- [3] Calhoun M. *Microscopic Anatomy of the Digestive System*. Iowa: Iowa State College Press, Ames; 1954.
- [4] Ziswiler V, Farner DS. Digestion and digestive system. In: Farner D, King J, editors. *Avian Biology*. London: Academic Press, London; 1972. p. 343-430.
- [5] Bennett T. Peripheral and autonomic nervous systems. In: Farner DS, King JR, editors. *Avian Biology*. New York: Academic Press, New York; 1974. p. 1-77.
- [6] Bartlet AL. Actions of putative transmitters in the chicken vagus nerve/oesophagus and Remak nerve/rectum preparations. *Br J Pharmacol Chemother*. 1974;51:549-558.
- [7] Takewaki T, Ohashi H, Okada T. Non-cholinergic and non-adrenergic mechanism in the contraction and relaxation of the chicken rectum. *Jap J Pharmac*. 1977;27:105-115.
- [8] Baumgart DC, Dignass AU. Intestinal barrier function. *Curr Opin Clin Nutr Metab Care*. 2002;5:685-694.
- [9] Hughes RJ. An integrated approach to understanding gut function and gut health of chickens. *Asia Pac J Clin Nutr*. 2005;14:S27.
- [10] Yegani M, Korver DR. Review Factors Affecting Intestinal Health in Poultry. *Poult Sci* [Internet]. 2008;87(10):2052-2063. Available from: <http://dx.doi.org/10.3382/ps.2008-00091>
- [11] Jiang Z. Gene and expression analysis of secretory mucins and trefoil factor(s) in the intestinal mucosa of chicken. PhD Thesis Purdue University; 2011.
- [12] Johansson ME V, Hansson GC. Immunological aspects of intestinal mucus and mucins. *Nat Rev Immunol* [Internet]. 2016;16(10):639-649. Available from: <http://dx.doi.org/10.1038/nri.2016.88>
- [13] Johansson MEV, Sjövall H, Hansson GC. The gastrointestinal mucus system in health and disease. *Nat Rev Gastroenterol Hepatol* [Internet]. 2013;10(6):352-361. Available from: <http://dx.doi.org/10.1038/nrgastro.2013.35>
- [14] Bansil R, Turner BS. The biology of mucus: Composition, synthesis and organization. Vol. 124, *Advanced Drug Delivery Reviews*. Elsevier B.V.; 2018. p. 3-15.
- [15] Hollingsworth MA, Swanson BJ. Mucins in cancer: Protection and control of the cell surface. *Nat Rev Cancer*. 2004;4(1):45-60.
- [16] Hill HD, Reynolds JA, Hill RL. Purification, composition, molecular weight, and subunit structure of ovine submaxillary mucin. *J Biol Chem*. 1977;252(11):3791-3798.
- [17] Lang T, Hansson GC, Samuelsson T. An inventory of mucin genes in the chicken genome shows that the mucin domain of Muc13 is encoded by multiple exons and that ovomucin is part of a locus of related gel-forming mucins. *BMC Genomics*. 2006;7:1-10.
- [18] Lai SK, Wang YY, Wirtz D, Hanes J. Micro- and macrorheology of mucus. Vol. 61, *Advanced Drug Delivery Reviews*. Elsevier; 2009. p. 86-100.

- [19] Berkes J, Viswanathan VK, Savkovic SD. Intestinal epithelial responses to enteric pathogens: effects on the tight junction barrier, ion transport, and inflammation. *Gut* [Internet]. 2003;52:439-451. Available from: www.gutjnl.com
- [20] Png CW, Lindén SK, Gilshenan KS, Zoetendal EG, McSweeney CS, Sly LI, et al. Mucolytic bacteria with increased prevalence in IBD mucosa augment in vitro utilization of mucin by other bacteria. *Am J Gastroenterol*. 2010;105(11):2420-2428.
- [21] Sonnenburg JL, Xu J, Leip DD, Chen CH, Westover BP, Weatherford J, et al. Glycan foraging in vivo by an intestine-adapted bacterial symbiont. *Science* (80-). 2005;307(5717):1955-9.
- [22] Linden SK, Sutton P, Karlsson NG, Korolik V, McGuckin MA. Mucins in the mucosal barrier to infection. *Mucosal Immunol*. 2008;1(3):183-197.
- [23] Kashyap PC, Marcobal A, Ursell LK, Smits SA, Sonnenburg ED, Costello EK, et al. Genetically dictated change in host mucus carbohydrate landscape exerts a diet-dependent effect on the gut microbiota. *Proc Natl Acad Sci U S A*. 2013;110(42):17059-17064.
- [24] Sommer F, Adam N, Johansson MEV, Xia L, Hansson GC, Bäckhed F. Altered mucus glycosylation in core 1 O-glycan-deficient mice affects microbiota composition and intestinal architecture. *PLoS One*. 2014;9(1).
- [25] Thim L. Trefoil peptides: from structure to function. *Cell Mol Life Sci*. 1997;53.
- [26] Tran CP, Cook GA, Yeomans ND, Thim L, Giraud AS. Trefoil peptide TFF2 (spasmolytic polypeptide) potentially accelerates healing and reduces inflammation in a rat model of colitis. *Gut*. 1999;44(5):636-642.
- [27] Babyatsky MW, DeBeaumont M, Thim L, Podolsky DK. Oral trefoil peptides protect against ethanol- and indomethacin-induced gastric injury in rats. *Gastroenterology*. 1996;110(2):489-497.
- [28] Wong WM, Poulosom R, Wright NA. Trefoil peptides. *Gut*. 1999;44(6):890-895.
- [29] Taupin D, Podolsky DK. Trefoil factors: Initiators of mucosal healing. *Nat Rev Mol Cell Biol*. 2003;4(9):721-732.
- [30] Baus-Loncar M, Kayademir T, Takaishi S, Wang T. Trefoil factor family 2 deficiency and immune response. *Cell Mol Life Sci*. 2005;62(24):2947-2955.
- [31] Cook GA, Familiari M, Thim L, Giraud AS. The trefoil peptides TFF2 and TFF3 are expressed in rat lymphoid tissues and participate in the immune response. *FEBS Lett*. 1999 Jul 30;456(1):155-159.
- [32] Dubeykovskaya Z, Dubeykovskiy A, Solal-Cohen J, Wang TC. Secreted trefoil factor 2 activates the CXCR4 receptor in epithelial and lymphocytic cancer cell lines. *J Biol Chem*. 2009;284(6):3650-3662.
- [33] Thim L, Madsen F, Poulsen SS. Effect of trefoil factors on the viscoelastic properties of mucus gels. *Eur J Clin Invest*. 2002;32(7):519-527.
- [34] Specian RD, Oliver MG. Functional biology of intestinal goblet cells. *Am J Physiol - Cell Physiol*. 1991;260(2 29-2):83-93.
- [35] Yang Q, Bermingham NA, Finegold MJ, Zoghbi HY. Requirement of Math1 for secretory cell lineage commitment in the mouse intestine. *Science* (80-). 2001;294(5549):2155-8.
- [36] Ye DZ, Kaestner KH. Foxa1 and Foxa2 Control the Differentiation of

- Goblet and Enteroendocrine L- and D-Cells in Mice. *Gastroenterology*. 2009 Dec 1;137(6):2052-2062.
- [37] Nexø E, Poulsen SS, Hansen SN, Kirkegaard P, Olsen PS. Characterisation of a novel proteolytic enzyme localised to goblet cells in rat and man. *Gut*. 1984;25(6):656-664.
- [38] Lo DD. Vigilance or Subversion? Constitutive and Inducible M Cells in Mucosal Tissues. *Trends Immunol* [Internet]. 2018;39(3):185-95. Available from: <https://doi.org/10.1016/j.it.2017.09.002>
- [39] Barton ES, Forrest JC, Connolly JL, Chappell JD, Liu Y, Schnell FJ, et al. Junction adhesion molecule is a receptor for reovirus. *Cell*. 2001;104(3):441-451.
- [40] Clark MA, Hirst BH, Jepson MA. M-cell surface $\beta 1$ integrin expression and invasin-mediated targeting of *Yersinia pseudotuberculosis* to mouse Peyer's patch M cells. *Infect Immun*. 1998;66(3):1237-1243.
- [41] Glomski IJ, Piris-Gimenez A, Huerre M, Mock M, Goossens PL. Primary involvement of pharynx and Peyer's patch in inhalational and intestinal anthrax. *PLoS Pathog*. 2007;3(6):0699-0708.
- [42] Sakhony OS, Rossy B, Gusti V, Pham AJ, Vu K, Lo DD. M cell-derived vesicles suggest a unique pathway for trans-epithelial antigen delivery. *Tissue Barriers*. 2015;3(1):1-2.
- [43] Gassler N. Paneth cells in intestinal physiology and pathophysiology. *World J Gastrointest Pathophysiol*. 2017;8(4):150-160.
- [44] Wang L, Li J, Li J, Li RX, Lv CF, Li S, et al. Identification of the Paneth cells in chicken small intestine. *Poult Sci*. 2016;95(7):1631-1635.
- [45] Deckx RJ, Vantrappen GR, Parein MM. Localization of lysozyme activity in a Paneth cell granule fraction. *Biochim Biophys Acta*. 1967;139(1):204-207.
- [46] Ouellette AJ, Miller SI, Henschen AH, Selsted ME. Purification and primary structure of murine cryptdin-1, a Paneth cell defensin. *FEBS Lett*. 1992;304(2-3):146-148.
- [47] Peterson LW, Artis D. Intestinal epithelial cells: Regulators of barrier function and immune homeostasis. *Nat Rev Immunol* [Internet]. 2014;14(3):141-153. Available from: <http://dx.doi.org/10.1038/nri3608>
- [48] Clevers HC, Bevins CL. Paneth cells: Maestros of the small intestinal crypts. *Annu Rev Physiol*. 2013;75:289-311.
- [49] Mowat AM, Agace WW. Regional specialization within the intestinal immune system. *Nat Rev Immunol*. 2014;14(10):667-685.
- [50] Pasupuleti M, Schmidtchen A, Malmsten M. Antimicrobial peptides: Key components of the innate immune system. *Crit Rev Biotechnol*. 2012;32(2):143-171.
- [51] Evans EW, Beach GG, Wunderlich J, Harmon BG. Isolation of antimicrobial peptides from avian heterophils. *J Leukoc Biol*. 1994;56(5):661-665.
- [52] Cuperus T, Coorens M, van Dijk A, Haagsman HP. Avian host defense peptides. *Dev Comp Immunol* [Internet]. 2013;41(3):352-369. Available from: <http://dx.doi.org/10.1016/j.dci.2013.04.019>
- [53] Zhang G, Sunkara LT. Avian antimicrobial host defense peptides: From biology to therapeutic applications. *Pharmaceuticals*. 2014;7(3):220-247.
- [54] Zasloff M. Antimicrobial peptides of multicellular organisms:

My perspective. *Adv Exp Med Biol.* 2002;1117(January):3-6.

[55] Takahashi D, Shukla SK, Prakash O, Zhang G. Structural determinants of host defense peptides for antimicrobial activity and target cell selectivity. *Biochimie* [Internet]. 2010;92(9):1236-1241. Available from: <http://dx.doi.org/10.1016/j.biochi.2010.02.023>

[56] Wang G. Human antimicrobial peptides and proteins. *Pharmaceuticals.* 2014;7(5):545-594.

[57] Brogden KA. Antimicrobial peptides: Pore formers or metabolic inhibitors in bacteria? *Nat Rev Microbiol.* 2005;3(3):238-250.

[58] Jenssen H, Hamill P, Hancock REW. Peptide antimicrobial agents. *Clin Microbiol Rev.* 2006;19(3):491-511.

[59] Pálffy R, Gardlík R, Behuliak M, Kadasi L, Turna J, Celec P. On the physiology and pathophysiology of antimicrobial peptides. *Mol Med.* 2009;15(1-2):51-59.

[60] Nicolas P. Multifunctional host defense peptides: Intracellular-targeting antimicrobial peptides. *FEBS J.* 2009;276(22):6483-6496.

[61] Evans EW, Beach FG, Moore KM, Jackwood MW, Glisson JR, Harmon BG. Antimicrobial activity of chicken and turkey heterophil peptides CHP1, CHP2, THP1, and THP3. *Vet Microbiol.* 1995 Dec 1;47(3-4):295-303.

[62] Harwig SSL, Swiderek KM, Kokryakov VN, Tan L, Lee TD, Panyutich EA, et al. Gallinacins: cysteine-rich antimicrobial peptides of chicken leukocytes. *FEBS Lett.* 1994;342(3):281-285.

[63] Milona P, Townes CL, Bevan RM, Hall J. The chicken host peptides, gallinacins 4, 7, and 9 have antimicrobial activity against *Salmonella* serovars.

Biochem Biophys Res Commun. 2007 Apr 27;356(1):169-174.

[64] Derache C, Labas V, Aucagne V, Meudal H, Landon C, Delmas AF, et al. Primary structure and antibacterial activity of chicken bone marrow-derived β -defensins. *Antimicrob Agents Chemother.* 2009;53(11):4647-4655.

[65] Hervé-Grépinet V, Réhault-Godbert S, Labas V, Magallon T, Derache C, Lavergne M, et al. Purification and characterization of avian β -defensin 11, an antimicrobial peptide of the hen egg. *Antimicrob Agents Chemother.* 2010;54(10):4401-4408.

[66] Lee MO, Jang HJ, Rengaraj D, Yang SY, Han JY, Lamont SJ, et al. Tissue expression and antibacterial activity of host defense peptides in chicken. *BMC Vet Res* [Internet]. 2016;12(1):1-9. Available from: <http://dx.doi.org/10.1186/s12917-016-0866-6>

[67] Higgs R, Lynn DJ, Cahalane S, Alaña I, Hewage CM, James T, et al. Modification of chicken avian β -defensin-8 at positively selected amino acid sites enhances specific antimicrobial activity. *Immunogenetics.* 2007;59(7):573-580.

[68] Higgs R, Lynn DJ, Gaines S, McMahan J, Tierney J, James T, et al. The synthetic form of a novel chicken β -defensin identified in silico is predominantly active against intestinal pathogens. *Immunogenetics.* 2005;57(1-2):90-98.

[69] Van Dijk A, Veldhuizen EJA, Kalkhove SIC, Tjeerdsma-Van Bokhoven JLM, Romijn RA, Haagsman HP. The β -defensingallinacin-6 is expressed in the chicken digestive tract and has antimicrobial activity against food-borne pathogens. *Antimicrob Agents Chemother.* 2007;51(3):912-922.

- [70] Bommineni YR, Dai H, Gong YX, Soulages JL, Fernando SC, DeSilva U, et al. Fowlicidin-3 is an α -helical cationic host defense peptide with potent antibacterial and lipopolysaccharide-neutralizing activities. *FEBS J*. 2007;274(2):418-428.
- [71] Goitsuka R, Chen CLH, Benyon L, Asano Y, Kitamura D, Cooper MD. Chicken cathelicidin-B1, an antimicrobial guardian at the mucosal M cell gateway. *Proc Natl Acad Sci U S A*. 2007;104(38):15063-15068.
- [72] van Dijk A, Molhoek EM, Veldhuizen EJA, Bokhoven JLMT van, Wagendorp E, Bikker F, et al. Identification of chicken cathelicidin-2 core elements involved in antibacterial and immunomodulatory activities. *Mol Immunol*. 2009;46(13):2465-2473.
- [73] Xiao Y, Cai Y, Bommineni YR, Fernando SC, Prakash O, Gilliland SE, et al. Identification and functional characterization of three chicken cathelicidins with potent antimicrobial activity. *J Biol Chem*. 2006;281(5):2858-2867.
- [74] Oh D, Shin SY, Lee S, Kang JH, Kim SD, Ryu PD, et al. Role of the hinge region and the tryptophan residue in the synthetic antimicrobial peptides, cecropin A(1-8)-magainin 2(1-12) and its analogues, on their antibiotic activities and structures. *Biochemistry*. 2000;39(39):11855-11864.
- [75] Xiao Y, Herrera AI, Bommineni YR, Soulages JL, Prakash O, Zhang G. The central kink region of fowlicidin-2, an α -helical host defense peptide, is critically involved in bacterial killing and endotoxin neutralization. *J Innate Immun*. 2009;1(3):268-280.
- [76] Molhoek EM, van Dijk A, Veldhuizen EJA, Dijk-Knijnenburg H, Mars-Groenendijk RH, Boele LCL, et al. Chicken cathelicidin-2-derived peptides with enhanced immunomodulatory and antibacterial activities against biological warfare agents. *Int J Antimicrob Agents*. 2010;36(3):271-274.
- [77] Giangaspero A, Sandri L, Tossi A. Amphipathic α helical antimicrobial peptides.. *Eur J Biochem*. 2001;268(21):5589-5600.
- [78] Shin SY, Kang JH, Jang SY, Kim Y, Kim KL, Hahm KS. Effects of the hinge region of cecropin A(1-8)-magainin 2(1-12), a synthetic antimicrobial peptide, on liposomes, bacterial and tumor cells. *Biochim Biophys Acta - Biomembr*. 2000;1463(2):209-218.
- [79] Yang YR, Jiang YB, Yin QQ, Liang H De, She RP. Chicken intestine defensins activated murine peripheral blood mononuclear cells through the TLR4-NF- κ B pathway. *Vet Immunol Immunopathol*. 2010;133(1):59-65.
- [80] Yang D, Liu Z, Tewary P, Chen Q, de la Rosa G, Oppenheim J. Defensin Participation in Innate and Adaptive Immunity. *Curr Pharm Des*. 2007;13(30):3131-3139.
- [81] Zhang H hua, Yang X mei, Xie Q mei, Ma J yun, Luo Y na, Cao Y chang, et al. The potent adjuvant effects of chicken β -defensin-1 when genetically fused with infectious bursal disease virus VP2 gene. *Vet Immunol Immunopathol [Internet]*. 2010;136(1-2):92-7. Available from: <http://dx.doi.org/10.1016/j.vetimm.2010.02.018>
- [82] Macpherson AJS, Maloy KJ. Adaptive immunity in the gastrointestinal tract. In: *Immunological Aspects of Gastroenterology*. 2001. p. 35-53.
- [83] Vervelde L, Jeurissen SHM. Postnatal development of intra-epithelial leukocytes in the chicken digestive tract: phenotypical characterization in situ. *Cell Tissue Res*. 1993;274(2):295-301.

- [84] Lillehoj HS, Min W, Dalloul RA. Recent progress on the cytokine regulation of intestinal immune responses to *Eimeria*. *Poult Sci* [Internet]. 2004;83(4):611-623. Available from: <http://dx.doi.org/10.1093/ps/83.4.611>
- [85] Imhof BA, Dunon D, Courtois D, Luhtala M, Vainio O. Intestinal CD8 $\alpha\alpha$ and CD8 $\alpha\beta$ Intraepithelial Lymphocytes Are Thymus Derived and Exhibit Subtle Differences in TCR β Repertoires. *J Immunol*. 2000;165(12):6716-6722.
- [86] Egan CE, Maurer KJ, Cohen SB, Mack M, Simpson KW, Denkers EY. Synergy between intraepithelial lymphocytes and lamina propria T cells drives intestinal inflammation during infection. *Mucosal Immunol*. 2011;4(6):658-670.
- [87] Brandtzaeg P, Farstad IN, Johansen FE, Morton HC, Norderhaug IN, Yamanaka T. The B-cell system of human mucosae and exocrine glands. *Immunol Rev*. 1999;171(1):45-87.
- [88] Everett M Lou, Palestrant D, Miller SE, Bollinger RR, Parker W. Immune exclusion and immune inclusion: A new model of host-bacterial interactions in the gut. *Clin Appl Immunol Rev*. 2004;4(5):321-332.
- [89] Slack E, Balmer ML, Fritz JH, Hapfelmeier S. Functional flexibility of intestinal IgA -broadening the fine line. *Front Immunol*. 2012;3(MAY):1-10.
- [90] Johansen FE, Brandtzaeg P. Transcriptional regulation of the mucosal IgA system. *Trends Immunol*. 2004;25(3):150-157.
- [91] Mantis N, Rol N, Corthésy B. Secretory IgA's Complex Roles in Immunity and Mucosal Homeostasis in the Gut. *Mucosal Immunol* [Internet]. 2011;6:603-611. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3624763/pdf/nihms412728.pdf>
- [92] Liévin-Le Moal V, Servin AL. The front line of enteric host defense against unwelcome intrusion of harmful microorganisms: Mucins, antimicrobial peptides, and Microbiota. *Clin Microbiol Rev*. 2006;19(2):315-337.
- [93] Mantis NJ, Forbes SJ. Secretory IgA: Arresting microbial pathogens at epithelial borders. *Immunol Invest*. 2010;39(4-5):383-406.
- [94] Mazanec MB, Nedrud JG, Kaetzel CS, Lamm ME. A three-tiered view of the role of IgA in mucosal defense. *Immunol Today*. 1993;14(9):430-435.
- [95] Fernandez MI, Pedron T, Tournebize R, Olivo-Marin JC, Sansonetti PJ, Phalipon A. Anti-inflammatory role for intracellular dimeric immunoglobulin A by neutralization of lipopolysaccharide in epithelial cells. *Immunity*. 2003;18(6):739-749.
- [96] Brisbin JT, Gong J, Sharif S. Interactions between commensal bacteria and the gut-associated immune system of the chicken. *Anim Heal Res Rev*. 2008;9(1):101-110.
- [97] Honda K, Takeda K. Regulatory mechanisms of immune responses to intestinal bacteria. *Mucosal Immunol*. 2009;2(3):187-196.
- [98] Snel J, Bakker MH, Heidt PJ. Quantification of antigen-specific immunoglobulin A after oral booster immunization with ovalbumin in mice mono-associated with segmented filamentous bacteria or *Clostridium innocuum*. *Immunol Lett*. 1997;58(1):25-28.
- [99] Yurong Y, Ruiping S, Shimin Z, Yibao J. Effect of probiotics on intestinal

mucosal immunity and ultrastructure of cecal tonsils of chickens. *Arch Anim Nutr.* 2005;59(4):237-246.

[100] Haghghi HR, Gong J, Gyles CL, Hayes MA, Zhou H, Sanei B, et al. Probiotics stimulate production of natural antibodies in chickens. *Clin Vaccine Immunol.* 2006;13(9):975-980.

[101] Zhang Q, Eicher SD, Applegate TJ. Development of intestinal mucin 2, IgA, and polymeric Ig receptor expressions in broiler chickens and Pekin ducks. *Poult Sci [Internet].* 2015;94(2):172-180. Available from: <http://dx.doi.org/10.3382/ps/peu064>

[102] Friedman A, Elad O, Cohen I, Bar Shira E. The gut associated lymphoid system in the post-hatch chick: Dynamics of maternal IgA. *Isr J Vet Med.* 2012;67(2):75-81.

[103] Bar-Shira E, Cohen I, Elad O, Friedman A. Role of goblet cells and mucin layer in protecting maternal IgA in precocious birds. *Dev Comp Immunol [Internet].* 2014;44(1):186-194. Available from: <http://dx.doi.org/10.1016/j.dci.2013.12.010>

[104] Snoeck V, Peters IR, Cox E. The IgA system: a comparison of structure and function in different species. *Vet Res.* 2006;37:455-467.

[105] Kaetzel CS. The polymeric immunoglobulin receptor: Bridging innate and adaptive immune responses at mucosal surfaces. *Immunol Rev.* 2005;206:83-99.

[106] Phalipon A, Corthésy B. Novel functions of the polymeric Ig receptor: Well beyond transport of immunoglobulins. *Trends Immunol.* 2003;24(2):55-58.

[107] Johansen FE, Kaetzel CS. Regulation of the polymeric immunoglobulin receptor and IgA transport: New advances in

environmental factors that stimulate pIgR expression and its role in mucosal immunity. *Mucosal Immunol.* 2011;4(6):598-602.

[108] Blanch VJ, Piskurich JF, Kaetzel CS. Cutting edge: Coordinate regulation of IFN regulatory factor-1 and the polymeric Ig receptor by proinflammatory cytokines. *J Immunol.* 1999;162(3):1232-1235.

[109] Schjerven H, Brandtzaeg P, Johansen F-E. Mechanism of IL-4-Mediated Up-Regulation of the Polymeric Ig Receptor: Role of STAT6 in Cell Type-Specific Delayed Transcriptional Response. *J Immunol.* 2000;165(7):3898-3906.

[110] Bar-Shira E, Sklan D, Friedman A. Establishment of immune competence in the avian GALT during the immediate post-hatch period. *Dev Comp Immunol.* 2003;27(2):147-157.

[111] Lammers A, Wieland WH, Kruijt L, Jansma A, Straetemans T, Schots A, et al. Successive immunoglobulin and cytokine expression in the small intestine of juvenile chicken. *Dev Comp Immunol [Internet].* 2010;34(12):1254-1262. Available from: <http://dx.doi.org/10.1016/j.dci.2010.07.001>

[112] Bruno MEC, Frantz AL, Rogier EW, Johansen FE, Kaetzel CS. Regulation of the polymeric immunoglobulin receptor by the classical and alternative NF- κ B pathways in intestinal epithelial cells. *Mucosal Immunol.* 2011;4(4):468-478.

[113] Cant JP, McBride BW, Croom WJ. The Regulation of Intestinal Metabolism and Its Impact on Whole Animal Energetics. *J Anim Sci.* 1996;74(10):2541-2553.

[114] Applegate T. Influence of Phytogenics on the Immunity of Livestock and Poultry. In: Steiner T, editor. *Phytogenics in Animal Nutrition.* Nottingham, United Kingdom:

Nottingham University Press; 2009. p. 39-59.

[115] Thornburn CC, Willcox JS. The caeca of the domestic fowl and digestion of the crude fibre complex: II. Experiments in vivo with fistulated birds, and the artificial and isolated caecum in vitro. *Br Poult Sci*. 1965;6(1):33-43.

[116] Józefiak D, Rutkowski A, Martin SA. Carbohydrate fermentation in the avian ceca: A review. *Anim Feed Sci Technol*. 2004;113(1-4):1-15.

[117] Rinttilä T, Apajalahti J. Intestinal microbiota and metabolites- Implications for broiler chicken health and performance. *J Appl Poult Res [Internet]*. 2013;22(3):647-658. Available from: <http://dx.doi.org/10.3382/japr.2013-00742>

[118] Bortoluzzi C, Rochell SJ, Applegate TJ. Threonine, arginine, and glutamine: Influences on intestinal physiology, immunology, and microbiology in broilers. *Poult Sci [Internet]*. 2018;97(3):937-945. Available from: <http://dx.doi.org/10.3382/ps/pex394>

[119] Bortoluzzi C, Fernandes JIM, Doranalli K, Applegate TJ. Effects of dietary amino acids in ameliorating intestinal function during enteric challenges in broiler chickens. *Anim Feed Sci Technol [Internet]*. 2020;262(September):114383. Available from: <https://doi.org/10.1016/j.anifeedsci.2019.114383>

[120] Gilbert MS, Ijssennagger N, Kies AK, van Mil SWC. Protein fermentation in the gut; implications for intestinal dysfunction in humans, pigs, and poultry. *Am J Physiol - Gastrointest Liver Physiol*. 2018;315(2):G159-G170.

[121] Bortoluzzi C, Vieira BS, Applegate TJ. Influence of Dietary Zinc, Copper, and Manganese on the

Intestinal Health of Broilers Under *Eimeria* Challenge. *Front Vet Sci*. 2020;7(January):1-7.

[122] Patra AK, Amasheh S, Aschenbach JR. Modulation of Gastrointestinal Barrier and Nutrient Transport Function in Farm Animals by Natural Plant Bioactive Compounds – A Comprehensive Review. 2018;8398.

[123] Patra AK, Amasheh S, Aschenbach JR. Modulation of gastrointestinal barrier and nutrient transport function in farm animals by natural plant bioactive compounds—A comprehensive review. *Crit Rev Food Sci Nutr [Internet]*. 2019;59(20):3237-66. Available from: <https://doi.org/10.1080/10408398.2018.1486284>

[124] Applegate TJ, Troche C. Influence of probiotics on intestinal structure and barrier functionality of poultry. In: Abdelrahman WHA, Mohlml M, editors. *Probiotics in Poultry Production*. 5MEnterpri ed. Sheffield, England; 2014. p. 51-69.

[125] Patra AK. Influence of Plant Bioactive Compounds on Intestinal Epithelial Barrier in Poultry. *Mini-Reviews Med Chem*. 2020;20: 566-577.

[126] Montagne L, Pluske JR, Hampson DJ. A review of interactions between dietary fibre and the intestinal mucosa, and their consequences on digestive health in young non-ruminant animals. *Anim Feed Sci Technol*. 2003;108(1-4):95-117.

[127] Bach Knudsen KE. The nutritional significance of “dietary fibre” analysis. *Anim Feed Sci Technol*. 2001;90(1-2):3-20.

[128] Kumar V, Sinha AK, Makkar HPS, de Boeck G, Becker K. Dietary Roles of Non-Starch Polysachharides in Human Nutrition: A Review. *Crit Rev Food Sci Nutr*. 2012;52(10):899-935.

- [129] Marquardt RR, Brenes A, Zhang Z, Boros D. Use of enzymes to improve nutrient availability in poultry feedstuffs. *Anim Feed Sci Technol.* 1996;60(3-4):321-330.
- [130] Perry GC. Effects of Non-starch polysaccharide on Avian Gastrointestinal Function. In: *Avian Gut Function in Health and Disease*. Oxon: CABI; 2006. p. 159-170.
- [131] Shakouri MD, Iji PA, Mikkelsen LL, Cowieson AJ. Intestinal function and gut microflora of broiler chickens as influenced by cereal grains and microbial enzyme supplementation. *J Anim Physiol Anim Nutr (Berl)*. 2009;93(5):647-658.
- [132] Korver DR. Overview of the immune dynamics of the digestive system. *J Appl Poult Res [Internet]*. 2006;15(1):123-135. Available from: <http://dx.doi.org/10.1093/japr/15.1.123>
- [133] Crisol-Martínez E, Stanley D, Geier MS, Hughes RJ, Moore RJ. Sorghum and wheat differentially affect caecal microbiota and associated performance characteristics of meat chickens. *PeerJ*. 2017;2017(3):1-20.
- [134] Field CJ, McBurney MI, Massimino S, Hayek MG, Sunvold GD. The fermentable fiber content of the diet alters the function and composition of canine gut associated lymphoid tissue. *Vet Immunol Immunopathol.* 1999;72(3-4):325-341.
- [135] Cox CM, Stuard LH, Kim S, McElroy AP, Bedford MR, Dalloul RA. Performance and immune responses to dietary β -glucan in broiler chicks. *Poult Sci [Internet]*. 2010;89(9):1924-1933. Available from: <http://dx.doi.org/10.3382/ps.2010-00865>
- [136] Cox CM, Sumners LH, Kim S, McElroy AP, Bedford MR, Dalloul RA. Immune responses to dietary β -glucan in broiler chicks during an *Eimeria* challenge. *Poult Sci [Internet]*. 2010;89(12):2597-2607. Available from: <http://dx.doi.org/10.3382/ps.2010-00987>
- [137] Schley PD, Field CJ. The immune-enhancing effects of dietary fibres and prebiotics. *Br J Nutr.* 2002;87(S2):S221-S230.
- [138] Roberfroid M. Prebiotics: The concept revisited. *J Nutr.* 2007;137(3).
- [139] Koh A, De Vadder F, Kovatcheva-Datchary P, Bäckhed F. From dietary fiber to host physiology: Short-chain fatty acids as key bacterial metabolites. *Cell.* 2016;165(6):1332-1345.
- [140] Liao X, Shao Y, Sun G, Yang Y, Zhang L, Guo Y, et al. The relationship among gut microbiota, short-chain fatty acids, and intestinal morphology of growing and healthy broilers. *Poult Sci [Internet]*. 2020;99(11):5883-95. Available from: <https://doi.org/10.1016/j.psj.2020.08.033>
- [141] Song J, Li Q, Li P, Liu RR, Cui H, Zheng M, et al. The effects of inulin on the mucosal morphology and immune status of specific pathogen-free chickens. *Poult Sci [Internet]*. 2018;97(11):3938-3946. Available from: <http://dx.doi.org/10.3382/ps/pey260>
- [142] Gibson GR. Fibre and effects on probiotics (the prebiotic concept). *Clin Nutr Suppl.* 2004;1(2):25-31.
- [143] Sanchez JI, Marzorati M, Grootaert C, Baran M, Van Craeyveld V, Courtin CM, et al. Arabinoxylan-oligosaccharides (AXOS) affect the protein/carbohydrate fermentation balance and microbial population dynamics of the Simulator of Human Intestinal Microbial Ecosystem. *Microb Biotechnol.* 2009;2(1):101-113.
- [144] Jha R, Berrocoso JFD. Dietary fiber and protein fermentation in the

intestine of swine and their interactive effects on gut health and on the environment: A review. *Anim Feed Sci Technol* [Internet]. 2016;212:18-26. Available from: <http://dx.doi.org/10.1016/j.anifeedsci.2015.12.002>

[145] Bach Knudsen KE, Hedemann MS, Lærke HN. The role of carbohydrates in intestinal health of pigs. *Anim Feed Sci Technol* [Internet]. 2012;173(1-2):41-53. Available from: <http://dx.doi.org/10.1016/j.anifeedsci.2011.12.020>

[146] Iqbal A, Qudoos A, Çetingül IS, Rizwan S, Shah A. Importance of dietary fiber in poultry nutrition Importance of Dietary Fiber in Poultry Nutrition. *J Anim Sci Prod*. 2019;(January):21-9.

[147] Jiménez-Moreno E, González-Alvarado JM, González-Serrano A, Lázaro R, Mateos GG. Effect of dietary fiber and fat on performance and digestive traits of broilers from one to twenty-one days of age. *Poult Sci*. 2009;88(12):2562-2574.

[148] Jiménez-Moreno E, Chamorro S, Frikha M, Safaa HM, Lázaro R, Mateos GG. Effects of increasing levels of pea hulls in the diet on productive performance, development of the gastrointestinal tract, and nutrient retention of broilers from one to eighteen days of age. *Anim Feed Sci Technol*. 2011;168(1-2):100-112.

[149] Sacranie A, Svihus B, Denstadli V, Moen B, Iji PA, Choct M. The effect of insoluble fiber and intermittent feeding on gizzard development, gut motility, and performance of broiler chickens. *Poult Sci* [Internet]. 2012;91(3):693-700. Available from: <http://dx.doi.org/10.3382/ps.2011-01790>

[150] Roberts SA, Xin H, Kerr BJ, Russell JR, Bregendahl K. Effects of dietary fiber and reduced crude protein on ammonia emission from

laying-hen manure. *Poult Sci* [Internet]. 2007;86(8):1625-1632. Available from: <http://dx.doi.org/10.1093/ps/86.8.1625>

[151] van Krimpen MM, Kwakkel RP, van der Peet-Schwering CMC, den Hartog LA, Verstegen MWA. Effects of nutrient dilution and nonstarch polysaccharide concentration in rearing and laying diets on eating behavior and feather damage of rearing and laying hens. *Poult Sci* [Internet]. 2009;88(4):759-773. Available from: <http://dx.doi.org/10.3382/ps.2008-00194>

[152] Hartini S, Choct M, Hinch G, Kocher A, Nolan J V. Effects of light intensity during rearing and beak trimming and dietary fiber sources on mortality, egg production, and performance of ISA Brown laying hens. *J Appl Poult Res* [Internet]. 2002;11(1):104-10. Available from: <http://dx.doi.org/10.1093/japr/11.1.104>

[153] De Maesschalck C, Eeckhaut V, Maertens L, De Lange L, Marchal L, Nezer C, et al. Effects of Xylo-oligosaccharides on broiler chicken performance and microbiota. *Appl Environ Microbiol*. 2015;81(17):5880-5888.

[154] Zhao PY, Li HL, Mohammadi M, Kim IH. Effect of dietary lactulose supplementation on growth performance, nutrient digestibility, meat quality, relative organ weight, and excreta microflora in broilers. *Poult Sci*. 2016;95(1):84-89.

[155] Meijer K, De Vos P, Priebe MG. Butyrate and other short-chain fatty acids as modulators of immunity: What relevance for health? *Curr Opin Clin Nutr Metab Care*. 2010;13(6):715-721.

[156] Li B, Leblois J, Taminiau B, Schroyen M, Beckers Y, Bindelle J, et al. The effect of inulin and wheat bran on intestinal health and microbiota in the early life of broiler chickens.

Poult Sci [Internet]. 2018;97(9):3156-3165. Available from: <http://dx.doi.org/10.3382/ps/pey195>

[157] Hutsko SL, Meizlisch K, Wick M, Lilburn MS. Early intestinal development and mucin transcription in the young poult with probiotic and mannan oligosaccharide prebiotic supplementation. Poult Sci [Internet]. 2016;95(5):1173-1178. Available from: <http://dx.doi.org/10.3382/ps/pew019>

[158] Totton SC, Farrar AM, Wilkins W, Bucher O, Waddell LA, Wilhelm BJ, et al. The effectiveness of selected feed and water additives for reducing *Salmonella* spp. of public health importance in broiler chickens: A systematic review, meta-analysis, and meta-regression approach. Prev Vet Med. 2012;106(3-4):197-213.

[159] Kollanoor JA, Venkitanarayanan K. Use of fatty acids in controlling enteric pathogens. In: Cherian G, Poureslami R, editors. Fats and fatty acids in poultry nutrition and health. Nottingham University Press; 2013.

[160] Gill PA, van Zelm MC, Muir JG, Gibson PR. Review article: short chain fatty acids as potential therapeutic agents in human gastrointestinal and inflammatory disorders. Aliment Pharmacol Ther. 2018;48(1):15-34.

[161] Vinolo MAR, Rodrigues HG, Nachbar RT, Curi R. Regulation of Inflammation by Short Chain Fatty Acids. Nutrients. 2011;(3):858-876.

[162] Guilloteau P, Martin L, Eeckhaut V, Ducatelle R, Zabielski R, Van Immerseel F. From the gut to the peripheral tissues: The multiple effects of butyrate. Nutr Res Rev. 2010;23(2):366-384.

[163] Van Immerseel F, Russell JB, Flythe MD, Gantois I, Timbermont L, Pasmans F, et al. The use of organic

acids to combat *Salmonella* in poultry: A mechanistic explanation of the efficacy. Avian Pathol. 2006;35(3):182-188.

[164] Huyghebaert G, Ducatelle R, Immerseel F Van. An update on alternatives to antimicrobial growth promoters for broilers. Vet J [Internet]. 2011;187(2):182-188. Available from: <http://dx.doi.org/10.1016/j.tvjl.2010.03.003>

[165] Song B, Li H, Wu Y, Zhen W, Wang Z, Xia Z, et al. Effect of microencapsulated sodium butyrate dietary supplementation on growth performance and intestinal barrier function of broiler chickens infected with necrotic enteritis. Anim Feed Sci Technol [Internet]. 2017;232:6-15. Available from: <http://dx.doi.org/10.1016/j.anifeedsci.2017.07.009>

[166] Van Immerseel F, Boyen F, Gantois I, Timbermont L, Bohez L, Pasmans F, et al. Supplementation of coated butyric acid in the feed reduces colonization and shedding of *Salmonella* in poultry. Poult Sci [Internet]. 2005;84(12):1851-1856. Available from: <http://dx.doi.org/10.1093/ps/84.12.1851>

[167] Van Immerseel F, Fievez V, De Buck J, Pasmans F, Martel A, Haesebrouck F, et al. Microencapsulated short-chain fatty acids in feed modify colonization and invasion early after infection with *Salmonella enteritidis* in young chickens. Poult Sci [Internet]. 2004;83(1):69-74. Available from: <http://dx.doi.org/10.1093/ps/83.1.69>

[168] Byrd JA, Hargis BM, Caldwell DJ, Bailey RH, Herron KL, McReynolds JL, et al. Effect of lactic acid administration in the drinking water during preslaughter feed withdrawal on *Salmonella* and *Campylobacter* contamination of broilers. Poult Sci [Internet]. 2001;80(3):278-283. Available from: <http://dx.doi.org/10.1093/ps/80.3.278>

- [169] Lawhon SD, Maurer R, Suyemoto M, Altier C. Intestinal short-chain fatty acids alter Salmonella typhimurium invasion gene expression and virulence through BarA/SirA. *Mol Microbiol.* 2002;46(5):1451-1464.
- [170] Sunkara LT, Jiang W, Zhang G. Modulation of Antimicrobial Host Defense Peptide Gene Expression by Free Fatty Acids. *PLoS One.* 2012;7(11).
- [171] Hinton M, Linton AH. Control of salmonella infections in broiler chickens by the acid treatment of their feed. *Vet Rec.* 1988;123:416-421.
- [172] McHan F, Shotts EB. Effect of feeding selected short-chain fatty acids on the in vivo attachment of Salmonella typhimurium in chick ceca. *Avian Dis.* 1992;36:139-142.
- [173] Skånseng B, Kaldhusdal M, Moen B, Gjevre AG, Johannessen GS, Sekelja M, et al. Prevention of intestinal *Campylobacter jejuni* colonization in broilers by combinations of in-feed organic acids. *J Appl Microbiol.* 2010;109(4):1265-1273.
- [174] Baltić B, Starčević M, Dordević J, Mrdović B, Marković R. Importance of medium chain fatty acids in animal nutrition. *IOP Conf Ser Earth Environ Sci.* 2017;85:012048.
- [175] Vandeplas S, Dubois Dauphin R, Beckers Y, Thonart P, Théwis A. Salmonella in chicken: Current and developing strategies to reduce contamination at farm level. *J Food Prot.* 2010;73(4):774-785.
- [176] Vasudevan P, Marek P, Nair MKM, Annamalai T, Darre M, Khan M, et al. In vitro inactivation of Salmonella enteritidis in autoclaved chicken cecal contents by caprylic acid. *J Appl Poult Res [Internet].* 2005;14(1):122-125. Available from: <http://dx.doi.org/10.1093/japr/14.1.122>
- [177] Van Immerseel F, De Buck J, Boyen F, Bohez L, Pasmans F, Volf J, et al. Medium-chain fatty acids decrease colonization and invasion through hila suppression shortly after infection of chickens with Salmonella enterica serovar enteritidis. *Appl Environ Microbiol.* 2004;70(6):3582-3587.
- [178] Kollanoor-Johny A, Mattson T, Baskaran SA, Amalaradjou MAR, Hoagland TA, Darre MJ, et al. Caprylic acid reduces salmonella Enteritidis populations in various segments of digestive tract and internal organs of 3-and 6-week-old broiler chickens, therapeutically. *Poult Sci [Internet].* 2012;91(7):1686-1694. Available from: <http://dx.doi.org/10.3382/ps.2011-01716>
- [179] Johny AK, Baskaran SA, Charles AS, Amalaradjou MAR, Darre MJ, Khan MI, et al. Prophylactic supplementation of caprylic acid in feed reduces salmonella enteritidis colonization in commercial broiler chickst. *J Food Prot.* 2009;72(4):722-727.
- [180] Solis de los Santos F, Hume M, Venkitanarayanan K, Donoghue AM, Hanning I, Slavik MF, et al. Caprylic acid reduces enteric *Campylobacter* colonization in market-aged broiler chickens but does not appear to alter cecal microbial populations. *J Food Prot.* 2010;73(2):251-257.