

1 **Understanding country-specific determinants of stillbirth using household surveys – the case of**
2 **Afghanistan**

3

4 Aliko Christou^{1*}, Michael J Dibley¹, Mohammad Hafiz Rasooly², Adela Mubasher³, Sayed Murtaza
5 Sadat Hofiani², Mohammad Khakerah Rashidi⁴, Patrick J. Kelly¹, Camille Raynes-Greenow¹

6

7 ¹Sydney School of Public Health, Faculty of Medicine and Health, The University of Sydney, Sydney,
8 NSW Australia

9 ²Afghanistan National Public Health Institute, Ministry of Public Health, Kabul, Afghanistan

10 ³World Health Organisation, Kabul, Afghanistan

11 ⁴Management Sciences for Health, Kabul, Afghanistan

12

13

14 *Corresponding author

15 Aliko Christou

16 Sydney School of Public Health

17 Room 121C, Edward Ford Building A27

18 The University of Sydney

19 Sydney, NSW 2006

20

21 Telephone: +61293516816

22 Mobile: +61435809757

23 Email: alichristou@gmail.com

24

25

26

27

28

29 **Abstract**

30

31 **Background:** Stillbirth rates in Afghanistan have declined little in the past decade with no data
32 available on key risk factors. Healthcare utilisation and maternal complications are important factors
33 influencing pregnancy outcomes but rarely captured for stillbirth in national surveys from low- and
34 middle-income countries. The 2010 Afghanistan Mortality Survey (AMS) is one of few surveys with
35 this information.

36

37 **Methods:** We used data from the 2010 AMS that included a full pregnancy history and verbal
38 autopsy. Our sample included the most recent live birth or stillbirth of 13,834 women aged 12-49
39 years in the three years preceding the survey. Multivariable Poisson regression was used to identify
40 socio-demographic, maternal, and healthcare utilisation risk factors for stillbirth.

41

42 **Results:** The risk of stillbirth was increased among women in the Central highlands (aRR:3.01,
43 95%CI:1.35, 6.70) and of Nuristani ethnicity (aRR:9.15, 95%CI: 2.95, 28.74). Women that didn't
44 receive antenatal care had three times increased risk of stillbirth (aRR:3.03, 95% CI:1.73, 5.30), while
45 high-quality antenatal care was important for reducing the risk of intrapartum stillbirth. Bleeding,
46 infection, headache, and reduced fetal movements were antenatal complications strongly associated
47 with stillbirth. Reduced fetal movements in the delivery period increased stillbirth risk by almost
48 seven (aRR:6.82, 95%CI:4.20,11.10). Facility births had a higher risk of stillbirths overall (aRR:1.55,
49 95%CI:1.12, 2.16), but not for intrapartum stillbirths.

50

51 **Conclusions:** Targeted interventions are needed to improve access and utilisation of services for
52 high-risk groups. Early detection of complications through improved quality of antenatal and obstetric
53 care is imperative. We demonstrate the potential of household surveys to provide country-specific
54 evidence on stillbirth risk factors for LMICs where data is lacking.

55

56 **Key words:** stillbirth, fetal death, perinatal death, Afghanistan, low- and middle-income country, risk
57 factor, household surveys, Demographic and Health Survey

58

59

60

61

62

63

64

65

66

67

68

69

70

71

72

73

74

75

76

77

78

79

80

81

82

83

84 **Background**

85

86 A major challenge for stillbirth prevention in low- and middle-income countries (LMIC),
87 where the largest burden lies, is the lack of adequate data to identify and quantify major risk factors at
88 the national level (1). Existing studies have been predominantly with women who have had contact
89 with the formal health care system (2), and while prospective, population-based studies such as those
90 from demographic surveillance sites in LMICs are increasing (3, 4), there are no national-level data
91 for many countries (1). This lack of data on country-specific risk factors makes it challenging to direct
92 attention to stillbirth at a national level and for countries to prioritise programmatic and policy areas
93 for action to reduce stillbirths.

94

95 In 2009, Afghanistan was among the top ten nations accounting for almost two-thirds of the
96 global stillbirth burden, and by 2016 little improvement was observed (5, 6). The annual reduction in
97 stillbirths between 2000 and 2015 in Afghanistan was only 1.9% (6). The stillbirth rate remains high
98 at 27 per 1000 births – six times that of high-income settings, yet there are no published studies to
99 understand stillbirths in this context. Stillbirths have not been a public health priority in Afghanistan
100 partly because of the absence of evidence on the major factors contributing to these deaths. The UN’s
101 2016 Global Strategy for Women’s, Children’s and Adolescent Health now includes reduction in the
102 stillbirth rate as a core indicator, and the 2014 Every Newborn Action Plan set the first-ever targets to
103 reduce stillbirths to 12 per 1000 births by 2030 which was endorsed by 190 countries, including
104 Afghanistan (7). It is therefore, both timely and crucial to investigate stillbirths in this high-burden
105 country.

106

107 The 2010 Afghanistan Mortality Survey (AMS) was a modified, special Demographic and
108 Health Survey (DHS) and one of a few nationally-representative surveys conducted in a LMIC in the
109 last ten years that collected health service utilisation data for stillbirths and also included a verbal
110 autopsy (8). The country’s unique and diverse socio-cultural, linguistic, and geographic characteristics
111 in addition to the current complex humanitarian situation, makes the need for context-specific data

112 imperative (Box 1). The objective of this study was to identify key maternal, obstetric and health-care
113 utilisation factors associated with stillbirth in Afghanistan, and to demonstrate the potential of a
114 modified DHS survey to provide country-specific evidence on risk factors for stillbirth if applied in
115 other LMICs.

116

117 **[Insert Box 1: Afghanistan country context and health situation]**

118

119

120

121

122

123

124

125

126

127

128

129

130

131

132

133

134

135

136

137

138

139

140

141

142

143

144

145

146

147 **Methods**

148

149 **Data sources**

150 Data for this analysis are from the 2010 Afghanistan Mortality Survey (9). This was the
151 country's first nationally representative household survey and is currently the only national,
152 population-based survey that has collected data on women's health service utilisation for stillbirth.
153 The survey adopted a two-stage sampling design based on the 2011 Population and Housing Census
154 preparatory frame from the Central Statistics Organisation. The design produced a sample
155 representative at the country level for rural and urban areas, and for the North, Central and South
156 geographical domains that are regroupings of eight geographical regions (Figure 1). The rural areas of
157 Kandahar, Helmand, and Zabul provinces in the South were not surveyed for security reasons.
158 Overall, the survey covered 87% of the population; the 13% not surveyed belonged mostly to the
159 South zone (9).

160

161 **[Insert Figure 1]**

162

163 We used data from three questionnaires in the AMS survey; the household, women's, and
164 verbal autopsy (VA) questionnaire, based on the DHS model questionnaires developed by the DHS
165 program and adapted for Afghanistan. The women's questionnaire collected information from ever-
166 married women aged 12-49 years including background characteristics and a complete pregnancy
167 history which captured all pregnancies and their outcomes in a woman's lifetime. Among women that
168 gave birth in the preceding five years, the women's questionnaires captured maternal health care
169 utilisation including antenatal, delivery and post-natal care for the mother's last live birth or stillbirth.
170 The VA questionnaire was completed for each death that occurred in the preceding three years.

171

172 In total, 22,351 households were interviewed, which included 47,848 women aged 12-49
173 years, yielding a response of 98%. We limited our analysis to all women's births within the last three
174 years, giving a base of 17,215 births. We merged data from the VA with the women's and household

175 data so that selected variables not available in the pregnancy histories for stillbirths could be included
176 (fetal sex, multiple pregnancy, and timing of the stillbirth). We further restricted our sample to
177 mothers' most recent birth, giving a sample of 13,844 women/births (13,528 live births, 316
178 stillbirths) then corrected any misclassification between miscarriages, stillbirths or early neonatal
179 deaths using the VA data. This gave a final sample of 13834 births (13,523 live births, 311 stillbirths).
180 Details on this procedure is available in the Appendix.

181

182 **Study variables**

183

184 *Dependent variable: pregnancy outcome*

185 Our main outcome variable was pregnancy outcome for the mothers' most recent pregnancy
186 and was coded as stillbirth or live birth (see Appendix for detail). We used the definition of stillbirth
187 to be a late fetal death at ≥ 28 weeks' gestation as recommended by WHO for international
188 comparisons. The 2010 AMS recorded gestational age in months so we used seven months or more as
189 our cut off. We defined intrapartum stillbirths as those stillbirths where the mother reported no signs
190 of skin maceration based on the VA data.

191

192 *Independent variables and analytical framework*

193 We included individual, household, and community-level explanatory variables based on
194 those identified in the literature as having an important effect on stillbirth, and availability in the 2010
195 AMS dataset (1, 10) (see Appendix). To guide the analysis, we developed an analytical framework by
196 adapting existing frameworks (11, 12). This framework mapped explanatory variables according to
197 proximity to the outcome as distal, intermediate and proximal determinants (Figure A2) and
198 represented three defined time periods - pre-conception, pregnancy, and childbirth.

199

200

201

202

203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229

Statistical analysis

All analyses were performed using STATA/SE version 14.2. For the binary outcome, stillbirth, we used Poisson regression models with a log link function to estimate relative risks. All models were weighted using sample weights to account for the complex survey design and adjusted standard errors were used to obtain Wald test p-values and 95% confidence intervals.

We fitted univariable models and built three multivariable regression models to examine the association between stillbirth and the explanatory variables. We applied a sequential approach (13) based on the three stages of pregnancy (Figure A2): model 1 included variables from the pre-conception period (community level, socio-economic, environmental & maternal factors); model 2 included factors related to the pregnancy period (antenatal care and pregnancy complications), having adjusted for the variables from stage 1; and model 3 included factors related to the delivery period (complications during the delivery period and delivery care), having adjusted for the variables from the first two stages. In the first model, no p-value criterion was used for including variables, but for subsequent models (model 2 and model 3) only variables with $p \leq 0.20$ from the previous model were included into the next stage. Wealth, maternal age and education, fetal sex, and multiple gestation were considered important factors and were retained in the models regardless of their p-values. All other variables were removed one at a time, starting with the highest p-value, until only those that had $p \leq 0.05$ remained. Multi-collinearity was checked using variance inflation factors. Area under the curve (AUC) and calibration plots were used to assess model performance. We used the same model building approach for identifying independent risk factors for intrapartum stillbirth.

230

231 **Results**

232 We included 13,834 births, of which 311 were stillbirths and 13,523 were live births (23
233 stillbirths per 1000 total births) (Table A2). Most women resided in rural areas (80.9%), were married
234 (99.5%), and had no formal education (89.4%). First-time mothers comprised 16% of the sample,
235 however, fertility was high with over 40% of women having at least five children prior to the index
236 pregnancy. Approximately 5% had experienced a previous pregnancy loss. Over one-third (36.0%) of
237 women had not received ANC for their last birth, while 16% had the recommended four or more
238 visits. Quality of ANC was generally low, with most women receiving less than five of the nine
239 recommended services. Only one-third of births took place at a health facility with a skilled birth
240 attendant, and less than 2% of births were caesarean. Almost two-thirds (60.6%) of stillbirths
241 occurred during the intrapartum period; although the timing was unknown for nearly 20%. The most
242 frequent maternal conditions during pregnancy were headaches, possible hypertension or infection,
243 and bleeding. Common complications in the delivery period were headaches, blurry vision, possible
244 hypertension or infection, excessive bleeding, prolonged labour/malpresentation. About 1% of women
245 in the pregnancy and delivery periods reported reduced fetal movements (Table A2).

246

247 Disparities in stillbirth rates across the eight geographical regions were high, ranging from 13
248 per 1000 births in the Northern regions to over 40 per 1000 births in the Central Highlands (Table 1).
249 In the univariate analysis, region of residence, ethnicity and maternal age were strongly associated
250 with stillbirth, but wealth quintile and education were not. First and higher order pregnancies,
251 multiple gestation, previous pregnancy loss, and not receiving ANC, were all associated with stillbirth
252 (Table 1).

253

254 **[Insert Table 1]**

255

256 Table 2 shows the multivariable results of factors associated with stillbirth. Factors associated
257 with stillbirth in the pre-pregnancy period (model 1) were region of residence, ethnicity, previous

258 pregnancy loss, and nulliparity. In particular, women in the Central highlands and South-Eastern
259 regions had twice the risk of stillbirth and women of Nuristani ethnicity were ten times more likely to
260 experience stillbirth. Once pregnant, taking into account utilisation of ANC and pregnancy
261 complications, region of residence was no longer associated with stillbirth, but ethnicity remained,
262 with Nuristani women having over nine times increased risk of stillbirth (model 2). Not receiving any
263 ANC during pregnancy increased the likelihood of stillbirth by almost three times, while women that
264 experienced possible infection, bleeding, and headache during their pregnancy had approximately
265 twice the risk of stillbirth, while women with reduced fetal movement were almost four times more
266 likely to have a stillbirth. Factors independently associated with stillbirth in the delivery period
267 (model 3) were, again, region of residence, ethnicity, previous pregnancy loss, first and multiple
268 pregnancies, not receiving ANC, and giving birth in a health facility. The same pregnancy
269 complications were predictive of stillbirth, except that the effect of reduced or no fetal movement as a
270 pregnancy complication was reduced. This is likely due to the inclusion of reduced or no fetal
271 movement as a delivery complication in the final model, which was now the factor with the highest
272 relative risk, increasing risk of stillbirth by nearly seven times. Across all models Nuristani women
273 consistently had a higher risk of stillbirth being over 12 times that of the Tajik population. There was
274 no difference in stillbirth across wealth quintiles or levels of maternal education or age after
275 accounting for all other factors. Models were well calibrated (Figures A5) and discrimination
276 improved from model 1 to model 3 (see AUC in Table 2).

277

278 **[Insert Table 2]**

279

280 The sub-group analysis on intrapartum stillbirth showed that being of Nuristani or Pashai
281 ethnicity, nulliparous women, multiple pregnancies, receiving no or low-quality ANC, or
282 experiencing possible infection or headache during pregnancy increased the risk of stillbirth. Reduced
283 or no fetal movement during the delivery period was also a strong predictor, whereas reduced or no
284 fetal movement as a pregnancy complication and giving birth in health facility were no longer
285 associated with intrapartum stillbirth once other factors were taken into account (Table 3, Figure A6).

286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313

[Insert Table 3]

We examined health care access disaggregated by ethnicity and region of residence to understand the disparities in stillbirth observed in access between different ethnic groups and geographic regions (Table A3, Figures A3-A4) and found that Nuristani women had the largest proportion of women that did not receive any ANC (89%) and the lowest levels of skilled birth attendance (2.5%) for their last pregnancy. The highest proportion of women receiving low quality ANC were residents of the Capital and Northern region (Figure A3). The South-Eastern, Western and Central highlands regions had the most women who did not receive any ANC. Skilled birth attendance was highest in the capital and lowest in the Central highlands, North-Eastern and Western regions (Figure A4).

314

315 **Comment**

316

317 *Principal findings*

318 Our analysis of the 2010 Afghanistan Mortality Survey has highlighted several socio-
319 demographic, health service utilisation, and maternal conditions that increase stillbirth in Afghan
320 women, some of which are modifiable and can inform and prioritise programmatic focus for future
321 stillbirth prevention in the country. Determinants of stillbirth in Afghanistan included residing in the
322 Central highlands, being of Nuristan ethnicity, not receiving ANC, and experiencing bleeding,
323 possible infection or headache during pregnancy. Reduced or no fetal movements during the delivery
324 period and giving birth in a health facility were also strongly associated with stillbirth. Factors
325 associated with intrapartum stillbirths differed slightly and included being of Nuristan or Pashai
326 ethnicity, utilisation and quality of ANC, possible infection or headache during pregnancy, and
327 reduced fetal movements in the delivery period. Women with first or multiple pregnancies, and
328 previous pregnancy loss also had increased risk of intrapartum stillbirth. These findings offer an
329 evidence-base to integrate efforts into health service delivery programmes focused on maternal,
330 perinatal and newborn survival, as well as future national health policies where until now, no such
331 information was available. We also demonstrate how DHS surveys can be adapted to generate more
332 data to understand the underlying factors driving stillbirths in other LMIC settings. DHSs

333

334 The overall stillbirth rate of 23 per 1000 total births is lower than adjusted rates reported for
335 Afghanistan in 2009 (29.3 per 1000) and in 2015 (26.7 per 1000) from the Lancet series which
336 accounted for under-reporting (1). Intrapartum stillbirths constituted almost two-thirds of stillbirths in
337 our study and is consistent with findings from other LMICs (3). Within-country variations in stillbirth
338 risk have been observed in many countries, as have ethnic differences (14, 15). Nuristani people are a
339 minority group that reside predominantly in the Eastern part of Afghanistan (Nuristan province) and
340 the low levels of healthcare utilisation may explain the extremely high rates of stillbirth. The 2015
341 Afghanistan DHS also found only 1% of births in Nuristan province were in a health facility, and this

342 province had the lowest levels of ANC utilisation across the country (11%). Exacerbating the
343 situation is that the East is a high-intensity conflict zone and one of the poorest regions in the country.
344 For intrapartum stillbirths, both Nuristani and Pashai women had higher risk of stillbirth. Pashai
345 women also reside in the East, where high levels of conflict could have compromised access and
346 quality of health services. The 2010 AMS did not report mortality rates according to province or
347 ethnicity; however, the 2015 DHS reported provincial level mortality rates which showed that
348 Nuristan province had the highest infant and under-five child mortality rates nation-wide (123 and
349 170 per 1000 live births respectively, compared to 45 and 55 per 1000 live births nationally) (16). The
350 high stillbirth rates in this group appears to reflect the pattern in regional disparities in other mortality
351 rates.

352

353 Geographic disparities underlie maternal and child mortality, morbidity, and healthcare seeking
354 in Afghanistan (17). The high rates of stillbirth among women in the Central highlands are likely due
355 to lack of access and availability of health services, as these areas are characterised by mountainous
356 terrain often isolated by snow. This region experiences scarcities in medical supplies due to poor
357 transport infrastructure and security concerns, and a shortage of medical doctors willing to work there.
358 Women from the South-Eastern region had higher risk of stillbirth in the initial multivariable model
359 until adjustment with antenatal and delivery care variables, indicating the importance of health service
360 utilisation in this area. High levels of conflict would likely limit access and availability of services in
361 this area.

362

363 The diverse geographical terrain with concentrated ethnic groups in specific regions, combined
364 with insecurity will require tailored approaches to reach these hard-to-reach, high-risk women. Tappis
365 et. al. in their study examining coverage of intrapartum care in selected areas of Afghanistan also
366 identified the importance of context-specific service delivery models to ensure women in high conflict
367 areas can access services (18). A major barrier to ensuring facility deliveries in some parts of the
368 country was the inability to travel at night along major roads because of insecurity. Delivering health
369 services to remote and mountainous areas is challenging especially in the context of insecurity, but

370 strategies which strengthen the role of local community health workers and task shifting can be
371 effective. A revised primary health care service delivery model, currently under development by the
372 Ministry of Public Health may provide an opportunity to integrate alternative approaches to facilitate
373 reductions in stillbirth.

374

375 Mothers who did not receive ANC were three times more likely to experience stillbirth and
376 while quality did not appear to make a difference for all stillbirths, it did matter for intrapartum
377 stillbirths. This suggests having any ANC is important for preventing stillbirths, but that quality and
378 content of care may be critical for identifying and managing maternal conditions early that could lead
379 to childbirth complications and intrapartum stillbirth. Overall, ANC utilisation was very low, and we
380 measured quality according to whether the mother received any of the nine checks, not necessarily,
381 the adequacy of the service or the initiation of treatment. Our measurement method may partly
382 explain the absence of an overall effect of quality of ANC for stillbirths. These downstream factors
383 are important to consider when assessing the effectiveness of ANC on stillbirth (19). Further
384 investigation is needed to examine the quality of care provided and adherence to recommended advice
385 among women. Our analysis showed that areas that achieved higher coverage of ANC (i.e. the Capital
386 and Northern regions) actually had a higher proportion of women receiving lower quality of ANC.
387 Ensuring adequate and high-quality ANC is one of the simplest and most cost-effective recommended
388 interventions to reduce stillbirths (20). Efforts to strengthen ANC are in progress where the Afghan
389 government is administering a maternal and child health handbook that contains information on safe
390 pregnancy, childbirth, and childcare to each pregnant woman and documents details of visits. It will
391 be important to record the services received, pregnancy progress, and results from any screening tests
392 in this handbook.

393

394 We identified several pregnancy conditions that were associated with stillbirth and are
395 preventable. Signs of infection and antepartum bleeding were important determinants in our study and
396 are well-established risks. Effective interventions exist for treating malaria and syphilis to reduce
397 stillbirth (21), and while malaria is endemic in some of the semi-arid Eastern and northern provinces

398 in Afghanistan, syphilis and HIV prevalence in Afghanistan is generally very low and limited to high
399 risk groups such as injecting drug users and sex workers (22, 23). Further research is needed to
400 identify common infections contributing to stillbirth in this setting. Hirose et al.(24) identified that
401 care-seeking delays in Afghanistan were higher among women experiencing severe infections
402 compared to other complications with more concerning symptoms, so it would be important to ensure
403 early detection and management of both bleeding and infections by educating women and family
404 members on the urgency of care-seeking for symptoms. Headaches during pregnancy were also a
405 strong risk factor of both stillbirth and intrapartum stillbirth and likely a sign of pre-eclampsia or
406 pregnancy-induced hypertension, which are known risk factors for stillbirth. Ensuring that ANC
407 includes blood pressure checks and appropriate management will be critical for reducing
408 complications that lead to stillbirth. Reduced fetal movements have rarely been examined in low-
409 income countries but is a known risk factor for stillbirth (25). Of all delivery complications, reduced
410 fetal movements was one of the strongest determinants for both stillbirth and intrapartum stillbirth in
411 our study. It would be important to ensure women understand the need to act upon any perceived
412 reduction or change in fetal movements, and that during the intrapartum period movements are
413 closely monitored.

414

415 Variations exist on the effect of delivery location on stillbirth with some studies showing an
416 increase in risk (26) while others indicate a protective effect (3). We found facility births had
417 increased odds of stillbirth overall, but for intrapartum stillbirth place of birth had no effect. Referral
418 bias, delays in care-seeking, or quality of care may account for these findings. The absence of an
419 association with intrapartum stillbirths is likely related to the quality of care or care-seeking delays. A
420 study examining delays in care-seeking in Afghanistan showed substantial departure and decision
421 making delays among pregnant women with life-threatening conditions (27). Concerns regarding
422 quality of intrapartum care in maternity hospitals in Afghanistan have also been documented (28).
423 Ballard et al. (29) in their Ethiopian study also found that women with an intrapartum emergency
424 were twice as likely to give birth in a health facility and that facility births did not reduce stillbirth
425 risk, suggesting the three delays was at play here (30).

426

427 We could not include mode of delivery in our multivariable analysis, but caesarean births
428 showed a high positive association with stillbirth in the univariate results. A study of over 50,000
429 births in Kabul hospitals identified high rates of stillbirth in caesareans done for obstructed labour,
430 malpresentation and uterine rupture, which are preventable with timely intervention (31). We did not
431 have data on indication for caesarean but an assessment of 78 first line referral facilities in
432 Afghanistan found 88% of caesarean births were emergencies (32), so it is likely most were
433 unplanned. The ideal caesarean rate to observe reductions in intrapartum stillbirth is between 5-10%
434 (33), but here we found it was under 2% and more recent national data reports a rate of only 3% (16),
435 therefore, improving access to caesarean would be important to prevent stillbirths in Afghanistan.

436

437 Male babies have an increased risk of stillbirth (34); however, we only found a slightly
438 increased risk among intrapartum stillbirths but not in the analysis of the full sample of stillbirths.
439 This may have been affected by the skewed sex ratio among stillbirths in our sample which had
440 almost 20% more male babies to female. This is higher than the usual 10% elevated risk of stillbirth
441 in male babies (34). The 2010 AMS (9) and 2015 Afghanistan DHS (16) both identified under-
442 reporting of neonatal and under-five child female deaths, as have other household surveys from
443 Afghanistan (35). This might suggest that under-reporting could also be a problem with female
444 stillborn deaths and partly contributing to the overall under-estimate of the true stillbirth burden.
445 Under-reporting of stillbirths can occur due to social, cultural or other factors including stigma or
446 blame towards the mother or other consequences that might preclude disclosure (36). Further
447 investigation into these issues is needed for Afghanistan.

448

449 *Strengths of the study*

450 A key strength of this study is the use of a large nation-wide population sample to identify
451 risk factors for stillbirth. In addition, this survey collected a comprehensive range of socio-
452 demographic, maternal and fetal characteristics, maternal complications and health care utilisation
453 factors for stillbirths which are not usually available in similar household surveys in LMICs.

454

455 *Limitations of the data*

456 There are several limitations to this study that should be considered. Although the 2010 AMS
457 was a national survey, there was an underrepresentation of the South because of highly insecure areas
458 that were not surveyed. Concerns about the accuracy of maternal and child mortality measures from
459 this survey have been noted (37) and highlight the challenges with collecting reliable data in conflict
460 zones. While we acknowledge this limitation, this is currently the only data source in the country with
461 information to enable understanding of key determinants of stillbirth. Although an updated DHS
462 survey was subsequently conducted, it did not capture health service utilisation, maternal or fetal
463 factors for stillbirth, which precludes the kind of analysis reported in this paper.

464

465 Collecting information on pregnancy histories is challenging in low-income settings, and
466 stillbirths are known to be under-estimated by about 30% when collected through household surveys
467 (5). The overall stillbirth rate of 22.5 per 1000 births in our study is low given the high levels of
468 neonatal and maternal mortality in the country. It is possible these estimates have been affected by
469 under-reporting and the data quality concerns raised with the 2010 AMS survey. The exclusion of
470 some rural areas of the South zone of Afghanistan during sampling because of security reasons also
471 meant that the survey covered only 66% of the South (94% of urban and 63% of rural areas were
472 sampled) and so many stillbirths from rural areas would not have been included. The lower stillbirth
473 rates observed in the Southern region (17.8 per 1000 births) is likely to have been affected by under-
474 sampling of rural areas of the south. Medical terminations are illegal in Afghanistan, so women may
475 report these as stillbirths or omit them entirely which might affect the total number of pregnancies and
476 stillbirths reported. Misclassification of stillbirths and early neonatal deaths is an issue with household
477 surveys, but we have minimised this by using data from the VA. The reliability of using skin
478 appearance to determine the timing of stillbirth may lead to an overestimate of intrapartum stillbirth
479 (38). Due to the small number of antepartum stillbirths we were not able to model antepartum
480 stillbirth risk factors separately to compare with the intrapartum stillbirth risk factors. Several known
481 risk factors were not captured, and we could not adjust for them (i.e. consanguinity, maternal

482 nutrition, distance to health facility, and care-seeking delays). Exposures related to the armed conflict
483 including chemicals and radiation, are known to increase the risk of stillbirth (39), but we had no
484 measurement of these exposures. Finally, it is possible there was under-reporting of the self-reported
485 maternal complications due to recall bias.

486

487 *Interpretation*

488 We provide for the first-time the major risk factors associated with stillbirth in Afghanistan,
489 where there was previously a complete absence of evidence to inform future interventions and
490 prevention efforts. Evidence-based interventions to prevent stillbirth exist (20) and their
491 implementation should be a priority for Afghanistan. We outline some recommendations in Box 2.
492 This study also demonstrates it is feasible to rapidly produce a comprehensive analysis of stillbirth
493 determinants for other LMICs if appropriate DHS data was available. To achieve this outcome would
494 require some modification to the standard DHS questionnaire format to include a full pregnancy
495 history as opposed to a live birth history (8), as well as the inclusion of stillbirth in collection of
496 information on women's health care utilisation and maternal complications during pregnancy and
497 childbirth. The 2010 AMS provides a model from which future household surveys can be adapted to
498 collect better data for stillbirth.

499

500 *Conclusions*

501 Countries affected by conflict and instability account for the largest burden of stillbirths (40),
502 but strategies to improve reproductive outcomes in these areas have not received sufficient global
503 attention and is urgently needed. Development assistance and international focus on Afghanistan has
504 declined recently as the security situation has worsened, and gains in maternal and child health are at
505 risk of deteriorating. To accelerate reductions in stillbirth, concerted efforts and commitment by the
506 government and international donors are needed to invest in prioritising implementation of
507 interventions to reduce stillbirth. Evaluating different approaches to overcome challenges in the
508 access and utilisation of care during pregnancy and childbirth to ensure services can reach the most

509 hard-to-reach women where the majority of stillbirths occur, will be imperative for future stillbirth
510 reduction in Afghanistan.

511 **[Insert Box 2: Recommendations for the prevention and reduction of stillbirth in Afghanistan]**

512

513

514

515

516

517

518

519

520

521

522

523

524

525

526

527

528

529

530

531

532

533

534

535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576

Declaration of interests

The authors declare no competing interests.

Acknowledgements

This study had no direct funding sources. During the completion of this work AC was supported by an Australian Postgraduate Award funded through the Australian Commonwealth Government and CRG was supported by a National Health and Medical Research Council Career Development Fellowship (1087062). We would like to thank Mr Rafiqul Haider for preparation of the map, and Ms Alison Gibberd for helpful discussions on the data analysis. We also acknowledge and thank the DHS programme for providing access to the 2010 Afghanistan Mortality Survey dataset.

577
578
579
580

581 **References**

- 582
- 583 1. Lawn JE, Blencowe H, Waiswa P, Amouzou A, Mathers C, Hogan D, et al. Stillbirths: rates,
584 risk factors, and acceleration towards 2030. *The Lancet*. 2016;387(10018):587-603.
- 585 2. Ashish K, Wrammert J, Ewald U, Clark RB, Gautam J, Baral G, et al. Incidence of
586 intrapartum stillbirth and associated risk factors in tertiary care setting of Nepal: a case-control study.
587 *Reproductive Health*. 2016;13(1):103.
- 588 3. McClure EM, Saleem S, Goudar SS, Moore JL, Garces A, Esamai F, et al. Stillbirth rates in
589 low-middle income countries 2010-2013: a population-based, multi-country study from the Global
590 Network. *Reproductive health*. 2015;12(Suppl 2):S7.
- 591 4. Kujala S, Waiswa P, Kadobera D, Akuze J, Pariyo G, Hanson C. Trends and risk factors of
592 stillbirths and neonatal deaths in Eastern Uganda (1982-2011): A cross-sectional, population-based
593 study. *Tropical Medicine & International Health*. 2016:n/a-n/a.
- 594 5. Cousens S, Blencowe H, Stanton C, Chou D, Ahmed S, Steinhardt L, et al. National, regional,
595 and worldwide estimates of stillbirth rates in 2009 with trends since 1995: a systematic analysis. *The*
596 *Lancet*. 2011;377(9774):1319-30.
- 597 6. Blencowe H, Cousens S, Jassir FB, Say L, Chou D, Mathers C, et al. National, regional, and
598 worldwide estimates of stillbirth rates in 2015, with trends from 2000: a systematic analysis. *Lancet*
599 *Glob Health*. 2016;4(2):e98-e108.
- 600 7. WHO. Every Newborn Action Plan: Progress Report May 2015. Geneva: World Health
601 Organization,; 2015.
- 602 8. Christou A, Dibley MJ, Raynes-Greenow C. Beyond counting stillbirths to understanding
603 their determinants in low- and middle-income countries: a systematic assessment of stillbirth data
604 availability in household surveys. *Tropical Medicine & International Health*. 2017;22(3):294-311.
- 605 9. ANPHI MOPH, CSO, ICF Macro, IIMMR, WHO (EMRO). Afghanistan Mortality Survey
606 2010. Calverton, Maryland, USA: Afghan National Public Health Institute, Ministry of Public Health
607 (ANPHI/MoPH), Central Statistics Organization (CSO), ICF Macro, Indian Institute of Health
608 Management Research (IIMMR), World Health Organization Regional Office for the Eastern
609 Mediterranean (WHO/EMRO),; 2011.
- 610 10. Aminu M, Unkels R, Mdegela M, Utz B, Adaji S, van den Broek N. Causes of and factors
611 associated with stillbirth in low- and middle-income countries: a systematic literature review. *BJOG:*
612 *An International Journal of Obstetrics & Gynaecology*. 2014;121:141-53.

- 613 11. Stanton C. Perinatal mortality in the Phillippines: an invesigation into the use of demographic
614 survey data for the sutdy of perinatal mortality and its determinants. Baltimore, Maryland: The Johns
615 Hopkins University; 1996.
- 616 12. Mosley WH, Chen LC. An analytical framework for the study of child survival in developing
617 countries. *Population and development review*. 1984;10:25-45.
- 618 13. Victora CG, Huttly SR, Fuchs SC, Olinto MT. The role of conceptual frameworks in
619 epidemiological analysis: a hierarchical approach. *Int J Epidemiol*. 1997;26(1):224-7.
- 620 14. Lee AC, Mullany LC, Tielsch JM, Katz J, Khattry SK, LeClerq SC, et al. Community-based
621 stillbirth rates and risk factors in rural Sarlahi, Nepal. *International Journal of Gynecology &
622 Obstetrics*. 2011;113(3):199-204.
- 623 15. Carvalho TS, Pellanda LC, Doyle P. Stillbirth prevalence in Brazil: an exploration of regional
624 differences. *Jornal de Pediatria*. 2017.
- 625 16. CSO, MOPH, ICF. Afghanistan Demographic and Health Survey 2015. Kabul, Afghanistan:
626 Central Statistics Organization (CSO); 2017.
- 627 17. Akseer N, Bhatti Z, Rizvi A, Salehi AS, Mashal T, Bhutta ZA. Coverage and inequalities in
628 maternal and child health interventions in Afghanistan. *BMC Public Health*. 2016;16(2):797.
- 629 18. Tappis H, Koblinsky M, Winch PJ, Turkmani S, Bartlett L. Context matters: Successes and
630 challenges of intrapartum care scale-up in four districts of Afghanistan. *Glob Public Health*.
631 2016;11(4):387-406.
- 632 19. Pai M, Schumacher SG, Abimbola S. Surrogate endpoints in global health research: still
633 searching for killer apps and silver bullets? *BMJ Glob Health*. 2018;3(2):e000755.
- 634 20. Bhutta ZA, Yakoob MY, Lawn JE, Rizvi A, Friberg IK, Weissman E, et al. Stillbirths: what
635 difference can we make and at what cost? *Lancet*. 2011;377(9776):1523-38.
- 636 21. Ishaque S, Yakoob MY, Imdad A, Goldenberg RL, Eisele TP, Bhutta ZA. Effectiveness of
637 interventions to screen and manage infections during pregnancy on reducing stillbirths: a review.
638 *BMC Public Health*. 2011;11(3):S3.
- 639 22. Todd CS, Ahmadzai M, Atiqzai F, Miller S, Smith JM, Ghazanfar SAS, et al. Seroprevalence
640 and correlates of HIV, syphilis, and hepatitis B and C virus among intrapartum patients in Kabul,
641 Afghanistan. *BMC Infectious Diseases*. 2008;8(1):119.
- 642 23. Todd CS, Nasir A, Stanekzai MR, Fiekert K, Rasuli MZ, Vlahov D, et al. Prevalence and
643 correlates of HIV, syphilis, and hepatitis B and C infection and harm reduction program use among

- 644 male injecting drug users in Kabul, Afghanistan: A cross-sectional assessment. *Harm Reduction*
645 *Journal*. 2011;8(1):22.
- 646 24. Hirose A, Borchert M, Cox J, Alkozai AS, Filippi V. Determinants of delays in travelling to
647 an emergency obstetric care facility in Herat, Afghanistan: an analysis of cross-sectional survey data
648 and spatial modelling. *BMC Pregnancy and Childbirth*. 2015;15(1):14.
- 649 25. Heazell AEP, Warland J, Stacey T, Coomarasamy C, Budd J, Mitchell EA, et al. Stillbirth is
650 associated with perceived alterations in fetal activity – findings from an international case control
651 study. *BMC Pregnancy and Childbirth*. 2017;17(1):369.
- 652 26. Ha YP, Hurt LS, Tawiah-Agyemang C, Kirkwood BR, Edmond KM. Effect of
653 Socioeconomic Deprivation and Health Service Utilisation on Antepartum and Intrapartum Stillbirth:
654 Population Cohort Study from Rural Ghana. *PLOS ONE*. 2012;7(7):e39050.
- 655 27. Hirose A, Borchert M, Niksear H, Alkozai AS, Cox J, Gardiner J, et al. Difficulties leaving
656 home: A cross-sectional study of delays in seeking emergency obstetric care in Herat, Afghanistan.
657 *Social Science & Medicine*. 2011;73(7):1003-13.
- 658 28. Kim Y-M, Zainullah P, Mungia J, Tappis H, Bartlett L, Zaka N. Availability and quality of
659 emergency obstetric and neonatal care services in Afghanistan. *International Journal of Gynecology*
660 *& Obstetrics*. 2012;116(3):192-6.
- 661 29. Ballard K, Belete Z, Kinfu H, Tadesse M, Amin M, Atnafu H. The effect of prenatal and
662 intrapartum care on the stillbirth rate among women in rural Ethiopia. *International Journal of*
663 *Gynecology & Obstetrics*. 2016;133(2):164-7.
- 664 30. Thaddeus S, Maine D. Too far to walk: maternal mortality in context. *Soc Sci Med*.
665 1994;38(8):1091-110.
- 666 31. Guidotti RJ, Kandasamy T, Betrán AP, Merialdi M, Hakimi F, Van Look P, et al. Monitoring
667 perinatal outcomes in hospitals in Kabul, Afghanistan: The first step of a quality assurance process.
668 *The Journal of Maternal-Fetal & Neonatal Medicine*. 2009;22(4):285-92.
- 669 32. Kim YM, Tappis H, Zainullah P, Ansari N, Evans C, Bartlett L, et al. Quality of caesarean
670 delivery services and documentation in first-line referral facilities in Afghanistan: a chart review.
671 *BMC Pregnancy Childbirth*. 2012;12:14.
- 672 33. Goldenberg RL, McClure EM, Bann CM. The relationship of intrapartum and antepartum
673 stillbirth rates to measures of obstetric care in developed and developing countries. *Acta Obstet*
674 *Gynecol Scand*. 2007;86(11):1303-9.
- 675 34. Mondal D, Galloway TS, Bailey TC, Mathews F. Elevated risk of stillbirth in males:
676 systematic review and meta-analysis of more than 30 million births. *BMC Medicine*. 2014;12(1):220.

- 677 35. Viswanathan K, Becker S, Hansen PM, Kumar D, Kumar B, Niayesh H, et al. Infant and
678 under-five mortality in Afghanistan: current estimates and limitations. Bull World Health Organ.
679 2010;88(8):576-83.
- 680 36. Haws RA, Mashasi I, Mrisho M, Schellenberg JA, Darmstadt GL, Winch PJ. "These are not
681 good things for other people to know": how rural Tanzanian women's experiences of pregnancy loss
682 and early neonatal death may impact survey data quality. Soc Sci Med. 2010;71(10):1764-22.
- 683 37. Hill K. Gains in Afghan Health: Too Good to Be True? 2012 [Available from:
684 http://www.cgdev.org/doc/blog/Gains_in_Afghan_Health_CGD_Talk_pdf.pdf.
- 685 38. Gold KJ, Abdul-Mumin AR, Boggs ME, Opare-Addo HS, Lieberman RW. Assessment of
686 "fresh" versus "macerated" as accurate markers of time since intrauterine fetal demise in low-income
687 countries. Int J Gynaecol Obstet. 2014;125(3):223-7.
- 688 39. Keasley J, Blickwedel J, Quenby S. Adverse effects of exposure to armed conflict on
689 pregnancy: a systematic review. BMJ Global Health. 2017;2(4).
- 690 40. Wise PH, Darmstadt GL. Confronting stillbirths and newborn deaths in areas of conflict and
691 political instability: a neglected global imperative. Paediatrics and international child health.
692 2015;35(3):220-6.
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709

710 **Tables and Boxes**

711

712 **Box 1: Afghanistan country context and health situation**

713

Afghanistan is a culturally rich nation located in south-central Asia sharing borders with six different countries, the longest being with Pakistan. The country's 34 provinces comprise a diverse range of ethnicities, languages, and geographic terrains. It is mostly a mountainous landscape with the Hindu Kush mountain range dividing the country from the northeast to the southwest into three distinct regions – the mountainous central highlands, the south-west plateaus characterised by deserts, and the smaller and most fertile northern plains. The current population is estimated to be approximately 30 million. The country is one of the least developed nations in the world, ranked 169 out of 188 nations on the human development index in 2015. About one third (37%) of the population lives below the poverty line and this has remained unchanged since 2007-08.

Afghanistan has a very young population structure with 48% aged under 15 years, and average life expectancy of only 60 years. Fertility rates are high with an average of 5.3 children in 2015 – a slight increase from 2010 rate of 5.1 (9, 16). Adult literacy rates remain low at 31%, particularly among females (males 45%; females 17%).

Afghanistan has faced over four decades of ongoing conflict, unstable governance and population displacement which continues. In 2016, the conflict led to the displacement of over half a million people, more than half of whom were children, and an unexpected influx of over one million Afghan refugees and returnees from Pakistan. It is estimated that over nine million people have limited or no access to essential health services, straining an already weak and recovering health system. The impact of the conflict on access to health services and health education for women and their families is therefore, particularly challenging.

Since its release from Taliban rule in 2002, immense efforts have been made by the Afghan government and international community to repair and strengthen the health system. Although rates of maternal and child deaths continue to be some of the highest in the world, there have been some encouraging improvements; maternal mortality has declined from 1600 deaths per 100,000 in 2002 to 327 per 100,000 in 2010, however, a 2013 analysis suggests these rates may be inaccurate and could be around 885 per 100 000 live births. Recent estimates for under five child mortality suggest around 70 deaths per 1000 live births and a neonatal mortality rate of 40 per 1000 live births. In 2015, 18% of women received the recommended four or more ANC visits and 50% reported attendance of a skilled birth provider at their most recent birth, an increase from 34% in 2010 (9). Despite these improvements, there remains inadequate access to, and utilisation of, ANC and quality obstetric care services (28), with stark inequities in access between urban and rural areas and across regions (17). Health system challenges exist around sufficient numbers of female health care providers and the costs of health services and treatment. There are also additional contextual challenges and social and cultural norms surrounding women's low levels of autonomy and education that directly impact on care-seeking delays and child health outcomes (27).

714

715

716

717

718

719

720

721

722

Box 2: Recommendations for the prevention and reduction of stillbirth in Afghanistan**Health systems strengthening and health service delivery**

- Improved coverage and monitoring of content of ANC for the early identification and management of high-risk pregnancies and early referral.
- The high number of intrapartum stillbirths indicates a need for improved quality and timely management of childbirth complications. Ensuring birth attendants at all levels of the health system are adequately trained and have the skills and resources available to manage complications will be essential.
- Increasing the availability and access to timely caesarean sections for high-risk pregnancies and minimising delays at the facility level.
- The highest burden of stillbirth in the country falls in the Central highlands and among minority ethnolinguistic groups which will require specialised attention and targeted strategies.

Community-based education and mobilisation

- Improve community awareness and education on key danger signs during pregnancy and childbirth that need immediate action.
- Sensitisation of community specifically about stillbirths and their prevention will also be important but will require additional strategies to overcome barriers and delays in care-seeking.

Further research

- Identification of the leading infections that may be contributing to stillbirth in Afghanistan require further research and understanding.
- Improved understanding of bottlenecks and barriers at the health facility level in regard to the prevention of stillbirth.
- Assessment of the quality of antenatal and intrapartum care provided at the various levels of health facilities.
- Development of strategies to strengthen referral linkages and facilitate referral and reduce care-seeking delays at the community level.

**Commitment to stillbirth targets in national health strategies and policies
& continued data collection on stillbirth**

- Afghanistan's current National Health Policy for 2015-2020 and Reproductive, Maternal, Newborn Child and Adolescent Health (RMNCAH) Strategy for 2017-2021 do not include targets for stillbirth reduction. National commitment in future policies and strategies to the recommended targets agreed upon as part of the 2014 Every Newborn Action Plan and endorsed at the World Health Assembly will direct national attention, prioritisation and funding towards reducing stillbirths.
- Future national population-based surveys should include a full pregnancy history similar to the 2010 Afghanistan Mortality Survey to ensure ongoing data availability on the key risk factors for stillbirths. This survey data will assist with tracking progress towards meeting the global target of 12 stillbirths per 1000 births by 2030 and identifying key areas of need for interventions.

727 **Table 1. Univariable results of factors associated with stillbirth for women's most recent birth in the**
 728 **preceding three years, Afghanistan 2010**
 729

	Stillbirths	All births	Stillbirth rate per 1000	Unadjusted RR [95% CI]
	N (%)	N (%)		
Total pregnancy outcomes (weighted)	311 (2.2)	13 834 ^a (100)	22.5	
COMMUNITY LEVEL				
Residence				
Urban	49 (15.7)	2636 (19.1)	18.5	1.00 [Reference]
Rural	262 (84.3)	11198 (80.9)	23.4	1.26 [0.90, 1.77]
Region				
North-Eastern	34 (10.9)	2081 (15.0)	16.3	1.00 [Reference]
Northern	28 (9.0)	2145 (15.5)	13.0	0.80 [0.45, 1.42]
Western	37 (12.0)	1841 (13.3)	20.3	1.25 [0.72, 2.16]
Central Highland	20 (6.4)	430 (3.1)	46.6	2.86 [1.34, 6.12]
Capital	58 (18.6)	2635 (19.1)	21.9	1.35 [0.82, 2.21]
Eastern	79 (26.5)	2472 (17.8)	32.1	1.97 [0.94, 4.16]
Southern	16 (5.2)	906 (6.6)	17.8	1.09 [0.62, 1.92]
South-Eastern	39 (12.5)	1324 (9.6)	29.3	1.80 [1.12, 2.88]
Ethnicity¹				
Tajik	76 (24.3)	4386 (31.7)	17.2	1.00 [Reference]
Pashtun	127 (40.9)	5992 (43.4)	21.2	1.23 [0.90, 1.69]
Hazara	30 (9.5)	1125 (8.1)	26.3	1.53 [0.90, 2.60]
Uzbek	24 (7.6)	1218 (8.8)	19.4	1.13 [0.66, 1.94]
Nuristan	35 (11.3)	189 (1.4)	186.0	10.80 [3.67, 31.77]
Pashai	10 (3.1)	318 (2.3)	30.2	1.75 [0.86, 3.57]
Other (Baloch/Turkmen/Other)	10 (3.3)	595 (4.3)	16.8	1.00 [0.43, 2.32]
SOCIO-ECONOMIC & ENVIRONMENTAL				
Wealth quintile				
Lowest	58 (18.5)	2828 (20.4)	20.4	1.07 [0.70, 1.762]
Second	76 (24.5)	2817 (20.4)	27.0	1.41 [0.99, 2.03]
Middle	75 (24.0)	2757 (19.9)	27.1	1.47 [1.01, 2.12]
Fourth	51 (16.4)	2736 (19.8)	18.7	0.98 [0.66, 1.45]
Highest	52 (16.6)	2696 (19.5)	19.1	1.00 [Reference]
Marital status				
Currently married	311 (100.0)	13769 (99.5)	-	-
Previously married	0 (0.7)	65.4 (0.5)	-	-
Maternal education				
No education or Madrassa	289 (93.1)	12372 (89.4)	23.4	1.59 [0.99, 2.55]
Any education ^b	22 (6.9)	1463 (10.6)	14.8	1.00 [Reference]
Source of drinking water²				
Improved water source	150 (48.3)	7653 (55.4)	19.6	1.00 [Reference]
Unimproved water source	161 (51.7)	6158 (44.6)	26.1	1.33 [0.95, 1.86]
Sanitation facility³				
Improved sanitation facility	117 (37.5)	5043 (36.5)	23.1	1.00 [Reference]
Unimproved sanitation facility/other	194 (62.5)	8781 (63.5)	22.1	0.96 [0.59, 1.56]
Fuel used for cooking⁴				
Clean fuel/no food cooked in house	49 (15.8)	2783 (20.2)	17.7	1.00 [Reference]
Solid fuel/other	262 (84.2)	11024 (79.8)	7.6	1.34 [0.94, 1.91]
MATERNAL & FETAL CHARACTERISTICS				
Sex of baby⁵				
Female	124 (42.0)	6280 (45.4)	19.8	1.00 [Reference]
Male	172 (58.0)	7538 (54.6)	22.8	1.15 [0.88, 1.51]
Pregnancy type⁶				
Singleton	285 (96.8)	13684 (99.0)	20.9	1.00 [Reference]
Multiple	9 (3.2)	133 (1.0)	70.2	3.37 [1.62, 6.98]
Maternal age				
12-18	22 (7.1)	1209 (8.7)	18.3	0.99 [0.53, 1.87]
19-24	92 (29.6)	5013 (36.2)	18.4	1.00 [Reference]
25-34	134 (43.0)	5666 (41.0)	23.6	1.28 [0.88, 1.87]
35+	63 (20.2)	1947 (14.1)	32.3	1.76 [1.26, 2.45]
Pregnancy order				
First pregnancy	58 (18.5)	2165 (15.7)	26.6	1.64 [1.10, 2.45]
2 nd -4 th pregnancy	98 (31.5)	6046 (43.7)	16.2	1.00 [Reference]
≥5th pregnancy	156 (50.0)	5623 (40.6)	27.7	1.71 [1.30, 2.24]
Pregnancy interval^f				
First pregnancy	58 (18.5)	2165 (15.7)	26.6	1.35 [0.89, 2.04]
<18 months	42 (13.5)	1664 (12.0)	25.2	1.28 [0.81, 2.05]
18-58 months	181 (58.0)	9174 (66.3)	19.7	1.00 [Reference]
≥59 months	31 (10.0)	831 (6.0)	37.3	1.90 [1.25, 2.87]
Previous adverse pregnancy outcome				
No prior adverse outcome ^d	270 (86.8)	13080 (94.6)	20.6	1.00 [Reference]
Yes	41 (13.2)	754 (5.5)	54.4	2.63 [1.87, 3.71]

	Stillbirths	All births	Stillbirth rate per 1000	Unadjusted RR [95% CI]
	N (%)	N (%)		
Total pregnancy outcomes (weighted)	311 (2.2)	13 834 ^a (100)	22.5	
ANTENATAL CARE				
Number of ANC visits^{c,7}				
None	145 (47.3)	4969 (36.2)	29.1	1.58 [0.98, 2.55]
1	43 (14.2)	1912 (13.9)	22.7	1.23 [0.74, 2.04]
2-3	76 (24.9)	4575 (33.3)	16.7	0.90 [0.60, 1.36]
4 or more	42 (13.7)	2272 (16.6)	18.4	1.00 [Reference]
Timing of first ANC visit^{c,8}				
First trimester	47 (15.4)	2569 (18.7)	18.5	1.00 [Reference]
Second trimester	61 (19.9)	3721 (27.1)	16.5	0.89 [0.60, 1.34]
Third trimester	55 (17.8)	2459 (17.9)	22.3	1.21 [0.77, 1.89]
No ANC	145 (46.9)	4969 (36.2)	29.1	1.58 [0.99, 2.52]
ANC provider^{c,9}				
Trained provider ^c	160 (51.5)	8413 (60.9)	19.0	1.00 [Reference]
Untrained provider ^f	6 (2.0)	432 (3.1)	14.1	0.74 [0.29, 1.89]
No ANC	145 (46.5)	4969 (36.0)	29.1	1.53 [1.07, 2.18]
Place of ANC^c				
Health facility/clinic	142 (45.5)	7694 (55.6)	18.4	1.00 [Reference]
Home/multiple providers/other	25 (7.9)	1171 (8.5)	21.1	1.14 [0.63, 2.07]
No ANC	145 (46.5)	4969 (35.9)	29.1	1.58 [1.11, 2.26]
ANC components^{c,g}				
Weighed ¹⁰	51 (16.4)	3481 (25.3)	14.6	0.58 [0.42, 0.80]
Blood pressure taken ¹¹	151 (48.6)	7932 (57.5)	19.0	0.70 [0.50, 0.98]
Urine sample taken ¹²	66 (21.2)	2920 (21.2)	22.5	1.00 [0.74, 1.36]
Blood sample taken ¹³	72 (23.1)	2742 (19.9)	26.2	1.21 [0.90, 1.63]
Given/bought iron tablets ¹⁴	103 (33.1)	5290 (38.3)	19.4	0.80 [0.56, 1.13]
Took intestinal parasite drugs ¹⁵	11 (3.7)	580 (4.2)	19.7	0.87 [0.41, 1.85]
Told signs of pregnancy complications ¹⁶	56 (18.1)	2888 (20.9)	19.2	0.83 [0.59, 1.17]
Told where to go for complications ¹⁷	47 (15.2)	2416 (17.5)	19.6	0.85 [0.58, 1.24]
Received 2+ tetanus injections ¹⁸	113 (36.4)	6868 (49.8)	16.4	0.58 [0.42, 0.80]
ANC quality score^h				
Low (0-5)	139 (45.3)	7115 (51.9)	19.5	1.40 [0.86, 2.29]
High (6-9)	23 (7.4)	1636 (11.9)	13.9	1.00 [Reference]
No ANC	145 (47.3)	4969 (36.2)	29.1	2.10 [1.34, 3.29]
PREGNANCY COMPLICATIONS				
Headache				
No	258 (83.0)	12102 (87.5)	21.3	1.00 [Reference]
Yes	53 (17.0)	1733 (12.5)	30.6	1.45 [1.00, 2.09]
Blurry vision				
No	279 (89.7)	12891 (93.2)	21.6	1.00 [Reference]
Yes	32 (10.3)	943 (6.8)	34.1	1.58 [1.05, 2.38]
Bleeding or spotting				
No	285 (91.7)	13300 (96.1)	21.4	1.00 [Reference]
Yes	26 (8.3)	534 (3.9)	48.4	2.26 [1.41, 3.60]
Probable hypertensionⁱ				
No	278 (89.5)	12827 (92.7)	21.7	1.00 [Reference]
Yes	33 (10.5)	1007 (7.3)	32.5	1.50 [0.97, 2.32]
Probable infection^j				
No	279 (89.7)	13079 (94.5)	21.3	1.00 [Reference]
Yes	32 (10.3)	755 (5.5)	42.6	2.00 [1.35, 2.96]
Anaemia or thin/weak blood				
No	291 (93.7)	13152 (4.9)	22.2	1.00 [Reference]
Yes	20 (6.3)	682 (4.9)	28.8	1.30 [0.83, 2.03]
Reduced or no fetal movement				
No	299 (96.1)	13684 (98.9)	21.8	1.00 [Reference]
Yes	12 (3.9)	150 (1.1)	81.5	3.73 [1.99, 7.94]
Too early contractions				
No	294.8 (94.8)	13424 (97.0)	22.0	1.00 [Reference]
Yes	16.3 (5.2)	410 (3.0)	39.8	1.81 [0.83, 3.95]
Abdominal pain				
No	276 (88.6)	12275 (88.7)	22.5	1.00 [Reference]
Yes	35 (11.4)	1559 (11.3)	22.7	1.01 [0.64, 1.59]
Fainted/unconsciousness				
No	304 (97.8)	13595 (98.3)	22.4	1.00 [Reference]
Yes	7 (2.2)	240 (1.7)	28.0	1.25 [0.59, 2.66]
DELIVERY CARE				
Birth attendant¹⁹				
Skilled provider ^c	112 (2.2)	4965 (36.1)	22.5	1.00 [Reference]
Unskilled provider ^f	189 (60.9)	8488 (61.7)	22.3	1.01 [0.70, 1.45]
No one	10 (3.2)	306 (2.2)	32.6	1.47 [0.71, 3.04]
Delivered in health facility²⁰				
No	201 (64.8)	9108 (66.0)	19.4	1.00 [Reference]

	Stillbirths	All births	Stillbirth rate per 1000	Unadjusted RR [95% CI]
	N (%)	N (%)		
Total pregnancy outcomes (weighted)	311 (2.2)	13 834 ^a (100)	22.5	
Yes	109 (35.2)	4702 (34.0)	23.0	1.05 [0.73, 1.51]
Mode of delivery^{c,21}				
Vaginal	263 (86.0)	12867 (94.2)	20.5	1.00 [Reference]
Caesarean section	19 (6.3)	238 (1.7)	80.6	3.95 [2.02, 7.69]
Instrumental (forceps or vacuum)	24 (7.8)	560 (4.1)	42.5	2.12 [1.33, 3.37]
COMPLICATIONS IN DELIVERY PERIOD				
Headache				
No	192 (61.7)	9023 (65.2)	19.4	1.00 [Reference]
Yes	119 (38.3)	4811 (34.8)	23.6	1.16 [0.82, 1.66]
Blurry vision				
No	238 (76.6)	11390 (82.3)	19.8	1.00 [Reference]
Yes	73 (23.4)	2444 (17.7)	25.6	1.42 [1.01, 1.99]
Excessive bleeding^c				
No	217 (69.7)	11907 (86.1)	17.1	1.00 [Reference]
Yes	94 (30.3)	1927 (13.9)	43.6	2.69 [2.07, 3.50]
Probable hypertension				
No	240 (77.2)	11594 (83.8)	20.7	1.00 [Reference]
Yes	71 (22.8)	2240 (16.2)	31.6	1.53 [1.05, 2.22]
Probable infection				
No	237 (76.1)	12039 (87.0)	19.7	1.00 [Reference]
Yes	74 (23.9)	1796 (13.0)	41.5	2.11 [1.59, 2.81]
Prolonged/obstructed labour/malpresentation				
No	271 (87.1)	12844 (92.8)	19.1	1.00 [Reference]
Yes	40 (12.9)	990 (7.2)	43.4	1.96 [1.33, 2.89]
Water broke too early				
No	280 (90.2)	13005 (94.0)	20.0	1.00 [Reference]
Yes	31 (9.8)	829 (6.0)	34.9	1.71 [1.03, 2.85]
Reduced or no fetal movement				
No	283 (91.1)	13678 (98.9)	18.9	1.00 [Reference]
Yes	28 (8.9)	157 (1.1)	185.	8.56 [5.51, 13.3]
Lower abdominal pain				
No	182 (58.5)	9568 (69.2)	19.0	1.00 [Reference]
Yes	129 (41.5)	4267 (30.8)	30.3	1.59 [1.24, 2.04]
Fainting/unconsciousness				
No	289 (93.0)	13202 (95.4)	21.9	1.00 [Reference]
Yes	22 (7.0)	632 (4.6)	34.7	1.58 [0.97, 2.58]

730 **Abbreviation:** ANC- antenatal care; RR- risk ratio

731 **Footnotes**

732 ^aN= 13 834 unless otherwise indicated

733 ^b Any education refers to any primary, secondary or higher level of education

734 ^c These variables were not included in the multivariable analyses. ANC variables not included due to multi-collinearity with quality of ANC.

735 Delivery assistant was not included due to collinearity with place of delivery. Mode of delivery not included as these are procedures might have occurred after the outcome. Severe bleeding during labour was not included as it was unknown if this was pre or post-partum

736 haemorrhage and may have occurred after the outcome.

737 ^d Includes first pregnancies

738 ^e Skilled/trained provider refers to doctor, nurse or midwife

739 ^f Unskilled/untrained provider refers to traditional birth attendant (TBA), Community health worker (CHW), relative or friend

740 ^g Reference category are those that did not receive the intervention

741 ^h ANC quality score calculated by number of components received out of a total of 9 components (1- weight taken, 2- blood pressure taken, 3- blood sample taken, 4- urine sample taken, 5- informed signs of pregnancy complications, 6- informed where to seek care for complications, 7- received 2+ tetanus injections, 8- received iron/folic acid, and 9- received anti-helminths)

742 ⁱ Probable hypertension was based on mother's report of convulsions/fits/shaking/eclampsia/pre-eclampsia and/or swelling/oedema

743 ^j Probable infection was based on mother's report of high fever and/or foul-smelling vaginal discharge

744 Missing values (unweighted observations): ¹ n=11; ² n=24 ³ n=9, ⁴ n=23, ⁵ n=16, ⁶ n=49, ⁷ n=122, ⁸ n=118, ⁹ n=22, ¹⁰ n=58, ¹¹ n=48

745 ¹² n=65, ¹³ n=69, ¹⁴ n=20, ¹⁵ n=133, ¹⁶ n=49, ¹⁷ n=10, ¹⁸ n=133, ¹⁹ n=78, ²⁰ n=29, ²¹ n=230

746

Table 2. Multivariable results of factors associated with stillbirth for women's most recent birth in the preceding three years, Afghanistan 2010

	Model 1: PRE-PREGNANCY Community+ socio-economic + maternal factors	Model 2: PREGNANCY Community + socioeconomic + maternal + pregnancy complications + ANC + biological	Model 3: DELIVERY TIME Community + socioeconomic + maternal + pregnancy complications + ANC + delivery care + delivery complications + biological
Factors	aRR [95%CI]	aRR [95%CI]	aRR [95%CI]
Region			
North-Eastern	1.00 [Reference]		1.00 [Reference]
Northern	0.70 [0.38, 1.26]		0.68 [0.38, 1.24]
Western	1.52 [0.84, 2.75]		1.20 [0.66, 2.20]
Central highlands	2.72 [1.16, 6.36]		3.01 [1.35, 6.70]
Capital	1.27 [0.71, 2.27]		1.14 [0.59, 2.29]
Eastern	1.31 [0.69, 2.49]		1.16 [0.51, 2.02]
Southern	1.20 [0.62, 2.34]		1.01 [0.45, 1.91]
South-Eastern	1.97 [1.10, 3.53]		1.59 [0.87, 2.89]
Ethnicity			
Tajik	1.00 [Reference]	1.00 [Reference]	1.00 [Reference]
Pashtun	1.07 [0.71, 1.60]	1.22 [0.86, 1.74]	1.07 [0.72, 1.60]
Hazara	1.13 [0.63, 2.03]	1.49 [0.88, 2.55]	1.03 [0.61, 1.75]
Uzbek	1.62 [0.85, 3.09]	1.21 [0.70, 2.09]	1.63 [0.86, 3.11]
Nuristani	10.39 [3.39, 31.86]	9.22 [3.49, 24.34]	9.15 [2.95, 28.74]
Pashai	1.65 [0.79, 3.45]	1.95 [0.94, 4.07]	1.78 [0.72, 4.37]
Baloch/Turkmen/Other	1.04 [0.50, 2.16]	0.92 [0.37, 2.27]	1.02 [0.48, 2.15]
Wealth Index			
Poorest	1.05 [0.65, 1.67]	0.96 [0.61, 1.49]	1.08 [0.64, 1.80]
Poorer	1.13 [0.74, 1.74]	1.08 [0.71, 1.64]	1.26 [0.77, 2.07]
Middle	1.10 [0.74, 1.62]	1.02 [0.69, 1.60]	1.16 [0.77, 1.75]
Richer	0.87 [0.58, 1.31]	0.87 [0.58, 1.30]	0.94 [0.62, 1.41]
Richest	1.00 [Reference]	1.00 [Reference]	1.00 [Reference]
Maternal education			
No education/Madrassa	1.59 [0.93, 273]	1.41 [0.83, 2.40]	1.50 [0.87, 2.58]
Any education	1.00 [Reference]	1.00 [Reference]	1.00 [Reference]
Previous pregnancy loss			
Yes ^b	2.61 [1.74, 3.91]	2.57 [1.77, 3.75]	2.43 [1.65, 3.59]
Maternal age (years)			
12-18	0.79 [0.40, 1.56]	0.80 [0.41, 1.59]	0.82 [0.42, 1.60]
19-24	1.00 [Reference]	1.00 [Reference]	1.00 [Reference]
25-34	1.33 [0.90, 1.96]	1.32 [0.86, 2.01]	1.36 [0.89, 2.08]
≥35	1.65 [1.02, 2.66]	1.58 [0.96, 2.62]	1.62 [0.99, 2.64]
Pregnancy order			
1st pregnancy	2.18 [1.46, 325]	2.33 [1.56, 3.47]	2.27 [1.52, 3.38]
2 nd -4 th pregnancy	1.00 [Reference]	1.00 [Reference]	1.00 [Reference]
≥5 th pregnancy	1.32 [0.92, 1.89]	1.35 [0.93, 1.96]	1.37 [0.95, 1.97]
ANC Quality Index			
High (6-9)		1.00 [Reference]	1.00 [Reference]
Low (0-5)		1.50 [0.94, 2.41]	1.56 [0.96, 2.53]
No ANC		2.77 [1.67, 4.61]	3.03 [1.73, 5.30]
Antepartum complication: Probable infection^c			
Yes ^b		2.25 [1.36, 3.09]	1.94 [1.29, 2.92]
Antepartum complication: Bleeding or spotting			
Yes ^b		2.25 [1.45, 3.49]	1.90 [1.19, 3.04]
Antepartum complication: Reduced or no fetal movement			
Yes		3.71 [1.94, 7.12]	2.06 [1.06, 3.97]
Antepartum complication: Headache			
Yes ^b		1.70 [1.23, 2.35]	1.67 [1.20, 2.33]
Delivery complication: Reduced or no fetal movement			
Yes ^b			6.82 [4.20, 11.10]
Delivered in health facility			
Yes ^b			1.55 [1.12, 2.16]
Sex of baby			
Female		1.00 [Reference]	1.00 [Reference]
Male		1.17 [0.89, 1.54]	1.16 [0.88, 1.52]
Multiple pregnancy			
Yes ^b		3.01 [1.60, 75.67]	3.19 [1.75, 5.80]
Area under the Curve (AUC)	0.66	0.69	0.73

Abbreviations: ANC – antenatal care, aRR – adjusted risk ratio, CI – confidence interval

Footnotes:

^a N presented is the weighted population and includes all cases with complete data (13 393 live births & 290 stillbirths).

^b Reference category for variables with yes/no responses is the “No” category

° Probable infection: if mother reported having symptoms of high fever and/or foul-smelling vaginal discharge

Table 3. Multivariable results of factors associated with intrapartum stillbirths for women's most recent birth in the preceding three years, Afghanistan 2010

	Model 1: PRE-PREGNANCY Community + socio-economic + environmental + maternal	Model 2: PREGNANCY PERIOD Community + socioeconomic + maternal + pregnancy complications + ANC + biological	Model 3: DELIVERY TIME Community + socioeconomic + maternal + pregnancy complications + ANC + delivery care + delivery complications + biological
Independent variables	aRR [95% CI]	aRR (95% CI)	aRR (95% CI)
N=13 577^a			
Ethnicity			
Tajik	1.00 [Reference]	1.00 [Reference]	1.00 [Reference]
Pashtun	1.25 [0.82, 1.90]	1.21 [0.80, 1.84]	1.17 [0.77, 1.78]
Hazara	1.21 [0.62, 2.38]	1.19 [0.61, 2.35]	1.17 [0.61, 2.27]
Uzbek	1.48 [0.83, 2.66]	1.59 [0.89, 2.86]	1.52 [0.82, 2.87]
Nuristani	12.55 [4.08, 38.66]	11.32 [3.71, 34.52]	11.13 [3.56, 34.80]
Pashai	2.81 [1.42, 5.56]	3.11 [1.50, 6.47]	2.92 [1.28, 6.64]
Baloch/Turkmen/Other	0.55 [0.20, 1.50]	0.57 [0.21, 1.57]	0.57 [0.21, 1.59]
Wealth index			
Poorest	0.99 [0.57, 1.70]	0.90 [0.53, 1.55]	0.89 [0.51, 1.53]
Poorer	1.08 [0.65, 1.79]	1.06 [0.64, 1.78]	1.04 [0.62, 1.77]
Middle	1.22 [0.72, 2.06]	1.19 [0.71, 1.98]	1.22 [0.73, 2.05]
Richer	0.84 [0.50, 1.42]	0.84 [0.52, 1.39]	0.85 [0.51, 1.53]
Richest	1.00 [Reference]	1.00 [Reference]	1.00 [Reference]
Maternal education			
No education/madrassa	1.89 [0.98, 3.66]	1.67 [0.86, 3.24]	1.70 [0.87, 3.32]
Any education	1.00 [Reference]	1.00 [Reference]	1.00 [Reference]
Maternal age			
12-18	0.80 [0.31, 2.06]	0.80 [0.31, 2.03]	0.83 [0.33, 2.12]
19-24	1.00 [Reference]	1.00 [Reference]	1.00 [Reference]
25-34	1.37 [0.86, 2.22]	1.34 [0.84, 2.15]	1.37 [0.86, 2.18]
≥35	1.65 [0.94, 2.92]	1.56 [0.87, 2.80]	1.61 [0.91, 2.87]
Pregnancy order			
1st pregnancy	2.10 [1.20, 3.70]	2.19 [1.25, 3.86]	2.19 [1.24, 3.88]
2 nd -4 th pregnancy	1.00 [Reference]	1.00 [Reference]	1.00 [Reference]
≥5 th pregnancy	1.29 [0.84, 1.99]	1.33 [0.86, 2.06]	1.34 [0.86, 2.07]
Previous pregnancy loss			
Yes ^b	2.91 [1.79, 4.72]	2.98 [1.87, 4.75]	2.91 [1.82, 4.65]
Quality of ANC			
High (6-9)		1.00 [Reference]	1.00 [Reference]
Low (0-5)		2.18 [1.04, 4.60]	2.17 [1.03, 4.57]
No ANC		3.55 [1.60, 7.88]	3.33 [1.56, 7.32]
Antepartum complication: Probable infection^c			
Yes ^b		2.02 [1.13, 3.62]	1.96 [1.09, 3.52]
Antepartum complication: Bleeding or spotting			
Yes		2.04 [1.06, 3.92]	
Antepartum complication: Headache			
Yes ^b		1.63 [1.05, 2.52]	1.63 [1.05, 2.52]
Delivery complication: Reduced or no fetal movements			
Yes ^b			8.15 [4.68, 14.18]
Sex of baby			
Female		1.00 [Reference]	1.00 [Reference]
Male		1.50 [1.02, 2.22]	1.51 [1.02, 2.22]
Multiple pregnancy			
Yes ^b		4.89 [2.12, 11.30]	4.96 [2.19, 11.24]
Area under the curve	0.65	0.70	0.72

Abbreviations: ANC – antenatal care, aRR – adjusted risk ratio, CI – confidence interval.

Footnotes:

^a N presented is the weighted population and includes all cases with complete data (13 393 live births and 184 stillbirths).

^b Reference category for variables with yes/no responses is the “No” category

^c Probable infection: if mother reported having symptoms of high fever and/or foul-smelling vaginal discharge