

This is a postprint version of the following published document:

González-Pérez, Marina; Gómez-Bombarelli, Rafael; Pérez-Prior, M. Teresa; Arenas-Valgañón, Jorge; García-Santos, M. Pilar; Calle, Emilio; Casado, Julio (2014). Alkylating Potential of Styrene Oxide: Reactions and Factors Involved in the Alkylation Process. *Chemical Research in Toxicology*, 27(10), 1853-1859.

DOI: <https://doi.org/10.1021/tx5002892>


© 2014 American Chemical Society.

# 1 Alkylating Potential of Styrene Oxide: Reactions and Factors 2 Involved in the Alkylation Process

3 Marina González-Pérez, Rafael Gómez-Bombarelli,<sup>‡</sup> M. Teresa Pérez-Prior,<sup>§</sup> Jorge Arenas-Valgañón,  
4 M. Pilar García-Santos, Emilio Calle, and Julio Casado\*

5 Departamento de Química Física, Facultad de Ciencias Químicas, Universidad de Salamanca, Plaza de los Caídos 1-5, E-37008  
6 Salamanca, Spain

7 **ABSTRACT:** The chemical reactivity of styrene-7,8-oxide (SO),  
8 an alkylating agent with high affinity for the guanine-N7  
9 position and a probable carcinogen for humans, with 4-(*p*-  
10 nitrobenzyl)pyridine (NBP), a trap for alkylating agents with  
11 nucleophilic characteristics similar to those of DNA bases, was  
12 investigated kinetically in water/dioxane media. UV-vis  
13 spectrophotometry and ultrafast liquid chromatography were  
14 used to monitor the reactions involved. It was found that in the  
15 alkylation process four reactions occur simultaneously: (a) the  
16 formation of a  $\beta$ -NBP-SO adduct through an  $S_N2$  mechanism;  
17 (b) the acid-catalyzed formation of the stable  $\alpha$ -NBP-SO adduct  
18 through an  $S_N2'$  mechanism; (c) the base-catalyzed hydrolysis of  
19 the  $\beta$ -adduct, and (d) the acid-catalyzed hydrolysis of SO. At 37.5  
20 °C and pH = 7.0 (in 7:3 water/dioxane medium), the values of the respective reaction rate constants were as follows:  $k_{\text{alk}\beta} = (2.1$   
21  $\pm 0.3) \times 10^{-4} \text{ M}^{-1} \text{ s}^{-1}$ ,  $k_{\text{alk}\alpha} = (1.0 \pm 0.1) \times 10^{-4} \text{ M}^{-1} \text{ s}^{-1}$ ,  $k_{\text{hyd}}^{\text{AD}} = (3.06 \pm 0.09) \times 10^{-6} \text{ s}^{-1}$ , and  $k_{\text{hyd}} = (4.2 \pm 0.9) \times 10^{-6} \text{ s}^{-1}$ .  
22 These values show that, in order to determine the alkylating potential of SO, none of the four reactions involved can be  
23 neglected. Temperature and pH were found to exert a strong influence on the values of some parameters that may be useful to  
24 investigate possible chemico-biological correlations (e.g., in the pH 5.81–7.69 range, the fraction of total adducts formed  
25 increased from 24% to 90% of the initial SO, whereas the adduct lifetime of the unstable  $\beta$ -adduct, which gives an idea of the  
26 permanence of the adduct over time, decreased from 32358 to 13313 min). A consequence of these results is that the conclusions  
27 drawn in studies addressing alkylation reactions at temperatures and/or pH far from those of biological conditions should be  
28 considered with some reserve.



## 29 ■ INTRODUCTION

30 The reactivity of styrene-7,8-oxide (SO) has been the subject of  
31 intense research from 1900 to the present, mainly due to its  
32 industrial applications and biological activity. This oxirane is  
33 widely used as a reactive plasticizer or diluent for epoxy resins,  
34 in the production of phenethyl alcohol and styrene glycol and  
35 its derivatives, and as a precursor for cosmetics, surface  
36 coatings, and agricultural and biological chemicals.<sup>1,2</sup> It is also  
37 the main *in vivo* mutagenic metabolite of styrene, a compound  
38 extensively used in the production of plastic, rubber, fiberglass,  
39 pipes, and food containers.<sup>3</sup> Thus, humans and other living  
40 organisms are exposed to SO, whose carcinogenic, neurotoxic,  
41 and mutagenic effects are well-known.<sup>4</sup>

42 SO is a potent alkylating agent with the capacity to form  
43 adducts *in vivo* and *in vitro* with DNA bases, mainly at the N7-  
44 position of guanine.<sup>5</sup> It has been classified as probable  
45 carcinogen for humans (2A IARC).<sup>2</sup>

46 In contrast to other epoxides that only react with  
47 nucleophiles through their less hindered carbon, SO can  
48 achieve this through the two electrophilic carbons of the  
49 oxirane ring, the primary ( $\beta$ ) and the secondary ( $\alpha$ ),<sup>5–10</sup> to  
50 form the corresponding adducts.

Regarding the chemical reactivity and biological effectiveness  
51 of SO with different nucleophiles, including DNA bases, there  
52 are abundant references in the literature.<sup>6,9,11–16</sup> In spite of this,  
53 many authors overlook some reactions that may be involved in  
54 the alkylation mechanism and, hence, in the alkylation efficacy  
55 of SO, such as (a) the potential formation of more than one  
56 adduct between SO and the alkylation substrate<sup>5–10,17–20</sup> and  
57 (b) the solvolysis reactions of the SO and of the adducts  
58 formed.<sup>19</sup> Accordingly, in the present study the chemical  
59 reactivity and alkylating potential of SO were investigated in  
60 depth. The molecule 4-(*p*-nitrobenzyl)pyridine (NBP), which  
61 has several advantages as an alkylation substrate,<sup>21</sup> was used as a  
62 nucleophilicity model for DNA. 63


## ■ EXPERIMENTAL PROCEDURES

64 Styrene oxide (99%), 4-(*p*-nitrobenzyl)pyridine (98%), and triethyl-  
65 amine (Et<sub>3</sub>N 99%) were purchased from Sigma-Aldrich (Steinheim,  
66 Germany). 1,4-Dioxane was from Panreac (Barcelona, Spain). 67

**Caution:** Styrene oxide is hazardous and should be handled  
68 carefully. 69

**Received:** July 15, 2014

## Scheme 1. Alkylation Mechanism of NBP by SO



70 To obtain deionize water, a Milli Q-Gradient (Millipore) system  
71 was used. Absorbance measurements were made with a Shimadzu UV-  
72 2401-PC spectrophotometer with a thermoelectric six-cell holder  
73 temperature control system ( $\pm 0.1$  °C). For pH measurements, a  
74 Metrohm 827 pH-meter with a glass electrode was used. The reaction  
75 temperature was kept constant ( $\pm 0.05$  °C) with a Lauda Ecoline  
76 RE120 thermostat. All kinetic runs were performed in triplicate.

77 The alkylation reaction of NBP by SO was carried out (i) in an  
78 excess of NBP to inhibit the formation of bis-alkylation adducts (1:2  
79 NBP:SO) and to work in pseudo-first-order conditions and (ii) in  
80 water/dioxane medium, to render the NBP soluble. Alkylation  
81 mixtures were prepared by adding 1.0 mL of 0.005–0.03 M epoxide  
82 stock solution (SO in dioxane) to 100 mL of NBP solution (0.01–0.02  
83 M) in water/dioxane medium. Acetate, phosphate, and borate buffers  
84 were used when needed to adjust the pH of the alkylation mixture.

85 Two different complementary techniques were used to investigate  
86 the formation and characterization of the SO-NBP adducts kinetically:

87 **Vis Spectrophotometry.** At different times, aliquots (2.4 mL) of  
88 the alkylation mixture (SO + NBP in a water/dioxane medium) were  
89 collected and placed in a cuvette containing 0.6 mL of  $\text{Et}_3\text{N}$  to induce  
90 the deprotonation of the uncolored adducts ( $\text{AD}_{\text{un}\alpha}$  and  $\text{AD}_{\text{un}\beta}$ ),  
91 yielding the SO-NBP colored adducts ( $\text{AD}_\alpha$  and  $\text{AD}_\beta$ ; Scheme 1),  
92 whose absorbance was measured at  $\lambda = 560$  nm, where only both  
93 adducts absorb. Detailed experimental conditions are given in the  
94 figure and table legends. Rate constants were obtained by nonlinear  
95 regression analysis of the absorbance/time data.

96 **HPLC.** A gradient-controlled Shimadzu ultrafast liquid chromatog-  
97 raphy system equipped with a diode array detector (UFLC-DAD) was  
98 used to monitor the formation of each adduct. Chromatographic  
99 separation was achieved with a Mediterranean Sea C18 column (25 cm  
100  $\times$  1 cm, 5  $\mu\text{m}$ ). Mobile phase A was acetate buffer (pH = 4.75 and 0.1  
101 M), and mobile phase B was acetonitrile. 10% B was held for 5 min,  
102 after which a gradient from 10% B up to 16% B was run over 8 min.  
103 Then, a gradient of up to 80% B was run over 1 min. 80% B was held  
104 for 14 min. The flow rate was set at 1.0 mL  $\text{min}^{-1}$ , and 200  $\mu\text{L}$  of the  
105 alkylation mixture was injected.

106 The hydrolysis reaction of SO was monitored by UV spectropho-  
107 tometry. Aliquots of 50–200  $\mu\text{L}$  of the stock solution of the epoxide in  
108 dioxane (0.085 M) were added to a thermostated cuvette containing  
109 3.0 mL of the reaction mixture ( $\text{HClO}_4$  + water + dioxane) at a  
110 constant pH. To convert the pH-meter output obtained in water/  
111 dioxane mixtures into hydrogen ion concentrations, it is necessary to  
112 use a correction factor.<sup>22–24</sup> However, with the solutions studied in  
113 this work (7:3 water:dioxane and  $[\text{HClO}_4] = 0.1\text{--}0.02$  M), this  
114 correction can be neglected.<sup>24</sup>

115 Absorbance at  $\lambda = 226$  nm, where the difference between the signals  
116 of SO and its hydrolysis product (styrene glycol, SG)<sup>16</sup> is maximum,  
117 was measured until no change was observed.


## RESULTS AND DISCUSSION

118

**Reaction Mechanism.** In order to determine the reactions, 119  
besides SO hydrolysis, involved in the alkylation mechanism, 120  
UFLC-DAD analysis was performed. The results obtained 121  
revealed the formation of two SO–NBP adducts with different 122  
retention times (see Scheme 1). This result is in agreement 123  
with their distinctive fragmentation patterns, obtained by 124  
analysis of the alkylation mixtures by mass spectrometry in an 125  
excess of NBP ( $m/z = 335, 317, 304$  for  $\alpha$ -adduct, and  $m/z =$  126  
 $335, 121$  for  $\beta$ -adduct), as well as with previous results obtained 127  
with other nucleophiles.<sup>6,13,19,20,25</sup> 128

Assignment of the UFLC-DAD peaks was performed by 129  
comparison of their retention times, UV–vis spectra, and 130  
kinetic profiles with those of the  $\beta$ -adduct formed between 131  
NBP and *p*-nitrostyrene oxide, a substituted styrene oxide 132  
whose alkylating potential has been studied by us previously.<sup>26</sup> 133  
The adducts identified were  $\alpha$ -NBP–SO ( $\text{AD}_{\text{un}\alpha}$ ),  $t_R = 11.0$  134  
min, and  $\beta$ -NBP–SO ( $\text{AD}_{\text{un}\beta}$ ),  $t_R = 8.5$  min (see Scheme 1). 135  
These two types of adduct also result in alkylation at the N-7 136  
position in the reaction between SO and guanosine, 137  
deoxyguanosine, and the single- or double-stranded DNA *in* 138  
*vitro* under physiological conditions.<sup>5,7,8,18</sup> 139

Figure 1 shows the profile of a typical kinetic run with 140  
UFLC-DAD, carried out by monitoring the area of the  $\text{AD}_{\text{un}\alpha}$  141




**Figure 1.** Kinetic profiles obtained by monitoring the variation in the  
UFLC-DAD peak area,  $\text{AD}_{\text{un}\beta}$  (gray circles) and  $\text{AD}_{\text{un}\alpha}$  (black circles),  
over time in 7:3 (v:v) water/dioxane media.  $T = 37.5$  °C; pH = 7.50;  
 $[\text{SO}]_0 = 1.7 \times 10^{-4}$  M;  $[\text{NBP}]_0 = 2.0 \times 10^{-2}$  M;  $\lambda = 254$  nm.

and  $\text{AD}_{\text{un}\beta}$  peaks over time. As can be observed, while the 142  
 $\text{AD}_{\text{un}\alpha}$  signal remains fairly stable along the reaction time, 143  
 $\text{AD}_{\text{un}\beta}$  undergoes hydrolysis. 144

It should be pointed out that depending on the reaction rates 145  
involved in the NBP alkylation process, two different 146

Scheme 2. Reactions Involved in NBP Alkylation by SO in Aquo–Organic Media



147 approaches can be used: (i) considering only the alkylation  
 148 reaction; (ii) taking into account all the reactions occurring  
 149 simultaneously. Since abundant instances of the use of a poor  
 150 mechanistic framework to determine the alkylating potential of  
 151 SO can be found in the literature,<sup>5–10,17–20</sup> in our case the  
 152 second approach was applied (Scheme 2) and included the  
 153 following: (a) the formation of the  $\alpha$ -NBP–SO and  $\beta$ -NBP–  
 154 SO adducts; (b) hydrolysis of the  $\beta$ -adduct (the  $\alpha$ -adduct  
 155 remained stable along the reaction); and (c) SO hydrolysis.  
 156 Equations 1–3 can be readily deduced from the mechanism  
 157 depicted in Scheme 2. In these equations,  $k_{\text{alk}}$  is the global  
 158 alkylation rate constant, defined as the sum of the second-order  
 159 rate constants for alkylation through the  $\alpha$  and  $\beta$  carbons,  $k_{\text{alk}\alpha}$   
 160 and  $k_{\text{alk}\beta}$ , respectively. Since NBP is present in a large excess, its  
 161 concentration can be assumed to be constant, i.e.  $[\text{NBP}] =$   
 162  $[\text{NBP}]_0$ .

$$163 \quad r = \frac{d[\text{AD}_{\text{un}\alpha}]}{dt} = k_{\text{alk}\alpha}[\text{NBP}][\text{SO}] \quad (1)$$

$$164 \quad r = \frac{d[\text{AD}_{\text{un}\beta}]}{dt} = k_{\text{alk}\beta}[\text{NBP}][\text{SO}] - k_{\text{hyd}}^{\text{AD}}[\text{AD}_{\text{un}\beta}] \quad (2)$$

$$165 \quad r = \frac{-d[\text{SO}]}{dt} = [\text{SO}](k_{\text{alk}}[\text{NBP}] + k_{\text{hyd}}) \quad (3)$$

166 Rate constants were determined using the NBP test<sup>21</sup> (see  
 167 Scheme 1) by monitoring the variation in absorbance at  $\lambda =$   
 168 560 nm (see above).

169 Absorbance at time  $t$  ( $A_{\text{AD}}$ ) can be expressed as the sum of  
 170 the absorbances of the colored  $\beta$ - and  $\alpha$ -adducts:

$$171 \quad A_{\text{AD}} = A_{\text{AD}\beta} + A_{\text{AD}\alpha} \quad (4)$$

172 The absorbance of each adduct at time  $t$  was obtained from  
 173 eqs 1 and 2 by applying the Lambert–Beer law to the  
 174 respective integrated rate equations in terms of concentration:

$$175 \quad A_{\text{AD}\alpha} = \frac{\epsilon_{\alpha} k_{\text{alk}\alpha} [\text{NBP}] [\text{SO}]_0}{k_{\text{alk}} [\text{NBP}] + k_{\text{hyd}}} (1 - e^{-(k_{\text{alk}} [\text{NBP}] + k_{\text{hyd}}) t}) \quad (5)$$

$$A_{\text{AD}\beta} = \frac{\epsilon_{\beta} k_{\text{alk}\beta} [\text{NBP}] [\text{SO}]_0}{(k_{\text{alk}} [\text{NBP}] + k_{\text{hyd}}) - k_{\text{hyd}}^{\text{AD}}} (e^{-(k_{\text{hyd}}^{\text{AD}}) t} - e^{-(k_{\text{alk}} [\text{NBP}] + k_{\text{hyd}}) t}) \quad (6) \quad 176$$

Equation 4 can thus be written as follows: 177

$$A_{\text{AD}} = \frac{a}{b - c} (e^{-ct} - e^{-bt}) + \frac{d}{b} (1 - e^{-bt}) \quad (7) \quad 178$$

with  $a$ ,  $b$ ,  $c$ , and  $d$  being parameters obtained by the nonlinear  
 179 fitting of the  $A_{\text{AD}}/t$  data and defined as 180


$$a = k_{\text{alk}\beta} [\text{NBP}] [\text{SO}]_0 \epsilon_{\beta} l; \quad b = k_{\text{alk}} [\text{NBP}] + k_{\text{hyd}}; \\ c = k_{\text{hyd}}^{\text{AD}}; \quad d = k_{\text{alk}\alpha} [\text{NBP}] [\text{SO}]_0 \epsilon_{\alpha} l \quad (8) \quad 181$$

The goodness of the data fit to eq 7, where the parameters  $a$   
 182 and  $d$  include a first-order reaction dependence on  $[\text{SO}]$  and  $183$   
 $[\text{NBP}]$ , while the parameter  $b$  includes a first-order dependence  
 184 on  $[\text{NBP}]$ , reveals that the formation of the  $\alpha$ - and  $\beta$ -adducts  
 185 occurs through  $\text{S}_{\text{N}}2'$  and  $\text{S}_{\text{N}}2$  mechanisms, as expected.<sup>6</sup> This,  
 186 together with the preference of SO for the N7-position of  
 187 guanine over other nucleophile positions of the DNA  
 188 bases,<sup>7,8,13,18,27</sup> suggests that NBP is a suitable model to  
 189 investigate the reactivity of SO with DNA.<sup>21</sup> 190

The good fitting of the experimental  $A_{\text{AD}}/t$  data obtained  
 191 under neutral conditions (Figure 2) to eq 7 supports the  
 192 proposed mechanism. 193

**Determination of the Kinetic Parameters Involved in  
 the NBP Alkylation Process.** As regards the global alkylation  
 195 rate constant,  $k_{\text{alk}}$  (eq 3), the value of this was obtained from  
 196 the slope of the  $b/[\text{NBP}]$  plot ( $b = k_{\text{alk}} [\text{NBP}] + k_{\text{hyd}}$ ),  $k_{\text{alk}} =$   
 197  $(3.1 \pm 0.4) \times 10^{-4} \text{ M}^{-1} \text{ s}^{-1}$  (37.5 °C, pH 7.0). Equation 9 was  
 198 used to determine  $k_{\text{alk}\beta} = (2.1 \pm 0.3) \times 10^{-4} \text{ M}^{-1} \text{ s}^{-1}$ , and  
 199 hence  $k_{\text{alk}\alpha} = (1.0 \pm 0.1) \cdot 10^{-4} \text{ M}^{-1} \text{ s}^{-1}$  (37.5 °C, pH 7.0),  
 200 assuming the ratio  $k_{\text{alk}\alpha}/k_{\text{alk}\beta} = 33/67$  (obtained previously for  
 201 the reaction of SO with deoxyguanosine in water/ethanol,  
 202 50:50 (v:v), at 37.0 °C)<sup>19</sup> for the alkylation of NBP by SO at  
 203 37.5 °C. This value was accepted, since it is known that in  
 204 protic solvents the  $k_{\text{alk}\alpha}/k_{\text{alk}\beta}$  ratio is solvent-independent<sup>11</sup> and  
 205 is similar for the alkylation of substrates analogous to 206





**Figure 2.** Experimental  $A_{AD}/t$  kinetic profiles in the formation of the SO–NBP adducts in 7:3 water/dioxane media (black circles), nonlinear fitting to eq 7 (dashed line), and the corresponding profiles calculated from eqs 5 and 6 for the  $\alpha$ -NBP–SO (black line) and  $\beta$ -NBP–SO (gray line) adducts, respectively.  $\lambda = 560$  nm;  $T = 37.5$  °C; pH = 7.0;  $[NBP]_0 = 1.99 \times 10^{-2}$  M;  $[SO]_0 = 7.69 \times 10^{-5}$  M.

207 guanosine<sup>18–20</sup> by *p*-substituted styrene oxides at the same  
208 temperature.

$$k_{alk\beta} = \frac{k_{alk}}{(k_{alk\alpha}/k_{alk\beta}) + 1} \quad (9)$$

210 With the values of the alkylation rate constants and those of  
211 the parameters  $a$  and  $d$ , the values of  $\varepsilon_\beta$  and  $\varepsilon_\alpha$  ( $\lambda = 560$  nm)  
212 were obtained (eqs 10 and 11):  $\varepsilon_\beta = (8 \pm 1) \times 10^3$  M<sup>-1</sup> cm<sup>-1</sup>  
213 and  $\varepsilon_\alpha = (40 \pm 8) \times 10^3$  M<sup>-1</sup> cm<sup>-1</sup>, respectively.

$$\varepsilon_\beta = \frac{a}{k_{alk\beta}I[SO]_0[NBP]} \quad (10)$$

$$\varepsilon_\alpha = \frac{d}{k_{alk\alpha}I[SO]_0[NBP]} \quad (11)$$

216 Regarding the SO hydrolysis reaction, its rate constant was  
217 obtained: (i) indirectly, by using the values of parameters  $b$  and  
218  $k_{alk}$  (eq 12), and (ii) directly, by spectrophotometric measure-  
219 ments (see Experimental Procedures).

$$k_{hyd} = b - k_{alk}[NBP] \quad (12)$$


221 Since the absorbance,  $A$ , of the hydrolysis reaction mixture at  
222 time  $t$  is the sum of the resulting styrene glycol (SG)<sup>16</sup> and  
223 epoxide contributions, eq 13 can be written.

$$A = I[SO]_0(\varepsilon_{SG} + (\varepsilon_{SO} - \varepsilon_{SG})e^{-k_{hyd}t}) \quad (13)$$

225 Excellent fitting of the experimental data to eq 13 was  
226 obtained (Figure 3a). In addition, the SO hydrolysis rate  
227 constants determined from eq 13 revealed a high degree of  
228 convergence with respect to those calculated from the  $b$   
229 parameter (Figure 3b). This supports the methodology used to  
230 determine the reaction rate constants involved in the alkylation  
231 process.

232 The values of the  $\beta$ -NBP–SO adduct hydrolysis rate  
233 constant,  $k_{hyd}^{AD}$  (see Table 1), are given directly by parameter  $c$   
234 (eq 8).

235 The values of the fitting parameters  $a$ ,  $b$ ,  $c$ , and  $d$ , as well as  
236 those of  $k_{alk\beta}$ ,  $k_{alk\alpha}$ ,  $k_{hyd}$ , and  $k_{hyd}^{AD}$  were used to determine some  
237 associated kinetic parameters that may be useful to investigate  
238 possible correlations between chemical reactivity and biological  
239 activity. Three such useful parameters are the fraction of the  
240 alkylating agent that eventually forms the  $\alpha$ - and  $\beta$ -adducts,  $f_\alpha$   
241 and  $f_\beta$ , respectively (eqs 14 and 15), and the adduct lifetime,  
242 AL. The sum  $f = f_\alpha + f_\beta$  ranges from 0 to 1, with 0 meaning that



**Figure 3.** (a) Kinetic profile of a typical kinetic run for SO hydrolysis (experimental data (circles) and fitting to eq 13 (continuous line)). (b) Variation in the SO hydrolysis rate constant with the acidity of the medium in 7:3 (v:v) water/dioxane media obtained directly (circles) and indirectly (squares) at 25.0 °C (gray circles and squares) and 37.5 °C (black circles and squares), respectively.

**Table 1.** Influence of Temperature on the Rate Constants of the Reactions Involved in the Alkylation of NBP by SO and Values of Some of the Associated Kinetic Parameters<sup>a</sup>

$T$ (°C)	25.0	27.5	30.0	32.5	35.0	37.5
$10^4 k_{alk\beta}$ (M <sup>-1</sup> s <sup>-1</sup> )	0.7	0.9	1.1	1.3	1.5	2.1
$10^4 k_{alk\alpha}$ (M <sup>-1</sup> s <sup>-1</sup> )	0.2	0.3	0.5	0.4	0.7	1.1
$10^4 k_{alk}$ (M <sup>-1</sup> s <sup>-1</sup> )	0.9	1.2	1.6	1.7	2.2	3.2
$10^6 k_{hyd}$ (s <sup>-1</sup> )	1.0	0.8	0.8	0.8	2.8	4.2
$10^6 k_{hyd}^{AD}$ (s <sup>-1</sup> )	0.5	0.6	1.0	1.5	1.9	3.1
$f_\beta$	0.50	0.57	0.54	0.62	0.42	0.40
$f_\alpha$	0.13	0.18	0.24	0.19	0.19	0.20
$f$	0.63	0.75	0.79	0.81	0.61	0.60
AL $_\beta$ (min)	17377	15209	8419	6810	3622	2185

<sup>a</sup>7:3 (v:v) water/dioxane media; pH = 7.0.

all the initial epoxide is hydrolyzed and with 1 meaning that the  
243 epoxide reacts exclusively with the NBP, yielding alkylation  
244 adducts.<sup>21</sup> These parameters have been correlated previously  
245 with the mutagenic potency of different compounds.<sup>21,26,28–31</sup>  
246

247 The adduct lifetime gives an idea of the permanence of the  
248 adduct over time. Thus, it is only meaningful for the unstable  $\beta$ -  
249 adduct. AL $_\beta$  is defined as the area under the curve  $[AD]_\beta/t$  per  
250 unit of alkylating agent concentration (eq 16).<sup>21,26,28–31</sup>  
251 Consequently, it can be calculated from the area under the  
252 curve  $A_{AD\beta}/t$  (i.e., Figure 2), as shown in eq 16. Calculation of  
253 this parameter is based on the rate constants of the three  
254 reactions involved,  $k_{hyd}$ ,  $k_{alk}$ , and  $k_{hyd}^{AD}$ . As may be seen from eq  
255 16, AL $_\beta$  is also related to the alkylating efficacy,  $f_\beta$  (i.e., the  
256 higher the  $f_\beta$  value and the lower the adduct hydrolysis rate  
257 constant, the higher the AL $_\beta$ ).

$$f_\alpha = \frac{k_{alk\alpha}[NBP]}{(k_{alk}[NBP] + k_{hyd})} = \frac{k_{alk\alpha}[NBP]}{b} \quad (14)$$

$$f_\beta = \frac{k_{alk\beta}[NBP]}{(k_{alk}[NBP] + k_{hyd})} = \frac{k_{alk\beta}[NBP]}{b} \quad (15)$$

$$AL_\beta = \frac{\int_{t=0}^{t=\infty} [AD]_\beta dt}{[SO]_0} = \frac{\int_{t=0}^{t=\infty} A_{AD\beta} dt}{\varepsilon_\beta I [SO]_0} \\ = \frac{k_{alk\beta}[NBP]}{(k_{alk}[NBP] + k_{hyd})k_{hyd}^{AD}} = \frac{f_\beta}{k_{hyd}^{AD}} = \frac{f_\beta}{c} \quad (16)$$

261 **Influence of Temperature on the Kinetic Parameters**  
 262 **Involved in the Alkylation Process.** Table 1 shows the  
 263 values of the above parameters determined in the 25.0–37.5 °C  
 264 range.

265 As can be observed, whereas the values of  $f_\alpha$  and  $f_\beta$  do not  
 266 show any clear trend with temperature,  $AL_\beta$  is strongly  
 267 influenced by it, being higher at lower temperatures. This  
 268 means that the stability of the  $\beta$ -adduct is higher at lower  
 269 temperatures; that is, a greater accumulation of the  $\beta$ -adduct  
 270 occurs with time.

271 It can also be seen that the  $k_{alk\beta}/k_{alk\alpha}$  ratio is inversely related  
 272 to temperature, showing an increase from 67/33 to 78/22 with  
 273 a decrease in temperature from 37.5 to 25.0 °C. This trend,  
 274 which seems to be common for the reaction of SO with  
 275 nucleophiles similar to NBP in protic solvents,<sup>6</sup> highlights the  
 276 relevant role played by temperature on the alkylating capacity  
 277 of SO. This result also suggests that any conclusions drawn  
 278 about the chemico-biological correlations established from  
 279 alkylation reactions with styrene oxide at temperatures far  
 280 from those of biological conditions should be taken with some  
 281 reserve.

282 The activation parameters calculated with the Arrhenius and  
 283 Eyring–Wynne–Jones equations,  $E_{a\beta} = (64 \pm 4) \text{ kJ mol}^{-1}$ ;  $E_{a\alpha}$   
 284  $= (99 \pm 5) \text{ kJ mol}^{-1}$ ;  $\Delta G_{\beta(25.0\text{ °C})}^\ddagger = (96 \pm 8) \text{ kJ mol}^{-1}$ ; and  
 285  $\Delta G_{\alpha(25.0\text{ °C})}^\ddagger = (99 \pm 19) \text{ kJ mol}^{-1}$ , revealed that the formation  
 286 of the activated intermediate state of the  $\alpha$ -adduct is enthalpy-  
 287 controlled whereas that of the  $\beta$ -adduct is not.

288 **Influence of the Acidity of the Medium on the Kinetic**  
 289 **Parameters Involved in the Alkylation Process.** A series of  
 290 experiments was performed in media of different acidities in the  
 291 5.0 < pH < 8.0 range, because experimental limitations did not  
 292 allow the reaction rate constants from eqs 7 and 8 to be  
 293 determined outside this range. (Since at pH < 5.0 the values of  
 294 the hydrolysis rate constant are equal to or higher than those of  
 295 the pseudo-first-order alkylation rate constant, calculation of  
 296 the individual values is not possible. At pH > 7.7 some  
 297 unidentified byproducts appear.) For the reaction at the  $\beta$   
 298 carbon of SO, it was observed that the alkylation rate constant  
 299 was not subject to any influence of pH. The  $\beta$ -NBP–SO adduct  
 300 was significantly hydrolyzed in neutral aquo-organic media  
 301 (Table 2), whereas the hydrolysis of the  $\beta$ -adduct is basic-  
 302 catalyzed and can be considered negligible in acid aquo-organic  
 303 media (Table 2), as occurs with other oxiranes.<sup>26,30</sup> In contrast,  
 304 alkylation through the  $\alpha$  carbon is acid-catalyzed (Table 2),

with the protonated transition state probably being favored by 305  
 the positive charge stabilization on that atom,<sup>9,32</sup> such that the 306  
 $\alpha$ -NBP–SO adduct formed remains stable along the reaction 307  
 (Figure 2). It should be noted that although consideration of 308  
 the stabilities of the adducts formed by SO is important, in the 309  
 literature they are usually overlooked.<sup>7,27,33</sup> 310

The hydrolysis of styrene oxide is also influenced by the 311  
 acidity of the medium. In the pH 2–8 range, the rate constant 312  
 $k_{hyd}$  can be expressed as the sum of the spontaneous and 313  
 catalytic hydrolysis rate constants (eq 17). 314

$$k_{hyd} = k_{hyd}^o + k_{H^+}[H^+] \quad (17) \quad 315$$

In 7:3 water/dioxane medium the values of  $k_{hyd}$  at 37.5 and 316  
 25.0 °C are given by the expressions  $k_{hyd37.5} = (2.5 \pm 0.8) \times$  317  
 $10^{-6} + (18 \pm 1) [H^+] \text{ (s}^{-1}\text{)}$  and  $k_{hyd25.0} = (1.0 \pm 0.1) \times$  318  
 $10^{-6} + (6.5 \pm 0.5) [H^+] \text{ (s}^{-1}\text{)}$  (Figure 3b). 319

The reaction order with respect to  $[H^+]$ , determined in pH < 320  
 5 acidic media at 25.0 and 37.5 °C, was found to be one. 321

The importance of the concurrent SO hydrolysis in the study 322  
 of NBP alkylation by the epoxide is such that at pH < 5 we 323  
 observed that the value of the above parameter  $b = k_{alk}[NBP] +$  324  
 $k_{hyd}$  gave the value of  $k_{hyd}$  directly because it was significantly 325  
 higher than that of  $k_{alk}[NBP]$ . In spite of this, in the literature it 326  
 is unusual to find studies that take SO hydrolysis into account 327  
 (see above). 328

As can be observed (Table 2), all the associated kinetic 329  
 parameters are clearly influenced by the acidity of the medium, 330  
 except  $f_\alpha$ . This means that a small fraction of the initial SO, 331  
 around 9–13%, is converted into  $\alpha$ -adduct regardless of the pH 332  
 of the medium. Additionally, the fraction of  $\beta$ -adduct formed is 333  
 substantially influenced by pH and is higher than that of the  $\alpha$ - 334  
 adduct. Thus, the fraction of total adducts formed increases 335  
 from 24% to 90% of the initial SO in the pH 5.81–7.69 range. 336  
 This means that in acid media SO hydrolysis is predominant, 337  
 whereas in neutral–basic media adduct formation predom- 338  
 inates, particularly that of the  $\beta$ -adduct (the  $k_{alk\beta}/k_{alk\alpha}$  ratio 339  
 increases from 63/37 to 93/7 due to an increase in the pH of 340  
 the medium from 5.8 to 7.7). This is important because it could 341  
 be related to the alkylating activity of styrene oxide in the 342  
 neutral biological medium. The results highlight the importance 343  
 of choosing the appropriate kinetic approach and of considering 344  
 all the reactions involved in the alkylation process. Neglecting 345  
 them would lead to mistaken conclusions about the SO 346  
 alkylating potential. Despite this, scrutiny of the literature 347  
 shows how frequently this is the case;<sup>5–10,17–20</sup> that is, when 348  
 using the NBP test or another spectroscopic method, the 349  
 increase in absorbance at a fixed time is considered to be a 350  
 measure of the reactivity or alkylating potential of SO<sup>34–37</sup> (this 351  
 alkylating potential has sometimes even been used to establish 352  
 correlations between chemical and biological activity).<sup>34,36</sup> 353

Another interesting result is that although the  $f_\beta$  value 354  
 declines with the acidity of the medium,  $\beta$ -adduct stability 355  
 clearly increases with it. 356


A consequence of the above is the strong influence of the 357  
 acidity of the medium on the reactivity of styrene oxide as an 358  
 alkylating agent, which underscores the importance of this 359  
 analysis in the search for possible chemico-biological correla- 360  
 tions. 361

**Influence of the Composition of the Reaction** 362  
**Medium on the Alkylation Process.** Figure 4a shows the 363 f4  
 strong dependence of  $k_{alk\beta}$  and  $k_{alk\alpha}$  on the dielectric constant, 364  
 $D$ , of the reaction medium in different water/dioxane mixtures, 365

Table 2. Influence of pH on the Rate Constants of the Reactions Involved in the Alkylation of NBP by SO and Associated Kinetic Parameters<sup>a</sup>

pH	5.29	5.81	6.57	6.95	7.27	7.69
$10^5 k_{alk\beta} \text{ (M}^{-1} \text{ s}^{-1}\text{)}$	7.30	7.15	7.09	6.98	7.29	7.20
$10^5 k_{alk\alpha} \text{ (M}^{-1} \text{ s}^{-1}\text{)}$		4.27	1.94	1.81	0.71	0.58
$10^5 k_{alk} \text{ (M}^{-1} \text{ s}^{-1}\text{)}$		11.42	9.03	8.79	8.00	7.78
$10^6 k_{hyd} \text{ (s}^{-1}\text{)}$	20.89	7.08	2.03	1.05	0.89	0.19
$10^7 k_{hyd}^{AD} \text{ (s}^{-1}\text{)}$		0.79	2.59	4.81	8.44	10.43
$f_\beta$	0.07	0.15	0.37	0.50	0.59	0.83
$f_\alpha$		0.09	0.10	0.13	0.06	0.07
$f$		0.24	0.47	0.63	0.65	0.90
AL (min)		32358	23874	17377	11635	13313

<sup>a</sup>7:3 water/dioxane media;  $T = 25.0\text{ °C}$ .



**Figure 4.** (a) Variation in the alkylation rate constants (37.5 °C) with the dielectric constant of the reaction medium, and (b) isokinetic correlations in the formation of both the  $\alpha$ - and  $\beta$ -adducts (black squares and black circles, respectively) in media of different composition.

with the alkylation rate increasing upon increasing  $D$ . This behavior, consistent with a reaction between polar molecules in which the activated complex is more polar than the reagents,<sup>38</sup> supports the proposed mechanism.

As is known,<sup>39</sup> the existence of an isokinetic relationship can serve to defend the argument that the reactions of a series share a common mechanism. The members may differ in the identities of a functional group, the length of a side chain, the composition of the solvent, and so on. In this work we considered the solvent composition, that is, the percentage of dioxane in the reaction medium. A mathematical formulation of the isokinetic effect is the linear relationship between two series of  $\log k_{alk}$  values measured at two temperatures  $T_1$  and  $T_2$ :  $\log k_{alk}(T_2) = a + b \log k_{alk}(T_1)$ .

It should be noted that the meaning of the isokinetic relationship is the existence of a *compensation effect* between the values of the enthalpy,  $\Delta H^\ddagger$ , and entropy of activation,  $\Delta S^\ddagger$ , such that the Gibbs free energy of activation,  $\Delta G^\ddagger$ , is approximately constant. The results shown in Figure 4b support the idea of common mechanisms of formation of both the  $\alpha$ - and  $\beta$ -SO-NBP adducts, respectively bimolecular nucleophilic  $S_N2'$  and  $S_N2$  substitutions, across the whole composition range of the solvent mixture.

## CONCLUSIONS

- NBP alkylation reactions through the  $\alpha$  and  $\beta$  carbons of SO occur via  $S_N2'$  and  $S_N2$  mechanisms, respectively.
- Neglecting some of the reactions involved in the alkylation process would lead to mistaken conclusions about the alkylating potential of SO.
- Temperature and pH are crucial factors because: (a) The lifetime of the unstable  $\beta$ -adduct is strongly influenced by temperature, and hence, the accumulation of this adduct is greater at lower temperatures; (b) The hydrolysis of the  $\beta$ -NBP-SO adduct is basic-catalyzed. The formation of the  $\alpha$ -NBP-SO adduct is acid-catalyzed, with the  $\alpha$ -NBP-SO adduct remaining stable along the reaction. The hydrolysis of SO is acid-catalyzed.
- Any conclusions drawn in studies addressing alkylation reactions at temperature and/or pH far from those found in biological conditions should be considered with some reserve when searching for chemico-biological correlations.
- The existence of an isokinetic relationship between alkylation rate constants in different water/dioxane ratios reveals a compensation effect that supports the idea of

common mechanisms of formation of both adducts across the whole composition range of the solvent mixture.

## AUTHOR INFORMATION

### Corresponding Author

\*Phone: (+34) 923 294486. Fax: (+34) 923 294574. E-mail: jucali@usal.es.

### Present Addresses

<sup>‡</sup>(R.G.-B.) Department of Chemistry and Chemical Biology, Harvard University, Cambridge, MA 02138, USA.

<sup>§</sup>(T.P.-P.) Departamento de Ciencia e Ingeniería de Materiales e Ingeniería Química, Universidad Carlos III de Madrid Avenida de la Universidad 30, E-28911 Leganés, Madrid, Spain.

### Funding

Thanks are given to the Spanish Ministerio de Economía y Competitividad and the European Regional Development Fund (Project CTQ2010-18999) for supporting the research reported in this article. M.G.-P. and J.A.-V. thank the Junta de Castilla y León-FEDER Funds for Ph.D. grants, and R.G.-B. thanks the Spanish Ministerio de Educación for a Ph.D. fellowship.

### Notes

The authors declare no competing financial interest.

## ACKNOWLEDGMENTS

The authors thank Dr. J. Bolton and the reviewers of the manuscript for their valuable comments.

## ABBREVIATIONS

$A_{AD}$ , sum of the absorbances of the colored  $\beta$ - and  $\alpha$ -adducts at time  $t$ ;  $AD_\omega$  colored  $\alpha$ -NBP-SO adduct;  $AD_\beta$  colored  $\beta$ -NBP-SO adduct;  $AD_{un\omega}$  uncolored  $\alpha$ -NBP-SO adduct;  $AD_{un\beta}$  uncolored  $\beta$ -NBP-SO adduct; AL, adduct lifetime;  $D$ , dielectric constant;  $f_\alpha$  and  $f_\beta$ , fraction of the alkylating agent that eventually forms the  $\alpha$ - and  $\beta$ -adducts, respectively; NBP, 4-(*p*-nitrobenzyl)pyridine; SG, styrene glycol; SO, styrene-7,8-oxide; UFLC-DAD, ultrafast liquid chromatography system equipped with a diode array detector

## REFERENCES

- NTP. (2011) Styrene-7,8-oxide, In *Report on Carcinogens*, pp 391–396, U.S. Department of Health and Human Services PHS, Morrisville, NC.
- IARC. (1994) Some industrial chemicals, In *Monographs on the Evaluation of Carcinogenic Risks to Humans*, pp 321–347, WHO, Lyon.
- NTP. (2011) Styrene, In *Report on Carcinogens*, pp 383–391, U.S. Department of Health and Human Services PHS, Morrisville, NC.
- Cruzan, G., Bus, J., Hotchkiss, J., Sura, R., Moore, C., Yost, G., Banton, M., and Sarang, S. (2013) Studies of styrene, styrene oxide and 4-hydroxystyrene toxicity in CYP2F2 knockout and CYP2F1 humanized mice support lack of human relevance for mouse lung tumors. *Regul. Toxicol. Pharmacol.* 66, 24–29.
- Latif, F., Moschel, R. C., Hemminki, K., and Dipple, A. (1988) Styrene oxide as a stereochemical probe for the mechanism of aralkylation at different sites on guanosine. *Chem. Res. Toxicol.* 1, 364–369.
- Laird, R. M., and Parker, R. E. (1961) Mechanism of epoxide reactions 0.4. Reactions of benzylamine with a series of *m*- and *p*-substituted styrene oxides in ethanol. *J. Am. Chem. Soc.* 83, 4277–4281.
- Koskinen, M., Vodičková, L., Vodička, P., Warner, S., and Hemminki, K. (2001) Kinetics of formation of specific styrene oxide adducts in double-stranded DNA. *Chem.-Biol. Interact.* 138, 111–124.



- 471 (8) Koskinen, M., and Plná, K. (2000) Specific DNA adducts induced  
472 by some mono-substituted epoxides in vitro and in vivo. *Chem.-Biol.*  
473 *Interact.* 129, 209–229.
- 474 (9) Parker, R. E., and Isaacs, N. S. (1959) Mechanisms of epoxide  
475 reactions. *Chem. Rev.* 59, 737–799.
- 476 (10) Swern, D., Billen, G. N., and Knight, H. B. (1949) Chemistry of  
477 epoxy compounds. VIII. Reaction of allyl alcohol with unsymmetrical  
478 oxirane compounds. An electronic interpretation. *J. Am. Chem. Soc.* 71,  
479 1152–1156.
- 480 (11) Parker, R. E., and Rockett, B. W. (1965) The mechanism of  
481 epoxide reactions. Part VIII. The effect of solvent variation on the  
482 reaction of benzylamine with 1,2-epoxyethylbenzene. *J. Chem. Soc.*,  
483 2569–2579.
- 484 (12) Nelis, H. J. C. F., and Sinsheimer, J. E. (1981) A sensitive  
485 fluorimetric procedure for the determination of aliphatic epoxides  
486 under physiological conditions. *Anal. Biochem.* 115, 151–157.
- 487 (13) Kumar, R., Vodicka, P., Peltonen, K., and Hemminki, K. (1997)  
488 32P-postlabelling analysis of isomeric 7-alkylguanine adducts of  
489 styrene oxide. *Carcinogenesis* 18, 407–414.
- 490 (14) Reeve, W., and Christoffel, I. (1950) The reaction of styrene  
491 oxide with methanol. *J. Am. Chem. Soc.* 72, 1480–1483.
- 492 (15) Laird, R. M., and Parker, R. E. (1965) The mechanism of  
493 epoxide reactions. Part X. The reactions of (epoxyethyl)benzene with  
494 m-chloro-, 3,4-dimethyl-, and o-(hydroxymethyl)-benzylamine. *J.*  
495 *Chem. Soc.*, 4784–4788.
- 496 (16) Blumenstein, J. J., Ukachukwu, V. C., Mohan, R. S., and Whalen,  
497 D. L. (1993) Effects of para-substituents on the mechanisms of  
498 solvolysis of styrene oxides. *J. Org. Chem.* 58, 924–932.
- 499 (17) Agarwal, S. C., Van Duuren, B. L., and Kneip, T. J. (1979)  
500 Detection of epoxides with 4-(p-Nitrobenzyl) pyridine. *Bull. Environ.*  
501 *Contam. Toxicol.* 23, 825–829.
- 502 (18) Barlow, T., and Dipple, A. (1998) Aralkylation of guanosine  
503 with para-substituted styrene oxides. *Chem. Res. Toxicol.* 11, 44–53.
- 504 (19) Sugiura, K., and Goto, M. (1981) Mutagenicities of styrene  
505 oxide derivatives on bacterial test systems: Relationship between  
506 mutagenic potencies and chemical reactivity. *Chem.-Biol. Interact.* 35,  
507 71–91.
- 508 (20) Tamura, N., Takahashi, K., Shirai, N., and Kawazoe, Y. (1982)  
509 Studies on chemical carcinogens. XXI. Quantitative structure-  
510 mutagenicity relationship among substituted styrene oxides. *Chem.*  
511 *Pharm. Bull. (Tokyo)* 30, 1393–1400.
- 512 (21) Gómez-Bombarelli, R., González-Pérez, M., Calle, E., and  
513 Casado, J. (2012) Potential of the NBP method for the study of  
514 alkylation mechanisms: NBP as a DNA-model. *Chem. Res. Toxicol.* 25,  
515 1176–1191.
- 516 (22) Agrawal, Y. K. (1973) Correction factors for the glass electrode  
517 in aqueous dioxan. *Talanta* 20, 1354–1356.
- 518 (23) Irving, H. M. N. H., and Mahnot, U. S. (1968) pH-meter  
519 corrections for titrations in mixtures of water and dioxan. *J. Inorg. Nucl.*  
520 *Chem.* 30, 1215–1220.
- 521 (24) Van Uitert, L. G., and Fernelius, W. C. (1954) Determination of  
522 thermodynamic equilibrium constants in mixed solvents. *J. Am. Chem.*  
523 *Soc.* 76, 5887–5888.
- 524 (25) Savela, K., Hesso, A., and Hemminki, K. (1986) Character-  
525 ization of reaction products between styrene oxide and deoxynucleo-  
526 sides and DNA. *Chem.-Biol. Interact.* 60, 235–246.
- 527 (26) González-Pérez, M., Gómez-Bombarelli, R., Pérez-Prior, M. T.,  
528 Manso, J. A., Céspedes-Camacho, I. F., Calle, E., and Casado, J. (2011)  
529 Reactivity of p-nitrostyrene oxide as an alkylating agent. A kinetic  
530 approach to biomimetic conditions. *Org. Biomol. Chem.* 9, 7016–7022.
- 531 (27) Vodicka, P., Koskinen, M., Arand, M., Oesch, F., and Hemminki,  
532 K. (2002) Spectrum of styrene-induced DNA adducts: the relationship  
533 to other biomarkers and prospects in human biomonitoring. *Mutat.*  
534 *Res., Rev. Mutat. Res.* 511, 239–254.
- 535 (28) Pérez-Prior, M. T., Gómez-Bombarelli, R., González-Pérez, M.,  
536 Manso, J. A., García-Santos, M. P., Calle, E., and Casado, J. (2010)  
537 Reactivity of the mutagen 1,4-dinitro-2-methylpyrrole as an alkylating  
538 agent. *J. Org. Chem.* 75, 1444–1449.
- (29) Pérez-Prior, M. T., Gómez-Bombarelli, R., González-Pérez, M.,  
Manso, J. A., García-Santos, M. P., Calle, E., and Casado, J. (2009) 539  
Sorbate-nitrite interactions: acetonitrile oxide as an alkylating agent. 540  
*Chem. Res. Toxicol.* 22, 1320–1324. 542
- (30) González-Pérez, M., Gómez-Bombarelli, R., Arenas-Valgañón, J.,  
Pérez-Prior, M. T., García-Santos, M. P., Calle, E., and Casado, J. 543  
(2012) Connecting the chemical and biological reactivity of epoxides. 544  
*Chem. Res. Toxicol.* 25, 2755–2762. 546
- (31) Arenas-Valgañón, J., Gómez-Bombarelli, R., González-Pérez, M.,  
González-Jiménez, M., Calle, E., and Casado, J. (2012) Taurine–nitrite 547  
interaction as a precursor of alkylation mechanisms. *Food Chem.* 134, 548  
986–991. 550
- (32) Kirkovsky, L. I., Lermontov, S. A., Zavorin, S. I., Sukhozhenko, I. 551  
I., Zavel'sky, V. I., Thier, R., and Bolt, H. M. (1998) Hydrolysis of 552  
genotoxic methyl-substituted oxiranes: Experimental kinetic and 553  
semiempirical studies. *Environ. Toxicol. Chem.* 17, 2141–2147. 554
- (33) Koskinen, M., Calebiro, D., and Hemminki, K. (2000) Styrene 555  
oxide-induced 2'-deoxycytidine adducts: implications for the muta- 556  
genicity of styrene oxide. *Chem.-Biol. Interact.* 126, 201–213. 557
- (34) Kim, J. H., and Thomas, J. J. (1992) Use of 4-(nitrobenzyl)- 558  
pyridine (4-NBP) to test mutagenic potential of slow-reacting 559  
epoxides, their corresponding olefins, and other alkylating-agents. 560  
*Bull. Environ. Contam. Toxicol.* 49, 879–885. 561
- (35) Hemminki, K., Heinonen, T., and Vainio, H. (1981) Alkylation 562  
of guanosine and 4-(p-nitrobenzyl)-pyridine by styrene oxide 563  
analogues in vitro. *Arch. Toxicol.* 49, 35–41. 564
- (36) Hemminki, K., and Falck, K. (1979) Correlation of mutagenicity 565  
and 4-(para-nitrobenzyl)-pyridine alkylation by epoxides. *Toxicol. Lett.* 566  
4, 103–106. 567
- (37) Rosman, L. B., Beylin, V. G., Gaddamidi, V., Hooberman, B. H.,  
and Sinsheimer, J. E. (1986) Mutagenicity of para-substituted alpha- 568  
methylstyrene oxide derivatives with *Salmonella*. *Mutat. Res.* 171, 63– 569  
70. 571
- (38) Connors, K. A. (1990) *Chemical Kinetics: The Study of Reaction* 572  
*Rates in Solution*, VCH, New York. 573
- (39) Exner, O. (1988) *Correlation Analysis of Chemical Data*, Plenum, 574  
New York. 575