

Title	The association between low pH value and unfavorable neurological outcome among the out-of-hospital cardiac arrest patient treated by extracorporeal CPR: Sensitivity analysis
Author(s)	Okada, Yohei; Kiguchi, Takeyuki; Kitamura, Tetsuhisa; Iwami, Taku
Citation	Journal of Intensive Care. 8 P.53
Issue Date	2020-07-22
Text Version	publisher
URL	http://hdl.handle.net/11094/78372
DOI	10.1186/s40560-020-00470-3
rights	<p>This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated in a credit line to the data.</p>
Note	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

Osaka University

LETTER TO THE EDITOR

Open Access



The association between low pH value and unfavorable neurological outcome among the out-of-hospital cardiac arrest patient treated by extra-corporeal CPR: sensitivity analysis

Yohei Okada^{1,2}, Takeyuki Kiguchi^{3,4}, Tetsuhisa Kitamura⁵ and Taku Iwami^{1,3*}

Abstract

This is the response to the comment from Dr. Romain Jouffroy and his colleague, on the manuscript “Association between low pH and unfavorable neurological outcome among out-of-hospital cardiac arrest patients treated by extracorporeal CPR: a prospective observational cohort study in Japan”. We performed sensitivity analysis based on the comment from them. It indicated that the results of primary analysis were robust even in considering their criticism.

To the editor,

We would like to thank Dr. Romain Jouffroy and his team for their interest and giving some suggestions for our research to investigate the association between pH value and neurological outcome among the out-of-hospital cardiac arrest (OHCA) patients treated by extra-corporeal cardiopulmonary resuscitation (ECPR) [1, 2].

First, Dr. Romain and his team suggested that the time duration from OHCA occurrence to the blood test or no flow duration should be in the analysis as a covariate. Basically, we believe that it is unnecessary to adjust the low or no flow time duration to estimate the association between the pH value and outcome, because we think pH value is the representative of the duration and the quality of the resuscitation as described in the original

article [1], and we assumed that the duration was not a confounder but a “parent” of the association between pH value and outcome in the causal inference. Further, they also suggested that categorization of the pH value to three groups could not reflect the pathophysiology accurately, and it should be treated as a continuous variable. In the primary analysis of the original article, the reason why we categorized the pH values was that it may be easy to interpret the association. However, we agree with the limitation that the categorization may not be able to represent the pathophysiology in more detail.

According to their suggestion, we performed the sensitivity analysis in which the time duration from call to blood test was added as a covariate, and the pH value was treated as continuous variable in the logistic model and indicated the association between the initial pH value and neurological outcome using the restricted cubic spline curve (Fig. 1). The detail of the method was described in additional file 1. This figure showed that the adjusted odds ratio for favorable neurological outcome was constantly low in the area of pH value less than 7.0, and it gradually increased in higher than 7.0.

This reply refers to the comment available at <https://doi.org/10.1186/s40560-020-00461-4>.

* Correspondence: iwami.taku.8w@kyoto-u.ac.jp

¹Department of Preventive Services, School of Public Health, Kyoto University, Kyoto, Japan

³Kyoto University Health Service, Yoshida Honmachi, Sakyo, Kyoto 606-8501, Japan

Full list of author information is available at the end of the article



© The Author(s). 2020 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.




Fig. 1 The results of sensitivity analysis. X-axis: Initial pH value before implementation of the extracorporeal cardio-pulmonary resuscitation (ECPR). Y-axis: The adjusted odds ratio and 95% confidence interval (CI) for favorable neurological outcome referred to median value in the pH ≥ 7.03 group (pH 7.13) were adjusted by sex (men, women), age (< 65, 65–74 and ≥ 75 years), witnessed by bystander, CPR by bystander, pre-hospital initial cardiac rhythm (shockable, non-shockable), cardiac rhythm on hospital arrival (shockable, non-shockable and ROSC) and the time from call to blood test. The method in detail was described in additional file 1

This result was almost the same as our primary analysis. Therefore, we believe that the result in primary analysis would be robust even though their concerns were taken into account.

Second, we understood their criticism that the residual confounding in in-hospital phase might have exist. However, even if such an unmeasured in-hospital confounding would have existed, we believe that the association in our results would be robust. It is because the association was so obvious, and in order to change the result, it would be required to assume enough strong association between the potential unmeasured confounders and outcomes, based on the recent idea for unmeasured confounding [3, 4]; however, association between in-hospital factors (e.g., tracheal intubation or drug administration) and outcomes are generally small compared to pre-hospital factors [5, 6].

Third, we agree with the concern that the blood test results including pH value might influence on the decision-making of the implementation of extracorporeal cardio-pulmonary resuscitation, and it could be a selection bias as mentioned in limitation part [1]. However, this potential selection would not change the results of this study. Because we assumed that most cases influenced by the selection were decided to withdraw the ECPR due to low

pH value, these patients would tend to die in the emergency department regardless of introduction of the ECPR. Therefore, we believe that it could not change the association of low pH value with unfavorable outcome.

Fourth, we understand their argument that the prevalence of return of spontaneous circulation in the tertile 1–3 groups were different; however, it was adjusted by logistic regression model in the primary analysis. Thus, it may not be a problem to interpret the result.

In conclusion, the result of sensitivity analysis based on the suggestion from Dr. Romain Jouffroy [2] indicated that the results of primary analysis were robust even in considering their criticism. We really appreciate for their suggestion on our analysis and giving the important opportunity to show the result of sensitivity analysis. We hope this process helps better understanding for readers.

Supplementary information

Supplementary information accompanies this paper at <https://doi.org/10.1186/s40560-020-00470-3>.

Additional file 1. Method.

Acknowledgements

We appreciate all members and institutions of the CRITICAL study for their contribution.

Authors' contributions

YO performed the conceptualization, visualization, formal analysis, methodology and writing the original draft. Takeyuki K performed the conceptualization, writing, reviewing and editing. Tetsuhisa K performed the methodology and supervision. Taku I performed the methodology, writing, reviewing and editing and supervision. All authors approved the final manuscript.

Funding

This study was supported by a scientific research grant from the Ministry of Education, Culture, Sports, Science and Technology of Japan (15H05006 and 19K09393). Funding sources had no role in the study design, analysis and interpretation of the data and writing of the paper.

Availability of data and materials

Not applicable

Ethics approval and consent to participate

The Ethics Committee of Kyoto University and each participating institution approved this study protocol (R1045), and written informed consent was waived.

Consent for publication

Not applicable

Competing interests

The authors declare that they have no competing interests.

Author details

¹Department of Preventive Services, School of Public Health, Kyoto University, Kyoto, Japan. ²Department of Primary Care and Emergency Medicine, Graduate School of Medicine, Kyoto University, Kyoto, Japan. ³Kyoto University Health Service, Yoshida Honmachi, Sakyo, Kyoto 606-8501, Japan. ⁴Critical Care and Trauma Center, Osaka General Medical Center, Osaka, Japan. ⁵Division of Environmental Medicine and Population Sciences, Department of Social and Environmental Medicine, Graduate School of Medicine, Osaka University, Osaka, Japan.

Received: 13 July 2020 Accepted: 15 July 2020

Published online: 22 July 2020

References

- Okada Y, Kiguchi T, Irisawa T, Yoshiya K, Yamada T, Hayakawa K, Noguchi K, Nishimura T, Ishibe T, Yagi Y, et al. Association between low pH and unfavorable neurological outcome among out-of-hospital cardiac arrest patients treated by extracorporeal CPR: a prospective observational cohort study in Japan. *Journal of Intensive Care*. 2020;8(1):34.
- Jouffroy R, Vivien B: Association between low pH and unfavorable neurological outcome among out-of-hospital cardiac arrest patients treated by extracorporeal CPR: do not dismiss confounders! *Journal of Intensive Care* 2020, 8(1):42.
- VanderWeele TJ, Ding P. Sensitivity analysis in observational research: introducing the E-value. *Ann Intern Med*. 2017;167(4):268–74.
- Haneuse S, VanderWeele TJ, Arterburn D. Using the E-value to assess the potential effect of unmeasured confounding in observational studies. *JAMA*. 2019;321(6):602–3.
- Iwami T, Nichol G, Hiraide A, Hayashi Y, Nishiuchi T, Kajino K, Morita H, Yukioka H, Ikeuchi H, Sugimoto H, et al. Continuous improvements in "Chain of Survival" increased survival after out-of-hospital cardiac arrests. *Circulation*. 2009;119(5):728–34.
- Stiell IG, Wells GA, Field B, Spaitte DW, Nesbitt LP, De Maio VJ, Nichol G, Cousineau D, Blackburn J, Munkley D, et al. Advanced cardiac life support in out-of-hospital cardiac arrest. *New England Journal of Medicine*. 2004;351(7):647–56.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

