

Supplemental Online Content

Weiss EM, Olszewski AE, Guttman KF, et al. Parental factors associated with the decision to participate in a neonatal clinical trial. *JAMA Netw Open*. 2021;4(1):e2032106. doi:10.1001/jamanetworkopen.2020.32106

eTable. Perception of Severity of Childs' Illness and Sarnat Classification

This supplemental material has been provided by the authors to give readers additional information about their work.

eTable. Perception of Severity of Childs' Illness and Sarnat Classification

I very much or moderately ...	Moderate ^a	Severe	Odds Ratio (95% CI) ^b
... thought my infant's illness would affect his or her life ^b	154 (69%)	33 (72%)	1.5 (0.7, 3.4)
... thought my infant was sick	166 (75%)	41 (89%)	3.2 (1.2, 11)
... was concerned about my infant's illness	213 (96%)	45 (98%)	-- ^c
... understood my infant's illness	108 (49%)	19 (41%)	0.8 (0.4, 1.6)
... thought my infant's illness was a serious condition	201 (92%)	44 (96%)	4.7 (0.9, 87)
... thought my infant's illness would have long term effects	149 (67%)	37 (80%)	2.4 (1.0, 6.0)
I thought my infant was likely to die at birth ^c	105 (49%)	28 (61%)	1.6 (0.8, 3.4)

^aThe columns are Sarnat severity staging of participant's infant.

^bThe odds ratio has adjusted for Medicaid status, household income, and race.

^cNot calculated because only a single participant with an infant in the severe category did not choose "very much" or "moderately."