



VU Research Portal

Another look at estimating parameters in systems of ordinary differential equations via regularization

Vujacic, I.; Mahmoudi, S.M.; de Wit, E.

published in

Proceedings of the 60th World Statistics Congress of the International Statistical Institute, ISI2015
2015

document version

Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

citation for published version (APA)

Vujacic, I., Mahmoudi, S. M., & de Wit, E. (2015). Another look at estimating parameters in systems of ordinary differential equations via regularization. In *Proceedings of the 60th World Statistics Congress of the International Statistical Institute, ISI2015* The International Statistical Institute.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.


- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl



Another look at estimating parameters in systems of ordinary differential equations via regularization

Ivan Vujačić*

VU University Amsterdam, Amsterdam, The Netherlands - i.vujacic@vu.nl

Seyed Mahdi Mahmoudi

University of Groningen, Groningen, The Netherlands - s.m.mahmoudi@rug.nl

Ernst Wit

University of Groningen, Groningen, The Netherlands - e.c.wit@rug.nl

Abstract

We consider estimation of parameters in systems of ordinary differential equations (ODEs). This problem is important because many processes in various fields of science are modelled by a system of ODEs. Since the system usually contains unknown parameters it is of interest to estimate them. The problem is approached from the viewpoint of M -estimation. In general, for a given parameter the true solution of the system is unavailable, therefore any M -criterion function is necessarily defined via an approximation of the solution. We define an approximation by viewing the system of ODEs as an operator equation and exploiting the connection with regularization theory. Combining the introduced regularized solution with the M -criterion function provides a general framework for estimating parameters in ODEs, which can handle partially observed systems. If the M -criterion function is the log-likelihood then suitable regularized solutions yield estimators which are consistent and asymptotically efficient. Also, connections with the generalized profiling procedure are made.

Keywords: M -estimation; Quasisolution; Generalized Tikhonov regularizer; Asymptotic efficiency.

1. Introduction

Consider the system of ordinary differential equations of the form

$$\begin{cases} \mathbf{x}'(t) = \mathbf{f}(\mathbf{x}(t), t; \boldsymbol{\theta}), & t \in [0, T], \\ \mathbf{x}(0) = \boldsymbol{\xi}, \end{cases} \quad (1)$$

where $\mathbf{x}(t)$ takes values in \mathbb{R}^d , $\boldsymbol{\xi}$ in $\Xi \subset \mathbb{R}^d$, $\boldsymbol{\theta}$ in $\Theta \subset \mathbb{R}^p$ and \mathbf{f} is known function. Given the values of $\boldsymbol{\xi}$ and $\boldsymbol{\theta}$, we denote the solution of (1) by $\mathbf{x}(t; \boldsymbol{\theta}, \boldsymbol{\xi})$. Let us assume that a process is modelled by ODEs (1) with unknown parameters $\boldsymbol{\xi}_0$ and $\boldsymbol{\theta}_0$. For simplicity, assume that we have noisy observations $y_i(t_j)$, $j = 1, \dots, n$ of the first $1 \leq d_1 \leq d$ states $x_i(t; \boldsymbol{\theta}_0, \boldsymbol{\xi}_0)$, $i = 1, \dots, d_1$ at time points $t_j \in [0, T]$, $j = 1, \dots, n$:

$$y_i(t_j) = x_i(t_j; \boldsymbol{\theta}_0, \boldsymbol{\xi}_0) + \varepsilon_i(t_j), \quad i = 1, \dots, d_1; j = 1, \dots, n. \quad (2)$$

The problem is to estimate $\boldsymbol{\theta}_0$ from the data \mathbf{Y} , where $\mathbf{Y} = (y_i(t_j))_{ij}$ denotes the matrix that contains all the observations. A general way to estimate the unknown parameter is to minimize some known function M_n of the parameter and the data; the obtained estimator is called an M -estimator (Van der Vaart, 2000). In the problem we consider M_n depends on the solution of the ODE system. In general the solution is not available, therefore its approximation has to be used. Approximation can be deterministic or stochastic; those used in the literature include:

1. Numerical solution given by a numerical ODE solver (Xue et al., 2010).
2. Classical smoothers like cubic splines, kernel estimators, regression splines, local polynomials, step function estimators (Varah, 1982; Brunel et al., 2008; Gugushvili and Klaassen, 2012; Dattner and Klaassen, 2013; Vujačić et al., 2014).

3. Specially designed smoothers that use ODEs model, like model based smoothing (Ramsay et al., 2007) or reproducing kernel Hilbert space based smoothers (González et al., 2014).

In this paper, we generalize and combine some of the ideas described above and estimate the parameter as follows:

$$1. \hat{\mathbf{x}} = \underset{\mathbf{x} \in \mathcal{X}_m}{\operatorname{argmin}} \mathcal{T}_{\alpha, \gamma}(\mathbf{x}), \quad (3)$$

$$2. \hat{\boldsymbol{\theta}}_n = \underset{\boldsymbol{\theta} \in \Theta}{\operatorname{argmin}} M_n(\boldsymbol{\theta} | \hat{\mathbf{x}}, \mathbf{Y}). \quad (4)$$

Here $\mathcal{T}_{\alpha, \gamma}$ is a functional with parameters $\alpha \geq 0$ and $\gamma \geq 0$, which is optimized over some finite-dimensional subspace \mathcal{X}_m of the solution space. M_n is a criterion function to be optimized, for example the log-likelihood criterion. The main goal of the paper is to show how to define the functional $\mathcal{T}_{\alpha, \gamma}$ by using regularization theory. We will call $\mathcal{T}_{\alpha, \gamma}$ generalized Tikhonov functional and its minimizer a generalized Tikhonov regularizer. The proposed framework can handle fully and partially observed systems and is not based on numerical integration of the system. The trade-off is that the numerical integration is substituted with numerical optimization. Some issues related to the proposed methodology that are also present in similar methods are selection of α , γ and the dimension m of the space \mathcal{X}_m .

The rest of the paper is organized as follows. A review of the regularization theory is provided in Section 2. In Section 3 the described theory is used to define generalized Tikhonov functional for the ODE system. Section 4 contains theoretical results for the estimator defined by (3) and (4) with M -criterion function being the log-likelihood. Some comparison with the generalized profiling procedure of Ramsay et al. (2007) is also provided. The final section contains discussion.

2. Ill-posed problems, quasisolutions and regularization

Let $F: \mathcal{X} \rightarrow \mathcal{Y}$ where \mathcal{X}, \mathcal{Y} are linear normed spaces and consider the operator equation

$$F(x) = y, \quad (5)$$

$x \in \mathcal{X}, y \in \mathcal{Y}$. The problem (5) is *well-posed in the sense of Hadamard* on the pair of normed spaces \mathcal{X} and \mathcal{Y} if the solution of (5) exists, it is unique and it is continuous with respect to y . The problem(5) is *ill-posed* on the pair of normed spaces \mathcal{X} and \mathcal{Y} if at least one of the three well-posedness conditions does not hold.

Equation (5) can be solved on the set $S \subset \mathcal{X}$ by finding the minimum of the *objective functional*

$$\mathcal{J}(x) = \|F(x) - y\|^2, \quad (6)$$

on S . This idea dates back to the works of A. M. Legendre and K. Gauss from the beginning of the 19th century, who proposed the least squares method for solving systems of linear algebraic equations. (Kabanikhin, 2011) In this regard, Ivanov (Tikhonov and Arsenin, 1977, Sec 1.2) introduced a concept of *quasisolution* (or pseudo solution or least squares solution) of equation (5) on $S \subset \mathcal{X}$ — quasisolution is any minimizer of (6) on S . Numerical optimization of \mathcal{J} has to be implemented on some finite dimensional subspace of \mathcal{X} . Moreover, the finite dimensional subspace of \mathcal{X} has to be chosen from the family of subspaces $\mathcal{X}_1 \subset \mathcal{X}_2 \subset \dots$ such that their union is dense in \mathcal{X} . This is minimal assumption for the sequence of minimum norm minimizers x_n of (6) on \mathcal{X}_n to converge to the minimum norm minimizer x of (6) on \mathcal{X} . When regularization is achieved by a finite dimensional approximation alone it is called *self-regularization* (Vainikko and Khyamarik, 1985) or *regularization by projection*. (Engl et al., 1996)

Tikhonov regularization involves further regularization by minimizing the so-called *Tikhonov functional*

$$\mathcal{T}_\alpha(x) = \mathcal{J}(x) + \alpha \Omega(x - x_0), \quad (7)$$

where x_0 is *trial solution*, $\alpha \geq 0$ is *regularization parameter* and Ω is *stabilizing functional* which is usually given by a norm or a semi-norm on \mathcal{X} . Stabilizing functional incorporates a priori information on the smoothness of the solution x . Some possible choices are the square of the norm $\Omega(x) = \|x\|^2$ or the total variation $\Omega(x) = V_0^T(x)$.

A priori information on values of the solution may be available, which allows to single out solutions which satisfy physical requirements. This information can be incorporated by adding an additional functional \mathcal{S} in (7) (Vasin and Ageev, 1995; Pöschl, 2008), which measures the closeness of the solution to the aforementioned a priori information:

$$\mathcal{T}_{\alpha,\gamma}(x) = \mathcal{J}(x) + \alpha\Omega(x - x_0) + \gamma\mathcal{S}(x), \quad (8)$$

where $\gamma \geq 0$ is the *penalty parameter*. In the noise free case, Vasin and Ageev (1995) treated information given via equality or inequality constraints; the functional \mathcal{S} was termed the penalty functional and the resulting method was called the generalized regularization method. In the presence of noise, Pöschl (2008) used the term "similarity functional" for \mathcal{S} and named the resulting method multi-modal Tikhonov regularization. We will use the terms *similarity functional* and *generalized Tikhonov regularization*. Thus, *generalized Tikhonov regularization* finds an approximation to the solution of (5) by minimizing (8) over some finite-dimensional subspace of \mathcal{X} . We will refer to the functional $\mathcal{T}_{\alpha,\gamma}(x)$ as *the generalized Tikhonov functional* and its minimizer as generalized Tikhonov regularizer.

3. Generalized Tikhonov regularization in ODE estimation setting

The problem (1) is well-posed. Indeed, the first two conditions of the definition of the well-posed problem in the sense of Hadamard follow from existence and uniqueness theorems for ordinary differential equations, and the third condition follows from the theorem of continuous dependence of solution on initial conditions and parameters (Agarwal and O'Regan, 2008). However, some states may not be observed in which case the initial conditions are not known. In this case the solution is not unique and the problem becomes ill-posed. Even if the initial conditions are known, non-uniqueness can still be introduced through finite dimensional approximation (Engl et al., 1996). Thus, to deal with this issue regularization can be employed. To this aim for system (1) without initial condition we define generalized Tikhonov regularizer by using the theory described in the previous section. For notational simplicity, we suppress the dependence on θ .

3.1 Finite dimensional approximation

The first step is the finite dimensional approximation; we assume that the solution of the system (1) lies in $(C^1[0, T])^d$. For simplicity we assume that each component of \mathbf{x} is approximated by an element from the same finite dimensional function space \mathcal{X}_m of dimension m with basis $\{h_1, \dots, h_m\}$. For any $x_i \in \mathcal{X}_m$, $i = 1, \dots, d$ we have:

$$x_i(t) = \sum_{k=1}^m \beta_{ik} h_k(t) = \beta_i^\top \mathbf{h}(t), \quad (9)$$

where $\beta_i = (\beta_{i1}, \dots, \beta_{im})^\top$ and $\mathbf{h}(t) = (h_1(t), \dots, h_m(t))^\top$. Commonly used basis functions are B-splines; they yield sequence of spaces \mathcal{X}_m^d whose union is dense in $(C^1[0, T])^d$ (see Lemma 1 in Section 4).

3.2 Objective functional

For fixed θ ODEs system (1) without initial condition is equivalent to the operator equation $F(\mathbf{x}) = \mathbf{0}$, where $F(\mathbf{x}(\cdot)) = \mathbf{x}'(\cdot) - \mathbf{f}(\mathbf{x}(\cdot), \cdot, \theta)$. The corresponding objective functional is

$$\mathcal{J}(\mathbf{x}) = \|\mathbf{x}' - \mathbf{f}(\mathbf{x}, \cdot, \theta)\|_{2, \mathbf{w}}^2, \quad (10)$$

where $\mathbf{w} = (w_1, \dots, w_d)$, $w_i > 0$ for $i = 1, \dots, d$ and $\|\mathbf{x}\|_{2, \mathbf{w}} = \sqrt{\sum_{i=1}^d w_i \int_0^T x_i^2(t) dt}$ is the norm on the space of d vector valued functions. Depending on the application other norms in \mathcal{J} can be chosen.

3.3 Stabilizing functional

Stabilizing functional enforces smoothing conditions on the solution \mathbf{x} . The common choice is the total curvature on $[0, T]$ measured by the integral $\int_0^T \{x''(t)\}^2 dt$. In regard to this measure one choice for stabilizing functional is

$$\Omega(\mathbf{x}) = \sum_{i=1}^d v_i \int_0^T \{x_i''(t)\}^2 dt \quad (11)$$

where v_i , $i = 1, \dots, d$ are nonnegative constants. Other choices are possible, like total variation $V_0^T(x)$.

3.4 Similarity functional

For simplicity let us assume that $d = 1$. Let $p(y(t)|x(t; \theta_0, \xi_0))$ be the distribution of the data based on

the model (2) and g be the true distribution of the data. The observations $y(t_i)$ represent the data for the problem of the estimation of $\boldsymbol{\theta}_0$ but they are a priori information for the problem of finding the solution $\boldsymbol{x}(t; \boldsymbol{\theta}_0, \boldsymbol{\xi}_0)$ of (1). The solution is measured with noise therefore the distribution of the data should be close to a priori distribution of the solution — distribution based on the model. One measure of closeness between distributions is the Kullback-Leibler (KL) divergence, which belongs to the class of f -divergences (Pöschl, 2008). Since g is not known the KL divergence can only be estimated from the data by approximating g with its empirical density:

$$KL(g(\cdot); p(\cdot|x(\cdot; \boldsymbol{\theta}_0, \boldsymbol{\xi}_0))) \approx -\frac{1}{n} \sum_{i=1}^n \log p(y(t_i)|x(t_i; \boldsymbol{\theta}_0, \boldsymbol{\xi}_0)).$$

Because the solution $x(t_i; \boldsymbol{\theta}_0, \boldsymbol{\xi}_0)$ is unknown by using the approximation $p(y(t_i)|x(t; \boldsymbol{\theta}_0, \boldsymbol{\xi}_0)) \approx p(y(t_i)|\hat{x}(t; \boldsymbol{\beta}))$ we obtain the similarity functional which for the system with d_1 observed components is

$$\mathcal{S}(\boldsymbol{x}) = -\sum_{i=1}^{d_1} \sum_{j=1}^n \log p(y_i(t_j)|\hat{x}_i(t_j; \boldsymbol{\beta})). \quad (12)$$

Here the scale $1/n$ is omitted because it can be subsumed in the penalty parameter γ which multiplies \mathcal{S} in (8). The weakness of this approach is that the employed approximation ignores the statistical uncertainty in the regularizer (Chkrebtii et al., 2013).

3.5 Generalized Tikhonov functional

From the previous subsections it follows that the generalized Tikhonov functional for the equation $F(\boldsymbol{x}) = \mathbf{0}$ is

$$\mathcal{T}_{\alpha, \gamma}(\boldsymbol{x}(\boldsymbol{\beta})) = \mathcal{J}(\boldsymbol{x}(\boldsymbol{\beta})) + \alpha \Omega(\boldsymbol{x}(\boldsymbol{\beta}) - \boldsymbol{x}_0) + \gamma \mathcal{S}(\boldsymbol{x}(\boldsymbol{\beta})), \quad (13)$$

where the functionals \mathcal{J} , Ω and \mathcal{S} are defined in (10), (11) and (12), respectively. The regularized solution is found by optimizing (13) over \mathcal{X}_m^d parametrized by $\boldsymbol{\beta} = (\boldsymbol{\beta}_1^\top, \dots, \boldsymbol{\beta}_d^\top)^\top$. This can be achieved by optimizing (13) with respect to $\boldsymbol{\beta}$ over \mathbb{R}^{dm} :

$$\hat{\boldsymbol{\beta}} = \underset{\boldsymbol{\beta} \in \mathbb{R}^{dm}}{\operatorname{argmin}} \mathcal{T}_{\alpha, \gamma}(\boldsymbol{x}(\boldsymbol{\beta})),$$

and applying (9).

4. Generalized Tikhonov regularizer with log-likelihood criterion and connection with the generalized profiling procedure

Combining generalized Tikhonov regularizer (13) with the log-likelihood criterion

$$M_n(\boldsymbol{\theta}) = -\sum_{i=1}^{d_1} \sum_{j=1}^n \log p(y_i(t_j)|\hat{x}_i(t_j; \boldsymbol{\theta})),$$

we obtain an estimator which is asymptotically efficient. The proof of asymptotic properties of the estimator follows like in Qi and Zhao (2010) with slight modifications. More specifically, let \boldsymbol{x}^o and \boldsymbol{x}^u denote the observed and unobserved part of \boldsymbol{x} , respectively and let $\boldsymbol{\xi}_0^o$ and $\boldsymbol{\xi}_0^u$ be their corresponding initial conditions. The following results, similar to Lemma 1, Theorem 3.2 and Theorem 3.3 of Qi and Zhao (2010), hold.

Lemma 1. *Under Assumption 2 of Qi and Zhao (2010), there exist a sequence of finite-dimensional subspaces \mathcal{X}_n of $C^1[0, T]$ such that for any compact subset Θ_0 of Θ and any compact subset Ξ_0 of Ξ , it holds*

$$\lim_{n \rightarrow \infty} r_n = 0,$$

where

$$r_n = \max \left[\sup_{(\boldsymbol{\theta}, \boldsymbol{\xi}) \in \Theta_0 \times \Xi_0} \inf_{\boldsymbol{w} \in \mathcal{X}_n, \boldsymbol{w}(0) = \boldsymbol{\xi}_0^o} \left\{ \|\boldsymbol{x}^o(\boldsymbol{\theta}, \boldsymbol{\xi}, \cdot) - \boldsymbol{w}\|_\infty \vee \left\| \frac{d\boldsymbol{x}^o}{dt}(\boldsymbol{\theta}, \boldsymbol{\xi}, \cdot) - \frac{d\boldsymbol{w}}{dt} \right\|_\infty \vee \left\| \frac{d^2\boldsymbol{x}^o}{dt^2}(\boldsymbol{\theta}, \boldsymbol{\xi}, \cdot) - \frac{d^2\boldsymbol{w}}{dt^2} \right\|_\infty \right\}, \right. \\ \left. \sup_{(\boldsymbol{\theta}, \boldsymbol{\xi}) \in \Theta_0 \times \Xi_0} \inf_{\boldsymbol{v} \in \mathcal{X}_n, \boldsymbol{v}(0) = \boldsymbol{\xi}_0^u} \left\{ \|\boldsymbol{x}^u(\boldsymbol{\theta}, \boldsymbol{\xi}, \cdot) - \boldsymbol{v}\|_\infty \vee \left\| \frac{d\boldsymbol{x}^u}{dt}(\boldsymbol{\theta}, \boldsymbol{\xi}, \cdot) - \frac{d\boldsymbol{v}}{dt} \right\|_\infty \vee \left\| \frac{d^2\boldsymbol{x}^u}{dt^2}(\boldsymbol{\theta}, \boldsymbol{\xi}, \cdot) - \frac{d^2\boldsymbol{v}}{dt^2} \right\|_\infty \right\} \right].$$

Theorem 1. *Let Assumptions 1-5 from Qi and Zhao (2010) hold. If $r_n \rightarrow 0$, $\alpha_n \rightarrow 0$ and $\gamma_n \rightarrow 0$, as $n \rightarrow \infty$ then*

$$\widehat{\boldsymbol{\theta}}_n - \boldsymbol{\theta}_0 = o_P(1).$$

Theorem 2. *Let Assumptions 1-6 from Qi and Zhao (2010) hold. If $r_n = o(n^{-1})$, $\alpha_n = o(n^{-2})$ and $\gamma_n = o(n^{-2})$ as $n \rightarrow \infty$ then the estimator $\widehat{\boldsymbol{\theta}}_n$ is asymptotically efficient.*

Model based smoother used in generalized profiling (Ramsay et al., 2007) can be viewed as a generalized Tikhonov regularizer. Indeed, the inner fitting criterion J defined in Ramsay et al. (2007) is

$$J(\mathbf{x}) = - \sum_{i=1}^{d_1} \sum_{j=1}^n \log p(y_i(t_j) | x_i(\boldsymbol{\theta})) + \sum_{i=1}^d \lambda_i \int_0^T \{x'_i(t) - f_i(\mathbf{x}(t), t, \boldsymbol{\theta})\}^2 dt. \quad (14)$$

With representation $\lambda_i = \lambda w_i$, where w_i , $i = 1, \dots, d$ are some constants, we obtain

$$J(\mathbf{x}) = \lambda \left\{ \frac{1}{\lambda} \mathcal{S}(\mathbf{x}) + \mathcal{J}(\mathbf{x}) \right\} = \lambda \mathcal{T}_{0,1/\lambda}(\mathbf{x}), \quad (15)$$

and consequently the model based smoother is a minimizer of $\mathcal{T}_{0,1/\lambda}$. It can be seen from (14) and (15) that the roles of the penalty terms and parameters in $\mathcal{T}_{0,1/\lambda}$ and J are reversed; relationship between the parameters is $\gamma = 1/\lambda$. For the solutions of the dynamic systems the fidelity to the ODEs is of the major concern and data term is of secondary importance (Gu, 2007). In regard to this, the regularization formulation seems more natural since the objective functional \mathcal{J} , which measures the fidelity to the ODEs, is not the penalty in $\mathcal{T}_{\alpha,\gamma}$ but the main term. On the other hand in J the ODE fidelity term is the penalty. The consequence of this is that to force the penalty to zero means that λ approaches ∞ , which leads to ill conditioning in the optimization (Biegler, 2007). This is avoided in the regularization formulation; here γ must approach zero.

5. Discussion

Generalized Tikhonov regularizer $\mathcal{T}_{\alpha,\gamma}$ is a broad framework because it allows one to incorporate various a priori information. It involves several layers of regularization, the first one being self-regularization (see Section 2). A practical issue is to determine the amount of regularization in $\mathcal{T}_{\alpha,\gamma}$. Table 1 gives a list of different regularizers for the ODEs estimation problem. The first four regularizers in the table all satisfy the conditions of Theorems 1 and 2 when $\alpha, \gamma \rightarrow 0$. Thus, they are all asymptotically efficient. The most sensible choices seem to be the model based smoother (Ramsay et al., 2007) and Ivanov's quasi solution. Indeed, the similarity functional \mathcal{S} is more informative than Ω since \mathcal{S} gives information on the values of the minimizer of \mathcal{J} while Ω narrows down the class of functions to which the minimizer of \mathcal{J} should belong. This class is already restricted through self-regularization and hence further regularization via Ω may not be necessary. This will be examined in simulation studies elsewhere.

Parameters	$\mathcal{T}_{\alpha,\gamma}(\mathbf{x})$	$\widehat{\mathbf{x}} = \operatorname{argmin}_{\mathbf{x} \in \mathcal{X}_m} \mathcal{T}_{\alpha,\gamma}(\mathbf{x})$
$\alpha > 0, \gamma > 0$	$\mathcal{J}(\mathbf{x}) + \alpha \Omega(\mathbf{x} - \mathbf{x}_0) + \gamma \mathcal{S}(\mathbf{x})$,	Generalized Tikhonov's regularizer
$\alpha = 0, \gamma = 0$	$\mathcal{J}(\mathbf{x})$,	Ivanov's quasi solution
$\alpha > 0, \gamma = 0$	$\mathcal{J}(\mathbf{x}) + \alpha \Omega(\mathbf{x} - \mathbf{x}_0)$,	Tikhonov's regularizer
$\alpha = 0, \gamma > 0$	$\mathcal{J}(\mathbf{x}) + \gamma \mathcal{S}(\mathbf{x})$	model based smoother of (Ramsay et al., 2007)
$\alpha = \infty, \gamma = 0$	$\mathcal{J}(\mathbf{x}_0) / \delta(\mathbf{x} - \mathbf{x}_0)$	trial solution \mathbf{x}_0

Table 1: Generalized Tikhonov regulariser and its special cases. The last row should be interpreted as $\mathcal{T}_{\alpha,0}(\mathbf{x}) \rightarrow \mathcal{J}(\mathbf{x}_0) / \delta(\mathbf{x} - \mathbf{x}_0)$ as $\alpha \rightarrow +\infty$, where δ is the Dirac's delta function.

Acknowledgements

The first author thanks Bartek Knapik for useful discussions on inverse and ill-posed problems and for providing useful references.

References

- Agarwal, R. P. and O'Regan, D. (2008). *An introduction to ordinary differential equations*. Springer Science & Business Media.
- Biegler, L. (2007). Discussion on the paper by Ramsay, Hooker, Campbell and Cao. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 69(5):776.
- Brunel, N. J. et al. (2008). Parameter estimation of ode's via nonparametric estimators. *Electronic Journal of Statistics*, 2:1242–1267.
- Chkrebtii, O. A., Campbell, D. A., Girolami, M. A., and Calderhead, B. (2013). Bayesian uncertainty quantification for differential equations. *arXiv preprint arXiv:1306.2365*.
- Dattner, I. and Klaassen, C. A. (2013). Estimation in systems of ordinary differential equations linear in the parameters. *arXiv preprint arXiv:1305.4126*.
- Engl, H. W., Hanke, M., and Neubauer, A. (1996). *Regularization of inverse problems*, volume 375. Springer.
- González, J., Vujačić, I., and Wit, E. (2014). Reproducing kernel hilbert space based estimation of systems of ordinary differential equations. *Pattern Recognition Letters*, 45:26–32.
- Gu, C. (2007). Discussion on the paper by Ramsay, Hooker, Campbell and Cao. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 69(5):780.
- Gugushvili, S. and Klaassen, C. A. J. (2012). \sqrt{n} -consistent parameter estimation for systems of ordinary differential equations: bypassing numerical integration via smoothing. *Bernoulli*, 18:1061–1098.
- Kabanikhin, S. I. (2011). *Inverse and ill-posed problems: theory and applications*, volume 55. Walter De Gruyter.
- Pöschl, C. (2008). *Tikhonov regularization with general residual term*. University Innsbruck.
- Qi, X. and Zhao, H. (2010). Asymptotic efficiency and finite-sample properties of the generalized profiling estimation of parameters in ordinary differential equations. *The Annals of Statistics*, 38(1):435–481.
- Ramsay, J. O., Hooker, G., Campbell, D., and Cao, J. (2007). Parameter estimation for differential equations: a generalized smoothing approach. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 69(5):741–796.
- Tikhonov, A. N. and Arsenin, V. Y. (1977). Solutions of ill-posed problems.
- Vainikko, G. M. and Khyamarik, U. (1985). Projection methods and self-regularization in ill-posed problems. *Izvestiya Vysshikh Uchebnykh Zavedenii. Matematika*, (10):3–17.
- Van der Vaart, A. W. (2000). *Asymptotic statistics*, volume 3. Cambridge university press.
- Varah, J. (1982). A spline least squares method for numerical parameter estimation in differential equations. *SIAM Journal on Scientific and Statistical Computing*, 3(1):28–46.
- Vasin, V. V. and Ageev, A. L. (1995). *Ill-posed problems with a priori information*, volume 3. Walter de Gruyter.
- Vujačić, I., Dattner, I., González, J., and Wit, E. (2014). Time-course window estimator for ordinary differential equations linear in the parameters. *Statistics and Computing*, pages 1–14.
- Xue, H., Miao, H., and Wu, H. (2010). Sieve estimation of constant and time-varying coefficients in nonlinear ordinary differential equation models by considering both numerical error and measurement error. *Annals of statistics*, 38(4):2351.