

1  
2  
3  
4 **Non-local acute stretching effects on strength performance in healthy young adults**  
5

6 **Authors:** David G Behm<sup>1</sup>, Shahab Alizadeh<sup>1</sup>, Ben Drury<sup>2</sup>, Urs Granacher<sup>3</sup>, Jason Moran<sup>4</sup>  
7  
8  
9

10 **Institutions:** <sup>1</sup> School of Human Kinetics and Recreation, Memorial University of  
11 Newfoundland, St. John's, Newfoundland and Labrador, Canada

12 <sup>2</sup> Department of Applied Sport Sciences, Hartpury University, UK

13 <sup>3</sup> Division of Training and Movement Science, University of Potsdam, Potsdam, Germany

14 <sup>4</sup> School of Sport, Rehabilitation and Exercise Sciences, University of Essex, Essex UK  
15  
16  
17  
18

19 **Corresponding Author:** David G Behm PhD

20 School of Human Kinetics and Recreation,  
21 Memorial University of Newfoundland,  
22 St. John's, Newfoundland and Labrador, Canada

23 [dbehm@mun.ca](mailto:dbehm@mun.ca)

24 Tel: 709-864-3408

25 Fax: 709-864-3979  
26  
27

28 **Running Title:** Global stretching effects on performance  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

50 Abstract: 249 words

51 Text 14 pages  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 **Abstract**  
5

6 *Background:* Static stretching (SS) can impair performance and increase range of motion of a non-  
7 exercised or non-stretched muscle respectively. An underdeveloped research area is the effect of  
8 unilateral stretching on non-local force output.  
9

10  
11  
12  
13  
14 *Objective:* The objective of this review was to describe the effects of unilateral SS on contralateral,  
15 non-stretched, muscle force and identify gaps in the literature.  
16

17  
18  
19 *Methods:* A systematic literature search following Preferred Reporting Items for Systematic  
20 Review and Meta-Analyses Protocols guidelines was performed according to prescribed inclusion  
21 and exclusion criteria. Weighted means and ranges highlighted the non-local force output response  
22 to unilateral stretching. The Physiotherapy Evidence Database scale was used to assess study risk  
23 of bias and methodological quality.  
24  
25

26  
27  
28  
29  
30  
31 *Results:* Unilateral stretching protocols from six studies involved  $6.3 \pm 2$  repetitions of  $36.3 \pm 7.4$   
32 seconds with  $19.3 \pm 5.7$  seconds recovery between stretches. The mean stretch-induced force  
33 deficits exhibited small magnitude effect sizes for both the stretched ( $-6.7 \pm 7.1\%$ ,  $d = -0.35$ : 0.01 to  
34  $-1.8$ ) and contralateral, non-stretched, muscles ( $-4.0 \pm 4.9\%$ ,  $d = -0.22$ : 0.08 to  $-1.1$ ). Control  
35 measures exhibited trivial deficits.  
36  
37

38  
39  
40  
41  
42  
43 *Conclusion:* The limited literature examining non-local effects of prolonged SS revealed that both  
44 the stretched and contralateral, non-stretched, limbs of young adults demonstrate small magnitude  
45 force deficits. However, the frequency of studies with these effects were similar with three  
46 measures demonstrating deficits, and four measures showing trivial changes. These results  
47 highlight the possible global (non-local) effects of prolonged SS. Further research should  
48 investigate effects of lower intensity stretching, upper versus lower body stretching, different age  
49 groups, incorporate full warm-ups, and identify predominant mechanisms among others.  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**Key Words:** flexibility; power; crossover; fatigue; mental fatigue; neural inhibition

**Abbreviations**

EMG: electromyography  
MVC: maximal voluntary contraction  
PEDro: Physiotherapy Evidence Database  
PIC: persistent inward currents  
PICOS: Population, Intervention, Comparison and Outcomes  
PRISMA-P: Preferred Reporting Items for Systematic review and Meta-Analysis Protocols  
ROM: range of motion  
SS: static stretching

## Introduction

A preponderance of published literature has reported that prolonged static stretching (SS: >60-s per muscle group) in isolation (without a full warm-up including prior and subsequent dynamic activities) can increase joint range of motion (ROM) and induce subsequent muscle strength (force), power, endurance, and sprint performance impairments in the stretched muscle or muscle groups (Behm and Chaouachi 2011; Behm et al. 2016a; 2020; Kay and Blazevich 2012). The mechanisms underlying SS-induced performance deficits have been attributed to both morphological and neural influences (Behm et al. 2020). Morphological mechanisms may include alterations in passive and active muscle force possibly indicating a reduction in the stiffness of the muscle's parallel elastic components (Kay and Blazevich 2009; Kay et al. 2015; Konrad et al. 2017a; 2017b), reduced myofibrillar  $\text{Ca}^{2+}$  sensitivity (i.e. due to metabolite accumulation) (Stephenson and Williams 1985; Sugi et al. 2013), and possibly changes in titin properties (Herzog 2014; Lee et al. 2007; Leonard and Herzog 2010). Neural mechanisms have been attributed to the exteroceptive reflex stimulation reducing excitatory sympathetic nervous activity (Guissard et al. 2001; Wu et al. 1999) and reduced strength of motoneurone dendrites persistent inward currents (PIC: (decreasing the amplification of central neural drive) (Trajano et al. 2017, 2020). Although stretch-induced neural inhibition is often hypothesised, a recent review by Behm and colleagues (Behm et al. 2020) found that SS effects on electromyographic (EMG) activity are conflicting, with both decrements and no changes reported, although when the EMG signal is normalised to the compound muscle action potential (M-wave) there are more consistent impairments. In addition, corticospinal excitability (Pulverenti et al. 2019) and cortical silent period (Pulverenti et al. 2020) are not substantially affected by stretching. Whereas morphological mechanisms can only apply to the stretched muscle (local effects), neural mechanisms can influence both local and non-

1  
2  
3  
4 local responses. Thus, in addition to the possible reductions in sympathetic nervous activity and  
5  
6 PIC strength, an alternative method of examining the extent of global or non-local neural effects  
7  
8 of SS would be to investigate crossover or non-local muscle performance responses.  
9

10  
11 There is evidence of non-local (contralateral homologous or heterologous muscles)  
12  
13 improvements in ROM following a bout of unilateral stretching (Behm et al. 2016b; 2019;  
14  
15 Caldwell et al. 2019a; 2019b; Ce et al. 2020; Chaouachi et al. 2017; Clark et al. 1999; De-la-Cruz-  
16  
17 Torres et al. 2020; Killen et al. 2019; Lima et al. 2014; Whalen et al. 2019; Wilke et al. 2016b;  
18  
19 Wilke et al. 2017). Global (whole body) mechanisms contributing to these non-local improvements  
20  
21 in ROM have been postulated to be related to psycho-physiological influences (i.e. increased  
22  
23 stretch or pain tolerance) (Magnusson et al. 1996; 1997) and the downregulation of sympathetic  
24  
25 excitatory nerve stimulation (Behm 2018). Prolonged SS (i.e. 30-s or more) can also reduce  
26  
27 motoneuron spindle reflexive (excitatory) activity (Avela et al. 1999; Guissard et al. 2001). Type  
28  
29 I and II muscle spindle afferents innervate spinal motoneurons (Prochazka and Ellaway 2012) and  
30  
31 project to the somatosensory and the primary motor cortex (Phillips et al. 1971; Rathelot and Strick  
32  
33 2009) and thus could have global consequences. The global impact of decreased sympathetic  
34  
35 nervous system stimulation and disfacilitation of excitatory spindle afferents (neural inhibition)  
36  
37 could potentially adversely affect the performance of non-stretched muscles (Behm 2018; Behm  
38  
39 et al. 2020).  
40  
41  
42  
43  
44  
45  
46  
47

48 There is evidence of non-local SS-induced isometric maximum voluntary contraction  
49  
50 (MVC) force (Behm et al. 2019), jump height (Marchetti et al. 2014) and muscle activation  
51  
52 (Marchetti et al. 2017) deficits following unilateral SS. However, these results contrast with those  
53  
54 that show no significant impairments of isokinetic torque following eight repetitions of 30-s each  
55  
56 of unilateral hip flexion static stretching (Chaouachi et al. 2017). Jelmini et al. (2018) reported no  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 change in handgrip force or EMG activity after three static stretching repetitions of 45-s each of  
5  
6 the finger flexors, but did find a 10.8% decrease in rate of force generation. In contrast, Caldwell  
7  
8 et al. (Caldwell et al. 2019a) found a near significant, small magnitude improvement ( $p=0.06$ ,  
9  
10  $d=0.22$ , 12.1%) in contralateral drop jump height and no change in ground contact time following  
11  
12 four repetitions of 30-s of unilateral SS. Hence, with this conflict in the literature it is not known  
13  
14 whether there is sufficient evidence to substantiate that unilateral SS generally has global  
15  
16 (contralateral or non-local) consequences (i.e. positive, negative or no significant effect) on  
17  
18 performance.  
19  
20  
21  
22

23  
24 The possibility of non-local force impairments from unilateral stretching may impact  
25  
26 training and rehabilitation. Sustained stretching of one muscle group not only improves ROM of  
27  
28 other non-stretched muscles (Behm et al. 2016b) but may also impair force and power. The  
29  
30 sequence of training may be a consideration since for example, lower body stretching could  
31  
32 negatively affect subsequent upper body strength and vice versa. The objective of this scoping  
33  
34 review was to investigate and summarize the literature on non-local stretching effects on force and  
35  
36 power (i.e. jump height) output to determine whether the prolonged SS-induced impairments  
37  
38 observed with stretched limbs are generally present in non-local, non-stretched, muscle groups. In  
39  
40 addition, we aimed to identify gaps in the literature to facilitate future lines of enquiry with regard  
41  
42 to this research area.  
43  
44  
45  
46  
47

## 48 **Methods**

### 49 *Search strategy*

50  
51  
52  
53 Literature searches following the Preferred Reporting Items for Systematic review and  
54  
55 Meta-Analysis Protocols (PRISMA-P) review guidelines was performed by two co-authors (SA  
56  
57 and DGB). These searches were conducted separately and independently using the PubMed,  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 Scopus, SPORTDiscus, Web of Science, Cochrane, and CINHAL Plus databases. The topic was  
5  
6 searched using a Boolean search strategy with the operator “OR” for the following title keywords:  
7  
8 stretch, stretching, flexibility, range of motion, non-local, unilateral, contralateral, ipsilateral,  
9  
10 crossover, or remote with the operator “AND” for terms such as performance, strength, force,  
11  
12 power and jump (Figure 1). Researchers’ personal computer databases were also examined for  
13  
14 related articles. Related peer-reviewed publications written in English, spanned the period from  
15  
16 1999 to November 2020. The search syntax was saved in PubMed so that we received regular  
17  
18 updates on the publication of further relevant studies. The selected articles were also cross  
19  
20 referenced by the authors to identify relevant studies that might have been overlooked in the  
21  
22 database search. In selecting studies for inclusion, a review of all relevant article titles was  
23  
24 conducted before an examination of article abstracts and, then, full published articles (Figure 1).  
25  
26  
27  
28  
29

### 30 31 *Data extraction*

32  
33 Data were extracted from gathered articles with a form created in Microsoft Excel. Where  
34  
35 required data were not clearly or completely reported, article authors were contacted for  
36  
37 clarification.  
38  
39

### 40 41 *Inclusion and exclusion criteria (study selection)*

42  
43 Articles investigating non-local stretching and performance measures were included  
44  
45 according to PICOS (Population, Intervention, Comparison and Outcomes) criteria (Table 1).  
46  
47

48 PLACE TABLE 1 AND FIGURE 1 (FLOW DIAGRAM) APPROXIMATELY HERE  
49

### 50 51 *Systematic review analysis*

52  
53 Means and standard deviations for a strength (force or torque) performance measure were  
54  
55 used to calculate an effect size ( $d = M_1 - M_2 / SD \text{ pooled}$ ) (Cohen 1988). Effect sizes descriptors  
56  
57 were trivial ( $<0.2$ ), small ( $0.2 - <0.5$ ), moderate ( $0.5 - <0.8$ ) and large ( $\geq 0.8$ ) (Cohen 1988). With  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 only six studies investigating the effects of unilateral stretching on non-local muscle performance  
5  
6 (Table 2), the number of studies for a full, meaningful meta-analysis was considered low. Hence,  
7  
8 weighted means ( $[\text{effect size for study \#1} \times (\text{number of participants}/\text{total participants}) + \text{effect}$   
9  
10  $\text{size for study \#2} \times (\text{number of participants}/\text{total participants}) + \dots] = \text{weighted mean of all}$   
11  
12 selected studies) were calculated to illustrate the average response.  
13  
14

15  
16 PLACE TABLE 2 APPROXIMATELY HERE  
17

### 18 *Assessment of risk of bias*

19  
20  
21 The Physiotherapy Evidence Database (PEDro) scale was used to assess the risk of bias  
22  
23 and methodological quality of the included studies (Table 3). This scale evaluates internal study  
24  
25 validity on a scale from 0 (high risk of bias) to 10 (low risk of bias). Two reviewers (DGB and  
26  
27 SA) independently rated each study. Any ratings that yielded different results were further  
28  
29 adjudicated by a third reviewer. This rating was then used in the risk of bias scale. A median score  
30  
31 of  $\geq 6$  represents the threshold for studies with a low risk of bias (Maher et al. 2003).  
32  
33  
34

35  
36 PLACE TABLE 3 APPROXIMATELY HERE  
37

### 38 **Results**

39  
40  
41 Table 2 illustrates the individual study characteristics. The six studies were acute  
42  
43 intervention studies with three studies involving males and three studies including both sexes. Five  
44  
45 of the six studies were published in 2017 or thereafter. Participant characteristics and stretch  
46  
47 intervention characteristics are provided in Tables 4 and 5 respectively. From the six studies, the  
48  
49 average SS-induced force deficit of the stretched muscle (weighted means) was a small magnitude  
50  
51  $(-6.7 \pm 7.1\%, d = -0.35; \text{range: } 0.01 \text{ to } -1.8)$ , whereas the contralateral, non-stretched, muscle also  
52  
53 showed a small magnitude effect size of  $(-4.0 \pm 4.9\%, d = -0.22; \text{range: } 0.08 \text{ to } -1.1)$ . The five  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



1  
2  
3  
4 control measures (no stretch intervention) from three studies (Behm et al. 2019, Caldwell et al.  
5  
6 2019a, Ce et al. 2020) exhibited a trivial strength deficit effect size of -0.03.  
7  
8

9 PLACE TABLES 4-5 APPROXIMATELY HERE  
10

## 11 **Discussion**

12  
13  
14 The major finding from this scoping review was that both the stretched and contralateral,  
15  
16 non-stretched limbs of the young adult participants in these studies experienced small magnitude,  
17  
18 SS-induced force decrements. This finding highlights the global (non-local) effects of prolonged,  
19  
20 unilateral, static stretching; in that the resultant force decrements experienced due to the stretching  
21  
22 of a limb do not necessarily manifest only in that limb.  
23  
24

25  
26 In accord with prior reviews, prolonged SS (>60 s per muscle group) without a proper  
27  
28 dynamic warm-up can induce small magnitude strength impairments (Behm 2018; Behm and  
29  
30 Chaouachi 2011; Behm et al. 2016a; 2020; Chaabene et al. 2019; Kay and Blazevich 2012) in the  
31  
32 stretched muscle. The typical stretching routine used in the studies in this review involved six  
33  
34 repetitions of 36 s each for a total duration of approximately 216 s (3 min: 36 s), which exceeds  
35  
36 the recommended maximum duration of 60 s of SS per muscle group. Furthermore, unlike the  
37  
38 typical sport or training scenario, none of the studies included in this review included a full  
39  
40 dynamic warm-up (prior aerobic activity and activity-specific dynamic activity following the  
41  
42 stretching) as suggested in prior reviews (Behm et al. 2016a; 2020). Although these results should  
43  
44 be expected for the stretched muscle based on the prolonged stretching interventions, this is the  
45  
46 first review to report small magnitude, force impairments in a non-local, non-stretched muscle as  
47  
48 a result of unilateral SS. Hence, the same SS routine characteristics that induce decrements in the  
49  
50 stretched muscle can also incur deficits in a non-stretched muscle, but due to different mechanisms.  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4           Although the average result indicated SS-induced non-local force impairments, an analysis  
5  
6 of the individual studies demonstrates variable responses. Behm et al. (2019) imposed eight SS  
7  
8 repetitions of 30 s each on the quadriceps and hamstrings and reported small magnitude (11.1%;  
9  
10  $d=-0.31$ ) contralateral MVC force deficits. Similarly, 5 x 45 s repetitions of knee extensors SS  
11  
12 generated large magnitude (-11.0%;  $d = -1.1$ ) impairments of knee extensors MVC force (Ce et al.  
13  
14 2020). In contrast, there were trivial (1.2%;  $d=0.08$ ), non-significant contralateral knee flexion  
15  
16 isokinetic torque changes following eight repetitions of 30-s each of unilateral hip flexion static  
17  
18 stretching (Chaouachi et al. 2017), trivial (-4.9%;  $d= -0.17$ ) knee flexion MVC force deficits  
19  
20 following 10 x 30 s SS of the hamstrings (Killen et al. 2019) and trivial (-4.2%;  $d= -0.15$ ) knee  
21  
22 flexion MVC force deficits with 4 repetitions of 30 s SS of the hamstrings (Caldwell et al. 2019a).  
23  
24 Jelmini et al. (2018) reported no change in handgrip force or EMG activity after three SS  
25  
26 repetitions of 45-s each of the finger flexors, but did report a small magnitude, 10.8% ( $d= -0.29$ )  
27  
28 decrease in rate of force generation.  
29  
30  
31  
32  
33

34  
35           Although jump performance (proxy measure of power) was not included in the present  
36  
37 weighted mean results, there is also variability in these findings. Ten SS of 30 s each of the upper  
38  
39 limbs (shoulder horizontal abduction) produced large magnitude deficits in countermovement  
40  
41 jump ground reaction forces (-25%;  $d = 1.41$ ) (Marchetti et al. 2014). In contrast, Caldwell et al.  
42  
43 (2019a) found a near significant, small magnitude improvement ( $p=0.06$ ,  $d=0.22$ , 12.1%) in  
44  
45 contralateral drop jump height following four repetitions of 30-s of unilateral SS.  
46  
47  
48  
49

50           In summary, although, the weighted means illustrated a small magnitude, non-local force  
51  
52 deficit, three measures indicated deficits in force output, while four measures showed only trivial  
53  
54 changes. The substantially larger effect size of the Ce et al. (2020) study, in concert with the small  
55  
56 magnitude impairments of the Behm et al. (2019) and Jelmini et al. (2018) studies counterbalanced  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 the four trivial measures resulting in an overall mean small magnitude force deficit. Further  
5  
6 research is necessary to strengthen the aggregate findings in the literature on the effect of different  
7  
8 types of unilateral stretching on non-local force as well as power and other physiological and  
9  
10 functional measures (i.e. sprint speed, sporting skills such as kicking and throwing).  
11  
12 Morphological mechanisms would not be a consideration of non-local force or power impairments  
13  
14 since the non-stretched muscle would not have experienced appreciable stress or strain. While  
15  
16 Halperin et al. (2015) suggested four possible mechanisms contributing to non-local muscle fatigue  
17  
18 including biochemical, biomechanical, neural or psycho-physiological factors, their relative  
19  
20 contribution to non-local stretching force and power impairments may not be similar.  
21  
22  
23  
24  
25

26 A fatigued muscle can accumulate a variety of metabolites such as potassium (alteration of  
27  
28 electrochemical gradients inhibiting muscle excitability (Juel 1986; Nordsborg et al. 2003)),  
29  
30 hydrogen ions (reduction in the force per cross bridge and myofibrillar Ca<sup>2+</sup> sensitivity (Fitts  
31  
32 2008)), inorganic phosphates (Debold et al. 2016), reactive oxygen species (Reid 2016) among  
33  
34 other metabolites. These metabolites dispersed from the exercised muscle can be transported to  
35  
36 non-exercised muscles throughout the body (Bangsbo et al. 1996; Halperin et al. 2014; Johnson et  
37  
38 al. 2014; Nordsborg et al. 2003) adversely affecting muscle contractility. However, a passive SS  
39  
40 routine is unlikely to generate metabolic expenditures that would result in a substantial  
41  
42 accumulation and global dispersal of force inhibiting metabolites (Colosio et al. 2020).  
43  
44  
45  
46  
47

48 Biomechanical factors concern the fatigue of core or trunk muscles that stabilise the body  
49  
50 when contracting a limb muscle. Decrements in core/trunk stability have been shown to  
51  
52 substantially impair limb force output (Anderson and Behm 2005; Behm and Anderson 2006;  
53  
54 Behm et al. 2010). This biomechanical mechanism is an unlikely factor as passive SS typically  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 does not induce considerable disruptive, fatiguing torques to the trunk that need to be countered  
5  
6  
7 by active stabilization of the trunk muscles.  
8

9         Neural inhibition has been postulated as a mechanism underlying stretch-induced  
10 performance impairments of the stretched limb (Behm 2018; Behm and Chaouachi 2011; Behm et  
11 al. 2016a; 2020) as well as a factor contributing to non-local muscle fatigue effects (Halperin et  
12 al. 2015). Using the interpolated twitch technique, voluntary activation of the contralateral  
13 quadriceps was reported to significantly decrease (-7%,  $d=-2.29$ ) following 4 x 45-s of unilateral  
14 static stretching (Ce et al. 2020). However, EMG changes of the non-stretched muscles following  
15 unilateral stretching is conflicting. There are reports of very large (-11 to -22%,  $d= 2.25-3.9$ ) (Ce  
16 et al. 2020) and small (-3.1%;  $d= -0.22$ ) (Marchetti et al. 2017) magnitude reductions in EMG  
17 activity during a MVC, which contrast with reports of no significant changes in MVC EMG  
18 activity (Behm 2019; Caldwell et al. 2019a; Jelmini et al. 2018; Killen et al. 2019). These  
19 difficulties in ascertaining EMG changes may be related to the curvilinearity of the force-EMG  
20 relationship with a plateau of EMG activity at higher force levels (Perry and Bekey 1981). In the  
21 literature, there are conflicting effects of stretching on EMG activity of the stretched muscle (Behm  
22 et al. 2020), as well as a lack of substantial corticospinal excitability (Budini et al. 2017; Budini et  
23 al. 2019; Pulverenti et al. 2019) and intracortical inhibition (silent period) (Opplert et al. 2020;  
24 Pulverenti et al. 2019; 2020) effects. The short post-SS duration of inhibitory reflex activity and  
25 lack of corticospinal excitability changes, suggest the influence of neural inhibition on non-local  
26 strength is not conclusive.  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51

52         However, according to the recent Behm et al. (2020) review, there may be two possible  
53 neural mechanisms associated with ROM increases of a stretched muscle; exteroceptive reflex  
54 stimulation, which can attenuate excitatory sympathetic nervous activity (Guissard et al. 2001; Wu  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 et al. 1999) and reductions in motoneurone dendrites persistent inward currents (PIC) (decreasing  
5  
6 the amplification of central neural drive (Trajano et al. 2017, 2020). Exteroceptive reflexes arising  
7  
8 from cutaneous receptors have polysynaptic innervations to motoneurons (Jenner and Stephens  
9  
10 1982; Kearney and Chan 1999), which can inhibit sympathetic nervous system excitation (Wu et  
11  
12 al. 1999), contributing to reductions in muscle tone, heart rate and blood pressure (van den Berg  
13  
14 and Cabri 1999; Wu et al. 1999). However, exteroceptive reflex activity only persists for several  
15  
16 seconds after stretching desists (Delwaide et al. 1981; Guissard et al. 2001). A similar response  
17  
18 duration has been observed with the attenuation of facilitatory reflex activity from proprioceptive  
19  
20 Ia and Ib afferents and thus their activity also seems to be too transient to persistently impair  
21  
22 subsequent strength and power output. Hence, while strength deficits of a stretched or non-  
23  
24 stretched muscle detected within seconds of a SS might be partially attributed to exteroceptive  
25  
26 reflex influence on the sympathetic nervous system, deficits in the stretched or non-stretched  
27  
28 muscles shortly thereafter would likely not be related to exteroceptive or proprioceptive reflexes.  
29  
30 However, monoamines from sympathetic nervous system activity such as noradrenaline can affect  
31  
32 motoneurone PIC activity that might impact non-local responses.

33  
34  
35  
36  
37  
38  
39  
40  
41 PICs can amplify (motoneurons fire at higher frequencies) and prolong synaptic input  
42  
43 from supraspinal and reflex pathways augmenting motor output 5-fold (Lee and Heckman 2000).  
44  
45 SS might affect PIC-dependent amplification of central drive to the stretched muscle, which would  
46  
47 diminish force production (Trajano et al. 2017, 2020). It has been found that SS-induced reductions  
48  
49 in PICs were fully recovered by 10 minutes post-SS (Trajano et al. 2014a). These durations of  
50  
51 impairment and recovery closely match the SS-induced deficits outlined in studies undertaken by  
52  
53 Pulverenti et al. (2019; 2020) and Trajano and colleagues (Trajano et al. 2014b). However, these  
54  
55 findings apply to the stretched muscle's performance and it is unknown whether the diminution of  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 the PICs would also be experienced by contralateral motoneurons affecting non-local muscle  
5  
6 strength. PIC activity is augmented by monoamines such as serotonin and noradrenaline (released  
7  
8 predominantly from sympathetic nerve fibres) (Lee and Heckman 2000). With a static stretching-  
9  
10 induced reduction in sympathetic activity (Inami et al. 2014, Thomas et al. 2021), there would be  
11  
12 a decrease in noradrenaline release globally, affecting non-local motoneurone PIC activity. In  
13  
14 summary, there are conflicting effects of stretching on EMG activity (Behm et al. 2020), a lack of  
15  
16 substantial corticospinal excitability (Budini et al. 2017; Budini et al. 2019; Pulverenti et al. 2019)  
17  
18 and intracortical inhibition (Opplert et al. 2020; Pulverenti et al. 2019; 2020) effects, which in  
19  
20 combination with the short post-SS duration of inhibitory reflex activity, suggest the effect of these  
21  
22 neural factors or measures may not be the primary mechanism but need further investigation. The  
23  
24 effects of unilateral stretching on contralateral motoneuron PIC activity may be a possible  
25  
26 mechanistic candidate but these PIC effects have not been examined in non-stretched muscles.  
27  
28  
29  
30  
31  
32

33  
34 Sustaining focus or concentration during prolonged SS would involve a level of cognitive  
35  
36 demand to maintain joint position and stretch intensity (Behm et al. 2020). Prolonged cognitive  
37  
38 activity can induce mental fatigue (impaired cognitive functioning), which can negatively affect  
39  
40 subsequent performance, principally with prolonged activities (Marcora et al. 2009; Pageaux et al.  
41  
42 2013; 2014). Mental fatigue can lead to the perception that an activity is more demanding, and  
43  
44 therefore an individual may succumb earlier or exert less effort (Marcora et al. 2009; Pageaux et  
45  
46 al. 2013; 2014). Mental fatigue would not be specific to a single muscle or muscle group. Halperin  
47  
48 et al. (2015) postulated that mental fatigue could be a major contributor to non-local muscle fatigue  
49  
50 as non-local muscle fatigue is more apparent with prolonged fatiguing tests that would require  
51  
52 sustained concentration. Although speculative, the concentration necessary to maintain a  
53  
54 prolonged SS to or near the point of discomfort could induce some mental fatigue that would  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 impair the ability of the individual to subsequently fully activate and co-ordinate the contraction  
5  
6 of multiple muscle groups (activate agonists, synergists and core stabilizers, and inhibit antagonists  
7  
8 muscles) to achieve maximal force output. Nonetheless, as there is a paucity of literature on non-  
9  
10 local stretching effects, more research is needed on both the non-local responses and the possible  
11  
12 mechanisms (i.e. neural inhibition and cortical fatigue).  
13  
14

15  
16 Another possible mechanism contributing to non-local force and power deficits may be  
17  
18 myofascial chains. Myofascial connective tissue is reported to furnish a network of myofascial  
19  
20 chains or meridians (Wilke et al. 2016a). Human (Norton-Old et al. 2013; van Wingerden et al.  
21  
22 1993; Vleeming et al. 1995) and animal (Schleip et al. 2012; Yahia et al. 1993) cadaveric studies  
23  
24 reveal that fascia can modify its stiffness and transfer stress to nearby structures. The myofascial  
25  
26 transfer of force is conducted through longitudinal (Benetazzo et al. 2011; Eng et al. 2014) and  
27  
28 spiral (oblique) (Myers 2001) lines as well as with transversal orientations to synergists and  
29  
30 antagonists (Huijing et al. 2007; Maas et al. 2001; 2005; Meijer et al. 2007). Improved acute  
31  
32 cervical ROM after hamstrings (Wilke et al. 2017) and gastrocnemius (Wilke et al. 2016b)  
33  
34 stretching was accredited to myofascial longitudinal and spiral lines. But the scope of these  
35  
36 transfer effects has been contested. Krause et al. (2016) proposed that myofascial strain is  
37  
38 transferred predominantly to neighbouring skeletal muscles. The myofascial literature typically  
39  
40 report force transmission that is distal to proximal or vice versa (i.e. stretch the plantar flexors  
41  
42 and increase cervical ROM). The unilateral, stretch-induced, non-local muscle force and power  
43  
44 impairments reported in this review occur primarily in contralateral muscles. Whereas  
45  
46 longitudinal or spiral chains may affect force output of neighbouring muscles or along a  
47  
48 longitudinal kinetic chain, they are unlikely to significantly contribute to contralateral limb  
49  
50 muscle impairments.  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4           Stretch-induced non-local force deficits can have implications for both training and  
5  
6 rehabilitation. Individuals should be aware that prolonged high intensity stretching of one muscle  
7  
8 group can not only increase ROM of other non-stretched muscles (Behm et al. 2016b) but may  
9  
10 also impair force and power. For example, with training or rehabilitation, extensive lower body  
11  
12 stretching prior to training could affect subsequent upper body force output (i.e., bench press or  
13  
14 shoulder press) or prolonged upper body stretching could negatively impact the ability to sprint,  
15  
16 jump or kick a ball. Stretch-induced impairments of a stretched muscle are trivial when less than  
17  
18 60 s of SS and a full warm-up is incorporated (Behm 2018; Behm and Chaouachi 2011; Behm et  
19  
20 al. 2016a; 2020; Chaabene et al. 2019; Kay and Blazevich 2012), but it is unknown if the same  
21  
22 parameters would nullify non-local stretching effects. These issues provoke the need for further  
23  
24 research.  
25  
26  
27  
28  
29

### 30 31 **Conclusions**

32  
33           The limited scope of literature examining non-local effects of prolonged SS reveal that  
34  
35 both the stretched and contralateral, non-stretched limbs of young adult participants demonstrate  
36  
37 on average, small magnitude, strength deficits. However, this assertion must be viewed with  
38  
39 caution as the frequency of these effects were nearly equally divided with three measures  
40  
41 demonstrating deficits, and four measures showing trivial changes. Possible mechanisms  
42  
43 underlying SS-induced non-local force deficits may involve a combination of factors such as a  
44  
45 decrease in sympathetic activation affecting overall central nervous system excitation and  
46  
47 motoneurone PIC activity as well as a mental energy deficit (cortical fatigue) adversely affecting  
48  
49 concentration to maximally activate and coordinate muscle contractions. Myofascial chains may  
50  
51 affect the force output of neighbouring muscles or more distant muscles located along a  
52  
53 longitudinal or spiral meridian. Unfortunately, there is insufficient evidence to identify a  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



1  
2  
3  
4 predominant mechanism(s). This review will hopefully act as a clarion to motivate researchers to  
5  
6 add and contribute to the area of non-local stretching effects as it has implications for training and  
7  
8 rehabilitation as well as insights into psychological and physiological mechanisms. Areas for  
9  
10 future investigation include the volume, intensity and type of stretching that might elicit non-local  
11  
12 force, power, speed and other performance impairments, whether similar to a stretched muscle;  
13  
14 would a full warm-up nullify possible decrements, upper versus lower body stretching effects,  
15  
16 possible differences associated with age, sex, trained state, muscle type, joint, and the underlying  
17  
18 mechanisms.  
19  
20  
21  
22  
23  
24  
25

## 26 **Compliance with Ethical Standards**

27  
28 Funding: Partial funding was provided by the Natural Science and Engineering Research Council  
29  
30 of Canada.  
31  
32

33  
34 Conflicts of Interest: The authors declare no conflicts of interest with the contents of this  
35  
36 manuscript.  
37

38  
39 Authorship Contributions: All authors contributed to the conception, writing and revisions to the  
40  
41 manuscript. DGB, SA, conducted the literature search and collected and collated the data. DGB  
42  
43 wrote the first draft of the manuscript. JM, BD and UG provided input into the analysis and revised  
44  
45 the original manuscript.  
46  
47

## 48 Data Availability Statement

49  
50 All pertinent data are provided in the listed tables (PEDro scale analysis, individual and mean  
51  
52 study characteristics) and figures (PRISMA flow chart).  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 **Table Legends**  
5

6 **Table 1:** PICOS selection criteria  
7

8 **Table 2:** Individual study characteristics  
9

10 **Table 3:** Physiotherapy Evidence Database (PEDro) scale used to assess the risk of bias and  
11 methodological quality of the included studies. PEDro Scale: Y: Yes, N: No  
12

13 **Table 4:** Mean study characteristics: Mean  $\pm$  standard deviation (participants, age) or number of  
14 studies (sex, trained state). Acronyms: RT: resistance trained, PA: physically active  
15

16 **Table 5:** Intervention characteristics: Mean  $\pm$  standard deviation or number of studies (static  
17 stretch intensity, tested muscle groups, dependent variables). Acronyms: MVC: maximal  
18 voluntary isometric contraction, POD: point of discomfort  
19

20  
21  
22  
23  
24 **Figure Legends**  
25

26 **Figure 1:** PRISMA flow chart illustrating different phases of the search and study selection using  
27 the following title based nomenclature: stretch, stretching, flexibility, range of motion, non-local,  
28 unilateral, contralateral, ipsilateral, crossover, or remote.  
29

30 **Figure 2:** Relative (%) change in force output following unilateral stretching.  
31 Abbreviations: Contra: contralateral (unilateral stretching of a muscle and testing of the  
32 contralateral muscle(s), Ipsi: ipsilateral (stretching and testing of the same muscle(s)).  
33

34 **Figure 3:** The effects of unilateral static stretching (SS) on subsequent non-local muscle  
35 performance.  
36

37 **A)** Mean characteristics of stretching protocols used in the reviewed studies.

38 **B)** Effects of unilateral static stretching (SS) on the stretched and contralateral non-stretched  
39 muscles.

40 **C)** Mechanisms potentially underpinning the SS-induced performance impairment. ES: mean  
41 effect size

42 **D)** Gaps in the literature to be investigated by future research.

43 Acronyms: CMJ: countermovement jump, DJ: drop jump, EMG: electromyography, GRF:  
44 ground reaction force  
45

## References

- Anderson K, and Behm DG (2005) The impact of instability resistance training on balance and stability. *Sports Med* 35(1):43-53. <http://www.ncbi.nlm.nih.gov/pubmed/15651912>
- Avela J, Kyriläinen H, and Komi PV (1999) Altered reflex sensitivity after repeated and prolonged passive muscle stretching. *J Appl Physiol* 86(4):1283-1291
- Bangsbo J, Madsen K, Kiens B, Richter EA (1996) Effect of muscle acidity on muscle metabolism and fatigue during intense exercise in man. *J Physiol* 495 ( Pt 2): 587-596. doi:10.1113/jphysiol.1996.sp021618
- Behm, DG (2018) *The Science and Physiology of Flexibility and Stretching: Implications and Applications in Sport Performance and Health*. Routledge Publishers, London, UK. Chapters 3-6
- Behm DG, Anderson,KG (2006) The role of instability with resistance training. *J Strength Cond Res* 20(3):716-722. doi:10.1519/R-18475.1
- Behm DG, Chaouachi A (2011) A review of the acute effects of static and dynamic stretching on performance. *Eur J Appl Physiol* 111(11):2633-2651. doi:10.1007/s00421-011-1879-2
- Behm DG, Drinkwater EJ, Willardson JM, Cowley PM (2010) The use of instability to train the core musculature. *Appl Physiol Nutr Metab* 35(1):91-108. doi:10.1139/H09-127
- Behm DG, Blazeovich AJ, Kay AD, McHugh M (2016a) Acute effects of muscle stretching on physical performance, range of motion, and injury incidence in healthy active individuals: a systematic review. *Appl Physiol Nutr Metab* 41(1):1-11. doi:10.1139/apnm-2015-0235
- Behm DG, Cavanaugh T, Quigley P, Reid JC, Nardi PS, Marchetti PH (2016b) Acute bouts of upper and lower body static and dynamic stretching increase non-local joint range of motion. *Eur J Appl Physiol* 116(1):241-249. doi:10.1007/s00421-015-3270-1
- Behm DG, Kay AD, Trajano GS, Blazeovich AJ (2020) Mechanisms underlying performance impairments following prolonged static stretching without a comprehensive warm-up. *Eur J Appl Physiol* DOI: 10.1007/s00421-020-04538-8
- Behm DG, Lau RJ, O'Leary JJ, Rayner MCP, Burton EA, Lavers L (2019) Acute effects of unilateral self-administered static stretching on contralateral limb performance. *Journal of Performance Health Research* 3(1):1-7. doi:DOI: 10.25036/jphr.2019.3.1.behm
- Benetazzo L, Bizzego A, De Caro R, Frigo G, Guidolin D, and Stecco C (2011) 3D reconstruction of the crural and thoracolumbar fasciae. *Surg Radiol Anat* 33(10):855-862 doi:10.1007/s00276-010-0757-7
- Budini F, Gallasch E, Christova M, Rafolt D, Rauscher AB, Tilp M (2017) One minute static stretch of plantar flexors transiently increases H reflex excitability and exerts no effect on corticospinal pathways. *Exper Physiol* 102(8): 901-910
- Budini F, Kemper D, Christova M, Gallasch E, Rafolt D, Tilp M (2019) Five minutes static stretching influences neural responses at spinal level in the background of unchanged corticospinal excitability. *J Musculoskel Neuronal Interactions* 19(1):30-36
- Caldwell SL, Bilodeau,RLS, Cox MJ, Peddle D, Cavanaugh T, Young JD, et al. (2019a) Unilateral hamstrings static stretching can impair the affected and contralateral knee extension force but improve unilateral drop jump height. *Eur J Appl Physiol* 119(9):1943-1949 doi:10.1007/s00421-019-04182-x
- Caldwell SL, Bilodeau, RLS, Cox MJ, Behm DG (2019b) Twice daily, self-administered band stretch training improves quadriceps isometric force and drop jump characteristics. *J Sport Sci Med* 18:544-551

1  
2  
3  
4 Ce E, Coratella G, Bisconti AV, Venturelli M, Limonta E, Doria C, et al. (2020) Neuromuscular  
5 versus Mechanical Stretch-induced Changes in Contra- versus Ipsilateral Muscle. *Med Sci Sports*  
6 *Exerc.* doi:10.1249/MSS.0000000000002255  
7  
8 Chaabene H, Behm DG, Negra Y, Granacher U (2019) Acute Effects of Static Stretching on  
9 Muscle Strength and Power: An Attempt to Clarify Previous Caveats. *Front Physiol* 10:1468-  
10 1472 doi:10.3389/fphys.2019.01468  
11  
12 Chaouachi A, Padulo J, Kasmi S, Ben Othmen, A, Chatra M, Behm DG (2017) Unilateral static  
13 and dynamic hamstrings stretching increases contralateral hip flexion range of motion. *Clin*  
14 *Physiol Funct Imaging* 37(1):23-29 doi:10.1111/cpf.12263  
15  
16 Clark S, Christiansen A, Hellman DF, Hugunin JW, Hurst KM (1999) Effects of ipsilateral  
17 anterior thigh soft tissue stretching on passive unilateral straight-leg raise. *J Orthop Sports Phys*  
18 *Ther* 29(1):4-12 doi:10.2519/jospt.1999.29.1.4  
19  
20 Cohen J (1988) *Statistical power analysis for the behavioural sciences*. L. Erbraum Associates,  
21 Hillside N.J. pp. 48-96  
22  
23 Colosio AL, Teso M, Pogliaghi S (2020) Prolonged static stretching causes acute, nonmetabolic  
24 fatigue and impairs exercise tolerance during severe-intensity cycling. *Appl Physiol Nutr Metab*  
25 45(8):902-910 doi:10.1139/apnm-2019-0981  
26  
27 Debold EP, Fitts RH, Sundberg CW, Nosek TM (2016) Muscle fatigue from the perspective of a  
28 single crossbridge. *Med Sci Sports Exerc* 48(11):2270-2280  
29  
30 De-la-Cruz-Torres B, Carrasco-Iglesias C, Minaya-Munoz F, Romero-Morales C (2020)  
31 Crossover effects of ultrasound-guided percutaneous neuromodulation on contralateral hamstring  
32 flexibility. *Acupunct Med*: doi:10.1177/0964528420920283  
33  
34 Delwaide PJ, Toulouse P, Crenna P (1981) Hypothetical role of long-loop reflex pathways. *Appl*  
35 *Neurophysiol* 44(1-3): 171-176 <http://www.ncbi.nlm.nih.gov/pubmed/7294778>  
36  
37 Eng CM, Pancheri FQ, Lieberman DE, Biewener AA, Dorfmann, L (2014) Directional  
38 differences in the biaxial material properties of fascia lata and the implications for fascia  
39 function. *Ann Biomed Eng* 42(6):1224-1237 doi:10.1007/s10439-014-0999-3  
40  
41 Fitts RH (2008) The cross-bridge cycle and skeletal muscle fatigue. *J Appl Physiol* 104(2):551-  
42 558 doi:10.1152/jappphysiol.01200.2007  
43  
44 Guissard N, Duchateau J, Hainaut K (2001) Mechanisms of decreased motoneurone excitation  
45 during passive muscle stretching. *Experimental Brain Research* 137:163-169  
46  
47 Halperin I, Aboodarda SJ, Behm DG (2014) Knee extension fatigue attenuates repeated force  
48 production of the elbow flexors. *Eur J Sport Sci* 14(8):823-829.  
49 doi:10.1080/17461391.2014.911355.  
50  
51 Halperin I, Chapman DW, Behm DG (2015) Non-local muscle fatigue: effects and possible  
52 mechanisms. *Eur J Appl Physiol* 115(10):2031-2048 doi:10.1007/s00421-015-3249-y  
53  
54 Herzog W (2014) The role of titin in eccentric muscle contraction. *J Exp Biol* 217(Pt 16):2825-  
55 2833 doi:10.1242/jeb.099127  
56  
57 Huijing PA, van de Langenberg RW, Meesters JJ, Baan GC (2007) Extramuscular myofascial  
58 force transmission also occurs between synergistic muscles and antagonistic muscles. *J*  
59 *Electromyogr Kinesiol* 17(6):680-689 doi:10.1016/j.jelekin.2007.02.005  
60  
61 Inami T, Shimizu T, Baba R, Nakagaki A (2014) Acute changes in autonomic nerve activity  
62 during passive static stretching. *Amer J Sports Sci Med* 2(4): 166-170.  
63  
64 Perry, J., and Bekey, G.A. 1981. EMG-force relationships in skeletal muscle. *CRC Critical*  
65 *Reviews in Biomedical Engineering*: 1-21.

1  
2  
3  
4 Jelmini JD, Cornwell A, Khodiguian N, Thayer J, Araujo AJ (2018) Acute effects of unilateral  
5 static stretching on handgrip strength of the stretched and non-stretched limb. *Eur J Appl Physiol*  
6 118(5):927-936 doi:10.1007/s00421-018-3810-6  
7  
8 Jenner JR, Stephens JA (1982) Cutaneous reflex responses and thier central nervous pathways  
9 studied in man. *J. Physiol.* 333:405-419  
10  
11 Johnson MA, Mills DE, Brown PI, Sharpe GR (2014) Prior upper body exercise reduces cycling  
12 work capacity but not critical power. *Med Sci Sports Exerc* 46(4):802-808.  
13 doi:10.1249/MSS.0000000000000159  
14  
15 Juel C (1986) Potassium and sodium shifts during in vitro isometric muscle contraction, and the  
16 time course of the ion-gradient recovery. *Pflugers Arch* 406(5):458-463.  
17 doi:10.1007/bf00583367  
18  
19 Kay AD, Blazevich AJ (2009) Isometric contractions reduce plantar flexor moment, Achilles  
20 tendon stiffness, and neuromuscular activity but remove the subsequent effects of stretch. *J Appl*  
21 *Physiol* 107(4):1181-1189 doi:10.1152/jappphysiol.00281.2009  
22  
23 Kay AD, Blazevich AJ (2012) Effect of acute static stretch on maximal muscle performance: a  
24 systematic review. *Med Sci Sports Exerc* 44(1):154-164 doi:10.1249/MSS.0b013e318225cb27  
25  
26 Kay AD, Husbands-Beasley J, Blazevich AJ (2015) Effects of Contract-Relax, Static Stretching,  
27 and Isometric Contractions on Muscle-Tendon Mechanics. *Med Sci Sports Exerc* 47(10):2181-  
28 2190 doi:10.1249/MSS.0000000000000632  
29  
30 Kearney RE, Chan CWY (1999) Reflex response of human arm muscles to cutaneous  
31 stimulation of the foot. *Brain Res* 170:214-217.  
32  
33 Killen BS, Zelizney KL, Ye X (2019) Crossover Effects of Unilateral Static Stretching and Foam  
34 Rolling on Contralateral Hamstring Flexibility and Strength. *J Sport Rehabil* 28(6):533-539  
35 doi:10.1123/jsr.2017-0356  
36  
37 Konrad A, Stafilidis S, Tilp M (2017a) Effects of acute static, ballistic, and PNF stretching  
38 exercise on the muscle and tendon tissue properties. *Scand J Med Sci Sports* 27(10):1070-1080  
39 doi:10.1111/sms.12725  
40  
41 Konrad A, Budini F, Tilp M (2017b) Acute effects of constant torque and constant angle  
42 stretching on the muscle and tendon tissue properties. *Eur J Appl Physiol* 117(8): 1649-1656  
43 doi:10.1007/s00421-017-3654-5  
44  
45 Krause F, Wilke J, Vogt L, Banzer W (2016) Intermuscular force transmission along myofascial  
46 chains: a systematic review. *J Anat* 228(6):910-918 doi:10.1111/joa.12464  
47  
48 Lee EJ, Joumaa V, Herzog W (2007) New insights into the passive force enhancement in skeletal  
49 muscles. *J Biomech* 40(4):719-727 doi:10.1016/j.jbiomech.2006.10.009  
50  
51 Lee RH, Heckman C (2000) Adjustable amplification of synaptic input in the dendrites of spinal  
52 motoneurons in vivo. *J Neurosci* 20(17):6734-6740  
53  
54 Leonard TR, Herzog W (2010) Regulation of muscle force in the absence of actin-myosin-based  
55 cross-bridge interaction. *Am J Physiol Cell Physiol* 299(1):C14-20  
56 doi:10.1152/ajpcell.00049.2010  
57  
58 Lima BN, Lucareli PR, Gomes WA, Silva JJ, Bley AS, Hartigan EH, et al. (2014) The acute  
59 effects of unilateral ankle plantar flexors static- stretching on postural sway and gastrocnemius  
60 muscle activity during single-leg balance tasks. *J Sports Sci Med* 13(3):564-570  
61 <http://www.ncbi.nlm.nih.gov/pubmed/25177183>  
62  
63 Maas H, Baan GC, Huijting PA (2001) Intermuscular interaction via myofascial force  
64 transmission: effects of tibialis anterior and extensor hallucis longus length on force transmission  
65

1  
2  
3  
4 from rat extensor digitorum longus muscle. *J Biomech* 34(7):927-940 doi:10.1016/s0021-  
5 9290(01)00055-0  
6  
7 Maas H, Meijer HJ, Huijing PA (2005) Intermuscular interaction between synergists in rat  
8 originates from both intermuscular and extramuscular myofascial force transmission. *Cells*  
9 *Tissues Organs* 181(1):38-50 doi:10.1159/000089967  
10  
11 Magnusson SP, Simonsen EB, Aagaard P, Sorensen H, Kjaer M (1996) A mechanism for altered  
12 flexibility in human skeletal muscle. *Journal of Physiology* 497 ( Pt 1):291-298 PM:8951730  
13  
14 Magnusson SP, Simonsen EB, Aagaard P, Boesen J, Johannsen F, Kjaer M (1997) Determinants  
15 of musculoskeletal flexibility: viscoelastic properties, cross-sectional area, EMG and stretch  
16 tolerance. *Scand J Med Sci Sports* 7(4):195-202. <https://www.ncbi.nlm.nih.gov/pubmed/9241023>  
17  
18 Maher CG, Sherrington C, Herbert RD, Moseley AM, Elkins M (2003) Reliability of the PEDro  
19 scale for rating quality of randomized controlled trials. *Phys Ther* 83(8):713-721  
20 <https://www.ncbi.nlm.nih.gov/pubmed/12882612>  
21  
22 Marchetti PH, Silva FH, Soares EG, Serpa EP, Nardi PS, Vilela Gode B, et al. (2014) Upper  
23 limb static-stretching protocol decreases maximal concentric jump performance. *J Sports Sci*  
24 *Med* 13(4): 945-950. <http://www.ncbi.nlm.nih.gov/pubmed/25435789>  
25  
26 Marchetti PH, Reis RG, Gomes WA, da Silva WA, Soares E.G, de Freitas FS, Behm DG (2017)  
27 Static-stretching of the pectoralis major decreases triceps brachii activation during a maximal  
28 isometric bench press. . *Gazzetta Medica Italiana* 176:212-218 DOI:10.23736/S0393-  
29 3660.17.03452-0  
30  
31 Marcora SM, Staiano W, Manning V (2009) Mental fatigue impairs physical performance in  
32 humans. *J Appl Physiol* 106(3):857-864 doi:10.1152/jappphysiol.91324.2008  
33  
34 Meijer HJ, Rijkeljkhuizen JM, Huijing PA (2007) Myofascial force transmission between  
35 antagonistic rat lower limb muscles: effects of single muscle or muscle group lengthening. *J*  
36 *Electromyogr Kinesiol* 17(6):698-707. doi:10.1016/j.jelekin.2007.02.006.  
37  
38 Myers TW (2001) *Anatomy trains: Myofascial meridians for manual and movement therapists.*  
39 Chirchill Livingstone Publishers, Edinburgh, Scotland, UK. pp. 73-157  
40  
41 Nordborg N, Mohr M, Pedersen LD, Nielsen JJ, Langberg H, Bangsbo J (2003) Muscle  
42 interstitial potassium kinetics during intense exhaustive exercise: effect of previous arm exercise.  
43 *Am J Physiol Regul Integr Comp Physiol* 285(1):R143-148 doi:10.1152/ajpregu.00029.2003  
44  
45 Norton-Old KJ, Schache AG, Barker PJ, Clark RA, Harrison SM, Briggs CA (2013) Anatomical  
46 and mechanical relationship between the proximal attachment of adductor longus and the distal  
47 rectus sheath. *Clin Anat* 26(4):522-530 doi:10.1002/ca.22116  
48  
49 Opplert J, Paizis C, Papitsa A, Blazevich AJ, Cometti C, Babault N (2020) Static stretch and  
50 dynamic muscle activity induce acute similar increase in corticospinal excitability. *PLoS One*  
51 15(3):e0230388. doi:10.1371/journal.pone.0230388  
52  
53 Pageaux B, Marcora SM, Lepers R (2013) Prolonged mental exertion does not alter  
54 neuromuscular function of the knee extensors. *Med Sci Sports Exerc* 45(12):2254-2264  
55 doi:10.1249/MSS.0b013e31829b504a  
56  
57 Pageaux B, Lepers R, Dietz KC, Marcora SM (2014) Response inhibition impairs subsequent  
58 self-paced endurance performance. *Eur J Appl Physiol* 114(5):1095-1105. doi:10.1007/s00421-  
59 014-2838-5  
60  
61 Perry J, Bekey GA (1981) EMG-force relationships in skeletal muscle. *CRC Critical Rev*  
62 *Biomedical Engin* 7(1):1-21.  
63  
64 Phillips C, Powell T, Wiesendanger M (1971) Projection from low-threshold muscle afferents of  
65 hand and forearm to area 3a of baboon's cortex. *The Journal of physiology* 217(2): 419-446

1  
2  
3  
4 Prochazka A, Ellaway P (2012) Sensory systems in the control of movement. *Comprehensive*  
5 *Physiol* 2:2615-2627  
6  
7 Pulverenti TS, Trajano GS, Kirk BJC, Blazevich AJ (2019) The loss of muscle force production  
8 after muscle stretching is not accompanied by altered corticospinal excitability. *Eur J Appl*  
9 *Physiol* 119(10):2287-2299 doi:10.1007/s00421-019-04212-8  
10  
11 Pulverenti TS, Trajano GS, Walsh A, Kirk BJC, Blazevich AJ (2020) Lack of cortical or Ia-  
12 afferent spinal pathway involvement in muscle force loss after passive static stretching. *J*  
13 *Neurophysiol* 123(5):1896-1906 doi:10.1152/jn.00578.2019  
14  
15 Rathelot JA, Strick PL (2009) Subdivisions of primary motor cortex based on cortico-  
16 motoneuronal cells. *Proceedings of the National Academy of Sciences* 106(3):918-923  
17  
18 Reid MB (2016) Reactive oxygen species as agents of fatigue. *Med Sci Sports Exerc*  
19 48(11):2239-2246 DOI: 10.1249/MSS.0000000000001006  
20  
21 Schleip R, Duerselen L, Vleeming A, Naylor IL, Lehmann-Horn F, Zorn A, et al. (2012) Strain  
22 hardening of fascia: static stretching of dense fibrous connective tissues can induce a temporary  
23 stiffness increase accompanied by enhanced matrix hydration. *J Bodyw Mov Ther* 16(1):94-100  
24 doi:10.1016/j.jbmt.2011.09.003  
25  
26 Stephenson DG, Williams DA (1985) Temperature-dependent calcium sensitivity changes in  
27 skinned muscle fibres of rat and toad. *J Physiol* 360:1-12 doi:10.1113/jphysiol.1985.sp015600  
28  
29 Sugi H, Abe T, Kobayashi T, Chaen S, Ohnuki Y, Saeki Y, et al. (2013) Enhancement of force  
30 generated by individual myosin heads in skinned rabbit psoas muscle fibers at low ionic strength.  
31 *PLoS One* 8(5):e63658 doi:10.1371/journal.pone.0063658  
32  
33 Thomas E, Bellafiore M, Gentile A, Paoli A, Palma A, Bianco A (2021) Cardiovascular  
34 responses to muscle stretching: A systematic review and meta-analysis. *Int J Sports Med*  
35 doi:10.1055/a-1312-7131.  
36  
37 Trajano GS, Nosaka K, Blazevich AJ (2017) Neurophysiological Mechanisms Underpinning  
38 Stretch-Induced Force Loss. *Sports Med* 47(8):1531-1541 doi:10.1007/s40279-017-0682-6  
39  
40 Trajano GS, Seitz LB, Nosaka K, Blazevich AJ (2014a) Can passive stretch inhibit motoneuron  
41 facilitation in the human plantar flexors? *J Apply Physiol* 117(12): 1486-1492  
42  
43 Trajano GS, Nosaka K, Seitz LB, Blazevich AJ (2014b) Intermittent stretch reduces force and  
44 central drive more than continuous stretch. *Med Sci Sports Exerc* 46(5): 902-910  
45 doi:10.1249/MSS.0000000000000185  
46  
47 Trajano GS, Taylor JL, Orssatto LBR, McNulty CR, Blazevich AJ (2020) Passive muscle  
48 stretching reduces estimates of persistent inward current strength in soleus motor units. *J Exp*  
49 *Biol* 223(Pt 21). doi:10.1242/jeb.229922.  
50  
51 van den Berg F, Cabri J (1999) *Angewandte Physiologie – Das Bindegewebe des*  
52 *Bewegungsapparates verstehen und beeinflussen*. Georg Thieme Verlag, Stuttgart, Germany. pp.  
53 17-48  
54  
55 van Wingerden JP, Vleeming A, Snijders CJ, Stoeckart R (1993) A functional-anatomical  
56 approach to the spine-pelvis mechanism: interaction between the biceps femoris muscle and the  
57 sacrotuberous ligament. *Eur Spine J* 2(3):140-144 doi:10.1007/BF00301411  
58  
59 Vleeming A, Pool-Goudzwaard AL, Stoeckart R, van Wingerden JP, Snijders CJ (1995) The  
60 posterior layer of the thoracolumbar fascia. Its function in load transfer from spine to legs. *Spine*  
61 20(7): 53-758 <https://www.ncbi.nlm.nih.gov/pubmed/7701385>  
62  
63 Whalen A, Farrell K, Roberts S, Smith H, Behm DG (2019) Topical Analgesic Improved or  
64 Maintained Ballistic Hip Flexion Range of Motion with Treated and Untreated Legs. *J Sports Sci*  
65 *Med* 18(3):552-558. <https://www.ncbi.nlm.nih.gov/pubmed/31427878>

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Wilke J, Krause F, Vogt L, Banzer W (2016a) What Is evidence-based about myofascial chains: A Systematic Review. *Arch Phys Med Rehabil* 97(3):454-461 doi:10.1016/j.apmr.2015.07.023

Wilke J, Niederer D, Vogt L, Banzer W (2016b) Remote effects of lower limb stretching: preliminary evidence for myofascial connectivity? *J Sports Sci* 34(22):2145-2148 doi:10.1080/02640414.2016.1179776


Wilke J, Vogt L, Niederer D, Banzer W (2017) Is remote stretching based on myofascial chains as effective as local exercise? A randomised-controlled trial. *J Sports Sci* 35(20):2021-2027 doi:10.1080/02640414.2016.1251606

Wu G, Ekedahl R, Stark B, Carlstedt T, Nilsson B, Hallin RG (1999) Clustering of Pacinian corpuscle afferent fibres in the human median nerve. *Exper Brain Res* 126: 399-409


Yahia LH, Pigeon P, DesRosiers EA (1993) Viscoelastic properties of the human lumbodorsal fascia. *J Biomed Eng* 15(5):425-429. doi:10.1016/0141-5425(93)90081-9



**Figure 1:** PRISMA flow chart illustrating different phases of the search and study selection using the following title based nomenclature: stretch, stretching, flexibility, range of motion, non-local, unilateral, contralateral, ipsilateral, crossover, or remote.



**Figure 2:** Relative (%) change in force output following unilateral stretching.  
Abbreviations: Contra: contralateral (unilateral stretching of a muscle and testing of the contralateral muscle(s)), Ipsi: ipsilateral (stretching and testing of the same muscle(s)).




**Figure 3:****A: Unilateral stretching protocols from this review:**

$6.3 \pm 2$  repetitions of  $36.3 \pm 7.4$  seconds with  $19.3 \pm 5.7$  seconds recovery between stretches.

**B**

Prolonged unilateral SS effects on stretched limb performance



↑ Range of motion (Behm et al. 2016, 2020)

↓ MVC force (6 studies)

ES: -0.36 (-0.01 to -1.8)

↓ Force, power, jump height, sprint speed, balance and reaction/movement time

Prolonged unilateral SS effects on contralateral, non-stretched limb performance



↑ Range of motion (Behm et al. 2020)

↓ MVC force (6 studies)

ES: -0.22 (0.08 to -1.1)

↓ MVC force: 3 studies, no change: 4 studies

↓ MVC EMG: 2 studies, no change: 4 studies

↓ CMJ GRF (ES = -1.4: 1 study)


↑ DJ Height (ES = 0.22: 1 study)

**C****– Possible**

– Cortical fatigue / Mental Energy Deficit

– ↓ Persistent inward currents  
– ↓ Exteroceptive reflex effects on sympathetic NS

– Myofascial chain effects on neighbouring and longitudinal / spiral oriented muscles.

**✗ Unlikely or ? Unclear**

✗ Corticospinal excitability  
✗ Cortical silent period (intracortical inhibition)

✗ Biomechanical (core or trunk stability)

✗ Biochemical (circulating metabolites)

✗ Morphological  
?

**D****Future research**

Effects of different stretching types, intensities, volumes, rest periods, muscles, upper versus lower body stretching, age groups (<18 and >30 years), sex, trained state, incorporate full warm-ups, on force, power, speed, functional movements and other measures as well as the predominant mechanisms.

**Table 1:** PICOS selection criteria

PICOS	Selection Criteria
Population	Only young healthy untrained, recreationally active, or trained individuals aged 18-30 years were recruited for the applicable studies.
Intervention	Static stretching programs
Comparator	Active/passive control
Outcomes	Measure of acute pre- and post-stretching of a homologous or heterologous muscle (MVC, isokinetic torque, drop jump height)
Study design	Randomized controlled or controlled trials

MVC: maximal voluntary contraction

**Table 2:** Individual study characteristics

Study	Age	M/F	Trained state	Stretched muscle group	Repetitions /Sets	Duration (s)	Volume (s)	Rest (s)	Intensity	Tested muscle group	%Δ
Behm et al. 2019	23.4±6.3	M:14	PA	Quadriceps Hamstring	8	30	240	-	100% POD	IL KE MVC	-6.9
										ILC KE MVC	-5.3
										CL KE MVC	-11.1
										CLC KE MVC	-5.1
Caldwell et al. 2019a	22.5±5.1	22/18	RT & RA	Hamstrings	4	30	120	15	100% POD	IL KE MVC	-8.1
										CL KE MVC	-4.2
Ce et al. 2020	22±3	M:21	H	KE	5	30	150	15	80%- 90% POD	IL KE MVC	-19.7
										CL KE MVC	-11
										ILC KE MVC	0
										CLC KE MVC	-0.1
Chaouachi et al. 2017	18±2	M:14	HTR	SS hip extensors and KF	8	30	240	20	POD	IL KF torque at 60°/s	-3.1
										CL KF torque at 60°/s	1.2
Chaouachi et al. 2017	18±2	M:14	HTR	DS hip extensors and KF	8	30	240	20	Full ROM	IL KF torque at 60°/s	1.5
										CL KF torque at 60°/s	1.3
Jelmini et al. 2018	25-27 ±6-10	15/15	NRT	Finger flexors	3	45	135	30	POD	IL HG force	-4.4
										IL HG RFD	-17.3
										CL HG force	0.4
										CL HG RFD	-10.8
Killen et al. 2017	M: 26±3 F:27±2	13/10	PA	Hamstrings	10	30	300	30	100% POD	CL hamstrings MVC	-4.9

CL: contralateral, CLC: contralateral control, DS: dynamic stretch, M: male, F: female, H: healthy, HG: handgrip, HRT: highly trained rowers, IL: ipsilateral, ILC: Ipsilateral control, KE: knee extensors, KF: knee flexors, MVC: maximum voluntary contraction, NRT: non-resistance trained, PA: physically active, POD: point of discomfort, RA: recreationally active, RT: resistance trained, RFD: rate of force development, SS: static stretch

**Table 3:** Physiotherapy Evidence Database (PEDro) scale used to assess the risk of bias and methodological quality of the included studies. PEDro Scale: Y: Yes, N: No

Study	1	2	3	4	5	6	7	8	9	10	11	Total
Behm et al. 2019	N	Y	N	Y	Y	N	N	N	Y	Y	Y	6
Caldwell et al. 2019	Y	Y	N	Y	N	N	N	N	N	Y	Y	5
Ce et al. 2020	Y	N	N	Y	N	N	N	N	N	N	Y	3
Chaouachi et al. 2017	N	Y	N	Y	N	N	N	Y	Y	Y	Y	6
da Silva et al. 2015	Y	N	N	Y	N	N	N	Y	N	Y	Y	5
Jelmini et al. 2018	Y	N	N	Y	Y	N	N	Y	Y	Y	Y	7
Killen et al. 2017	Y	Y	N	Y	N	N	N	Y	Y	Y	Y	7
Mean												5.6
Median												6
Mode												6

PEDro scale criteria

1. Eligibility criteria were specified.
2. Subjects were randomly allocated to groups
3. Allocation was concealed.
4. The groups were similar at baseline regarding the most important prognostic indicators.
5. There was blinding of all subjects.
6. There was blinding of all therapists/researchers who administered the therapy/protocol.
7. There was blinding of all assessors who measured at least one key outcome.
8. Measures of at least one key outcome were obtained from more than 85% of the subjects initially allocated to groups.
9. All subjects for whom outcome measures were available received the treatment or control condition as allocated or, where this was not the case, data for at least one key outcome was analysed by “intention to treat”.
10. The results of between-group statistical comparisons are reported for at least one key outcome.
11. The study provides both point measures and measures of variability for at least one key outcome.

**Table 4:** Mean study characteristics: Mean  $\pm$  standard deviation (participants, age) or number of studies (sex, trained state).  
 Acronyms: RT: resistance trained, PA: physically active

# of Participants	Age (years)	Sex	Trained State
20.6 $\pm$ 8.3	21.1 $\pm$ 3.3	Males: 7 Females: 3	PA : 2 RT+PA: 1 Healthy: 2 Rowers: 1 RT: 1

**Table 5:** Intervention characteristics: Mean  $\pm$  standard deviation or number of studies (static stretch intensity, tested muscle groups, dependent variables). Acronyms: MVC: maximal voluntary isometric contraction, POD: point of discomfort

Stretched muscle groups	Repetitions	Static stretch durations	Rest between repetitions	Static stretch intensity	Tested muscle groups	Dependent variables
Hamstrings: 2 Quadriceps and hamstrings: 2 Quadriceps: 1 Plantar flexors: 1 Finger flexors: 1	6.3 $\pm$ 2.0	36.3 $\pm$ 7.4	19.3 $\pm$ 5.7	Maximum 70-90% POD: 2	KE: 3 KF: 2 PF: 1 HG: 1	MVC: 5, Isokinetic Torque: 1, Drop Jump: 1

Conflicts of Interest: The authors declare no conflicts of interest with the contents of this manuscript.

Authorship Contributions: All authors contributed to the conception, writing and revisions to the manuscript. DGB, SA, conducted the literature search and collected and collated the data. DGB wrote the first draft of the manuscript. JM, BD and UG provided input into the analysis and revised the original manuscript.