

Association of methylenetetrahydrofolate reductase C677T CT gene polymorphism with a non-dipping blood pressure pattern in morbidly obese patients

 Rea Levicki^{1*},
 Martina Lovrić Benčić^{2,3},
 Martina Matovinović^{2,3},
 Lada Bradić²,
 Kristina Gašparović^{2,3},
 Tamara Božina³,
 Nada Božina^{2,3},
 Jadranka Sertić^{2,3},
 Bojan Jelaković^{2,3}

¹Požega General County Hospital, Požega, Croatia

²University Hospital Centre Zagreb, Zagreb, Croatia

³University of Zagreb School of Medicine, Zagreb, Croatia

KEYWORDS: MTHFR C677T polymorphisms, arterial hypertension, non-dipper.

CITATION: *Cardiol Croat.* 2021;16(1-2):38-39. | <https://doi.org/10.15836/ccar2021.38>

***ADDRESS FOR CORRESPONDENCE:** Rea Levicki, Opća županijska bolnica Požega, Osječka 107, HR-34000 Požega, Croatia. / Phone: +385-98-550-309 / E-mail: rlevicki@gmail.com

ORCID: Rea Levicki, <https://orcid.org/0000-0003-3687-1310> • Martina Lovrić Benčić, <https://orcid.org/0000-0001-8446-6120> • Martina Matovinović, <https://orcid.org/0000-0002-6325-7394> • Lada Bradić, <https://orcid.org/0000-0001-8296-699X> • Kristina Gašparović, <https://orcid.org/0000-0002-1191-4831> • Tamara Božina, <https://orcid.org/0000-0001-5883-7979> • Nada Božina, <https://orcid.org/0000-0001-6016-1699> • Jadranka Sertić, <https://orcid.org/0000-0001-9581-9489> • Bojan Jelaković, <https://orcid.org/0000-0002-2546-4632>

Introduction: Region near the gene encoding methylenetetrahydrofolate reductase (MTHFR) is associated with blood pressure. CT and TT genotype of the C677T MTHFR gene are more common in obese hypertensive patients with BMI>29 kg/m².^{1,2} The aim of this study was to prove the connection between C667T polymorphism of MTHFR gene and non-dipping pattern in 24h ambulatory blood pressure monitoring (ABPM) in obese patients.

Patients and Methods: We included 33 patients from a multidisciplinary weight management program in which genetic analysis on MTHFR gene polymorphism was tested and 24h ABPM was performed. Patients were divided into 3 groups: 12 patients with MTHFR C677T healthy genotype CC (9 W, 3 M, age 48.83±9.82y, BMI 44.39±9.69 kg/m²), 14 patients with MTHFR C677T heterozygous mutation CT (11 W, 3 M, age 47.93±8.74y, BMI 40.9±5.96 kg/m²), 7 patients with MTHFR C677T homozygous mutation TT (6 W, 1 M, age 43.43±12.22y, BMI 37.46±5.79 kg/m²). In each group 24h ABPM results were analyzed, and dipping status was determined according to percentage of night systolic blood pressure drop and divided into 4 groups: inverse dipper (<0%), non-dipper (0-10%), dipper (10-20%), extreme dipper (>20%).

Results: Patients with MTHFR C677T: CT had the lowest average night blood pressure drop (8.51±5.9%), comparing to MTHFR C677T: TT (10.17±4.72 %) and MTHFR C677T: CC (10.53± 6.07%). Patients with MTHFR C677T:CT (**Figure 1**) were the only group with inverse dipper pattern present, but also with significant non-dipping pattern (inverse dipper 7.1%, non-dipper 35.7%, dipper 57.1%) comparing to patients with MTHFR C677T:TT (non-dipper 28.6%, dipper 71.4%) (**Figure 2**). Patients with MTHFR C677T:CC (**Figure 3**) had the highest level of dipper pattern, but also extreme dipper pattern was present (non-dipper 25%, dipper 66.7%, extreme dipper 8.3%).

RECEIVED:
 December 4, 2020
ACCEPTED:
 December 18, 2020




FIGURE 1. Dipping status distribution in patients with MTHFR C677T heterozygous CT mutation.

