



Citation for published version:

Roberts, V, Joiner, R, Russell, C, Bradley, R, Bowles, M, Bowes, A & Nice, T 2019, 'Mind and Body: an early intervention group programme for adolescents with self-harm thoughts and behaviours', *Education and Health*, vol. 37, no. 2. <<http://sheu.org.uk/sheux/EH/eh372vr.pdf>>

Publication date:
2019

Document Version
Peer reviewed version

[Link to publication](#)

Publisher Rights
Unspecified

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Mind and Body: an early intervention group programme for adolescents with self-harm thoughts and behaviours

Accepted for publication in Health and Education

Author details:

Dr Veronica Roberts is an Independent Educational Psychologist and Research Associate at the University of Bath Psychology Department.

Professor Richard Joiner is Professor of Educational Psychology at the University of Bath's, Department of Psychology.

Claire Russell is an Honorary Lecturer in the University of Bath's, Department of Psychology

Dr Terrence Nice is a lecturer in psychotherapy at the Centre for Professional Practice, University of Kent.

Mark Bowles is the Director of The Training Effect. Contact details:

<http://www.thetrainingeffect.co.uk>

Aaron Bowes is Senior Project officer at The Training Effect, UK

For communication please email: v.roberts@bath.ac.uk and r.joiner@bath.ac.uk

Authors: Richard Joiner, Veronica Roberts, Claire Russell, Terence Nice, Mark Bowles, Aaron Bowes and Emma Howe

Mind and Body: an early intervention group programme for adolescents with self-harm thoughts and behaviours

Introduction

Mental health difficulties amongst children and young people (C&YP) in the UK are on the rise (NHS, 2018). In particular, emotional disorders have increased from 3.9% in 2004 to 5.8% in 2017 (NHS, 2018). In the same survey, self-harm and suicide attempts were reported by 25.5% of 11-16 year olds, and 46.8% of 17-19 year olds with a mental health condition (NHS, 2018). Such stark prevalence rates for those with a mental health condition, alongside the association between self-harm and suicide, identify a significant public health concern (Brent et al, 2013; Campbell, 2016; 2018). In addition, the survey found that 3% of 11-16 year olds *without* a disorder, had also self-harmed and considered suicide (NHS, 2018). Previous studies exploring community samples suggest that life-time prevalence rates for self-harm amongst adolescents is much higher - between 13-45% (Nock, 2010, p344). This suggests there is perhaps a 'hidden demographic' of adolescents who engage in self-harm but are unknown. This group is of particular concern due to the likelihood that they are *not* accessing appropriate support services. Indeed, research suggests that between a third and one-half of adolescents who self-harm do *not* seek help for this issue (Rowe et al, 2014; Deliberto & Nock, 2008).

Self-harm is a complex behaviour existing along a 'continuum of self-destructiveness' (Nock, 2010). At one end of the spectrum the individual's intent may be to end their own life, whilst at the other it may be to maintain it (Rae, 2016). It involves intentional and direct self-poisoning (e.g. overdose) and/or self-injury (e.g.

cutting) with methods being diverse and often conducted privately (Hawton et al, 2012).

The effective identification and treatment of this group therefore poses a considerable challenge and suggests that early, preventative and community-based interventions are needed. However, research into psychosocial interventions for adolescents, primarily focuses on higher-tier clinical support provided post hospital admission. The treatments considered most effective are intensive, long term, tend to support the family as well as the adolescent (Glen et al, 2015; Ougrin et al, 2015) and therefore have significant cost implications.

Research into community-based support for self-harm is limited but preliminary findings have suggested that programmes based within schools and/or that involve peers and non-familial supportive adults such as teachers, could be beneficial (Brent et al, 2013). This approach is further supported by evidence that interpersonal factors play a crucial role in self-harm and that the behaviour is strongly associated with peer relationship problems and peer victimisation (Giletta et al, 2012; Brunner et al, 2014). Having a friend who self-harms is also recognised as risk factor for self-harm behaviour (King and Merchant, 2008). Therefore, addressing self-harm within a peer group context has the potential of tackling key contributing social causes or maintenance factors. Preliminary evidence has suggested such programmes could be highly beneficial for adolescents (Nock, 2010, p355; Wood et al, 2001, p1247). Conversely though, concerns have also been raised regarding a 'social contagion' effect (Jarvi et al, 2013) potentially undermining the efficacy of group interventions. Therefore, further research into community-based early intervention group programmes is required.

Evaluation focus

This study evaluated Mind and Body (MAB), which is an early intervention group programme for adolescents at risk of, or engaging in, self-harm behaviours. It was delivered within secondary schools and community-based clinics by Addaction. This evaluation therefore contributes to the preliminary research into community-based interventions for self-harm in adolescents. Specifically, the programme was evaluated for its:

1. Identification, referral and support of young people at risk of, or engaging in self-harm behaviours
2. Impact on young people's awareness and management of thoughts, feelings and behaviours related to self-harm
3. Impact on mental health and emotional well-being.

Method

Students

8440 students (13-17 year olds) completed a screening survey. 1573 students (18.6% of the screened population), were identified at risk of self-harm and had one-to-one sessions with an Addaction professional to determine the programme's suitability. In addition, school staff could highlight students of concern and adolescents were also able to self-refer to the programme. 622 students participated in the programme between March 2016 and April 2017 (13.6% of the total screened population). We collected data from 299 young people who undertook and completed the programme. See Table 1 for a geographical break-down.


Table1: Screened and participating sample figures

Area	Number screened	Number identified as at risk	Number engaged in programme
Cornwall	914	203	107
Kent	6039	1115	421
Lancashire	1487	255	94
Totals:	8440	1573	622

Procedure

The MAB programme was delivered in three pilot sites across England: Kent, Cornwall and Lancashire within secondary schools and community centres, by Addaction professionals. We obtained ethical approval from the Psychology Department's ethics committee at the University of Bath. The parents were informed that the programme was being delivered in their child's school and they were told to notify the school if they did not want their child to participate. The programme involved 8 weekly group sessions and 3 one-to-one sessions (See Figure 1). Group sizes were between 6-8 students.

Figure 1: Programme structure and core themes



Baseline assessments were conducted one-to-one with the students prior to their participation in the programme and post-intervention.

In addition, we conducted semi-structured focus groups with key stakeholders at the start of the programme delivery and six months later in each area, except for Lancashire, where following an initial focus-group, feedback questionnaires were completed by stakeholders 6 months later.

Measures

i. Screening survey

We used a bespoke online screening survey developed by The Training Effect, to identify adolescents at risk of, or engaging in, self-harm. The questions explored emotions, life-focus, peers and family, alcohol, smoking, self-harm and school support.

ii Pre and post-intervention measures

We evaluated the effectiveness of the programme before and after the programme using measures of mental wellbeing and a self-harm risk assessment.

The Short Warwick Edinburgh Mental Well-being Scale (Stewart-Brown, 2009) a 7-item scale to assess and monitor students' mental health and well-being

The Strengths and Difficulties Questionnaire (SDQ) a 25-item questionnaire exploring emotions, behavioural conduct, attention/distractibility, peer relationships and prosocial behaviour; providing a quantitative measure of behavioural change (Goodman, 1997).

Self-Harm Risk Assessment – a 7-question protocol exploring the frequency of self-harm behaviour, thoughts of self-harm, nature of self-harm behaviours and presence of suicide ideation.

All the above measures are of known validity and reliability.

Quantitative analysis

The quantitative analysis we conducted compared students' pre- and post-intervention scores on the Short Warwick Edinburgh Mental Well-being Scale, the Strengths and Difficulties Questionnaire and the Self-Harm Risk Assessment. Due to the ordinal level of the questionnaire data, non-parametric statistics were conducted.

Qualitative analysis

We took notes during the focus groups and the transcripts were circulated to attendees for accuracy and perspective checks. Thematic analysis (Braun and Clarke, 2006) was conducted to explore core themes regarding the MAB programme delivery, impact and outcomes.

Findings

Quantitative findings

We found a significant reduction in the number of self-harm thoughts after the programme, for students who disclosed self-harm thoughts before the programme; 67% reported a reduction in self-harm thoughts after the programme, 24% reported an increase and for 9% there was no change (Wilcoxon signed ranks test, $z = 5.00$, $p < 0.05$). We found a similar pattern for self-harm actions: 64% of students reported a reduction in the number of self-harm actions; 27% reported an increase, and there was no change for 10% of students (Wilcoxon signed ranks test, $z = 2.44$, $p < 0.05$).

We compared the students' scores on the Short Warwick Edinburgh Mental Well-being Scale and the Strengths and Difficulties Questionnaire before and after the intervention (see Table 2).

Table 2: Changes in psychological well-being following participation in the MAB programme

	Deteriorated	Stayed the Same	Improved
Short Warwick Edinburgh Mental Well-Being Scale			
	21.4%	5.8%	72.8%
Strengths and Difficulties Questionnaire			
Emotional	11.8%	40.3%	48.5%
Conduct	13.6%	56.0%	30.5%
Hyperactivity	25.9%	43.2%	30.9%
Peer Problems	24.2%	42.8%	32.9%
Prosocial	18.0%	67.9%	14.0%

There was a significant improvement for students on the Short Warwick Edinburgh Mental Well-being Scale (Mann-Whitney, $z = 10.37$, $p < 0.05$), with 73% of students improving over

the course of the programme. In addition, significant improvements occurred on the Strength and Difficulties Questionnaire, for emotional problems (Mann-Whitney, $z = 7.42$, $p < 0.05$), conduct problems (Mann-Whitney, $z = 12.24$, $p < 0.05$) and peer problems (Mann-Whitney, $z = 2.45$, $p < 0.05$). There was no significant improvement in terms of hyperactivity (Mann-Whitney, $z = 1.28$, $p > 0.05$) or prosocial behaviour (Mann-Whitney, $z = 1.19$, $p > 0.05$).

We further explored whether there were any differences between those students who improved and those who did not (see Table 3). Those who improved on the Short Warwick Edinburgh Mental Well-being Scale, scored significantly lower before the programme, than those who did not improve (Kruskal-Wallis, Chi-squared = 27.79, $p < 0.05$). Those who improved on the Strengths and Difficulties Questionnaire had significantly higher ratings on emotional problems (Kruskal-Wallis, Chi-squared = 32.41, $p < 0.05$), conduct problems (Kruskal-Wallis, Chi-squared = 118.96, $p < 0.05$), hyperactivity (Kruskal-Wallis, Chi-squared = 66.59, $p < 0.05$), peer problems (Kruskal-Wallis, Chi-squared = 57.19, $p < 0.05$) and lower scores on prosocial behaviour (Kruskal-Wallis, Chi-squared = 116.70, $p < 0.05$). Therefore, students who benefitted the most from the programme were those who had the greatest difficulties before the programme.

Table 3: Students' mean pre-test scores and standard deviations

	Deteriorated	Stayed the Same	Improved
	Mean (SD)	Mean (SD)	Mean (SD)
Short Warwick Edinburgh Mental Well-being Scale			
	21.05 (3.18)	21.3 (2.95)	18.77 (3.77)
Strengths and Difficulties Questionnaire			
Emotional	2.21 (0.74)	3.18 (1.27)	3.38 (0.78)
Conduct	1.39 (0.56)	1.43 (0.92)	3.07 (0.85)

Hyperactivity	1.71 (0.77)	2.15 (1.33)	3.24 (0.80)
Peer Problems	1.92 (0.86)	2.79 (1.30)	3.43 (0.74)
Prosocial	2.59 (0.70)	1.19 (0.67)	1.41 (0.66)

Qualitative findings

We organised emergent themes from stakeholder focus groups around the key objectives of the MAB programme. Themes and the number of stakeholders contributing to the theme are summarised in Table 4.

Table 4. – Qualitative Themes from Stakeholder Focus Groups

Theme	No. of stakeholders	Cornwall	Kent	Lancs.
Identification, referral and support of young people at risk of, or engaging in self-harm behaviours				
<i>Under the radar</i>	10	3	4	3
<i>Filling a gap in services</i>	14	4	8	2
Young people’s awareness and management of thoughts, feelings and behaviours related to self-harm				
<i>Not alone</i>	5	-	3	2
<i>Reduced stigma and stereotyping</i>	10	-	7	3
<i>Increased awareness in staff</i>	11	1	8	2
<i>Increased support seeking</i>	7	2	4	1
<i>Independent coping strategies</i>	8	3	4	1

<i>Self-sustaining peer support</i>	10	2	7	1
Impact on students' mental health, emotional well-being and behaviour				
<i>Communication skills</i>	7	1	5	1
<i>Confidence and empowerment</i>	6	-	5	1
<i>Increased participation in school life</i>	6	2	4	-
Additional key themes				
<i>Skill of MAB practitioners</i>	15	4	8	3
<i>Parent and family support</i>	4	3	1	-

Identification, referral and support of young people at risk of, or engaging in self-harm behaviours

Under the radar

The screening survey was a powerful tool to identify 'at risk' students. In one stakeholder's view, it had saved the life of a student who had been completely unknown to them as at risk.

"It pulled out students who I would never have come across. They would never have got support...but doing that survey, it identified those students. It was a real eye opener."

Stakeholder, Lancashire

The high number of students identified was a surprise and led to concerns regarding how to support students who could not participate in MAB and did not meet the threshold for Child and Adolescent Mental Health Services (CAMHS).

Filling a gap in services

Stakeholders identified that the MAB programme bridged an important gap in services for students with self-harm behaviours who required support beyond school staff expertise, but did not meet the criteria for CAMHS involvement.

Some stakeholders also identified how the programme filled a gap in the *form* of service offered to young people. The ease of access due to the programme being school-based, flexibility/responsiveness of approach, and availability during the school holidays were cited as reasons the programme had been so effective for many of the students.

“The other advantage is that access to the MAB practitioners is on the young person’s terms.”

Stakeholder, Cornwall.

Young people’s awareness and management of thoughts, feelings and behaviours related to self-harm

Not alone

Stakeholders highlighted how the group sessions had increased students’ awareness that they were ‘not alone’ with their experiences, reducing their sense of isolation.

“Every single one of the students has acknowledged that there is support from the shared experience of their situations”

Stakeholder, Lancashire

Reduced stigma and stereotyping

Several stakeholders indicated that the programme led to a reduction in students' stereotypes and stigma around mental health. In some instances, this had led to increased mental health awareness within the school.

Increased awareness in school staff

Stakeholders' felt introductory assemblies had raised awareness for staff and students who attended, but that the impact could have been enhanced by presenting it to the whole-school and fostering a more supportive whole-school ethos. It was felt this would have helped overcome some teachers viewing students missing lessons for MAB as "a nuisance".

Increased support seeking

Several stakeholders observed increased communication between MAB students and school staff, including coded communications and seeking additional support from school personnel when needed.

"Some YP...have initiated a system where they wear wrist bands to identify risk feelings or vulnerabilities"

Stakeholder, Kent

Independent coping strategies

Stakeholders indicated that the students were now more able to independently and effectively use appropriate coping strategies. One stakeholder felt that the extended

time frame of the programme made it more realistic that young people would be able to develop and use these techniques.

Self-sustaining peer support

Stakeholders indicated that students had formed self-sustaining peer networks that provided support after the programme concluded. This was viewed as a particularly powerful and unanticipated outcome generated by the group format of the programme.

“I certainly think that the group aspect helps...young people are learning from one another without being judged.”

Stakeholder, Kent

“Because it’s a small group, the ethos is that the students will stick together. They may not be best friends but there is a support network based on shared experience”

Stakeholder, Lancashire

Impact on students’ mental health, emotional well-being and behaviour

Communication skills

Stakeholders reported that participating in MAB had supported students’ communication skills towards peers, school staff and/or their parents. One stakeholder felt this helped reduce their sense of isolation whilst others reported it had helped some students to talk to their parents.

“We do an exercise where we support young people to identify their communication styles. This really opens their eyes so that they can take different perspectives... Young people were able to change their communication styles.”

Stakeholder, Cornwall

Confidence and empowerment

Some stakeholders reported that many participating students had noticeably improved confidence. Four stakeholders felt that the group sessions provided a safe space to be ‘heard without judgement’ and that this had led to a lasting sense of empowerment.

Increased participation in school life

Some stakeholders had observed improvements in the students’ engagement and participation in school life including reduced absences, behavioural incidents and lateness, and improved attitude to learning. Two stakeholders reported some students had volunteered for events that previously they would not have put themselves forward for. One stakeholder reported that some students had improved so significantly that they were no longer on the school’s intervention list.

Additional key themes

Skill of MAB Practitioners

MAB practitioners’ skills were viewed as a key factor supporting the effectiveness of the programme. This included their ability to communicate clearly with school staff, their relationship with the students and flexibility in responding to the students’ needs within programme delivery. Their ability to do this was linked by stakeholders to the

level of prior experience and expertise practitioners had, as well as extensive training in the MAB programme and being external to the school organisation.

“Having an independent person seemed to promote honest and authentic disclosure.”

Stakeholder, Cornwall

“Because the practitioner was empathic, the students felt they could really trust her”

Stakeholder, Kent.

Areas for further development

A few stakeholders indicated that there was a potential gap in the provision of support to parents/families of students attending MAB.

“Parents are crying out for support when they don’t know what to do to help their children.”

Stakeholder, Cornwall

Discussion

The Mind and Body programme was developed in response to the need for preventative and early interventions for young people who engage in or, are at-risk of engaging in, self-harm behaviours. MAB aimed to support students in addressing thoughts and actions associated with self-harm and to increase mental well-being. The screening tool identified 1573 students at risk of or, engaging in, self-harm – 18.6% of the screened population. This included many “under the radar” students, who were not known by the school or local mental health services to be experiencing any difficulties. Follow-on interviews with all identified students indicated that this

level of identification was valid and therefore the screening tool was effective at identifying young people at risk of or, engaged in self-harm behaviours. The potential of this instrument for nationwide identification of young people at risk of self-harm is therefore worth further investigation.

Quantitative analysis indicated that the MAB programme was effective in improving mental well-being for the majority (72.8%) of students. In addition, significant improvements were found for students in emotional (48.5%), conduct (30.5%), and peer problems (32.9%). Furthermore, of students completing the self-harm risk assessment, 67% reported a reduction in self-harm thoughts and 63% reported a reduction in self-harm actions. These results are therefore supportive of the efficacy of the MAB programme for providing an effective school-based group intervention for students at risk of or, engaging in, self-harm behaviour.

Qualitative observations from stakeholders also supported the view that the MAB programme had a positive impact on young people's awareness, thoughts, feelings and behaviours relating to self-harm:

- (i) There was a reported increase in young people's appropriately seeking and accessing support from school staff.
- (ii) There was a reported increase in young people's coping skills/strategies
- (iii) There was a reduction in stigma about self-harm, aided by the group sharing experiences, understanding, empathy and compassion.
- (iv) There was an increase in young people's well-being and a broadening of their social networks.

- (v) There was also an observed improvement for some young people in their attendance and contribution in classes.

Given these findings regarding the impact of the MAB Programme, there is strong emerging evidence for the effectiveness of the programme.

However, whilst these findings are very promising, they need to be considered within the limitations of the evaluation. There was no control group and so it is not possible to show categorically that these improvements would not have happened without the programme. In addition, data was not collated for all of the students who completed the programme and so may not be fully representative. Furthermore, follow-up data is not available and so it is unclear whether these improvements were maintained over time. In addition, it is notable that a sizable minority of young people deteriorated during the course of the programme, and more exploration of the factors that may have contributed to their deterioration is needed.

In conclusion, both the screening tool and MAB programme demonstrate considerable promise in providing an effective way of identifying and supporting young people at risk of or, engaging in, self-harm behaviour, as a form of early and community-based intervention. However, further research is required to build upon the current evaluation's findings.

References

Braun, V. & Clarke, V. (2006). Using thematic analysis in psychology. *Qualitative Research in Psychology*, 3, pp77-101.

Brent, D.A., McMakin, D.L., Kennard, B.D., Goldstein, T.R., Mayes, T.L., Douaihy, A.B. (2013). Protecting adolescents from self-harm: a critical review of intervention studies. *Journal of the American Academy of Child and Adolescent Psychiatry*. 52(12), pp. 1260-1271.

Brunner, R., Kaess, M. Parzer, P., Fischer, G., Carli, V., Hoven, C.W., Wasserman, C., Sarchiapone, M., Resch, F., Apter, A., Balazs, J., Barzilay, S., Bobes, J., Corcoran, P., Cosmanm, D., Harins, C., Iosuec, M. Kahn, J., Keeley, H., Meszaros, G., Nemes, B., Podlogar, T., postuvan, V., Saiz, P.A., Sisask, M., Tubiana, A., Varnik, A. & Wasserman, D. (2014). Life-time prevalence and psychosocial correlates of adolescent direct self-injurious behaviour: a comparative study of findings in 11 European countries. *Journal of Child Psychology and Psychiatry*, 55(4), pp. 337-348.

Campbell, D. (2016). NHS figures show 'shocking' rise in self-harm among young. *The Guardian*, 23rd October. Accessed online on: 18/04/19 at:
<https://www.theguardian.com/society/2016/oct/23/nhs-figures-show-shocking-rise-self-harm-young-people>

Campbell, D. (2018). Fifth of 17 to 19-year-olds self-harm or attempt suicide. *The Guardian*, 22nd November. Accessed online on: 15/05/19 at:
<https://www.theguardian.com/society/2018/nov/22/concern-over-rise-in-suicide-attempts-among-young-women>

Deliberto, T.L. & Nock, M.K. (2008). An exploratory study of correlates, onset, and offset of nonsuicidal self-injury. *Archives of Suicide Research*.12, pp.219-31.

Giletta, M., Scholte, R.H., Engels, R.C., Ciairano, S. & Prinstein, M.J. (2012). Adolescent non-suicidal self-injury: a cross-national study of community samples from Italy, the Netherlands and the United States. *Psychiatry Research*, 197, pp.66-72.

Glen, C.R., Franklin, J.C. & Nock, M.K. (2015). Evidence-based psychosocial treatments for self-injurious thoughts and behaviors in youth. *Journal of Clinical Child and Adolescent Psychology*. 44(1), pp1537-4424.

Goodman R. (1997). The strengths and difficulties questionnaire: a research note. *Journal of Child Psychology and Psychiatry*. 38(5):581–586

Hawton, K., Saunders, K.E.A., O'Connor, R. (2012). Self-harm and suicide in adolescents. *The Lancet*. 379, pp.2373-2382.

Jacobson, C.M. & Gould, M. (2007). The epidemiology and phenomenology of non-suicidal self-injurious behaviour among adolescents: a critical review of the literature. *Archives of Suicide Research*. 11(2), pp1543-6136.

Jarvi, S., Jackson, B., Swenson, L. & Crawford, H. (2013). The impact of social contagion on non-suicidal self-injury: a review of the literature. *Archives of Suicide Research*, 17(1), pp. 1-19.

King, C.A., Merchant, C.R. (2008). Social and interpersonal factors relating to adolescent suicidality: a review of the literature. *Archives of Suicide Research*, 12, pp181-196.

NHS. (2018). Mental health of children and young people in England 2017. Published NHS Digital, 22nd November, 2018. Accessed online on: 15/05/19 at: <https://digital.nhs.uk/data-and-information/publications/statistical/mental-health-of-children-and-young-people-in-england/2017/2017>

Nock, M.K. (2009) Why do people hurt themselves? New insights into the nature and functions of self-injury. *Current Directions in Psychological Science*, 18, pp78-83.

Nock, M.K. (2010). Self-injury. *Annual Review of Clinical Psychology*. 6, pp. 339-363.

Ougrin, D., Tranah, T., Stahl, D., Moran, P. & Asarnow, J.R. (2015). Therapeutic interventions for suicide attempts and self-harm in adolescents: systematic review and meta-

analysis. *Journal of the American Academy of Child and Adolescent Psychiatry*, 54(2), pp97-107.

Rae, T. (2016). Introducing the topic of self-harm in schools: developing an educational and preventative support intervention. *Education and Health*, 34(2).

Rowe, S.L., French, R.S., Henderson, C., Ougrin, D., Slade, M. & Moran, P. (2014). Help-seeking behaviour and adolescent self-harm: a systematic review. *Australian and New Zealand Journal of Psychiatry*, 48(12), pp. 1083-95.

Stewart-Brown, S., Tennant, A., Tennant, R., Platt, S., Parkinson, J., & Weich, S. (2009). Internal construct validity of the Warwick-Edinburgh mental well-being scale (WEMWBS): a Rasch analysis using data from the Scottish health education population survey. *Health and quality of life outcomes*, 7(1), 15.

Wood, A., Trainor, G., Rothwell, J., Moore, A., Harrington, R. (2001). Randomized trial of group therapy for repeated deliberate self-harm in adolescents. *Journal of the American Academy of Child and Adolescent Psychiatry*. 40, pp. 1246-53.