

1 *Campylobacter pinnipediorum* subsp. *caledonicus* and *Campylobacter pinnipediorum*  
2 subsp. *pinnipediorum* recovered from abscesses in pinnipeds

3

4 Geoffrey Foster<sup>1\*</sup>, Johanna L. Baily<sup>2,10</sup>, Fiona Howie<sup>3</sup>, Andrew C. Brownlow<sup>1</sup>, Jaap A.  
5 Wagenaar<sup>4,5,6</sup>, Maarten J. Gilbert<sup>4,7</sup>, William G. Miller<sup>8</sup>, Barbara A. Byrne<sup>9</sup>, Kristin A.  
6 Clothier<sup>9</sup>, Todd Schmitt<sup>11</sup>, Tony Patterson<sup>1</sup>, Robert J. Reid<sup>1</sup> and Mark P. Dagleish<sup>2</sup>

7

8 <sup>1</sup> SRUC Veterinary Services, An Lochran, 10 Inverness Campus, Inverness, IV2  
9 5NA, Scotland, United Kingdom

10 <sup>2</sup> Moredun Research Institute, Pentlands Science Park, Bush Loan, Penicuik,  
11 Midlothian EH26 0PZ, Scotland, United Kingdom

12 <sup>3</sup> SRUC Veterinary Services, Pentlands Science Park, Bush Loan, Penicuik,  
13 Midlothian EH26 0PZ, Scotland, United Kingdom

14 <sup>4</sup> Department of Infectious Diseases and Immunology, Faculty of Veterinary  
15 Medicine, Utrecht University, Yalelaan 1, 3584CL, Utrecht, the Netherlands

16 <sup>5</sup> WHO Collaborating Center for *Campylobacter*/OIE Reference Laboratory for  
17 *Campylobacteriosis*, Yalelaan 1, 3584CL, Utrecht, The Netherlands

18 <sup>6</sup> Department of Bacteriology and Epidemiology, Wageningen Bioveterinary  
19 Research, Houtribweg 39, 8221 RA, Lelystad, the Netherlands

20 <sup>7</sup> Reptile, Amphibian and Fish Conservation Netherlands (RAVON), Toernooiveld  
21 1, 6525 ED, Nijmegen, the Netherlands

22 <sup>8</sup> Produce Safety and Microbiology Research Unit, Agricultural Research Service,  
23 United States Department of Agriculture, Albany, CA 94710, United States

24 <sup>9</sup> Pathology, Microbiology and Immunology, University of California, Davis, CA  
25 95616, United States

1 <sup>10</sup> Present address: Institute of Aquaculture, University of Stirling, Stirling, FK9 4LA,  
2 UK

3 <sup>11</sup>SeaWorld San Diego, Veterinary Services, San Diego, CA 92109, USA

4

5 Running title: *Campylobacter pinnipediorum* infections of seals

6

7 \* Corresponding author:

8 Mailing address:

9 SRUC Veterinary Services, An Lochran, 10 Inverness Campus, Inverness, IV2 5NB,  
10 Scotland, United Kingdom

11

12 \*Corresponding author: geoffrey.foster@sac.co.uk Phone: +441463243030. Fax:

13 +441463711103

1 ABSTRACT: *Campylobacter pinnipediorum* was described recently for isolates  
2 recovered from pinnipeds. The novel species was further split into two subspecies  
3 based on host and geography with *Campylobacter pinnipediorum* subsp.  
4 *pinnipediorum* recovered from otariid seals in California and *Campylobacter*  
5 *pinnipediorum* subsp. *caledonicus* recovered from phocid seals in Scotland. We report  
6 details of the infections of seven pinnipeds from which *C. pinnipediorum* was isolated:  
7 *C. pinnipediorum* subsp. *caledonicus* was isolated from two harbour seals (*Phoca*  
8 *vitulina*) and a single grey seal (*Halichoerus grypus*) and *C. pinnipediorum* subsp.  
9 *pinnipediorum* isolates from California sea lions (*Zalophus californianus*). Six of the  
10 isolates were recovered from samples collected at post-mortem investigation. In two  
11 of the Scottish seals and in three of the California seals, *C. pinnipediorum* was the  
12 sole bacterial isolate recovered from abscesses present and suggests they may have  
13 resulted from con- or intraspecific bite wounds.

## 1. INTRODUCTION

*Campylobacter* is a bacterial genus that has been associated with asymptomatic carriage and symptomatic infections in humans and other animals (Skirrow, 1994). In humans, campylobacteriosis is recognised as the most common cause of bacterial food poisoning in the UK, due mostly to *Campylobacter jejuni* and *Campylobacter coli* (Strachan and Forbes 2010, Tam et al. 2012). The aforementioned species have also been recognised in farm and companion animals as well as wildlife and are established zoonotic pathogens (Sheppard et al. 2009). In addition to the above, further *Campylobacter* spp. of veterinary importance include *Campylobacter fetus* subsp. *fetus*, which is associated with abortion in sheep and cattle, and reproductive disease in bovids due to *Campylobacter fetus* subsp. *venerealis* (Skirrow 1994).

Increasing microbiological investigations of wildlife have reported the isolation of *Campylobacter* spp., including strains infecting marine mammals. In seals inhabiting Scottish coastal waters, these have included *C. jejuni*, *C. coli*, *Campylobacter lari* (Baily et al. 2015) and *Campylobacter insulaenigrae* (Foster et al. 2004). First reported from 3 seals and a harbour porpoise, the latter has since been reported from seals in other regions of the world (Stoddard et al. 2007; Garcia-Pena et al. 2010, González et al. 2011) but also as a cause of human infections (Chua et al. 2007). Of late, isolates from harbour seals in the Netherlands were assigned to the novel species, *Campylobacter blaseri* (Gilbert et al. 2018).

Recently, a polyphasic analysis of *Campylobacter* isolates from pinnipeds resulted in the description of the novel, urease positive species, *Campylobacter pinnipediorum* (Gilbert et al. 2017). This novel species could be sub-divided further into two

1 subspecies based on phylogenetic analysis and catalase reaction. Notably, the two  
2 subspecies could also be differentiated by the animal hosts infected and their  
3 geographic location: *Campylobacter pinnipediorum* subsp. *pinnipediorum* was  
4 recovered from otariid seals in California and *Campylobacter pinnipediorum* subsp.  
5 *caledonicus* was recovered from phocid seals in Scotland. We report here the details  
6 of infections with *C. pinnipediorum*: *C. pinnipediorum* subsp. *caledonicus* in two  
7 harbour seals (*Phoca vitulina*) and a grey seal (*Halichoerus grypus*) in Scotland and  
8 *C. pinnipediorum* subsp. *pinnipediorum* in four California sea lions (*Zalophus*  
9 *californianus*) in California. Six of the isolates were isolated following post-mortem  
10 examination.

11

12

## 2. MATERIALS AND METHODS

13

### 2.1 Scottish seal cases

14 Pinnipeds that stranded in Scotland were reported under the Scottish Marine Animals  
15 Strandings Scheme (SMASS) funded by Marine Scotland as part of the UK  
16 Government's commitment to a number of international agreements and specifically  
17 under European Protected Species licence number 120139. Carcasses were  
18 transported to Scotland's Rural College (SRUC) Veterinary Services, Inverness for a  
19 post mortem examination performed according to a standard protocol (Dierauf, 1994).  
20 Selected tissues and gross lesions were sampled for microbiological and  
21 histopathological diagnoses. Cultures were made on Columbia sheep blood agar  
22 (CSBA) (Oxoid, Basingstoke, UK) and selectively for *Brucella* on Farrell's medium  
23 (FM) (Farrell 1974), incubated at 37 °C in air with 5 % added CO<sub>2</sub>. Selective cultures  
24 for *Campylobacter* and anaerobes incubated in microaerophilic and anaerobic

1 environments were also set up. All plates were examined for growth at frequent  
2 intervals up to 14 days.

3

4

## 2.2 California sea lion cases

5 Four California sea lions that live-stranded were rescued and cared for at animal  
6 rehabilitation centres in California. Three of the animals were cared for at SeaWorld in  
7 California under the NMFS Marine Mammal Health and Stranding Response Program  
8 Permit #932-1905 and the other at the Pacific Marine Mammal Center in California  
9 under the NMFS Marine Mammal Health and Stranding Response Program Permit  
10 #932-1489. The SeaWorld animals subsequently died and received post mortem  
11 investigations, which included sampling of abscesses. The sea lion at the Pacific  
12 Marine Mammal Center in California survived and abscess exudate was collected for  
13 cytology and bacterial culture on 5% sheep blood agar (Hardy Diagnostics, Santa  
14 Maria, CA) and incubated at 35°C in room air supplemented with 5% CO<sub>2</sub>.

15

16

## 3. RESULTS

17

### 3.1 Scottish cases

18 Case S1. A juvenile male harbour seal (M203/00) was found live-stranded at  
19 Lossiemouth in the Moray Firth (57°66'13.9"N 003°61'63.89"W) on 28/10/2000.  
20 There was a large poorly circumscribed sub-blubber abscess above the left  
21 shoulder, which contained dark red/brown pus. The left pre-scapular lymph node  
22 was markedly enlarged. The seal was also suffering from severe pneumonia and  
23 was euthanased on welfare grounds. Large numbers of *Campylobacter*-like bacteria  
24 were the only organisms observed in a Gram-stained smear from the abscess  
25 contents, but they were not seen in tissue smears from lung, liver, spleen, kidney or

1 brain. A pure growth of *Campylobacter pinnipediorum* subsp. *caledonicus* (+) was  
2 recovered from the abscess on CSBA after 10 days microaerophilic incubation at  
3 37°C and after 14 days in a capnophilic atmosphere at the same temperature.  
4 Microaerophilic cultures from brain, spleen and lung were negative and cultures  
5 using *Campylobacter*-selective media (charcoal cefoperazone deoxycholate agar  
6 [CCDA] and Blaser-Wang) were negative for these three tissues and the abscess.  
7 Additionally, *E. coli* was recovered from lung (+++), spleen (++) , mesenteric lymph  
8 node (++) , kidney (+) , liver (few) and in mixed growth from blood and intestine.  
9 Selective culture for *Brucella* spp. was negative for all tissues. Histological  
10 examination of the lung confirmed the presence of parasitic and suppurative  
11 broncho-interstitial pneumonia. Larvae were present in the airways with an  
12 associated eosinophilic and lymphocytic inflammatory response and suppurative  
13 infiltrates were associated with small numbers of adult worms in alveoli. The section  
14 of liver examined was diffusely congested and contained a focus of necrosis and  
15 suppurative inflammation. The thymus was depleted. The spleen was congested with  
16 evidence of extramedullary haematopoiesis and the kidney was congested.  
17 Histological examination of the brain showed a severe suppurative meningo-  
18 encephalitis, as denoted by the presence of large numbers of neutrophils, and protein  
19 rich oedema around blood vessels in the meninges and in the neuropil along with  
20 necrosis in the grey matter of the cerebral cortex and white matter of the internal  
21 capsule. These lesions are indicative of a bacterial aetiology, possibly with the  
22 production of bacterial toxins.

23

24 Case S2. A 3-4 week old female grey seal pup (M302/10) was found dead at  
25 Tynninghame Beach, East Lothian (56°02'49.53"N 002°59'84.13"W) on 09/12/2010. A

1 profuse growth of *C. pinnipediorum* subsp. *caledonicus* appeared initially as tiny  
2 colonies, which were possibly satellitic around other organisms as a mixed growth on  
3 chocolate agar incubated at 37°C in air plus 5% CO<sub>2</sub> from a lung abscess. Other  
4 bacteria isolated were *Streptococcus agalactiae*, *Arcanobacterium phocae* and a  
5 moderate growth of *Salmonella enterica* serovar Bovismorbificans. In addition, a  
6 profuse growth of an unidentified *Fusobacterium* sp. was recovered from anaerobic  
7 cultures. Interestingly, culture of non-abscessed lung tissue produced all the  
8 aforementioned organisms except for the *Campylobacter*. The only other tissues  
9 cultured were liver and kidney, which had profuse and moderate growths, respectively,  
10 of *S. Bovismorbificans*, a serovar encountered in grey seals in Scottish waters (Baily  
11 et al. 2016). Selective culture for *Brucella* was negative for all tissues.

12

13 Case S3. A juvenile female harbour seal (M341/11) was found dead at Findhorn in the  
14 Moray Firth, (57°66'13.9"N 003°61'63.89"W) on 27/11/2011. There was a large  
15 abscess within the axillary lymph node under the right scapula, which was confirmed  
16 histologically, but no overt evidence of a bite wound. A profuse pure growth of *C.*  
17 *pinnipediorum* subsp. *caledonicus* was recovered from the abscess in anaerobic  
18 culture. The only other bacterial isolate was *Brucella pinnipedialis* recovered as single  
19 colonies from lung, spleen and mesenteric lymph node and as a moderate growth  
20 alongside a sparse mixed growth of other organisms from the small intestine. No  
21 growth was obtained from the liver, kidney or brain. *Campylobacter* cultures of small  
22 intestine on Skirrow's agar were negative. Additional histopathology findings included  
23 mild drainage reactions in a distant lymph node and the spleen; mildly reactive hepatic  
24 sinusoids and mild broncho-interstitial pneumonia associated with adult parasites with  
25 gross morphology identical to *Parafilaroides* sp.



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24

### 3.2 California sea lion cases

Case C1. An emaciated female California sea lion pup (SW 130202), weighing 12 kg, live-stranded for the second time on 09/03/2013 and died at SeaWorld on 11/03/2013. A 9 X 5 X 3 cm abscess was found on the left pelvic limb surrounded by necrotic muscle and chylous peritoneal effusion containing 30 mL of opaque pink/red fluid. *Campylobacter pinnipediorum* subsp. *pinnipediorum* was isolated from the effusion along with mixed aerobic and anaerobic bacteria. The primary cause of death was likely sepsis due to the severe focally extensive necrotising myositis, peritonitis and emaciation.

Case C2. An emaciated juvenile female California sea lion (SW 130133), weighing 11.4 kg, died at SeaWorld on 15/02/2013. Open wounds were found on the lower jaw at the corner of the oral cavity and upper lip just behind the vibrissae. There was also a 6.0 X 3.5 cm abscess in the neck above the right shoulder joint which was full of opaque pink purulent fluid. Histologic examination of the abscess revealed numerous intact and degenerate neutrophils surrounded by abundant fibroplasia extending into the surrounding muscle and adipose tissues with occasional lymphoid aggregates. *Campylobacter pinnipediorum* subsp. *pinnipediorum* was isolated from the neck abscess, which was thought to possibly be the result of a bite wound from a conspecific sea lion. The sea lion received injectable antibiotics (ceftiofur), subcutaneous fluids, oral electrolytes and pinniped gruel formula for nutrition. The suspect bite wound abscess was deemed to be the cause of the poor body condition and death.

1 Case C3. An emaciated male California sea lion pup (SW 130167), weighing 13 kg,  
2 was admitted to SeaWorld on 03/03/2013 and died on 04/03/2013. The hypothermic  
3 sea lion was housed inside with thermal heating pad and treated supportively with  
4 subcutaneous fluids + 5% dextrose, dilute formula and oral electrolytes. Preliminary  
5 CBC (cell blood count) showed a low white blood cell count (5,230 cells/ $\mu$ L, 31%  
6 neutrophils, 18% band neutrophils, 27% lymphocytes, 16% monocytes), hematocrit  
7 (46%), and point-of-care I-stat showed hypoglycemia (27 mg/dL) and acidosis (pH  
8 7.1). At post-mortem, there was approx. 30 mL of opaque off-white pleural effusion  
9 with opaque orange fluid with proteinaceous particles. Both lungs were mottled with  
10 dark red regions at the lung periphery. The caudal left lung lobe was enlarged with the  
11 edge firm and dark red with clusters of off-white abscesses, from which *C.*  
12 *pinnipediorum* subsp. *pinnipediorum* was isolated in small numbers. Histologic  
13 examination demonstrated submassive pneumonia with necrosis of terminal  
14 bronchioles and alveoli and tissue replacement by degenerate neutrophils,  
15 proteinaceous oedema, and fibrin. No microorganisms were identified in any sections  
16 of the lung. Cause of death was sub-massive necrosis and pneumonia, pleural  
17 effusion and septic necrosis in the liver.

18

19 Case C4. A California sea lion (Z-11-06-01-057) began treatment at the Pacific Marine  
20 Mammal Center on 01/06/2011 and was released on 14/08/2011. Two abscesses  
21 were present: one on the right dorsolateral abdomen and one on the right thoracic  
22 area. The animal was also moderately underweight with several raised cutaneous  
23 nodules on the thoraco-abdominal dorsum. Two abscesses were opened, suppurative  
24 exudate removed, flushed and instilled with Techni-Care<sup>®</sup> antiseptic. Cytologic  
25 examination of the abscess exudate revealed masses of degenerate and/or necrotic

1 leukocytes, approximately 50:50% neutrophils:mononuclear cells with occasional  
2 scattered erythrocytes, no bacteria were definitively identified. The right dorsolateral  
3 abdominal abscess was aspirated, from which *C. pinnipediorum* subsp. *pinnipediorum*  
4 was the sole isolate. Treatment included flushing with salt and/or tap water,  
5 amoxicillin-clavulanic acid 17 mg/kg orally twice daily for 6 days. Wound care was  
6 terminated 12 days after admission and abscesses were noted to be healed 3 weeks  
7 after admission. The animal was released one month later weighing 44 kg.

8

### 9 **3.3 Characteristics of *Campylobacter pinnipediorum* isolates**

10 On CSBA, the isolates of both subspecies of *Campylobacter pinnipediorum* presented  
11 as  $\alpha$ -haemolytic beige, glossy, slightly raised and circular colonies with smooth  
12 margins. Growth was observed in microaerophilic and anaerobic environments and at  
13 25 °C but not in aerobic environments without added CO<sub>2</sub>, nor at 42 °C. Growth in air  
14 with 5% CO<sub>2</sub> varied with strain. All isolates were sensitive to cephalothin (30  $\mu$ g) and  
15 nalidixic acid (30  $\mu$ g). Positive reactions were obtained for urea hydrolysis, H<sub>2</sub>S  
16 production from triple sugar iron agar (TSI), nitrate production and oxidase but  
17 negative for hydrolysis of hippurate and indoxyl acetate. The aforementioned  
18 characteristics are shared between *C. pinnipediorum* subsp. *caledonicus* and *C.*  
19 *pinnipediorum* subsp. *pinnipediorum* and permitted their distinction from all other  
20 validated species of *Campylobacter* as well as urease-positive *Campylobacter lari*  
21 (UPTC) which is not, as yet, a defined taxon (Gilbert et. al. 2017; Miller et al. 2014).  
22 Differences in catalase reactions were observed between the Scottish and Californian  
23 strains, catalase-negative for *C. pinnipediorum* subsp. *caledonicus* and catalase-  
24 positive for *C. pinnipediorum* subsp. *pinnipediorum*. Extensive molecular testing,  
25 including whole genome sequencing, permitted further distinction between the two

1 subspecies of *C. pinnipediorum*. A full description of the characterisation methods and  
2 results is provided with the species description of *C. pinnipediorum* (Gilbert et al.  
3 2017).

4

5

#### 4. DISCUSSION

6 During the years 2000 to 2011, *Campylobacter pinnipediorum* subsp. *caledonicus* was  
7 isolated from the abscesses of two stranded juvenile harbour seals and a grey seal  
8 pup that stranded at different locations around the North Sea coast of Scotland. In  
9 each of these cases, a single abscess was the only site from which *C. pinnipediorum*  
10 subsp. *caledonicus* was recovered: one in the shoulder region, one in a limb and the  
11 other in the lung. *Campylobacter pinnipediorum* subsp. *pinnipediorum* was recovered  
12 from abscesses in four California sea lions during 2011 and 2013. In three of these  
13 abscesses *C. pinnipediorum* subsp. *pinnipediorum* was the only organism isolated. It  
14 is possible that other cases may have gone undetected, especially if culture under  
15 anaerobic or microaerophilic conditions was not performed.

16

17 It is interesting to compare these cases with findings from necropsies of eight harbour  
18 porpoises (*Phocoena phocoena*), which stranded in Scotland, England, Belgium and  
19 the Netherlands, following non-lethal attacks from grey seals (Foster et al. 2019). In  
20 all these cases, *Neisseria animaloris* was recovered from abscesses in skin and lung,  
21 similar to the findings for the seals with *C. pinnipediorum* reported here. Grey seals  
22 are known to attack harbour seals (van Neer et al. 2015), harbour porpoises  
23 (*Phocoena phocoena*) (Leopold et al., 2015) and their own species (Brownlow et al.  
24 2016). It is possible that the infections in seals follow a similar course to that of *N.*  
25 *animaloris* in porpoises, whereby the seals survive an initial assault only to die of

1 complications later, such as abscesses due to *C. pinnipediorum*, either at the site of  
2 puncture or following dissemination and abscessation in the lung. Indeed, a recent  
3 microbiome study found *C. pinnipediorum* is part of the grey and harbour seal oral  
4 microbiome, being particularly abundant in some grey seals, and was also detected in  
5 very low amounts in porpoise lesions as a result of bites from grey seals (Gilbert et al.  
6 2020). Seal lesions were not part of that study; however, the possibility remains that  
7 abscesses in seals become infected with *C. pinnipediorum* following traumatic injury  
8 from conspecific or other pinnipeds. Furthermore, all strains of *C. pinnipediorum*  
9 displayed urease activity, which may be related to a gastric niche and also shared  
10 100% 16S rRNA sequence homology with uncultured bacteria from the gastric  
11 microbiota of California sea lions (Bik et al. 2016). *Campylobacter pinnipediorum*  
12 subsp. *pinnipediorum* was also recovered from the rectal swabs of four sea lion pups  
13 and a yearling.

14

15 We suggest that the most likely pathogenesis for the infections presented in this study  
16 are abscessation, secondary to bite wounds, likely from pinnipeds. Although bite  
17 wounds were not observed, their absence was recorded only for the most recent case  
18 and it may be that they went unnoticed in the earlier cases or had resolved. The non-  
19 reporting of bite lesions may also reflect slow growth of *C. pinnipediorum* or the effects  
20 of other stressors on the pathogenicity of this bacterium. It is possible that smaller  
21 puncture wounds may partially heal or be difficult to identify but that *C. pinnipediorum*  
22 is inoculated into tissues, is able to thrive and replicate in the low-oxygen environment  
23 present in damaged or devitalized tissues, and produces the severe cellulitis and  
24 debilitation observed in these animals. We suggest, therefore, that abscesses and bite  
25 wounds in pinnipeds should be subjected to appropriate microbiological investigation

1 to determine if either *Campylobacter pinnipediorum* sub-species is present to help  
2 determine the origin of the wounds/abscesses which may help to determine treatments  
3 for live animals and possible ultimate causes of death in dead animals.

#### 6 **Acknowledgements**

7 The Scottish Marine Animal Stranding Scheme receives financial support from the  
8 Scottish Government Marine Directorate and the UK Department of Environment,  
9 Farming and Rural Affairs (Defra). MPD was funded by the Scottish Government.  
10 Three California sea lions at SeaWorld were contributed under SeaWorld Parks and  
11 Entertainment technical contribution number 2020-15.

#### 13 **References**

14 Baily JL, Foster G, Brown D, Davison NJ, Coia JE, Watson E, Pizzi R, Goldstone RJ,  
15 Willoughby K, Hall AJ, Dagleish MP (2016) *Salmonella* infection in grey seals  
16 (*Halichoerus grypus*), a marine sentinel species: pathogenicity and molecular typing  
17 of *Salmonella* strains compared with human and livestock isolates. Environ Microbiol  
18 18:1048-1087

19 Baily JL, Meric G, Bayliss S, Foster G, Moss, SE, Watson E, Pascoe B, Mikhail J, Pizzi  
20 R, Goldstone RJ, Smith DGE, Willoughby K, Hall AJ, Sheppard SK, Dagleish MP  
21 (2015) Evidence of land-sea transfer of the zoonotic pathogen *Campylobacter* to a  
22 wildlife marine sentinel species. Mol Ecol 24:208-221

23 Bik EM, Costello EK, Switzer AD, Callahan BJ, Holmes SP, Wells SR, Carlin KP,  
24 Jensen ED Venn-Watson S, Relman D (2016) Marine mammals harbor unique

1 microbiotas shaped by and yet distinct from the sea. Nat Commun  
2 7:doi:10.1038/ncomms10516

3 Brownlow A, Onoufriou A, Bishop J, Davison N, Thompson D (2016) Corkscrew seals:  
4 grey seal (*Halichoerus grypus*) infanticide and cannibalism may indicate the cause of  
5 spiral lacerations in seals. Plos One 11:e 0156464.

6 Chua K, Gurtler V, Montgomery J, Fraenkel M, Maeall BC, Grayson ML (2007)  
7 *Campylobacter insulaenigrae* causing septicaemia and enteritis. J Med Microbiol  
8 56:1565-1567

9 Dierauf LA (1994) Pinniped forensic necropsy and tissue collection guide, NOAA  
10 Technical Memorandum NMFS-OPR-94-3 US Department of Commerce, National  
11 Marine Fisheries Service, Silver Spring MD

12 Foster G, Holmes B, Steigerwalt AG, Lawson PA, Thorne P, Byrer DE, Ross HM, Xerry  
13 J, Thompson PM, Collins MD (2004) *Campylobacter insulaenigrae* sp. nov., isolated  
14 from marine mammals. Int J Syst Evol Microbiol 54:2369-2373

15 Foster G, Whatmore AM, Dagleish MP, Malnick H, Gilbert MJ, Begeman L, MacGregor  
16 S., Davison, NJ, Roest HJ, Jepson P, Howie F, Muchowski J, Brownlow AC,  
17 Wagenaar JA, Kik MJL, Deaville R, ten Doeschate M, Barley J, Hunter L, IJsseldijk  
18 (2019) *Neisseria animaloris* recovered from lesions in stranded harbor porpoises  
19 (*Phocoena phocoena*) that escaped from traumatic attacks by grey seals (*Halichoerus*  
20 *grypus*). Sci Rep s41598-019-50979-3

21 García-Peña FJ, Pérez-Boto D, Jiménez C, San Miguel E, Echeita A, Renqifo-Herrera  
22 C, García-Párraga D, Ortega-Mora LM, Pedraza-Diaz S (2010) Isolation and

1 characterisation of *Campylobacter* spp. from Antarctic fur seals (*Arctocephalus*  
2 *gazella*) at Deception Island, Antarctica. Appl Environ Microbiol 76:6013-6016

3 Gilbert MJ, Zomer AL, Timmerman AJ, Spaninks MP, Rubio-Garcia A, Rossen JW,  
4 Duim B, Wagenaar JA (2018) *Campylobacter blaseri* sp. nov. reported from harbor  
5 seals (*Phoca vitulina*). Int J Syst Evol Microbiol 68:1787-1794

6 Gilbert MJ, Miller WG, St Leger J, Chapman MH, Timmerman AJ, Foster G, Wagenaar  
7 JA (2017) *Campylobacter pinnipediorum* sp. nov. subsp. *pinnipediorum* subsp. nov. &  
8 *Campylobacter pinnipediorum* sp. nov. subsp. *caledonicus* subsp. nov., isolated from  
9 pinnipeds. Int J Syst Evol Microbiol 67:1961-1968

10 Gilbert MJ, IJsseldijk LL, Rubio-García A, Gröne A, Duim B, Rossen J, Zomer AL,  
11 Wagenaar JA (2020) After the bite: bacterial transmission from grey seals  
12 (*Halichoerus grypus*) to harbour porpoises (*Phocoena phocoena*). R Soc Open Sci  
13 7:192079

14 González M, Paz Villaneuva, M., DeBruyne L, Vandamme P, Fernández H (2011)  
15 *Campylobacter insulaenigrae*: first isolation report from South American sea lion  
16 (*Otaria flavescens*, Shaw, 1800). Braz J Microbiol 42:261-265

17 Leopold MF, Begeman L, van Bleijswijk JDL IJsseldijk LL, Witte HJ, Gröne A (2015a)  
18 Exposing the grey seal as a major predator of harbour porpoises. Proceed R Soc B  
19 282:20142429

20 Miller WG, Yee E, Chapman MH, Smith TPL, Bono JL, Huynh S, Parker CT,  
21 Vandamme P, Luong K, Korlach J (2014) Comparative genomics of the  
22 *Campylobacter lari* group. Genome Biol Evol 6:3252-3266



1 Sheppard SK, Dallas JF, Strachan NJ, MacRae M, McCarthy MD, Wilson DJ, Gormley  
2 FJ, Falush D, Ogden ID, Maiden MCJ, Forbes KJ (2009) *Campylobacter* genotyping  
3 to determine the source of human infection. Clin Infect Dis 48:1072-1078

4 Skirrow MB (1994) Diseases due to *Campylobacter*, *Helicobacter* and related bacteria.  
5 J.Comp Pathol 111:113-149

6 Stoddard RA, Miller WG, Foley JE, Lawrence J, Gulland FMD, Conrad PA, Byrne BA  
7 (2007) *Campylobacter insulaenigrae*: isolates from North American elephant seals  
8 (*Mirounga angustirostris*) in California. Appl Environ Microbiol 73:1729-1735

9 Strachan NJ, Forbes KJ (2010) The growing UK epidemic of campylobacteriosis.  
10 Lancet 376:665-667

11 Tam CC, Rodriguez LC, Viviani L, Dodds JP, Evans MR, Hunter PR, Gray JJ, Letley  
12 LH, Rait G, Tompkins DS, O'Brien S (2012) Longitudinal study of infectious intestinal  
13 disease in the UK (IID2 study): incidence in the community and presenting to general  
14 practice. Gut 61:69-77

15 Van Neer A, Jensen LF, Siebert U (2015) Grey seal (*Halichoerus grypus*) predation  
16 on harbor seals (*Phoca vitulina*) on the island of Helgoland, Germany. J Sea Res 97:1-  
17 4.