


## Article

## Seasonal and daytime variation in multiple immune parameters in humans: Evidence from 329,261 participants of the UK Biobank cohort

## Clocks and Calendars of the Immune System



Cathy Wyse,  
Grace O'Malley,  
Andrew N.  
Coogan, Sam  
McConkey, Daniel  
J. Smith

cathywyse@rcsi.com

**Highlights**

Seasonal outbreaks are common features of human infectious disease

Immune parameters varied across seasons and time of day in UK Biobank participants

Innate variability in immune parameters could influence human susceptibility to disease

Wyse et al., iScience 24,  
102255  
April 23, 2021 © 2021 The  
Authors.  
[https://doi.org/10.1016/  
j.isci.2021.102255](https://doi.org/10.1016/j.isci.2021.102255)

## Article

## Seasonal and daytime variation in multiple immune parameters in humans: Evidence from 329,261 participants of the UK Biobank cohort

Cathy Wyse,<sup>1,\*</sup> Grace O'Malley,<sup>1</sup> Andrew N. Coogan,<sup>2</sup> Sam McConkey,<sup>3</sup> and Daniel J. Smith<sup>4,5</sup>

## SUMMARY

Seasonal disease outbreaks are perennial features of human infectious disease but the factors generating these patterns are unclear. Here we investigate seasonal and daytime variability in multiple immune parameters in 329,261 participants in UK Biobank and test for associations with a wide range of environmental and lifestyle factors, including changes in day length, outdoor temperature and vitamin D at the time the blood sample was collected. Seasonal patterns were evident in lymphocyte and neutrophil counts, and C-reactive protein CRP, but not monocytes, and these were independent of lifestyle, demographic, and environmental factors. All the immune parameters assessed demonstrated significant daytime variation that was independent of confounding factors. At a population level, human immune parameters vary across season and across time of day, independent of multiple confounding factors. Both season and time of day are fundamental dimensions of immune function that should be considered in all studies of immuno-prophylaxis and disease transmission.

## INTRODUCTION

Annual cycles in vulnerability to infectious disease are an established feature of human epidemiology: most respiratory viruses cause winter-time infection and polio is principally a summer-time disease (Dowell and Shang Ho, 2004). Childhood infectious diseases (meningitis, mumps, pertussis, and varicella) (Shah et al., 2006), and many of the contagious diseases that affect domestic animals (Poljak et al., 2014) (Jactel et al., 1990) are seasonal as are relapses in autoimmune diseases (Harding et al., 2017; Mori et al., 2019). The factors that mediate this seasonality are poorly understood and circannual patterns are simply an assumed component of the dynamics of infectious diseases. In addition to seasonality, animals and humans are more susceptible to infectious disease during the resting phase of their daily cycle (Tognini et al., 2017), adding a further circadian dimension to disease vulnerability.

The axial and orbital rotations of the Earth generate predictable seasonal and daily rhythms of light and darkness. These conditions in turn generate circadian and seasonal oscillations in ambient temperature, food availability, predation, and risk of infection. Evolution has equipped animals with innate timing mechanisms, or "clocks", that synchronize physiology to these recurring periods of increased risk. The circadian clock is generated by a series of interconnected transcription–translation feedback loops that regulate the expression of a panel of clock-controlled genes (Takahashi, 2017). Most mammalian cells contain a molecular clock and overall rhythmicity is maintained by a master clock located in the suprachiasmatic nuclei of the hypothalamus, conferring time dependence on most physiological parameters through hormonal and neural signals (Takahashi, 2017). Mice deficient in the cryptochrome clock genes (*Cry1* and *Cry2*), show elevated proinflammatory cytokines (Narasimamurthy et al., 2012), and an autoimmune phenotype (Cao et al., 2017), while loss of clock function by deletion of the clock gene *Bmal1* was associated with augmented immunity against bacterial infection (Kitchen et al., 2020), all evidence for a close association between circadian timing mechanisms and immune function.

The mechanisms driving seasonality in humans are unclear, but in animals, the seasonal clock is generated by changes in thyroid hormones in the brain that respond to day length signaled by the pineal hormone melatonin (Wood and Loudon, 2014). The circadian clock is entrained by the 24-hr photoperiod, while the seasonal clock entrains to day length patterns in Northern latitudes and to seasonal patterns in rain

<sup>1</sup>School of Physiotherapy, Division of Population Health Sciences, Royal College of Surgeons in Ireland, Beaux Lane House, Mercer Street Lower, Dublin, Ireland

<sup>2</sup>Kathleen Lonsdale Institute for Human Health Research, Maynooth University, Maynooth, Kildare, Ireland

<sup>3</sup>Royal College of Surgeons in Ireland: University of Medicine and Health Science, Dublin, Ireland

<sup>4</sup>Institute of Health and Wellbeing, University of Glasgow, Glasgow, Scotland

<sup>5</sup>Lead contact

\*Correspondence: cathywyse@rcsi.com

<https://doi.org/10.1016/j.isci.2021.102255>



and food availability in tropical regions, where day length is constant (Bronson, 2009; Stevenson and Pendergast, 2015). This is analogous with aspects of seasonality of the human immune system, where viral infection and immune cell numbers are associated with day length (e.g. winter peak in influenza) in Northern clines and with climatic changes in tropical regions (Tamerius et al., 2011).

Together, the seasonal and circadian clock synchronize physiology in two dimensions of time, optimizing homeostasis by anticipating changes in the environment. For example, plants (Gangappa and Kumar, 2018), fish (Fortes-Silva et al., 2019), birds (Markowska et al., 2017) and mammals (Pawlak et al., 2005; Yellon and Tran, 2002) all align their immune defense with the time of day that pathogenic and physical challenge are most likely. This conservation across the biological kingdoms is strong evidence that temporal modulation of immune function is an ancient and fundamental mechanism that has evolved to optimize survival in variable environmental conditions.

Laboratory experiments corroborate epidemiological evidence of circadian and seasonal rhythms in disease susceptibility. For example, mice are more resilient to experimental inflammatory (Feigin et al., 1969; Halberg et al., 1960), infectious (Bellet et al., 2013; Edgar et al., 2016; Wongwiwat et al., 1972) and physical challenges (Cable et al., 2017) delivered at night (their active circadian phase) or in summer (Schulman and Kilbourne, 1963). Importantly, these daily cycles in vulnerability persist in constant conditions (photoperiod, temperature, or humidity) (Schulman and Kilbourne, 1963) (Shackelford and Feigin, 1973) and are absent in animals lacking a circadian clock (Curtis et al., 2015; Edgar et al., 2016), demonstrating clock-mediated regulation that is not driven by current environmental conditions. Similar to rodents, humans are more resistant to the effects of inflammatory or infectious challenge (Alamili et al., 2014) (Pollmächer et al., 1996) delivered during their active circadian phase (day time), or in summer (W M Lee et al., 2012) (Shadrin et al., 1977; Zykov and Sosunov, 1987).

Circulating white blood cell counts are known to oscillate across 24hr under basal conditions, reflecting distribution of cells between tissues and the periphery (Stenzinger et al., 2019). Importantly, these rhythms persist in constant conditions and are absent in animals with ablated clock function (Stenzinger et al., 2019), indicating that they are mediated via innate circadian timing mechanisms. The extensive data collection within UK Biobank represents an unprecedented opportunity to assess seasonal and time-of-day variation in levels of human immune parameters. Here, we provide evidence of endogenous seasonal and daytime variability in human immune function at a population level, and we demonstrate that these patterns are independent of a wide range of demographic, environmental, and lifestyle factors.

## RESULTS

The exclusion criteria for this study resulted in the removal of 173,275 study participants. The remaining cohort was mostly White (98%), with the other ethnic groups poorly represented (<2% participants). Summary data that describes the demography and lifestyle of the 329,261 participants that were eligible for inclusion are given in Table 1.

Mean values for WBC percentage and CRP were plotted against month and time of day for all participants, and annual and daily variation was evident on visual inspection and univariate analysis (Figures 1 and 2), but there was no seasonal or diurnal variation in the titer levels of any antigen (data not shown). The probability of seropositive status to any of the 20 antigens analysis was not associated with the month or time of day of analysis (data not shown). Summary data for white blood cell count, vitamin D, and CRP levels at all time points are shown in supplementary information (Tables S1–S4). Mean and 95% confidence intervals for monthly data with fitted cosinor models for lymphocyte, monocyte, neutrophil, and CRP are shown in Figures 1 and 2. Cosinor analysis showed that the seasonal patterns were statistically significant for a 12-month assumed periodicity for CRP and WBC counts (Table 2).


CRP levels were higher in the winter months, peaking in December, with lowest levels in July. The seasonal pattern of neutrophil counts was similarly higher in winter, peaking in Jan with lowest levels recorded in summer (July). The seasonal pattern of lymphocyte counts peaked in spring (March) and troughed in autumn (October) (; Figure 1). There was no significant seasonal pattern in monocyte counts.

Multiple linear regression was next applied to investigate whether the immune parameters were associated with day length and, if so, whether these were independent of other lifestyle and environmental factors that


**Table 1. Demographic and lifestyle characteristics**

Variable	(n = 329,261)
Age (years)	
Mean (SD)	55.8 (8.19)
Townsend deprivation index	
Mean (SD)	-1.55 (2.94)
Ethnicity	
White	306146 (91.0%)
Physical activity	
Mean (SD)	46.0 (62.8)
Sedentary behavior	
Mean (SD)	4.90 (2.24)
BMI	
Mean (SD)	26.8 (4.34)
Smoking	
Yes	31,063 (9.2%)
Chronotype	
More evening than morning	81,240 (24.1%)
Evening	23,185 (6.9%)
More morning than evening	105763 (31.4%)
Morning	78,226 (23.2%)

could confound associations via unrelated seasonality. There was some degree of correlation between outdoor temperature and day length, as expected, but the multicollinearity diagnostic indices, VIF (variance inflation factor) and tolerance did not reach the thresholds (>10 and <0.2, respectively) to indicate problematic multicollinearity (Table S5). CRP levels, as well as neutrophil and lymphocyte counts were found to be significantly associated with day length, independent of demographic, lifestyle, and



**Figure 1. Annual variation in total monocytes, neutrophils, lymphocytes, and CRP. Data are mean (bars) and 95% confidence intervals (boxes), with fitted cosinor curves (dotted line). Daylength is indicated by the box color gradients**



**Figure 2. Daytime variation in monocytes, neutrophils, lymphocytes, and CRP. Data are mean (bars) and 95% confidence intervals (boxes), with fitted segmented regression lines (dotted black lines). The color gradient represents mean zenith angle of the sun at each time point is given to indicate daylight**

environmental factors (Table 3) including outdoor temperature and vitamin D. The relationship between vitamin D and CRP was found to be dependent on BMI, and an interaction term to account for this effect was included in the CRP regression model. Interaction was also detected between vitamin D and sex for all WBC markers, and these interaction terms were added to the regression models (supplementary information Tables S5 and S6). In the fully adjusted model, neutrophil count and CRP showed significant negative associations with day length, while lymphocyte count was positively associated, as also shown in cosinor analysis. Monocyte count was not significantly associated with day length in the fully adjusted model (Model 3).

Segmented linear regression analysis of CRP and WBC counts over the daily time course showed significant daytime variation that was represented by segmental regression lines (Figure 2). The peaks and trough (break points) for each marker are shown in Table 4. All white blood cells showed significant daily variation, with counts lowest in the early morning and increasing as the day progressed. Neutrophil count reached a plateau at 3pm. CRP levels were highly variable, peaking at 1pm, and decreasing thereafter (Table 4).

Linear regression analysis demonstrated that the daytime changes in WBCs and CRP were in most cases independent of lifestyle and environmental factors (Table 5). The morning ascending segment of the CRP daily curve was the only section of any of the curves that did not retain statistical significance after adjustment. However the daily CRP curve showed a significant relationship with time of day for the later parts of the day, after the break point at 1pm.

## DISCUSSION

The human immune system is not constant over 24hr or across the seasons and the time of exposure to pathogens is an important consideration in determining risk of infection Here we report seasonal and day-time patterns in immune cells and inflammatory markers within a large sample of the UK population. Importantly, we demonstrate that these patterns are independent of multiple demographic, lifestyle and local

**Table 2. Parameters describing the amplitude, annual peak (acrophase), and mean value (mesor) predicted by fitted cosinor model**

Day length	Amplitude	Acrophase	Bathyphase	Mesor	p
Neutrophils (10 <sup>9</sup> /L)	0.80	Jan	Jul	4.10	<0.001
Lymphocytes (10 <sup>9</sup> /L)	0.03	April	Oct	1.93	<0.001
CRP (mg/L)	0.10	Jan	Jun	2.23	<0.001

Data are mean and sd.

environmental variables, supporting the existence of endogenous seasonal and daytime patterns in human immune parameters. These findings highlight the importance of future studies to understand the time dimensions of immune function and their implications for preventing and controlling outbreaks of infectious disease.

UK Biobank combines multiple blood, demographic, and environmental parameters in a large sample of the UK population that allowed us to generate novel information on the factors that mediate daytime and seasonal variations in peripheral immune cells. It was possible to adjust for multiple lifestyle-associated parameters that might be seasonal such as physical activity and sleep duration, and unlike previous studies, we were also able to adjust for the effects of vitamin-D and outdoor temperature. The timing of sample collection was randomized in this study, which allowed us to exclude the possibility of bias through self-selection of the time of sampling. Finally, this study contributes to current knowledge by demonstrating significant daytime and seasonal variability in immune cells in the biggest population sample reported to date.

The greatest seasonal and daytime changes in this study were seen in lymphocyte numbers, with high-amplitude variation over seasons and days. Lymphocytes were lower during the early parts of the day, increasing as the day progressed, consistent with previous reports that lymphocytes circulating in blood are lower during the respective active phase of humans (Ackermann et al., 2012) (He et al., 2018) (Lange et al., 2010) (Born et al., 1997) (Abo et al., 1981) and rodents (Druzd et al., 2017; He et al., 2018) (Suzuki et al., 2016). Interestingly, longer sleep times were associated with numbers of circulating white blood cells across multiple mammalian species (Preston et al., 2009), further supporting a relationship between the timing of activity and immune function. Circadian rhythms in the homing and egress of lymphocytes through the lymphatic system and other tissues underlies these diurnal changes of lymphocyte numbers in blood (Druzd et al., 2017; He et al., 2018) (Pick et al., 2019). Since the lymph nodes contain the interaction between lymphocytes and antigen, longer accumulation times increase antigen encounters and potentiate the adaptive immune response (Scheiermann et al., 2018) (Suzuki et al., 2016). Consequently, lymphocyte numbers in the periphery drop as trafficking to the tissues increases, along with increased tissue surveillance and resistance to infection (He et al., 2018) (Pick et al., 2019). Circadian rhythms of lymphocyte trafficking to the periphery are abolished by genetic ablation of clock function and persist in constant conditions (Deprés-Brummer et al., 1997; Druzd et al., 2017), confirming their regulation by the innate circadian clock in mice. These endogenous rhythms are associated with time-of-day dependent changes in adaptive

**Table 3. Associations between day length and CRP, lymphocyte, neutrophil, and monocyte count**

Day length	Model 1		Model 2		Model 3	
White blood cells (10 <sup>9</sup> /L)	-0.011***	[-0.013,-0.009]	-0.008***	[-0.011,-0.006]	-0.011***	[-0.015,-0.007]
Neutrophils (10 <sup>9</sup> /L)	-0.015***	[-0.017,-0.014]	-0.013***	[-0.015,-0.011]	-0.014***	[-0.018,-0.011]
Monocytes (10 <sup>9</sup> /L)	0.001***	[0.001,0.001]	0.001***	[0.001,0.001]	-0.000	[-0.001,-0.000]
Lymphocytes (10 <sup>9</sup> /L)	0.002***	[0.002,0.003]	0.003***	[0.002,0.004]	0.004***	[0.003,0.005]
CRP (mg/L)	-0.006***	[-0.007,-0.004]	-0.005***	[-0.006,-0.003]	-0.004***	[-0.007,-0.002]

Data are expressed as regression coefficients (B), with 95% confidence intervals in parentheses.

Model 1 was adjusted for age, sex, ethnicity, deprivation.

Model 2 was adjusted for Model 1 + BMI, physical activity, sedentary behavior, sleep duration, chronotype, shiftwork, smoking, alcohol.

Model 3 was adjusted for Model 2 + vitamin D, outdoor temperature, time of day, blood analyzer, and UK Biobank assessment center.

**Table 4. Segmented regression parameters showing predicted break points for each segment and regression coefficient for overall segmented linear model**

Time of day	Break point time of day (se)	p*
White blood cells (10 <sup>9</sup> /L)	14.34 (0.069)	<0.001
Neutrophils (10 <sup>9</sup> /L)	14.62 (0.040)	<0.001
Monocytes (10 <sup>9</sup> /L)	13.27 (0.160)	<0.001
Lymphocytes (10 <sup>9</sup> /L)	16.12 (0.180)	<0.001
CRP (mg/L)	12.71 (0.220)	<0.01

\*Davies' test was applied to test for significant differences of slopes between each segmented relationship.

immunity, including amplified response to induction of autoimmunity (experimental autoimmune encephalomyelitis), immunization (Suzuki et al., 2016) and viral infection (influenza) during the active phase (Druz et al., 2017). Thus the time of day that pathogenic challenge occurs affects the adaptive immune response generated days later, one of the mechanisms through which lymphocyte trafficking might modulate seasonal and circadian vulnerability to infection. In agreement with previous findings (Aguirre-Gamboa et al., 2016; Liu and Taioli, 2015), lymphocytes were positively associated with day length in our study; cell numbers were lower in autumn and peaked in spring. These seasonal variations in peripheral lymphocyte counts suggest that humans have some capacity for seasonal regulation of lymphocyte trafficking that could contribute to seasonality in susceptibility to infection.

Previous studies have reported seasonal and circadian patterns in antibody titers in humans (Rosenblatt et al., 1982), IgM (Dozier et al., 1997), often discovered serendipitously in the course of other investigations (Dozier et al., 1997). Leukocytes collected at different times of year showed decreased ex vivo response (thymidine incorporation, cytokine release) to activation in winter time in humans and rats (Boctor et al., 1989; Brock, 1983, 1987) (Amat and Torres, 1993), and diurnal patterns in antibody titers (Kurupati et al., 2017) and in ex vivo response to stimulation of PBMCs (Van Rood et al., 1991) (de Bree et al., 2020) have been reported in humans. This study in UK Biobank is the first to investigate circadian and seasonal patterns in antibody titers to common infectious agents at a population level. Despite our comparatively large sample size, we found no evidence for seasonal or daily variation in antibody titers or in the probability of testing immunopositive to any of the 20 antigens investigated in this study. However, the antibody response to vaccination or viral infection and subsequent decay is subject to wide variation between individuals (Antia et al., 2018), which is not accounted for by the cross-sectional design of the present study. Longitudinal experiments are required to establish if antibody titers vary by season or time of day and how this might impact on response to vaccination or infection. Daytime variation in antibody titers could confound studies of the efficacy of vaccination that use antibody response as an outcome variable (Kurupati et al., 2017) (Long et al., 2016a, 2016b), and future investigations within individuals and with multiple sample time points are warranted to understand basal variation in antibody titers.

Blood neutrophil counts were lowest in early morning in the UK Biobank participants, increasing thereafter to plateau after 3pm. Previous studies demonstrated comparable circadian rhythms in peripheral neutrophil counts that were low in the rest phase, and that increased over the active phase in both humans and mice (Ackermann et al., 2012) (Jilma et al., 1999). Neutrophils have a half-life less than 24hr, and circadian rhythmicity is regulated through clock-controlled oscillations in chemokine pathways that drive release of young cells in the active phase and clearance of aged neutrophils from the periphery in the resting phase (Adrover et al., 2019; Casanova-Acebes et al., 2013). These rhythms in neutrophil tissue migration were shown to underlie increased resistance to infection (*Candida albicans*), during the active phase in mice (Adrover et al., 2019) and to diurnal variation in bactericidal function ex vivo in human neutrophils (Ella et al., 2016). Neutrophil counts were negatively associated with day length in our study, in agreement with previous studies in humans living at temperate latitudes (Golding et al., 2015; Liu and Taioli, 2015). We extend these findings to demonstrate high peripheral neutrophil counts in winter time at a population level that were related to annual photoperiod, independent of participant lifestyle, local environmental conditions and vitamin D. In addition to total counts, previous studies have demonstrated seasonality of functional aspects of neutrophil immune function, including adhesive capacity, CD11b/CD18 expression and ROS production, resulting in augmented bactericidal properties of neutrophils collected in summer (Klink et al., 2012). The seasonal and daytime patterns in neutrophil count reported here, and

**Table 5. Associations between time of day and CRP, lymphocyte, neutrophil, and monocyte counts**

Day length	Model 1		Model 2		Model 3	
<b>WBCs (10<sup>9</sup>/L)</b>						
Segment 1	0.207***	[0.203,0.211]	0.199***	[0.195,0.203]	0.197***	[0.193,0.202]
Segment 2	0.078***	[0.073,0.082]	0.082***	[0.078,0.086]	0.082***	[0.078,0.087]
<b>Neutrophils (10<sup>9</sup>/L)</b>						
Segment 1	0.094**	[0.033,0.156]	0.095**	[0.030,0.159]	0.101**	[0.036,0.165]
Segment 2	0.268***	[0.243,0.293]	0.262***	[0.236,0.289]	0.259***	[0.233,0.285]
<b>Monocytes (10<sup>9</sup>/L)</b>						
Segment 1	0.085***	[0.048,0.122]	0.086***	[0.047,0.125]	0.085***	[0.046,0.124]
Segment 2	-0.206***	[-0.235,-0.177]	-0.199***	[-0.230,-0.168]	-0.195***	[-0.227,-0.164]
<b>Lymphocytes (10<sup>9</sup>/L)</b>						
Segment 1	0.051***	[0.050,0.052]	0.050***	[0.049,0.051]	0.050***	[0.049,0.051]
Segment 2	0.082***	[0.079,0.085]	0.085***	[0.082,0.088]	0.085***	[0.082,0.088]
<b>CRP (mg/L)</b>						
Segment 1	0.015	[-0.000,0.030]	0.009	[-0.007,0.025]	0.010	[-0.006,0.026]
Segment 2	-0.009*	[-0.017,-0.001]	-0.013***	[-0.021,-0.006]	-0.013**	[-0.021,-0.005]

Time of day is represented by 2 linear splines, to account for non-linear relationships with the independent variables.

Data are expressed as regression coefficients (B), with 95% confidence intervals in parentheses.

Model 1 was adjusted for age, sex, ethnicity, deprivation.

Model 2 was adjusted for BMI, physical activity, sedentary behavior, sleep duration, chronotype, smoking, alcohol.

Model 3 was adjusted for Model 2 + day length, blood analyzer and UK Biobank assessment center.

in previous studies support evidence from animal studies that time-dependent cycles of tissue migration could contribute to neutrophil-mediated resilience to infection during the active phase, and relative vulnerability to infection in winter time (Adrover et al., 2019; Casanova-Acebes et al., 2013).

Monocyte counts were lower in the morning compared to evening in UK Biobank participants, consistent with previous reports that monocytes increase during the active phase in mice (Nguyen et al., 2013) and humans (Born et al., 1997). The acute phase protein, CRP showed a weak daily pattern in this study, with levels higher in daytime, again corroborating previous reports of diurnal patterns of CRP in humans (Morris et al., 2017) (Chiriboga et al., 2009; Liu and Taioli, 2015; Rudnicka et al., 2007) The immune function of tissue macrophages was shown to be regulated by intrinsic circadian timing mechanisms that were independent of systemic glucocorticoid secretion (Keller et al., 2009). Circadian rhythms in the circulation and tissue migration of monocytes in mice were similarly regulated through an innate cell-intrinsic clock mechanism and their oscillation coincides with an enhanced inflammatory response when monocytes are decreasing at the beginning of the rest phase (Nguyen et al., 2013) and increased lethality of endotoxin challenge at this time (Halberg et al., 1960). In agreement, human volunteers show a heightened response to endotoxin challenge in the evening (Alamili et al., 2014; Pollmächer et al., 1996). This increased inflammatory response in the active phase might maximize innate immune defense at a time when pathogenic challenge is most likely (Curtis et al., 2014) but could also leave animals more vulnerable to the toxic effects of augmented inflammation.

Monocyte counts are higher in winter in some (Aguirre-Gamboa et al., 2016) but not all (Liu and Taioli, 2015) previous studies. There was no evidence of a seasonal pattern in UK Biobank participants, and monocytes were not associated with day length in the fully adjusted model. While peripheral counts are not always seasonal, monocyte function shows strong seasonality *ex vivo*, with an augmented proinflammatory response to activation in summer time (Myriantefs et al., 2003; ter Horst et al., 2016). We found a weak seasonal pattern in the acute phase protein, CRP in UK Biobank, with levels higher in the winter months. Peripheral CRP and other proinflammatory markers were higher in winter in many studies in humans (Sung, 2006) (Liu and Taioli, 2015), and this is thought to contribute to seasonal prevalence of cardiovascular disease (Sartini et al., 2017).

Seasonality of human viral infections is generally and intuitively thought to be driven by annual changes in temperature or humidity, but there is increasing evidence that innate variation in host disease susceptibility

is an important contributor. In support of this, many diseases are seasonal in tropical regions where temperature and humidity are constant (Bloom-Feshbach et al., 2013) (Tamerius et al., 2011). Furthermore, outbreaks of influenza occur annually and simultaneously at latitudes that are oceans apart (Lofgren et al., 2007) despite variations in local climatic conditions and human behavior. Recurrent seasonality is a feature of the epidemiology of infectious disease in animals that do not share human winter time behaviors such as increased time indoors, crowding or school terms. The prevalence of human respiratory viruses does not correspond with the prevalence of the respiratory disease they cause; remarkably, detection of viral infection is relatively low in the months that respiratory disease is highest (Lee et al., 2012).

Vitamin D is suspected to contribute to disease seasonality due to known associations with immune function and highly seasonal serum levels (Cannell et al., 2006), but this postulation is not corroborated by models that compared serum vitamin D with influenza transmission in population based studies (Shaman et al., 2011; Xu et al., 2016). Our findings in UK Biobank showed that seasonal changes in white blood cells and CRP were related to day length independent of vitamin D levels, in agreement with evidence that circulating vitamin D was not responsible for seasonality in the proinflammatory functions of human monocytes (ter Horst et al., 2016). Finally, evidence of widespread seasonal regulation of transcription of genes regulating immune function and of reversed expression patterns in Northern and Southern hemispheres strongly supports endogenous regulation of seasonality in human immune function (Dopico et al., 2015). Seasonality of human infectious disease may be driven by an endogenous circannual rhythmicity in host immunity that generates cycles of enhancement and suppression of immune function and windows of vulnerability to infection, as proposed by Dowell in 2001 (Dowell, 2001).

### Limitations of the study

This study in UK Biobank is the largest investigation of the seasonal and daytime patterns in human immune cells, inflammatory markers and antibody titers at population level, but our results are subject to many important limitations. Although UK Biobank represents a very large population sample of middle-aged UK adults, only 5.5% of those who received an invitation volunteered to take part. A broad range of socio-economic groups are represented but ethnic diversity is low (Sudlow et al., 2015). Furthermore, UK Biobank specifically recruited participants aged 40–69 years, so our findings may not apply to younger people. Some of the data that we analyzed were self-reported, including ethnicity, physical activity, health status and chronotype, and misclassification errors are possible. The participants denied chronic disease, but we cannot exclude the presence of acute infection at the time of assessment. The study design was cross-sectional, and a single blood sample was available from each participant so the influence of within subject variation cannot be assessed. We were unable to assess circadian patterns since there were no nighttime blood sample collections, and our results are limited to analysis of daytime variation. Nevertheless, the daily patterns we report in over 300,000 participants are consistent with the results of previous studies where blood was withdrawn at regular time points over 24hr under experimental conditions. We present results of total cell counts only, and further studies are required to investigate subtypes of lymphocytes and neutrophils. The immune parameters that we report are affected by a multitude of factors related to interactions between host, pathogen, and the environment. It is not possible to consider all of these in a population-based study, and the mechanisms driving the associations with day length and time of day that we report require investigation under controlled experimental conditions. Furthermore, the effect sizes we report are small, and likely to be of clinical significance for population-level disease control, rather than for the health of individuals. The strengths of this study are the large sample size and that the times of sample collection were randomly allocated to each participant. It is also a strength that we were able to investigate the effects of day length and time of day on immune parameters while adjusting for other factors thought to affect seasonal and daytime variability including physical activity, vitamin D, and outdoor temperature.

Seasonality in the epidemiology of infectious disease is considered to be generated by environment and pathogen related factors, and innate variability in host susceptibility to infection is rarely considered. Our findings of seasonal and daytime variability in multiple immune parameters in a large sample of the UK population under basal, free living conditions that were independent of environmental conditions, support the contribution of innate mechanisms to variability in disease susceptibility.

Future research should focus on whether elective restriction of human activity at times of increased vulnerability to infection through night time and winter curfews could control the spread of infectious disease by minimizing exposure to pathogens during susceptible periods. This is exactly the function that has driven

the evolution of temporal regulation of the immune system and harnessing this innate attribute could optimize our resilience to COVID-19 and future pandemics.

### Resource availability

#### Lead contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Cathy Wyse ([cathywyse@rcsi.com](mailto:cathywyse@rcsi.com)).

#### Materials availability

This study did not generate any new unique reagents.

#### Data and code availability

The data sets generated during this study are available at the UK Biobank repository, [www.ukbiobank.ac.uk](http://www.ukbiobank.ac.uk).

## METHODS

All methods can be found in the accompanying [Transparent Methods supplemental file](#).

## SUPPLEMENTAL INFORMATION

Supplemental information can be found online at <https://doi.org/10.1016/j.isci.2021.102255>.

## ACKNOWLEDGMENTS

The authors are grateful to the UK Biobank (Project ID 26209) and its participants for the data used in these studies. This study was funded by an MRC Mental Health Data Pathfinder Award (MC\_PC\_17217) to DJS. DJS also acknowledges support from a Lister Institute Prize Fellowship (2016-2021). GOM was supported by an RCSI Strategic Academic Recruitment (StAR) Fellowship. We are grateful to Claire Pallas for helpful comments on the manuscript.

## AUTHOR CONTRIBUTIONS

The study was conceived by CW and DS, data analysis was completed by CW. GOM, SMcC, and AC contributed to writing the manuscript and all authors approved the final manuscript.

## DECLARATION OF INTERESTS

The authors declare no competing interests.

Received: November 23, 2020

Revised: January 14, 2021

Accepted: February 25, 2021

Published: April 23, 2021

## REFERENCES

- Abo, T., Kawate, T., Itoh, K., and Kumagai, K. (1981). Studies on the bioperiodicity of the immune response. I. Circadian rhythms of human T, B, and K cell traffic in the peripheral blood. *J. Immunol.* *126*, 1360–1363.
- Ackermann, K., Revell, V.L., Lao, O., Rombouts, E.J., Skene, D.J., and Kayser, M. (2012). Diurnal rhythms in blood cell populations and the effect of acute sleep deprivation in healthy young men. *Sleep* *35*, 933–940.
- Adrover, J.M., del Fresno, C., Crainiciuc, G., Cuartero, M.I., Casanova-Acebes, M., Weiss, L.A., Huerga-Encabo, H., Silvestre-Roig, C., Rossaint, J., Cossio, I., et al. (2019). A neutrophil timer coordinates immune defense and vascular protection. *Immunity* *50*, 390–402.e10.
- Aguirre-Gamboa, R., Joosten, I., Urbano, P.C.M., van der Molen, R.G., van Rijssen, E., van Cranenbroek, B., Oosting, M., Smeekens, S., Jaeger, M., Zorro, M., et al. (2016). Differential effects of environmental and genetic factors on T and B cell immune traits. *Cell Rep.* *17*, 2474–2487.
- Alamili, M., Bendtzen, K., Lykkesfeldt, J., Rosenberg, J., and Gögenur, I. (2014). Pronounced inflammatory response to endotoxaemia during nighttime: a randomised cross-over trial. *PLoS One* *9*, e87413.
- Amat, J., and Torres, A. (1993). Circannual rhythm in the effects of stress on the humoral immune response of the rat. *Neurosci. Lett.* *160*, 190–192.
- Antia, A., Ahmed, H., Handel, A., Carlson, N.E., Amanna, I.J., Antia, R., and Slifka, M. (2018). Heterogeneity and longevity of antibody memory to viruses and vaccines. *PLoS Biol.* *16*, 1–15.
- Bellet, M.M., Deriu, E., Liu, J.Z., Grimaldi, B., Blaschitz, C., Zeller, M., Edwards, R.A., Sahar, S., Dandekar, S., Baldi, P., et al. (2013). Circadian clock regulates the host response to Salmonella. *Proc. Natl. Acad. Sci. U S A* *110*, 9897–9902.
- Bloom-Feshbach, K., Alonso, W.J., Charu, V., Tamerius, J., Simonsen, L., Miller, M.A., and

- Viboud, C. (2013). Latitudinal variations in seasonal activity of influenza and respiratory syncytial virus (RSV): a global comparative review. *PLoS One* 8, 3–4.
- Boctor, F.N., Charmy, R.A., and Cooper, E.L. (1989). Seasonal differences in the rhythmicity of human Male and female lymphocyte blastogenic responses. *Immunol. Invest.* 18, 775–784.
- Born, J., Lange, T., Hansen, K., Mölle, M., and Fehm, H.L. (1997). Effects of sleep and circadian rhythm on human circulating immune cells. *J. Immunol.* 158, 4454–4464.
- Brock, M.A. (1987). Age-related changes in circannual rhythms of lymphocyte blastogenic responses in mice. *Am. J. Physiol.* 252, 299–302.
- Brock, M.A. (1983). Seasonal rhythmicity in lymphocyte blastogenic responses of mice persists in a constant environment. *J. Immunol.* 130, 2586–2588.
- Bronson, F.H. (2009). Climate change and seasonal reproduction in mammals. *Philos. Trans. R. Soc. B Biol. Sci.* 27, 3331–3340.
- Cable, E.J., Onishi, K.G., and Prendergast, B.J. (2017). Circadian rhythms accelerate wound healing in female Siberian hamsters. *Physiol. Behav.* 171, 165–174.
- Cannell, J.J., Vieth, R., Umhau, J.C., Holick, M.F., Grant, W.B., Madronich, S., Garland, C.F., and Giovannucci, E. (2006). Epidemic influenza and vitamin D. *Epidemiol. Infect.* 134, 1129–1140.
- Cao, Q., Zhao, X., Bai, J., Gery, S., Sun, H., Lin, D.C., Chen, Q., Chen, Z., Mack, L., Yang, et al. (2017). Circadian clock cryptochrome proteins regulate autoimmunity. *Proc. Natl. Acad. Sci. U S A* 114, 12548–12553.
- Casanova-Acebes, M., Pitaval, C., Weiss, L.A., Nombela-Arrieta, C., Chèvre, R., A-González, N., Kunisaki, Y., Zhang, D., Van Rooijen, N., Silberstein, L.E., et al. (2013). Rhythmic modulation of the hematopoietic niche through neutrophil clearance. *Cell* 153, 1025.
- Chiriboga, D.E., Yunsheng, M., Li, W., Iii, E.J.S., Hébert, J.R., Merriam, P.A., Rawson, E.S., and Ockene, I.S. (2009). Seasonal and Sex variation of high-sensitivity C-reactive protein in healthy adults: a longitudinal study. *Clin. Chem.* 55, 313–321.
- Curtis, A.M., Bellet, M.M., Sassone-Corsi, P., and O'Neill, L.A. (2014). Circadian clock proteins and immunity. *Immunity* 40, 178–186, <https://doi.org/10.1016/j.immuni.2014.02.002>.
- Curtis, A.M., Fagundes, C.T., Yang, G., Palsson-McDermott, E.M., Wochal, P., McGettrick, A.F., Foley, N.H., Early, J.O., Chen, L., Zhang, H., et al. (2015). Circadian control of innate immunity in macrophages by miR-155 targeting Bmal1. *Proc. Natl. Acad. Sci. U S A* 112, 7231–7236.
- de Bree, L.C.J., Mourits, V.P., Koeken, V.A.C.M., Moorlag, S.J.C.F.M., Janssen, R., Folkman, L., Barreca, D., Krausgruber, T., Fife-Gernedl, V., Novakovic, B., et al. (2020). Circadian rhythm influences induction of trained immunity by BCG vaccination. *J. Clin. Invest.* 130, 5603–5617.
- Deprés-Brummer, P., Bourin, P., Pages, N., Metzger, G., and Lévi, F. (1997). Persistent T lymphocyte rhythms despite suppressed circadian clock outputs in rats. *Am. J. Physiol.* 273, 1891–1899.
- Dopico, X.C., Evangelou, M., Ferreira, R.C., Guo, H., Pekalski, M.L., Smyth, D.J., Cooper, N., Burren, O.S., Fulford, A.J., Hennig, B.J., et al. (2015). Widespread seasonal gene expression reveals annual differences in human immunity and physiology. *Nat. Commun.* 6, 7000.
- Dowell, S.F. (2001). Seasonal variation in host susceptibility and cycles of certain infectious diseases. *Emerg. Infect. Dis.* 7, 369–374.
- Dowell, S.F., and Shang Ho, M. (2004). Seasonality of infectious diseases and severe acute respiratory syndrome - what we don't know can hurt us. *Lancet Infect. Dis.* 4, 704–708.
- Dozier, M.M., Ratajczak, H.V., Sothorn, R.B., and Thomas, P.T. (1997). The influence of vehicle gavage on seasonality of immune system parameters in the B6C3F1 mouse. *Fundam. Appl. Toxicol.* 38, 116–122.
- Druz, D., Matveeva, O., Ince, L., Harrison, U., He, W., Schmal, C., Herzel, H., Tsang, A.H., Kawakami, N., Leliavski, A., et al. (2017). Lymphocyte circadian clocks control lymph node trafficking and adaptive immune responses. *Immunity* 46, 120–132.
- Edgar, R.S., Stangherlin, A., Nagy, A.D., Nicoll, M.P., Efstathiou, S., O'Neill, J.S., and Reddy, A.B. (2016). Cell autonomous regulation of herpes and influenza virus infection by the circadian clock. *Proc. Natl. Acad. Sci. U S A* 113, 10085–10090.
- Ella, K., Csépanyi-Kömi, R., and Káldi, K. (2016). Circadian regulation of human peripheral neutrophils. *Brain Behav. Immun.* 57, 209–221.
- Feigin, R.D., San Joaquin, V.H., Haymond, M.W., and Wyatt, R.G. (1969). Daily periodicity of susceptibility of mice to pneumococcal infection. *Nature* 224, 379–380.
- Fortes-Silva, R., Leme, F.D.O.P., Boaventura, T.P., Mendonça, H.C.P. De, Moreira, J.P.L., Cunha, P.H.H., and Luz, R.K. (2019). Daily rhythms of leukocytes populations, biochemical and enzymatic blood parameters in a carnivorous freshwater catfish (*Lophosilurus alexandri*). *Chronobiol. Int.* 36, 276–287.
- Gangappa, S.N., and Kumar, S.V. (2018). DET1 and COP1 modulate the coordination of growth and immunity in response to key seasonal signals in arabidopsis. *Cell Rep.* 25, 29–37.e3.
- Goldinger, A., Shakhbazov, K., Henders, A.K., McRae, A.F., Montgomery, G.W., and Powell, J.E. (2015). Seasonal effects on gene expression. *PLoS One* 10, e0126995.
- Halberg, F., Johnson, E.A., Brown, B.W., and Bittner, J.J. (1960). Susceptibility rhythm to *E. coli* endotoxin and bioassay. *Proc. Soc. Exp. Biol. Med.* 103, 142–144.
- Harding, K., Tilling, K., MacIver, C., Willis, M., Joseph, F., Ingram, G., Hirst, C., Wardle, M., Pickersgill, T., Ben-Shlomo, Y., and Robertson, N. (2017). Seasonal variation in multiple sclerosis relapse. *J. Neurol.* 264, 1059–1067.
- He, W., Holtkamp, S., Hergenhan, S.M., Kraus, K., de Juan, A., Weber, J., Bradfield, P., Grenier, J.M.P., Pelletier, J., Druz, D., et al. (2018). Circadian expression of migratory factors establishes lineage-specific signatures that guide the homing of leukocyte subsets to tissues. *Immunity* 49, 1175–1190.e7.
- Jactel, B., Espinasse, J., Viso, M., and Valiergue, H. (1990). An epidemiological study of winter dysentery in fifteen herds in France. *Vet. Res. Commun.* 14, 367–379.
- Jilma, B., Hergovich, N., Stohlawetz, P., Eichler, H.G., Bauer, P., and Wagner, O.F. (1999). Circadian variation of granulocyte colony stimulating factor levels in man. *Br. J. Haematol.* 106, 368–370.
- Keller, M., Mazuch, J., Abraham, U., Eom, G.D., Herzog, E.D., Volk, H.D., Kramer, A., and Maier, B. (2009). A circadian clock in macrophages controls inflammatory immune responses. *Proc. Natl. Acad. Sci. U S A* 106, 21407–21412.
- Kitchen, G.B., Cunningham, P.S., Poolman, T.M., Iqbal, M., Maidstone, R., Baxter, M., Bagnall, J., Begley, N., Saer, B., Hussell, T., et al. (2020). The clock gene Bmal1 inhibits macrophage motility, phagocytosis, and impairs defense against pneumonia. *Proc. Natl. Acad. Sci. U S A* 117, 1543–1551.
- Klink, M., Bednarska, K., Blus, E., Kielbik, M., and Sulowska, Z. (2012). Seasonal changes in activities of human neutrophils in vitro. *Inflamm. Res.* 61, 11–16.
- Kurupati, R.K., Kossenkoff, A., Kannan, S., Haut, L.H., Doyle, S., Yin, X., Schmader, K.E., Liu, Q., Showe, L., and Ertl, H.C.J. (2017). The effect of timing of influenza vaccination and sample collection on antibody titers and responses in the aged. *Vaccine* 35, 3700–3708.
- Lange, T., Dimitrov, S., and Born, J. (2010). Effects of sleep and circadian rhythm on the human immune system. *Ann. N. Y. Acad. Sci.* 1193, 48–59.
- Lee, W.M., Lemanske, R.F., Evans, M.D., Vang, F., Pappas, T., Gagnon, R., Jackson, D.J., and Gern, J.E. (2012). Human rhinovirus species and season of infection determine illness severity. *Am. J. Respir. Crit. Care Med.* 186, 886–891.
- Liu, B., and Taioli, E. (2015). Seasonal variations of complete blood count and inflammatory biomarkers in the US population - analysis of NHANES data. *PLoS One* 10, e0142382.
- Lofgren, E., Fefferman, N.H., Naumov, Y.N., Gorski, J., and Naumova, E.N. (2007). Influenza seasonality: underlying causes and modeling theories. *J. Virol.* 81, 5429–5436.
- Long, J.E., Drayson, M.T., Taylor, A.E., Toellner, K.M., Lord, J.M., and Phillips, A.C. (2016a). Morning vaccination enhances antibody response over afternoon vaccination: a cluster-randomised trial. *Vaccine* 34, 2679–2685.
- Long, J.E., Drayson, M.T., Taylor, A.E., Toellner, K.M., Lord, J.M., and Phillips, A.C. (2016b). Corrigendum to 'Morning vaccination enhances antibody response over afternoon vaccination: a cluster-randomised trial' (*Vaccine* (2016) 34(24) (2679–2685)). *Vaccine* 34, 4842.
- Markowska, M., Majewski, P.M., and Skwarko-Sońta, K. (2017). Avian biological clock – immune

- system relationship. *Dev. Comp. Immunol.* 66, 130–138.
- Mori, H., Sawada, T., Nishiyama, S., Shimada, K., Tahara, K., Hayashi, H., Kato, E., Tago, M., Matsui, T., and Tohma, S. (2019). Influence of seasonal changes on disease activity and distribution of affected joints in rheumatoid arthritis. *BMC Musculoskelet. Disord.* 20, 30.
- Morris, C.J., Purvis, T.E., Mistretta, J., Hu, K., and Scheer, F.A.J.L. (2017). Circadian misalignment increases C-reactive protein and blood pressure in chronic shift workers. *J. Biol. Rhythms* 32, 154–164.
- Myrianthefs, P., Karatzas, S., Venetsanou, K., Grouzi, E., Evagelopoulou, P., Boutzouka, E., Fildisis, G., Spiliotopoulou, I., and Baltopoulos, G. (2003). Seasonal variation in whole blood cytokine production after LPS stimulation in normal individuals. *Cytokine* 24, 286–292.
- Narasimamurthy, R., Hatori, M., Nayak, S.K., Liu, F., Panda, S., and Verma, I.M. (2012). Circadian clock protein cryptochrome regulates the expression of proinflammatory cytokines. *Proc. Natl. Acad. Sci. U S A* 109, 12662–12667.
- Nguyen, K.D., Fentress, S.J., Qiu, Y., Yun, K., Cox, J.S., and Chawla, A. (2013). Circadian gene Bmal1 regulates diurnal oscillations of Ly6C(hi) inflammatory monocytes. *Science* 341, 1483–1488.
- Pawlak, J., Majewski, P., Markowska, M., and Skwarło-Sońta, K. (2005). Season- and gender-dependent changes in the immune function of Siberian hamsters (*Phodopus sungorus*). *Neuroendocrinol. Lett.* 26, 55–60.
- Pick, R., He, W., Chen, C.S., and Scheiermann, C. (2019). Time-of-Day-Dependent trafficking and function of leukocyte subsets. *Trends Immunol.* 40, 524–537.
- Poljak, Z., Carman, S., and McEwen, B. (2014). Assessment of seasonality of influenza in swine using field submissions to a diagnostic laboratory in Ontario between 2007 and 2012. *Influenza Other Respi. Viruses* 8, 482–492.
- Pollmächer, T., Mullington, J., Korth, G., Schreiber, W., Hermann, D., Orth, A., Galanos, C., and Holsboer, F. (1996). Diurnal variations in the human host response to endotoxin. *J. Infect. Dis.* 174, 1040–1045.
- Preston, B.T., Capellini, I., McNamara, P., Barton, R.A., and Nunn, C.L. (2009). Parasite resistance and the adaptive significance of sleep. *BMC Evol. Biol.* 9, 9–17.
- Rosenblatt, L.S., Shifrine, M., Hetherington, N.W., Paglierioni, T., and MacKenzie, M.R. (1982). A circannual rhythm in rubella antibody titers. *J. Interdiscipl. Cycle Res.* 13, 81–88.
- Rudnicka, A.R., Rumley, A., Lowe, G.D.O., and Strachan, D.P. (2007). Diurnal, seasonal, and blood-processing patterns in levels of circulating fibrinogen, fibrin D-dimer, C-reactive protein, tissue plasminogen activator, and von Willebrand factor in a 45-year-old population. *Circulation* 115, 996–1003.
- Sartini, C., Barry, S.J.E., Whincup, P.H., Wannamethee, S.G., Lowe, G.D.O., Jefferis, B.J., Lennon, L., Welsh, P., Ford, I., Sattar, N., and Morris, R.W. (2017). Relationship between outdoor temperature and cardiovascular disease risk factors in older people. *Eur. J. Prev. Cardiol.* 24, 349–356.
- Scheiermann, C., Gibbs, J., Ince, L., and Loudon, A. (2018). Clocking in to immunity. *Nat. Rev. Immunol.* 18, 423–437.
- Schulman, J.L., and Kilbourne, E.D. (1963). Experimental transmission of influenza virus infection in mice. II. Some factors affecting the incidence of transmitted infection. *J. Exp. Med.* 118, 267–275.
- Shackelford, P.G., and Feigin, R.D. (1973). Periodicity of susceptibility to pneumococcal infection: influence of light and adrenocortical secretions. *Science* 182, 285–287.
- Shadrin, A.S., Marinich, I.G., and Taros Yu, .L. (1977). Experimental and epidemiological estimation of seasonal and climato geographical features of non specific resistance of the organism to influenza. *J. Hyg. Epidemiol. Microbiol. Immunol.* 21, 155–161.
- Shah, A.P., Smolensky, M.H., Burau, K.D., Cech, I.M., and Lai, D. (2006). Seasonality of primarily childhood and young adult infectious diseases in the United States. *Chronobiol. Int.* 23, 1065–1082.
- Shaman, J., Jeon, C.Y., Giovannucci, E., and Lipsitch, M. (2011). Shortcomings of vitamin D-based model simulations of seasonal influenza. *PLoS One* 6, 1–7.
- Steninger, M., Karpova, D., Unterrainer, C., Harenkamp, S., Wiercinska, E., Hoerster, K., Pfeffer, M., Maronde, E., and Bonig, H. (2019). Hematopoietic-extrinsic cues dictate circadian redistribution of mature and immature hematopoietic cells in blood and spleen. *Cells* 8, 1–22.
- Stevenson, T.J., and Prendergast, B.J. (2015). Photoperiodic time measurement and seasonal immunological plasticity. *Front. Neuroendocrinol.* 37, 76–88.
- Sudlow, C., Gallacher, J., Allen, N., Beral, V., Burton, P., Danesh, J., Downey, P., Elliott, P., Green, J., Landray, M., et al. (2015). UK biobank: an open access resource for identifying the causes of a wide range of complex diseases of middle and old age. *PLoS Med.* 12, e1001779.
- Sung, K.C. (2006). Seasonal variation of C-reactive protein in apparently healthy Koreans. *Int. J. Cardiol.* 107, 338–342.
- Suzuki, K., Hayano, Y., Nakai, A., Furuta, F., and Noda, M. (2016). Adrenergic control of the adaptive immune response by diurnal lymphocyte recirculation through lymph nodes. *J. Exp. Med.* 213, 2567–2574.
- Takahashi, J.S. (2017). Transcriptional architecture of the mammalian circadian clock. *Nat. Rev. Genet.* 18, 164–179.
- Tamerius, J., Nelson, M.I., Zhou, S.Z., Viboud, C., Miller, M.A., and Alonson, W.J. (2011). Global influenza seasonality: reconciling patterns across temperate and tropical regions. *Environ. Health Perspect.* 119, 439–445, <https://doi.org/10.1289/ehp.1002383>.
- ter Horst, R., Jaeger, M., Smeekens, S.P., Oosting, M., Swertz, M.A., Li, Y., Kumar, V., Diavatopoulos, D.A., Jansen, A.F.M., Lemmers, H., et al. (2016). Host and environmental factors influencing individual human cytokine responses. *Cell* 167, 1111–1124.e13.
- Tognini, P., Thaiss, C.A., Elinav, E., and Sassone-Corsi, P. (2017). Circadian coordination of antimicrobial responses. *Cell Host Microbe* 22, 185–192.
- Van Rood, Y., Goulmy, E., Blokland, E., Pool, J., Van Rood, J., and Van Houwelingen, H. (1991). Month-related variability in immunological test results; implications for immunological follow-up studies. *Clin. Exp. Immunol.* 86, 349–354.
- Wongwiwat, M., Sukapanit, S., Triyanond, C., and Sawyer, W.D. (1972). Circadian rhythm of the resistance of mice to acute pneumococcal infection. *Infect. Immun.* 5, 442–448.
- Wood, S., and Loudon, A. (2014). Clocks for all seasons: unwinding the roles and mechanisms of circadian and interval timers in the hypothalamus and pituitary. *J. Endocrinol.* 222, R39–R59.
- Xu, C., Fang, V.J., Perera, R.A.P.M., Kam, A.M.S., Ng, S., Chan, Y.H., Chan, K.H., Ip, D.K.M., Peiris, J.S.M., and Cowling, B.J. (2016). Serum 25-hydroxyvitamin D was not associated with influenza virus infection in children and adults in Hong Kong, 2009–2010. *J. Nutr.* 146, 2506–2512.
- Yellon, S.M., and Tran, L.T. (2002). Photoperiod, reproduction, and immunity in select strains of inbred mice. *J. Biol. Rhythm* 17, 65–75.
- Zykov, M.P., and Sosunov, A.V. (1987). Vaccination activity of live influenza vaccine in different seasons of the year. *J. Hyg. Epidemiol. Microbiol. Immunol.* 31, 453–459.

**iScience, Volume 24**

**Supplemental information**

**Seasonal and daytime variation in multiple  
immune parameters in humans: Evidence from 329,261  
participants of the UK Biobank cohort**

**Cathy Wyse, Grace O'Malley, Andrew N. Coogan, Sam McConkey, and Daniel J. Smith**

Table S1 Summary descriptive data for CRP, total white blood cells, monocytes, lymphocytes and neutrophils by month, mean [95% CI]. Related to Figure 1

Month	Jan (N=24283)	Feb (N=26731)	Mar (N=32217)	Apr (N=28549)	May (N=33871)	Jun (N=33883)	Jul (N=27606)	Aug (N=24426)	Sep (N=23422)	Oct (N=27710)	Nov (N=26992)	Dec (N=18161)
<b>CRP</b>												
mg/litre	2.337 (± 3.994)	2.279 (± 3.946)	2.264 (± 3.797)	2.266 (± 3.822)	2.232 (± 3.737)	2.133 (± 3.488)	2.152 (± 3.519)	2.122 (± 3.474)	2.101 (± 3.343)	2.260 (± 3.635)	2.257 (± 3.841)	2.392 (± 4.184)
<b>WBC</b>												
(10 <sup>9</sup> /litre)	6.792 (± 1.692)	6.769 (± 1.690)	6.831 (± 1.683)	6.772 (± 1.651)	6.708 (± 1.651)	6.690 (± 1.632)	6.630 (± 1.640)	6.627 (± 1.628)	6.650 (± 1.622)	6.699 (± 1.627)	6.775 (± 1.678)	6.725 (± 1.659)
<b>Neutrophils</b>												
(10 <sup>9</sup> /litre)	4.203 (± 1.324)	4.156 (± 1.315)	4.160 (± 1.294)	4.118 (± 1.275)	4.075 (± 1.279)	4.048 (± 1.254)	4.021 (± 1.267)	4.039 (± 1.254)	4.065 (± 1.255)	4.119 (± 1.279)	4.172 (± 1.305)	4.149 (± 1.300)
<b>Monocytes</b>												
(10 <sup>9</sup> /litre)	0.4514 (± 0.1679)	0.4499 (± 0.1655)	0.4668 (± 0.1592)	0.4665 (± 0.1568)	0.4626 (± 0.1562)	0.4675 (± 0.1554)	0.4644 (± 0.1535)	0.4524 (± 0.1540)	0.4624 (± 0.1524)	0.4633 (± 0.1544)	0.4621 (± 0.1594)	0.4586 (± 0.1669)
<b>Lymphocytes</b>												
(10 <sup>9</sup> /litre)	1.915 (± 0.5728)	1.946 (± 0.5948)	1.984 (± 0.6039)	1.966 (± 0.5856)	1.956 (± 0.5887)	1.950 (± 0.5764)	1.922 (± 0.5685)	1.918 (± 0.5732)	1.908 (± 0.5657)	1.903 (± 0.5618)	1.922 (± 0.5731)	1.906 (± 0.5822)

Table S2 Summary descriptive data for CRP, total white blood cells, monocytes, lymphocytes and neutrophils by time of day (hour), mean [95% CI]. Related to Figure 2

Time of Day	9 (N=20091)	10 (N=28642)	11 (N=34487)	12 (N=31376)	13 (N=31656)	14 (N=34257)	15 (N=32769)	16 (N=34022)	17 (N=31456)	18 (N=31409)	19 (N=13616)
<b>CRP</b>											
mg/litre	2.123 (± 3.538)	2.173 (± 3.597)	2.257 (± 3.838)	2.259 (± 3.614)	2.312 (± 3.853)	2.273 (± 3.826)	2.260 (± 3.725)	2.245 (± 3.772)	2.201 (± 3.778)	2.172 (± 3.700)	2.140 (± 3.535)
<b>WBC</b>											
(10 <sup>9</sup> /litre)	5.926 (± 1.514)	6.046 (± 1.546)	6.243 (± 1.564)	6.466 (± 1.579)	6.713 (± 1.616)	6.876 (± 1.622)	6.991 (± 1.608)	7.078 (± 1.635)	7.130 (± 1.616)	7.195 (± 1.624)	7.254 (± 1.639)
<b>Neutrophils</b>											
(10 <sup>9</sup> /litre)	3.640 (± 1.210)	3.717 (± 1.238)	3.829 (± 1.241)	3.978 (± 1.253)	4.159 (± 1.283)	4.271 (± 1.279)	4.340 (± 1.279)	4.339 (± 1.282)	4.295 (± 1.258)	4.256 (± 1.258)	4.231 (± 1.269)
<b>Monocytes</b>											
(10 <sup>9</sup> /litre)	0.4217 (± 0.1494)	0.4280 (± 0.1516)	0.4359 (± 0.1538)	0.4401 (± 0.1534)	0.4429 (± 0.1543)	0.4523 (± 0.1547)	0.4682 (± 0.1568)	0.4858 (± 0.1601)	0.4946 (± 0.1578)	0.5026 (± 0.1591)	0.5139 (± 0.1609)
<b>Lymphocytes</b>											
(10 <sup>9</sup> /litre)	1.654 (± 0.4742)	1.697 (± 0.4916)	1.776 (± 0.5167)	1.841 (± 0.5351)	1.904 (± 0.5525)	1.934 (± 0.5659)	1.963 (± 0.5606)	2.027 (± 0.5777)	2.107 (± 0.5852)	2.195 (± 0.6023)	2.263 (± 0.6292)

Table S3 Summary descriptive data for CRP, total white blood cells, monocytes, lymphocytes and neutrophils by month, median (interquartile range). Related to Figure 1

Month	Jan (N=24283)	Feb (N=26731)	Mar (N=32217)	Apr (N=28549)	May (N=33871)	Jun (N=33883)	Jul (N=27606)	Aug (N=24426)	Sep (N=23422)	Oct (N=27710)	Nov (N=26992)	Dec (N=18161)
<b>CRP</b>												
mg/litre	2.337 (3.994)	2.279 (3.946)	2.264 (3.797)	2.266 (3.822)	2.232 (3.737)	2.133 (3.488)	2.152 (3.519)	2.122 (3.474)	2.101 (3.343)	2.260 (3.635)	2.257 (3.841)	2.392 (4.184)
<b>WBC</b>												
(10 <sup>9</sup> /litre)	6.792 (1.692)	6.769 (1.690)	6.831 (1.683)	6.772 (1.651)	6.708 (1.651)	6.690 (1.632)	6.630 (1.640)	6.627 (1.628)	6.650 (1.622)	6.699 (1.627)	6.775 (1.678)	6.725 (1.659)
<b>Neutrophils</b>												
(10 <sup>9</sup> /litre)	4.203 (1.324)	4.156 (1.315)	4.160 (1.294)	4.118 (1.275)	4.075 (1.279)	4.048 (1.254)	4.021 (1.267)	4.039 (1.254)	4.065 (1.255)	4.119 (1.279)	4.172 (1.305)	4.149 (1.300)
<b>Monocytes</b>												
(10 <sup>9</sup> /litre)	0.4514 (0.1679)	0.4499 (0.1655)	0.4668 (0.1592)	0.4665 (0.1568)	0.4626 (0.1562)	0.4675 (0.1554)	0.4644 (0.1535)	0.4524 (0.1540)	0.4624 (0.1524)	0.4633 (0.1544)	0.4621 (0.1594)	0.4586 (0.1669)
<b>Lymphocytes</b>												
(10 <sup>9</sup> /litre)	1.915 (0.5728)	1.946 (0.5948)	1.984 (0.6039)	1.966 (0.5856)	1.956 (0.5887)	1.950 (0.5764)	1.922 (0.5685)	1.918 (0.5732)	1.908 (0.5657)	1.903 (0.5618)	1.922 (0.5731)	1.906 (0.5822)

Table S4 Summary descriptive data for CRP, total white blood cells, monocytes, lymphocytes and neutrophils by time of day (hour), median (interquartile range) Related to Figure 2

Time of Day	9 (N=20091)	10 (N=28642)	11 (N=34487)	12 (N=31376)	13 (N=31656)	14 (N=34257)	15 (N=32769)	16 (N=34022)	17 (N=31456)	18 (N=31409)	19 (N=13616)
<b>CRP</b>											
mg/litre	2.123 (3.538)	2.173 (3.597)	2.257 (3.838)	2.259 (3.614)	2.312 (3.853)	2.273 (3.826)	2.260 (3.725)	2.245 (3.772)	2.201 (3.778)	2.172 (3.700)	2.140 (3.535)
<b>WBC</b>											
(10 <sup>9</sup> /litre)	5.926 (1.514)	6.046 (1.546)	6.243 (1.564)	6.466 (1.579)	6.713 (1.616)	6.876 (1.622)	6.991 (1.608)	7.078 (1.635)	7.130 (1.616)	7.195 (1.624)	7.254 (1.639)
<b>Neutrophils</b>											
(10 <sup>9</sup> /litre)	3.640 (1.210)	3.717 (1.238)	3.829 (1.241)	3.978 (1.253)	4.159 (1.283)	4.271 (1.279)	4.340 (1.279)	4.339 (1.282)	4.295 (1.258)	4.256 (1.258)	4.231 (1.269)
<b>Monocytes</b>											
(10 <sup>9</sup> /litre)	0.4217 (0.149)	0.4280 (0.1516)	0.4359 (0.153)	0.4401 (0.1534)	0.4429 (0.1543)	0.4523 (0.1547)	0.4682 (0.1568)	0.4858 (0.1601)	0.4946 (0.1578)	0.5026 (0.1591)	0.5139 (0.1609)
<b>Lymphocytes</b>											
(10 <sup>9</sup> /litre)	1.654 (0.474)	1.697 (0.491)	1.776 (0.516)	1.841 (0.535)	1.904 (0.5525)	1.934 (0.5659)	1.963 (0.5606)	2.027 (0.5777)	2.107 (0.5852)	2.195 (0.6023)	2.263 (0.6292)

Table S5 Diagnostic tests for multicollinearity between predictor variables included in the multiple regression models. Related to Table 2.

Variable	VIF	SQRT VIF	Tolerance	R Squared
Daylength	2.76	1.66	0.3619	0.6381
Sex	1.09	1.04	0.9169	0.0831
Age	1.04	1.02	0.966	0.034
Ethnicity	1.08	1.04	0.922	0.078
Deprivation Index	1.09	1.04	0.9214	0.0786
BMI	1.1	1.05	0.912	0.088
Physical Activity	1.05	1.03	0.9514	0.0486
Sedentary Behaviour	1.09	1.04	0.9163	0.0837
Sleep Duration	1.02	1.01	0.9851	0.0149
Chronotype	1.03	1.01	0.9742	0.0258
Shiftwork	1.05	1.03	0.9489	0.0511
Smoking	1.04	1.02	0.9639	0.0361
Alcohol Frequency	1.04	1.02	0.9657	0.0343
Vitamin D	1.24	1.11	0.8069	0.1931
Outdoor Temperature	3.00	1.73	0.3323	0.6677
Time of Day	1.01	1	0.9932	0.0068
Blood Count Device	1	1	0.9972	0.0028
Assessment Centre	1.04	1.02	0.961	0.039
<b>Mean VIF</b>	<b>1.26</b>			

Table S6 Multiple regression analysis of associations between time of day and lymphocyte, neutrophil, monocyte counts and CRP. Related to Table 4 and Table 5.

Lymphocytes	Model 1	Model 2	Model 3
Segment 1	0.051*** [0.050,0.052]	0.050*** [0.049,0.051]	0.050*** [0.049,0.051]
Segment 2	0.082*** [0.079,0.085]	0.085*** [0.082,0.088]	0.085*** [0.082,0.088]
Sex	-0.120*** [-0.124,-0.116]	-0.155*** [-0.159,-0.150]	-0.154*** [-0.158,-0.150]
Age	0.002*** [0.001,0.002]	0.002*** [0.002,0.002]	0.002*** [0.002,0.002]
Ethnicity	0.047*** [0.044,0.049]	0.047*** [0.044,0.051]	0.044*** [0.041,0.047]
Social Deprivation	0.007*** [0.006,0.007]	0.000 [-0.000,0.001]	-0.000 [-0.001,0.000]
BMI		0.022*** [0.021,0.022]	0.022*** [0.021,0.022]
Physical Activity		-0.000*** [-0.000,-0.000]	-0.000*** [-0.000,-0.000]
Sedentary Behaviour		0.009*** [0.008,0.010]	0.008*** [0.007,0.009]
Sleep Duration		0.006*** [0.004,0.008]	0.006*** [0.004,0.008]
Chronotype		-0.015*** [-0.017,-0.012]	-0.015*** [-0.017,-0.012]
Smoking		0.346*** [0.339,0.353]	0.349*** [0.342,0.355]
Alcohol		0.007*** [0.005,0.008]	0.007*** [0.005,0.008]
Daylength			0.000 [-0.000,0.001]
Blood Analyser			-0.005*** [-0.006,-0.003]
Assessment Centre			0.004*** [0.004,0.005]
Observations	309220	275086	272393
R <sup>2</sup>	0.099	0.159	0.161

95% confidence intervals in brackets

\*  $p < 0.05$ , \*\*  $p < 0.01$ , \*\*\*  $p < 0.001$

<b>Neutrophils</b>	<b>Model 1</b>		<b>Model 2</b>		<b>Model 3</b>	
Segment 1	0.130***	[0.127,0.133]	0.126***	[0.123,0.128]	0.125***	[0.122,0.128]
Segment 2	-0.039***	[-0.043,-0.034]	-0.035***	[-0.040,-0.030]	-0.034***	[-0.039,-0.030]
Sex	0.037***	[0.028,0.046]	-0.021***	[-0.031,-0.011]	-0.022***	[-0.031,-0.012]
Age	-0.003***	[-0.003,-0.002]	-0.002***	[-0.003,-0.001]	-0.002***	[-0.003,-0.001]
Ethnicity	-0.153***	[-0.160,-0.147]	-0.146***	[-0.153,-0.139]	-0.151***	[-0.158,-0.144]
Social Deprivation	0.021***	[0.020,0.023]	0.008***	[0.007,0.010]	0.007***	[0.006,0.009]
BMI			0.035***	[0.034,0.036]	0.035***	[0.034,0.036]
Physical Activity			-0.000***	[-0.000,-0.000]	-0.000***	[-0.000,-0.000]
Sedentary Behaviour			0.017***	[0.015,0.019]	0.016***	[0.014,0.018]
Sleep Duration			0.025***	[0.020,0.030]	0.025***	[0.020,0.030]
Chronotype			-0.023***	[-0.028,-0.018]	-0.023***	[-0.028,-0.018]
Smoking			0.794***	[0.779,0.810]	0.797***	[0.781,0.813]
Alcohol			0.023***	[0.020,0.027]	0.023***	[0.020,0.027]
Daylength					-0.016***	[-0.018,-0.015]
Blood Analyser					0.003	[-0.001,0.007]
Assessment Centre					0.007***	[0.006,0.008]
Observations	308545		274483		271793	
R <sup>2</sup>	0.042		0.091		0.093	

95% confidence intervals in brackets

\*  $p < 0.05$ , \*\*  $p < 0.01$ , \*\*\*  $p < 0.001$

<b>CRP</b>	<b>Model 1</b>		<b>Model 2</b>		<b>Model 3</b>	
Segment 1	0.013*	[0.000,0.026]	0.004	[-0.009,0.017]	0.005	[-0.008,0.019]
Segment 2	-0.010*	[-0.019,-0.002]	-0.013**	[-0.022,-0.005]	-0.013**	[-0.021,-0.004]
Sex	-0.177***	[-0.204,-0.150]	-0.402***	[-0.431,-0.374]	-0.405***	[-0.434,-0.377]
Age	0.028***	[0.026,0.029]	0.025***	[0.024,0.027]	0.026***	[0.024,0.027]
Ethnicity	0.019	[-0.002,0.039]	0.003	[-0.018,0.024]	0.006	[-0.016,0.027]
Social Deprivation	0.054***	[0.050,0.059]	0.027***	[0.022,0.032]	0.029***	[0.024,0.034]
BMI			0.210***	[0.207,0.214]	0.210***	[0.207,0.214]
Physical Activity			-0.001***	[-0.001,-0.001]	-0.001***	[-0.001,-0.001]
Sedentary Behaviour			0.020***	[0.014,0.026]	0.020***	[0.014,0.027]
Sleep Duration			0.045***	[0.031,0.059]	0.045***	[0.031,0.059]
Chronotype			0.029***	[0.014,0.044]	0.029***	[0.014,0.044]
Smoking			0.718***	[0.671,0.765]	0.720***	[0.673,0.767]
Alcohol			0.014**	[0.004,0.024]	0.014**	[0.004,0.024]
Daylength					-0.017***	[-0.022,-0.013]
Assessment Centre					-0.005***	[-0.008,-0.003]
Observations	304000		270518		267872	
R <sup>2</sup>	0.006		0.069		0.069	

95% confidence intervals in brackets

\*  $p < 0.05$ , \*\*  $p < 0.01$ , \*\*\*  $p < 0.001$

<b>Monocytes</b>	<b>Model 1</b>		<b>Model 2</b>		<b>Model 3</b>	
Segment 1	0.005***	[0.005,0.006]	0.005***	[0.004,0.005]	0.005***	[0.004,0.005]
Segment 2	0.012***	[0.012,0.012]	0.012***	[0.012,0.013]	0.012***	[0.012,0.013]
Sex	0.070***	[0.069,0.071]	0.063***	[0.062,0.064]	0.063***	[0.062,0.064]
Age	0.002***	[0.002,0.002]	0.002***	[0.002,0.002]	0.002***	[0.002,0.002]
Ethnicity	-0.015***	[-0.016,-0.014]	-0.014***	[-0.015,-0.014]	-0.015***	[-0.016,-0.014]
Social Deprivation	0.001***	[0.001,0.001]	0.000**	[0.000,0.001]	0.000	[-0.000,0.000]
BMI			0.004***	[0.004,0.004]	0.004***	[0.004,0.004]
Physical Activity			-0.000***	[-0.000,-0.000]	-0.000***	[-0.000,-0.000]
Sedentary Behaviour			0.001***	[0.001,0.001]	0.001***	[0.001,0.001]
Sleep Duration			0.002***	[0.001,0.003]	0.002***	[0.002,0.003]
Chronotype			-0.000	[-0.001,0.000]	-0.000	[-0.001,0.000]
Smoking			0.050***	[0.048,0.052]	0.050***	[0.048,0.052]
Alcohol			-0.001***	[-0.002,-0.001]	-0.001***	[-0.002,-0.001]
Daylength					0.001***	[0.000,0.001]
Blood Analyser					-0.009***	[-0.010,-0.009]
Assessment Centre					0.001***	[0.001,0.001]
Observations	309207		275076		272380	
R <sup>2</sup>	0.091		0.115		0.120	

95% confidence intervals in brackets

\*  $p < 0.05$ , \*\*  $p < 0.01$ , \*\*\*  $p < 0.001$

Table S7 Multiple regression analysis of associations between daylength and lymphocyte, neutrophil, monocyte counts and CRP. Related to Table 2

<b>Neutrophils</b>	<b>Model 1</b>		<b>Model 2</b>		<b>Model 3</b>	
Daylength	0.002***	[0.002,0.003]	0.003***	[0.002,0.004]	0.004***	[0.003,0.005]
Sex	-0.116***	[-0.120,-0.111]	-0.131***	[-0.136,-0.125]	-0.129***	[-0.134,-0.124]
Age	0.001***	[0.001,0.002]	0.002***	[0.001,0.002]	0.001***	[0.001,0.001]
Ethnicity	0.056***	[0.053,0.059]	0.057***	[0.053,0.060]	0.043***	[0.039,0.047]
Social Deprivation	0.007***	[0.006,0.008]	0.001*	[0.000,0.002]	-0.001	[-0.002,0.000]
BMI			0.022***	[0.021,0.023]	0.021***	[0.020,0.022]
Physical Activity			-0.000***	[-0.000,-0.000]	-0.000***	[-0.000,-0.000]
Sedentary Behaviour			0.007***	[0.006,0.009]	0.007***	[0.006,0.008]
Sleep Duration			0.004**	[0.001,0.007]	0.004*	[0.001,0.006]
Chronotype			-0.007***	[-0.009,-0.004]	-0.014***	[-0.017,-0.011]
Shiftwork			-0.018***	[-0.025,-0.011]	-0.009*	[-0.016,-0.002]
Smoking			0.334***	[0.325,0.342]	0.329***	[0.321,0.338]
Alcohol			0.009***	[0.007,0.011]	0.008***	[0.006,0.010]
Vitamin D					-0.000***	[-0.000,-0.000]
Outdoor Temperature					-0.002***	[-0.003,-0.001]
Time of Day					0.057***	[0.057,0.058]
Blood Analyser					-0.005***	[-0.008,-0.003]
Assessment Centre					0.004***	[0.004,0.005]
Observations	308924		175498		161528	
R <sup>2</sup>	0.016		0.074		0.175	

95% confidence intervals in brackets

\*  $p < 0.05$ , \*\*  $p < 0.01$ , \*\*\*  $p < 0.001$

<b>Lymphocytes</b>	<b>Model 1</b>		<b>Model 2</b>		<b>Model 3</b>	
Daylength	-0.015***	[-0.017,-0.014]	-0.013***	[-0.015,-0.011]	-0.014***	[-0.018,-0.011]
Sex	0.027***	[0.018,0.036]	-0.121***	[-0.133,-0.108]	-0.127***	[-0.140,-0.115]
Age	-0.000	[-0.001,0.000]	-0.006***	[-0.007,-0.005]	-0.007***	[-0.008,-0.006]
Ethnicity	-0.141***	[-0.148,-0.134]	-0.139***	[-0.148,-0.131]	-0.168***	[-0.178,-0.159]
Social Deprivation	0.023***	[0.021,0.024]	0.005***	[0.003,0.007]	0.001	[-0.001,0.003]
BMI			0.039***	[0.038,0.040]	0.037***	[0.035,0.038]
Physical Activity			-0.000	[-0.000,0.000]	-0.000	[-0.000,0.000]
Sedentary Behaviour			0.018***	[0.015,0.020]	0.018***	[0.015,0.021]
Sleep Duration			0.023***	[0.017,0.029]	0.023***	[0.017,0.030]
Chronotype			-0.008*	[-0.015,-0.002]	-0.018***	[-0.025,-0.012]
Shiftwork			0.015	[-0.001,0.032]	0.031***	[0.015,0.048]
Smoking			0.773***	[0.754,0.793]	0.760***	[0.740,0.780]
Alcohol			0.020***	[0.015,0.024]	0.018***	[0.014,0.023]
Vitamin D					-0.003***	[-0.003,-0.002]
Outdoor Temperature					0.002	[-0.000,0.004]
Time of Day					0.070***	[0.068,0.072]
Blood Analyser					0.000	[-0.005,0.006]
Assessment Centre					0.007***	[0.006,0.008]
Observations	308250		175049		161112	
R <sup>2</sup>	0.008		0.062		0.093	

95% confidence intervals in brackets

\*  $p < 0.05$ , \*\*  $p < 0.01$ , \*\*\*  $p < 0.001$

<b>Monocytes</b>	<b>Model 1</b>		<b>Model 2</b>		<b>Model 3</b>	
Daylength	0.001***	[0.001,0.001]	0.001***	[0.001,0.001]	-0.000*	[-0.001,-0.000]
Sex	0.071***	[0.070,0.072]	0.058***	[0.056,0.059]	0.057***	[0.056,0.059]
Age	0.002***	[0.002,0.002]	0.001***	[0.001,0.001]	0.001***	[0.001,0.001]
Ethnicity	-0.013***	[-0.014,-0.013]	-0.013***	[-0.014,-0.012]	-0.016***	[-0.017,-0.015]
Social Deprivation	0.001***	[0.001,0.002]	0.000	[-0.000,0.000]	-0.000	[-0.000,0.000]
BMI			0.004***	[0.004,0.004]	0.004***	[0.004,0.004]
Physical Activity			-0.000	[-0.000,0.000]	-0.000	[-0.000,0.000]
Sedentary Behaviour			0.001***	[0.001,0.002]	0.001***	[0.001,0.002]
Sleep Duration			0.002***	[0.001,0.003]	0.002***	[0.001,0.002]
Chronotype			0.001*	[0.000,0.002]	-0.001	[-0.001,0.000]
Shiftwork			0.000	[-0.002,0.002]	0.001	[-0.000,0.003]
Smoking			0.048***	[0.046,0.050]	0.048***	[0.045,0.050]
Alcohol			-0.001***	[-0.002,-0.001]	-0.002***	[-0.002,-0.001]
Vitamin D					-0.000***	[-0.000,-0.000]
Outdoor Temperature					0.001***	[0.001,0.001]
Time of Day					0.010***	[0.009,0.010]
Blood Analyser					-0.010***	[-0.010,-0.009]
Assessment Centre					0.001***	[0.001,0.001]
Observations	308911		175441		161471	
R <sup>2</sup>	0.060		0.075		0.119	

95% confidence intervals in brackets

\*  $p < 0.05$ , \*\*  $p < 0.01$ , \*\*\*  $p < 0.001$

<b>CRP</b>	<b>Model 1</b>		<b>Model 2</b>		<b>Model 3</b>	
Daylength	-0.006***	[-0.007,-0.004]	-0.005***	[-0.006,-0.003]	-0.004***	[-0.007,-0.002]
Sex	-0.042***	[-0.050,-0.035]	-0.153***	[-0.162,-0.144]	-0.085***	[-0.108,-0.063]
Age	0.016***	[0.016,0.017]	0.012***	[0.012,0.013]	0.012***	[0.011,0.013]
Ethnicity	0.005	[-0.000,0.011]	-0.001	[-0.007,0.006]	-0.003	[-0.010,0.004]
Social Deprivation	0.020***	[0.019,0.022]	0.005***	[0.003,0.006]	0.005***	[0.003,0.007]
BMI			0.104***	[0.103,0.105]	0.100***	[0.097,0.102]
Physical Activity			-0.000***	[-0.000,-0.000]	-0.000***	[-0.000,-0.000]
Sedentary Behaviour			0.014***	[0.012,0.016]	0.014***	[0.012,0.016]
Sleep Duration			0.017***	[0.012,0.022]	0.016***	[0.011,0.021]
Chronotype			0.015***	[0.010,0.020]	0.016***	[0.011,0.021]
Shiftwork			0.050***	[0.038,0.062]	0.048***	[0.036,0.061]
Smoking			0.282***	[0.268,0.297]	0.281***	[0.267,0.296]
Alcohol			0.012***	[0.009,0.015]	0.012***	[0.009,0.016]
Vitamin D					-0.002**	[-0.004,-0.001]
Sex					0.000	[0.000,0.000]
Vitamin D # Sex					-0.001***	[-0.002,-0.001]
Vitamin D					0.000	[0.000,0.000]
BMI					0.000	[0.000,0.000]
Vitamin D # BMI					0.000***	[0.000,0.000]
Outdoor Temperature					-0.000	[-0.002,0.002]
Time of Day					-0.004***	[-0.005,-0.002]
Blood Analyser					-0.000	[-0.004,0.004]
Assessment Centre					-0.001**	[-0.002,-0.000]
Observations	303707		172476		161492	
R <sup>2</sup>	0.019		0.223		0.222	

95% confidence intervals in brackets

\*  $p < 0.05$ , \*\*  $p < 0.01$ , \*\*\*  $p < 0.001$

<b>WBC</b>	<b>Model 1</b>		<b>Model 2</b>		<b>Model 3</b>	
Daylength	-0.011***	[-0.013,-0.009]	-0.008***	[-0.011,-0.006]	-0.011***	[-0.015,-0.007]
Sex	0.005	[-0.007,0.016]	-0.181***	[-0.197,-0.166]	-0.186***	[-0.202,-0.171]
Age	0.002***	[0.002,0.003]	-0.003***	[-0.004,-0.002]	-0.005***	[-0.006,-0.004]
Ethnicity	-0.098***	[-0.107,-0.090]	-0.094***	[-0.105,-0.083]	-0.142***	[-0.153,-0.130]
Social Deprivation	0.033***	[0.031,0.035]	0.006***	[0.004,0.009]	0.001	[-0.002,0.003]
BMI			0.068***	[0.066,0.069]	0.064***	[0.062,0.066]
Physical Activity			-0.000*	[-0.000,-0.000]	-0.000**	[-0.000,-0.000]
Sedentary Behaviour			0.027***	[0.023,0.030]	0.027***	[0.024,0.030]
Sleep Duration			0.031***	[0.023,0.039]	0.030***	[0.022,0.038]
Chronotype			-0.013**	[-0.022,-0.005]	-0.033***	[-0.041,-0.025]
Shiftwork			-0.000	[-0.021,0.020]	0.027*	[0.006,0.047]
Smoking			1.223***	[1.198,1.248]	1.205***	[1.180,1.230]
Alcohol			0.028***	[0.022,0.033]	0.025***	[0.019,0.031]
Vitamin D					-0.003***	[-0.004,-0.003]
Outdoor Temperature					0.001	[-0.001,0.004]
Time of Day					0.142***	[0.139,0.144]
Blood Analyser					-0.016***	[-0.023,-0.009]
Assessment Centre					0.012***	[0.011,0.014]
Observations	309516		175753		161746	
R <sup>2</sup>	0.005		0.089		0.163	

95% confidence intervals in brackets

\*  $p < 0.05$ , \*\*  $p < 0.01$ , \*\*\*  $p < 0.001$

Table S8 Comparison of demographic and lifestyle characteristics of participants that self-reported good health and were included in this study, and the remainder of the UK Biobank database, that self-reported poor health. Related to Methods section and Table 1

	<b>Healthy</b> (N=329261)	<b>Diseased</b> (N=159904)
<b>Age (years)</b>		
Mean (SD)	55 ( $\pm$ 8)	58 ( $\pm$ 8)
<b>Sex</b>		
Female	185438 (56 %)	80199 (50 %)
Male	143823 (44 %)	79705 (50 %)
<b>Social Deprivation Index</b>		
Mean (SD)	-1.544 ( $\pm$ 2.945)	-0.8489 ( $\pm$ 3.296)
<b>Ethnicity</b>		
White	311054 (94 %)	150657 (94 %)
Asian	6369 (2 %)	3022 (2 %)
Black	4902 (1 %)	2817 (2 %)
Chinese	1190 (0 %)	302 (0 %)
Mixed	1929 (1 %)	931 (1 %)
Other	2849 (1 %)	1491 (1 %)
<b>Physical Activity</b>		
Mean (SD)	45.89 ( $\pm$ 62.72)	40.63 ( $\pm$ 62.25)
<b>Sedentary Behaviour</b>		
Mean (SD)	4.90 ( $\pm$ 2.24)	5.31 ( $\pm$ 2.50)
<b>BMI</b>		
Mean (SD)	26.82 ( $\pm$ 4.34)	28.64 ( $\pm$ 5.39)
<b>Smoker</b>		
No	296633 (90 %)	139513 (87 %)
Yes	31606 (10 %)	19615 (12 %)
<b>Chronotype</b>		
Evening	23586 (7 %)	15512 (10 %)
More evening than morning	82570 (25 %)	40817 (26 %)
More morning than evening	107374 (33 %)	46633 (29 %)
Morning	79704 (24 %)	38064 (24 %)

Table S9 Comparison of demographic and lifestyle characteristics of participants that were randomly selected by UK Biobank for inclusion in the study of blood levels of antibodies against 20 infectious agents, and the remainder of the UK Biobank database. Related to Table 1

	Not Antigen Tested (N=493224)	Antigen Tested* (N=9431)
<b>Age (years)</b>		
Mean (SD)	56 (± 8)	56 (± 8)
<b>Sex</b>		
Female	268121 (54 %)	5277 (56 %)
Male	224972 (46 %)	4153 (44 %)
<b>Social Deprivation Index</b>		
Mean (SD)	-1.292 (± 3.096)	-1.368 (± 3.052)
<b>Ethnicity</b>		
White	463829 (94 %)	8883 (94 %)
Asian	9687 (2 %)	195 (2 %)
Black	7920 (2 %)	141 (1 %)
Chinese	1538 (0 %)	36 (0 %)
Mixed	2905 (1 %)	53 (1 %)
Other	4481 (1 %)	78 (1 %)
<b>Physical Activity</b>		
Mean (SD)	43.94 (± 62.65)	43.85 (± 58.91)
<b>Sedentary Behaviour</b>		
Mean (SD)	5.02 (± 2.35)	5.01 (± 2.28)
<b>BMI</b>		
Mean (SD)	27.43 (± 4.80)	27.33 (± 4.80)
<b>Smoker</b>		
No	438167 (89 %)	8429 (89 %)
Yes	52028 (11 %)	949 (10 %)
<b>Chronotype</b>		
Evening	39375 (8 %)	736 (8 %)
More evening than morning	123899 (25 %)	2435 (26 %)
More morning than evening	154370 (31 %)	3009 (32 %)
Morning	118100 (24 %)	2271 (24 %)

\*Participants randomly selected for inclusion in serum antigen titre study

## Transparent Methods

### *Study Sample*

The study sample were participants of UK Biobank, a general population cohort study that recruited over half a million UK residents continuously between 2006 and 2010, at 22 assessment centres located across the UK ([www.ukbiobank.co.uk](http://www.ukbiobank.co.uk)). Eligible participants who lived within travelling distance of one of 22 UK assessment centres were identified through national health service patient registers and invited to participate by mail, resulting in a 5.5% response rate (Sudlow et al., 2015). The participants were aged between 37-73 years at the time of enrolment. Inclusion in the present study was restricted to participants who reported having no chronic disease at the time of recruitment. Participants provided full informed consent to participate in UK Biobank. This study was covered by the generic ethical approval for UK Biobank studies from the NHS National Research Ethics Service

### *Participant Measures*

Participants were invited to attend the assessment centre at a pre-booked provisional appointment time between 8am-7pm; they did not self-select the time of day of attendance but were free to reschedule if required. Baseline information was collected at the assessment centre using a questionnaire and an interview, and blood samples and physical measurements were taken. Information on the demographic status of the participants included age at baseline, sex (male/female), ethnicity (White, Black, Mixed, Chinese, Asian, Other). (large-scale and resource, 2007) Participants were self-categorised as morning or evening chronotype using the question, “Do you consider yourself to be: Definitely a morning person; More a morning than evening person; More an evening than a morning person; Definitely an evening person; or Don’t know”. The self-reported level and duration of usual physical activity was used to derive total physical activity, measured as metabolic equivalents (MET.hours/week). A proxy of sedentary behaviour was derived from the total number of self-reported hours spent driving, using a computer and watching television each day. Smoking status was self-reported and categorized as “never smoker,” “current smoker” and “former smoker.” The frequency and volume of alcohol intake were self-reported. Body mass index (BMI) was measured by trained UK Biobank staff using standardized methods and instruments. Habitual sleep duration was self-reported in hours per 24h. (large-scale and resource, 2007)

Blood samples were collected at the end of the assessment centre visit, and the time was immediately recorded on computerised system by swiping the unique barcode on the collection tube. Blood cell counts were performed within 24 hours using an LH750 haematology analyser (Coulter, Beckman Coulter, Brea, CA, USA) to determine the total number of white blood cells, plus the numbers of neutrophils, lymphocytes and monocytes, which were expressed as a percentage of the total white blood cell count. Vitamin D was measured in serum using a chemiluminescent direct competitive immunoassay (LIAISON XL; Diasorin, Italy). Serum C reactive protein (CRP) was measured using a high-sensitivity immunoturbidimetric assay

performed on a Beckman Coulter clinical chemistry analyser (AU5800 Immuno-turbidimetric, Beckman Coulter (UK)).

A subset of participants (n=9724) were chosen at random for assessment of blood levels of antibodies against 20 infectious agents (see tables S7-8 for details of antigens). Antibody levels were measured using a Luminex high-throughput platform following validation against gold-standard assays and independent reference sera. Results of analyses were expressed as median fluorescence intensity for each antigen, and seropositivity status based on suggested thresholds. Full details of the participant measures and analysis procedures are available at [www.ukbiobank.co.uk](http://www.ukbiobank.co.uk).

### *Environmental Variables*

Latitude and longitude were derived from the postcode of residence at 1km<sup>2</sup> resolution using Open Source Geographic Information System software (QGIS Open Source Geospatial Foundation Project, <http://qgis.osgeo.org>). These data were combined with information on the date of attendance at the assessment center to derive the length of daylight on that day for each participant using vectorial algorithms in R-software [R version] in the ‘*insol*’ package (*insol*: Solar Radiation). Daylight was approximated over the hours (9am-7pm) of the diurnal dataset by calculating the mean zenith angle of the sun at each time and assessment centre location using the R-package “*geolight*”. (Lisovski and Hahn, 2012) The derived data were verified using information provided by the Global Monitoring Division of the US Government National Oceanic and Atmospheric Administration. Outdoor temperature was averaged for the 3 weeks preceding the date of attendance from data provided by the UK Meteorological Office for the weather station nearest to each assessment centre. Monthly data was aggregated into seasons for the purposes of descriptive analysis with seasons defined as: Spring = February, March, April; Summer = May, June, July; Autumn = August, September, October; Winter = November, December, January.

### *Data Analysis*

Seasonal and daily variation were assessed by plotting mean values of white blood cell and CRP values against month or hour of sample collection, fitting models to describe annual and daily variation, and then investigating whether any variation was independent of confounding factors and directly related to day length. The sampling distribution for CRP was positively-skewed, and a logarithmic transformation was applied before regression analysis, but original data are shown in the descriptive data summaries.

Seasonal patterns were analyzed by fitting a linear regression model for each outcome of interest that included a sine and a cosine term of transformations of the time variable, taken as month:

$$Y_i = M + \beta \text{Cos}(2\pi t_i / 12) + \gamma \text{Sin}(2\pi t_i / 12) \quad (1)$$

Where  $Y$  is  $t$  is time (months), and  $M$ ,  $\beta$  and  $\gamma$  were predicted by regression, above. The acrophase ( $\Phi$ ) and amplitude ( $A$ ) was predicted using equations 2 and 3, with  $M$  predicted from equation (1) above.

$$A = (\beta^2 + \gamma^2)^{1/2} \quad (2)$$

$$\Phi = \tan^{-1}(-\gamma/\beta) \quad (3)$$

The intercept ( $M$ ) was the mean level of the curve and thus an estimate of the annual mean of each outcome variable. The amplitude ( $A$ ) was the distance from the mean to the acrophase or the nadir, providing an estimate of the magnitude of seasonality. The acrophase ( $\Phi$ ) is the peak  $x$  axis value of the curve, whereas the nadir is the trough. Seasonality was indicated by statistical significance of the estimated cosinor (sine and cosine) regression coefficients.

Variation of the markers over the daily time course of sample collection was modeled using linear methods since the absence of nighttime samples precluded assumption of circadian patterns. Although assessment centre appointments started at 8am, the blood sample was collected at the end of the 40-minute assessment, so the 8am time point was excluded due to small sample numbers at this time ( $n=1843$ ). The relationship between time of day and the immune parameters was represented by a series of linear regression lines connected at breakpoints where the slope of each line changed. This analysis was implemented using the R package “segmented” to predict the times of breakpoints during the test period for each analyte (Muggeo, 2003). The statistical significance of the segmented regression model was assessed using the Davies test to test the null hypothesis that a breakpoint does not exist, and that the difference in slope parameter ( $\psi$ ) of the segmented relationship is zero. The breakpoints and slopes of each segment indicate peaks and troughs in WBC and CRP levels over time, as well as the rate and direction of any changes.

If seasonal and daily variation were indicated, we next investigated if these patterns were related to day length, and to time of day, and if any relationships were independent of lifestyle and environmental factors. The daytime data were modelled as a series of linear splines to account for the non-linear relationships between time of day and the immune parameters. Three multiple linear regression models were run that included an increasing number of covariates and progressively adjusted for sociodemographic, disease, lifestyle and environmental (temperature and day length) factors, with results reported as point estimates and 95% confidence intervals.

Potential confounders included as covariables were age; sex; ethnicity; Townsend area-deprivation score; physical activity and sedentary behaviour; alcohol intake and smoking status; outdoor temperature; blood analyser; vitamin D; sleep duration; chronotype and UK Biobank assessment centre. Multicollinearity between the covariables was assessed using variance inflation factors (VIF) and tolerance factors, with values of  $VIF > 10$  taken to denote problematic collinearity.

All analyses were performed using R version 3.5, and Stata 14 statistical software (StataCorp LP) and values of  $p < 0.01$  were considered to represent statistical significance.

## Supplemental References

- Lisovski, S., Hahn, S., 2012. GeoLight - processing and analysing light-based geolocator data in R. *Methods Ecol. Evol.* 3, 1055-1059
- Muggeo, V.M.R., 2003. Estimating regression models with unknown break-points. *Stat. Med.* 22, 3055–3071.
- Sudlow, C., Gallacher, J., Allen, N., Beral, V., Burton, P., Danesh, J., Downey, P., Elliott, P., Green, J., Landray, M., Liu, B., Matthews, P., Ong, G., Pell, J., Silman, A., Young, A., Sprosen, T., Peakman, T., Collins, R., 2015. UK biobank: an open access resource for identifying the causes of a wide range of complex diseases of middle and old age. *PLoS Med* 12, e1001779.