### University of Montana

### ScholarWorks at University of Montana

UM Graduate Student Research Conference (GradCon)

### 1D Crustal Velocity Model for West-Central Montana

Courtenay Jo Duzet University of Montana, Missoula, cd164433@umconnect.umt.edu

Michael C. Stickney Montana Tech

Follow this and additional works at: https://scholarworks.umt.edu/gsrc

### Let us know how access to this document benefits you.

Duzet, Courtenay Jo and Stickney, Michael C., "1D Crustal Velocity Model for West-Central Montana" (2021). *UM Graduate Student Research Conference (GradCon)*. 13. https://scholarworks.umt.edu/gsrc/2021/stem\_poster/13

This Poster Presentation is brought to you for free and open access by ScholarWorks at University of Montana. It has been accepted for inclusion in UM Graduate Student Research Conference (GradCon) by an authorized administrator of ScholarWorks at University of Montana. For more information, please contact scholarworks@mso.umt.edu.

# 1D Crustal Seismic-Velocity Model for West-Central Montana

Courtenay J. Duzet<sup>1</sup>, Hilary R. Martens<sup>1</sup>, Michael C. Stickney<sup>2</sup>


1. University of Montana, Department of Geosciences, Missoula, MT

2. Montana Technological University, Montana Bureau of Mines and Geology, Butte, MT



### . Introduction & Motivation

- While 90% of earthquakes occur on plate boundaries, Montana is one 47°N of the most seismically active states in the country, despite being located in the continental interior (Martens, 2017).
- Western Montana is particularly active as it lies within a zone of distributed crustal deformation, influenced by tectonic-plate motion occurring off the coasts of Washington and Oregon, the Yellowstone hotspot volcano, and the Rocky Mountains (McMahon, et al. 2019).
- On July 6th, 2017, Lincoln, MT had a M5.8 event, the largest to occur since the 1959 M7.9 Hebgen Lake quake.
- This event created a prolific & still ongoing aftershock sequence, which has given us the perfect opportunity to create the first crustal seismic velocity model for west-central Montana.
- This will advance earthquake science in Montana and lead us to a better understanding of crustal structure & stress conditions.




# 2. University of Montana Seismic Network (UMSN)

- UM stations use digital, 3 component, broadband seismometers.
  - Components: Vertical, N-S, E-W
- A component from one of the UM stations can be seen in the figure below.
- This is a significant upgrade in technology compared to the short period, analog seismic stations used to create past velocity models, which only recorded one component data (vertical).



- Seismic stations are strategically placed around the epicenter of the 2017 earthquake to collect the best quality data (see figure above).
- Blue markers represent seismic stations, while the orange marker is the epicenter of the Lincoln event.
- Seismic data used to constrain our crustal seismic-velocity model are from the UMSN and the Montana Regional Seismic Network (MRSN).



## 3. Methods & VELEST


- We manually collect seismic data from UMSN and supplement with telemetered data from the MRSN.
  - Mainshock & aftershock data recorded over a 3 year period (2017-2020).
  - We select only the best seismic data to constrain the velocity model. Data are filtered based on azimuthal gap, epicentral distance, hypocenter depth, and number of recording stations.
- Once this has been done, we use the software program, VELEST, which simultaneously inverts the seismic data and parameters from several hundred well recorded earthquakes to produce a 1D velocity model (Kissling et al. 1994).
  - In order obtain effective results with VELEST, the following data inputs are required:
    - Seismic station coordinates & elevations.
    - Earthquake hypocenters & arrival times.
  - A reference velocity model to initiate the inversion process.




Seismic station at Bandy Ranch in Ovando, MT

# 4. The Crustal Seismic-Velocity Model

- The crustal seismic velocity model describes seismic-velocity as a function of depth.
- An integral tool for understanding the structure of the crust and for locating the origin of an earthquake.
- The last model for Western Montana was derived in 2003 and is comprised of 3 layers (Zeiler et. al,
  - However, based on the seismic data used to create it, it is most appropriate for Southwestern Montana.
- As we explore the model space, we have a handful of models that could potentially represent the seismic nature of the area.
- Our current models describe the west-central region's seismic velocities with 8 layers for an area of roughly 5,000 km<sup>2</sup>.
  - Greater detail for shallow depths of the upper crust





# 5. Summary & Outlook

- Western Montana is prone to infrequent, high magnitude earthquakes, providing rare opportunities to gather large quantities of high-quality seismic data in a particular region.
- The 2017 Lincoln event has provided a prime opportunity to collect quality seismic data that will allow us to create a much-needed crustal velocity model for this seismically active region of Montana.
- With the combination of a prolific aftershock sequence and upgraded seismometers, we are able to collect well constrained, high quality seismic data that can be used to create a velocity model that is able to describe crustal stress conditions & structure with greater accuracy.
- Not only will developing a new, regional crustal velocity model advance earthquake science in Montana, but this will also be the first model derived specifically for the west-central region of the state, as the current velocity model is most appropriate for southwestern Montana.

## Looking towards the future:

- Implement our modeling process beyond our study region, allowing us to derive velocity models for other areas of western Montana that do not yet have a region specific velocity model.
- Explore the idea of combining these 1D models to create a psuedo 3D model for Western Montana as a whole.

### References

Kissling, E., U. Kradolfer, and H. Maurer. "Program VELEST user's guide-Short Introduction." *Institute of Geophysics, ETH Zurich* (1995).

Martens, H. "Intraplate Earthquakes in Montana." Department of Geosciences, University of Montana (2017). http://hs.umt.edu/martens/research.php

McMahon, N. D., Aster, R. C., Yeck, W. L., Benz, H. M., Stickney, M. C., & Martens, H. R. (2019). Spatiotemporal analysis of the foreshock–mainshock–aftershock sequence of the 6 July 2017 M w 5.8 Lincoln, Montana, earthquake. Seismological Research Letters, 90(1), 131–139.

Zeiler, C. P., Stickney, M. C., & Speece, M. A. (2005). Revised velocity structure of western Montana. Bulletin of the Seismological Society of America, 95(2), 759–762.