

Andrews University

Digital Commons @ Andrews University

Faculty Publications

12-14-2012

Naturally Occurring Carboxypeptidase A6 Mutations: Effect on Enzyme Function and Association with Epilepsy

Matthew R. Sapio

Albert Einstein College of Medicine of Yeshiva University

Annick Salzmänn

Université de Genève

Monique Vessaz

Hôpitaux universitaires de Genève

Arielle Crespel

CHU Montpellier

Peter J. Lyons

Andrews University

See next page for additional authors

Follow this and additional works at: <https://digitalcommons.andrews.edu/pubs>



Part of the [Nervous System Diseases Commons](#)

Recommended Citation

Sapio, Matthew R.; Salzmänn, Annick; Vessaz, Monique; Crespel, Arielle; Lyons, Peter J.; Malafosse, Alain; and Fricker, Lloyd D., "Naturally Occurring Carboxypeptidase A6 Mutations: Effect on Enzyme Function and Association with Epilepsy" (2012). *Faculty Publications*. 1714.

<https://digitalcommons.andrews.edu/pubs/1714>

This Article is brought to you for free and open access by Digital Commons @ Andrews University. It has been accepted for inclusion in Faculty Publications by an authorized administrator of Digital Commons @ Andrews University. For more information, please contact repository@andrews.edu.

Authors

Matthew R. Sapiro, Annick Salzmann, Monique Vessaz, Arielle Crespel, Peter J. Lyons, Alain Malafosse, and Lloyd D. Fricker

Naturally Occurring Carboxypeptidase A6 Mutations

EFFECT ON ENZYME FUNCTION AND ASSOCIATION WITH EPILEPSY*[§]

Received for publication, August 27, 2012, and in revised form, October 23, 2012. Published, JBC Papers in Press, October 26, 2012, DOI 10.1074/jbc.M112.414094

Matthew R. Sapio[‡], Annick Salzmänn[§], Monique Vessaz[¶], Arielle Crespel^{||}, Peter J. Lyons^{**}, Alain Malafosse^{§¶}, and Lloyd D. Fricker^{‡¶¶¶}

From the [‡]Department of Neuroscience and the ^{¶¶}Department of Molecular Pharmacology, Albert Einstein College of Medicine, Bronx, New York 10461, the [§]Department of Psychiatry, University of Geneva, 1205 Geneva, Switzerland, the [¶]Department of Genetic Medicine and Laboratory, University Hospitals of Geneva, 1205 Geneva, Switzerland, the ^{||}Department of Neurology, Montpellier University Hospital, France, and the ^{**}Department of Biology, Andrews University, Berrien Springs, Michigan 49104

Background: Two mutations in the carboxypeptidase A6 (*CPA6*) gene were previously found in epilepsy patients.

Results: Many additional *CPA6* mutations were found. Some inactivated *CPA6* and were more frequently found in epilepsy patients than controls.

Conclusion: Several *CPA6* mutations greatly reduce enzyme activity, but the most frequently found mutations do not.

Significance: Mutations in *CPA6* are associated with rare cases of epilepsy.

Carboxypeptidase A6 (*CPA6*) is a member of the A/B subfamily of M14 metalloproteases that is expressed in brain and many other tissues during development. Recently, two mutations in human *CPA6* were associated with febrile seizures and/or temporal lobe epilepsy. In this study we screened for additional *CPA6* mutations in patients with febrile seizures and focal epilepsy, which encompasses the temporal lobe epilepsy subtype. Mutations found from this analysis as well as *CPA6* mutations reported in databases of single nucleotide polymorphisms were further screened by analysis of the modeled pro*CPA6* protein structure and the functional role of the mutated amino acid. The point mutations predicted to affect activity and/or protein folding were tested by expression of the mutant in HEK293 cells and analysis of the resulting *CPA6* protein. Common polymorphisms in *CPA6* were also included in this analysis. Several mutations resulted in reduced enzyme activity or *CPA6* protein levels in the extracellular matrix. The mutants with reduced extracellular *CPA6* protein levels showed normal levels of 50-kDa pro*CPA6* in the cell, and this could be converted into 37-kDa *CPA6* by trypsin, suggesting that protein folding was not greatly affected by the mutations. Interestingly, three of the mutations that reduced extracellular *CPA6* protein levels were found in patients with epilepsy. Taken together, these results provide further evidence for the involvement of *CPA6* mutations in human epilepsy and reveal additional rare mutations that inactivate *CPA6* and could, therefore, also be associated with epileptic phenotypes.

Epilepsy is one of the most common neurological disorders, affecting 1–2% of the population (1) with focal epilepsy (FE)²

* This work was supported, in whole or in part, by National Institutes of Health Grant DA-004494 (to L. D. F.)

[§] This article contains supplemental Tables S1 and S2 and Figs. S1–S7.

¹ To whom correspondence should be addressed: Dept. of Molecular Pharmacology, Albert Einstein College of Medicine, 1300 Morris Park Ave., Bronx, NY 10461. Tel.: 718-430-4225; Fax: 718-430-8922; E-mail: lloyd.fricker@einstein.yu.edu.

² The abbreviations used are: FE, focal epilepsy; *CPA6*, carboxypeptidase A6; ECM, extracellular matrix; FA, 3-(2-furyl) acryloyl-Phe-Phe; FS, febrile seizure; SNP, single nucleotide polymorphism; TLE, temporal lobe epilepsy.

accounting for ~60% of this prevalence (2). As proposed by the International League Against Epilepsy, FE includes temporal, frontal, parietal, or occipital lobe epilepsy, depending on the localization of the seizure discharge (3). Temporal lobe epilepsy (TLE) is the most frequent form of FE in adults, with hippocampal sclerosis being the most recurrent pathogenic feature of this illness (4). Moreover, 25% of patients with TLE and hippocampal sclerosis often show febrile seizures (FS) during childhood (5). Generally, FS are defined as seizures occurring during fever episodes without any central nervous system infection (3). FS are the most common convulsive events during early childhood (5), being seen in 5% of the general population (6). As suggested by segregation and linkage studies, FS (7) and TLE (8) could be considered as complex disorders with genetic predisposition. However, few genes have been reproducibly associated with FS (9) and TLE (10). This may be due to the clinical heterogeneity of the patients included in almost all studies. FS and TLE may also be caused by multiple rare and/or *de novo* mutations. This latter possibility is supported by the recent identification of two mutations in the carboxypeptidase A6 (*CPA6*) gene in a family with FS and TLE as well as in unrelated TLE patients (11).

CPA6 is an extracellular peptidase in the A/B subfamily of the M14 family of carboxypeptidases (12). *CPA6* is translated with an N-terminal signal peptide (residues 1–30), which routes it to the secretory pathway. Like most other members of its subfamily, *CPA6* is produced as an ~50-kDa inactive proenzyme, pro*CPA6* (residues 31–437), that is activated when cleaved by a furin-like enzyme into the 37-kDa form (residues 130–437) (13, 14). The propeptide found in most members of the A/B carboxypeptidase subfamily acts as an intramolecular chaperone, inhibiting enzymatic activity (15, 16). Upon secretion, *CPA6* binds to the extracellular matrix (ECM), where it is enzymatically active and cleaves bulky aliphatic and aromatic residues from the C termini of peptides and proteins (17). *CPA6* mRNA is most highly expressed in the olfactory bulb and epididymis of the adult mouse and is more broadly expressed in the developing forebrain and cerebellum as well as in developing

TABLE 1

Demographic and clinical characteristics of FE patients, FS patients, and Caucasian controls

HS, hippocampal sclerosis; NA, number of patients for whom no information is available

Subjects	n	Male	Female	Age at	Age at	HS	Personal	Familial history	Simple	Complex	Personal history
				inclusion \pm S.D.	onset \pm S.D.		history of FS	of FS/epilepsy	FS	FS	of epilepsy
		%	%				%	%	%	%	
FE patients	195	96 (49.2)	99 (50.8)	32.83 \pm 11.19 NA 3	14.37 \pm 9.22 NA 13	89 (46.8) NA 5	69 (35.6) NA 1	32 (17.5) NA 12	NA 195	NA 195	
FS patients	97	51 (52.6)	46 (47.4)	9.47 \pm 13.21 NA 6	1.68 \pm 1.73 NA 63			33 (84.6) NA 58	30 (56.6) NA 44	23 (43.4) NA 44	16 (21.1) NA 21
Controls	242	171 (70.7)	71 (29.3)	44.8 \pm 12.8							

skin, bone, and osteoblasts (18). CPA6 mRNA has been identified in the human central nervous system, notably the hippocampus (11).

One of the previously identified CPA6 mutations results in a substitution at alanine 270 (A270V) and appears to be linked to a familial autosomal recessive form of FS and TLE, whereas the other, a missense mutation at glycine 267 (G267R), is associated to unrelated TLE patients in the heterozygous state (11). Biochemical analysis revealed that the A270V mutation retains full carboxypeptidase activity but is present in ECM at about 40% that of the wild type (WT) level in the ECM. The G267R mutation was not detectable in the ECM. Although CPA6 has been shown to perform biologically significant cleavages on several peptides *in vitro* (13, 17), the function of CPA6 as well as the mechanism by which these mutations may lead to epilepsy remains unknown. Because both mutations reduce the level of CPA6 protein in the ECM, it remains unclear whether carboxypeptidase activity or the level of the protein is responsible for the observed effects of these mutations.

Given what is known about the G267R and A270V mutations, we hypothesized that loss of function of CPA6 could lead to seizures and epilepsy. Because other mutations in the CPA6 gene should be capable of reducing or eliminating the function of the protein, we searched for and found additional CPA6 mutations in FE and FS populations. Modeling of these mutations as well as other mutations reported in databases of single nucleotide polymorphisms (SNPs) suggested that many mutations in the CPA6 gene could alter structural integrity, carboxypeptidase activity, or both and thus could potentially be related to the disease. We analyzed these mutant proteins using a variety of methods to better understand their relationship to seizures and epilepsy.

EXPERIMENTAL PROCEDURES

Subjects—This study was approved by the ethics committees from the departments of neurology and pediatrics from the University Hospitals of Geneva. The FE group consisted of patients who suffered from non-lesional FE and lesional FE such as vascular malformation, cortical dysplasia, and nervous system tumor (Table 1). These unrelated Caucasian patients showed the following distribution of epilepsy syndromes already described in a previous paper (11): 138 patients with TLE (70.8%), 28 patients with undetermined focal epilepsy (15.4%), 12 patients with frontotemporal epilepsy (6.2%), 4 patients with parietal epilepsy (2.1%), 2 patients with temporo-parietal junction epilepsy (1.0%), 2 patients with parietoocipi-

tal epilepsy (1.0%), 2 patients with temporolateral epilepsy (1.0%), 2 patients with temporooccipital epilepsy (1.0%), 1 patient with parietotemporal epilepsy (0.5%), 1 patient with multifocal epilepsy (0.5%), and 1 patient with Rasmussen syndrome (0.5%). These patients suffered from a severe form of epilepsy with poor control of their seizures. Diagnosis was based on patient history, clinical examination, interictal, and ictal electroencephalography (EEG) analysis carried out with monitoring video-EEG and magnetic resonance imaging evaluation.

Unrelated Caucasian patients with FS were admitted to the pediatric emergency rooms at the University Hospitals of Geneva, Lausanne, and Neuchâtel (Switzerland) (Table 1). At that time, these patients were recruited following the Consensus statement (19). Simple FS were defined when the seizures were brief with a single seizure occurring during a febrile illness. If duration exceeded 15 min or if multiple seizures occurred during a single febrile illness or within the first 24 h period, they were defined as complex FS.

A healthy Caucasian control group (West European origin for at least two generations) was recruited from blood donors at the University Hospitals of Geneva (Table 1). Only unrelated blood donors without a personal and/or familial history of epilepsy and seizures were included.

Mutation and Single Nucleotide Polymorphism Screening—All patients and controls or their legal representatives gave their informed consent, and then venous blood was collected in EDTA tubes. DNA was extracted from peripheral blood leukocytes using the Nucleon BACC 2 kit (GE Healthcare). CPA6 exons were explored using a high resolution melt assay, which is a rapid, highly reproducible, and resolute process. Previous experiments have shown a sensitivity and specificity between 97 and 99% for PCR products up to 1000 bp (20, 21). We used this method followed by direct sequencing to confirm mutations and SNPs as described (11). PCR and high resolution melt assay conditions and primers sequences are available on request. For high resolution melt data analysis, we normalized melting curves for each sample by calculating the “line of best fit” between two normalization regions before and after the major fluorescence decrease representing the melting of the PCR product using the software provided by the Rotor-GeneTM 6000 (Corbett Research, Australia). This algorithm allows the direct comparison of samples that have different starting fluorescence levels with control samples, which were previously sequenced to determine genotype (22).

Carboxypeptidase A6 Mutations

Statistical Analysis—First, we checked Hardy-Weinberg equilibrium by using a χ^2 goodness-of-fit test. Then differences in allele frequencies between cases and controls were determined using UNPHASED software (Version 3.0.10) (23). The statistical power to detect association was estimated using the Genetic Power Calculator. For every calculation we used a genotype relative risk for a heterozygote of 2, for a homozygote of 3, and an additive model at α level = 0.05. Therefore, we found that for rs17343819, FE patients had 94% power to detect a risk allele G with 13% frequency in controls and a disease prevalence of 0.5% (1, 24). The power was 84% for FS patients with a disease prevalence of 2% (6). For rs17853192, FE patients had 82% power to detect no significant association with a risk allele G with 7% frequency in controls. The power was 68% for FS patients.

Protein Analysis and Modeling—CPA6 protein sequences were accessed from The National Center for Biotechnology Information. Sequence alignment was performed using ClustalW and analyzed using GeneDoc to quantify homology between human CPA6 and orthologs in other vertebrates. Modeling of CPA6 protein structure has been described previously (17). Mutations were modeled using Swiss PDB Viewer and PyMOL. Images were created using PyMOL.

Site-directed Mutagenesis—Point mutations were introduced into the pcDNA3.1(–)hCPA6-HAH6 plasmid (13) using PfuUltra II Hotstart DNA Polymerase (Agilent, Santa Clara, CA) or Pfu Turbo (Stratagene, La Jolla, CA) using the QuikChange mutagenesis protocol (Stratagene). The resulting cDNA constructs code for full-length human preproCPA6 protein with the hemagglutinin (HA) and hexahistidine (H6) epitopes at the C terminus (13). All mutations were confirmed by DNA sequence analysis.

Cell Culture and Analysis of CPA6 Mutant Proteins—HEK293T cells were cultured in Dulbecco's modified Eagle's medium supplemented with 10% fetal bovine serum and penicillin/streptomycin at 37 °C and 5% CO₂. Cells were transfected with human CPA6 cDNA using TransFectin (Bio-Rad) according to the manufacturer's instructions. The amount of plasmid used in transfection was adjusted so that expression levels of proCPA6 in cell lysates were equal when examined by Western blot (data not shown). At 48 h post-transfection, cells were harvested in phosphate-buffered saline (PBS) and centrifuged, and the pellet was sonicated in PBS solution containing 0.1% Nonidet P-40. Aliquots were analyzed on a denaturing polyacrylamide gel. After harvesting cells from the culture dish, the ECM remaining on the dish was washed several times with PBS to remove any remaining cells. ECM samples were analyzed for CPA6 activity by incubating for 2 h at 37 °C with 0.5 mM 3-(2-furyl) acryloyl-Phe-Phe (FA-Phe-Phe; Bachem, Bubendorf, Switzerland) in Tris buffer (pH 7.8) containing 150 mM NaCl. Reactions were stopped by transferring the soluble substrate to polypropylene tubes, separating it from the enzyme, which remained bound to the ECM. Solutions were transferred to cuvettes, and carboxypeptidase activity was detected by a decrease in absorbance at 336 nm as described previously (13). After incubation with FA-Phe-Phe, the same ECM samples were collected by the addition of SDS-PAGE sample buffer, fractionated on SDS-PAGE, and transferred to nitrocellulose.

Western blotting was performed according to standard protocol with mouse anti-HA (clone HA-7, Sigma; 1:5000 dilution) primary antibody and either IRdye800-conjugated anti-mouse secondary antibody (Rockland, Gilbertsville, PA; 1:3000 dilution) or horseradish peroxidase-conjugated anti-mouse secondary antibody (Cell Signaling, Danvers, MA; 1:2000). Images were obtained and quantified using the Li-COR Odyssey System (Lincoln, NE) or with enhanced chemiluminescence detection reagent.

In experiments where levels of CPA6 protein expression were compared between media and ECM fractions, cells were cultured in reduced volumes of media beginning at 24 h post-transfection to ensure detection of CPA6 protein in the media fractions. As a positive control, 400 μ g/ml heparin was added at 24 h post-transfection to solubilize CPA6 to show that CPA6 could be detected in the media when present. At 48 h post-transfection, media samples were collected, and cells were harvested in PBS, combined with SDS-PAGE sample buffer, and analyzed by Western blotting, as above.

In experiments testing the thermal stability of CPA6, ECM samples were washed to remove cells and cell fragments as described above. Samples were incubated in PBS at the indicated temperature for 1 h. After heating, PBS was removed, and samples were assayed for carboxypeptidase activity as described above. In experiments to determine if oxidation reduces the enzymatic activity of CPA6, a similar method was employed. In place of heating for 1 h, ECM samples were incubated with 2% hydrogen peroxide (w/w) for 30 min at 22 °C on a rotating platform (25). Samples were then washed several times with PBS to remove remaining hydrogen peroxide before assaying for carboxypeptidase activity.

In experiments where trypsin was used to digest proCPA6, cells were lysed as described above and incubated with trypsin for 1 h at 37 °C at a broad range of concentrations. A 1:1 concentration of trypsin represents a stock solution of 0.2 μ g/ μ l of trypsin in trypsin resuspension buffer (Promega, Fitchburg, WI), from which other concentrations were prepared by serial dilution.

RESULTS

Mutations and SNPs Screening—To screen for additional mutations in CPA6, an exonic high resolution melt analysis was performed. A previously unpublished missense mutation c.843A→G was found in a heterozygote FE patient (ET 174). This mutation changes a histidine at position 196 into an arginine (H196R) and was not found in 242 Caucasian controls. The parents of the individual harboring this mutation were not available. One FS patient (CF-29-0) was found with a putative missense heterozygous mutation (Table 2). This non-synonymous mutation replaces a proline at position 16 with a threonine (P16T). The patient's unaffected father (CF-29-2) was heterozygous, whereas his affected mother (CF-29-1) did not harbor the mutation (Table 2). The mutation was not found in 242 Caucasian controls.

The screening of CPA6 exons also revealed a published genetic variant, rs35993949 (c.875C→G) found in the heterozygous condition in one FE patient (Table 2, subject ET 158). The SNP rs35993949 affects a glutamine residue, which is

TABLE 2
CPA6 mutations in FE and FS patients and their available relatives

NA, data not available.

Subject	CF-29-0	CF-29-1 (mother of CF-29-0)	CF-29-2 (father of CF-29-0)	ET 174	ET 158 ^a
Sex	Male	Female	Male	Male	Male
Age	2 years	36 years	36 years	20 years	49 years
Age at onset	15 months	4 years		7 years	5 years
Simple FS	Yes	Yes	No	Yes	No
Complex FS	No	No	No	No	No
Afebrile seizures	No	No	No	Left temporal origin	Temporal origin
Familial FS Antecedent	Mother	No	No	Brother	Paternal grandfather
IRM	NA	NA		Hippocampal sclerosis	Cavernous malformation
CPA6 mutation	c.302C → A (p.P16T) CA	c.302C → A (p.P16T) CC	c.302C → A (p.P16T) CA	c.843A → G (p.H196R) AG	c.875C → G, (p.Q207E) CG
Alleles	Heterozygous	Homozygous for Pro-16 allele	Heterozygous	Heterozygous	Heterozygous ^b

^a Data from this patient were published previously (11).^b This patient also harbors the G267R mutation.

replaced by a glutamic acid residue (Q207E). The putative pathogenic G-allele was not found in 242 Caucasian controls. In a previous study we reported a heterozygous missense mutation (G267R) for patient ET 158 (11). Further analysis shows that patient ET 158 is compound heterozygous for the two pathogenic non-synonymous mutations c.875C → G (Q207E) and c.799G → A (G267R) on exons 6 and 8, respectively. As parents of ET 158 were not available, we could not determine whether these two mutations are *de novo*.

Three common allelic variations were examined. First, we checked Hardy-Weinberg equilibrium to test if both allele and genotype frequencies in our patient and control populations remained constant through generations (26). The rs17853192 (C → G → S173C) and rs17343819 (A → G → N249S) genotypic distributions were at Hardy-Weinberg equilibrium among FE and FS patients and controls (Table 3). The rs10957393 (T → C → F45L) genotypic distributions were at Hardy-Weinberg equilibrium among FS patients ($p = 0.39$) but at the limit in FE patients ($p = 0.03$) and in controls ($p = 0.02$).

The SNP rs17343819 showed a significant difference between FE patients and controls (Table 3). The G-allele displayed a higher frequency in FE patients (21.0%) than in unaffected cases (13.7%). This difference reached statistical significance with a p value of 0.0041. Moreover, the AA genotype appears to be protective against FE because controls show a higher frequency (74.0%) than FE patients (61.0%) ($p = 0.013$). No significant genotypic and allelic differences were found between FS patients and controls. For rs17853192 and rs10957393, no associations were found for FE or FS patients compared with controls.

Modeling of CPA6 Mutations within Predicted Protein Structure—In addition to the mutations described above in FE and FS patients, a large number of additional mutations in the CPA6 gene were reported in the National Center for Biotechnology Information SNP database (www.ncbi.nlm.nih.gov) and in the 1000 Genomes Project. We first examined all SNP mutations and selected those that were exonic and that resulted in a missense mutation (supplemental Table S1). These mutations were examined for conservation from human to zebrafish (supplemental Fig. S1) and across members of the carboxypeptidase A subfamily (supplemental Fig. S2). SNPs were selected based on a variety of factors including frequency of the allele, conser-

TABLE 3
Allele and genotype count in FE, FS patients, and controlsBold indicates statistically significant p values.

Genetic variation	Controls		
	<i>n</i>	FE <i>n</i>	FS <i>n</i>
	%	%	%
rs10957393 (T → C → p.F45L)			
Allele	466	390	192
T	378 (81.1)	312 (80.0)	165 (85.9)
C	88 (18.9)	78 (20.0)	27 (14.1)
P		0.681	0.132
Genotype	233	195	96
TT	148 (63.5)	120 (61.5)	70 (72.9)
TC	82 (35.2)	72 (36.9)	25 (26.0)
CC	3 (1.3)	3 (1.5)	1 (1.1)
P		0.904	0.252
rs17853192 (C → G → p.S173C)			
Allele	476	390	192
C	439 (92.6)	368 (94.4)	178 (91.8)
G	35 (7.3)	22 (5.6)	16 (8.2)
P		0.302	0.705
Genotype	238	195	96
CC	202 (85.2)	173 (88.7)	81 (83.5)
CG	35 (14.8)	22 (11.2)	16 (16.5)
GG	0 (0.0)	0 (0.0)	0 (0.0)
P		0.284	0.692
rs17343819 (A → G → p.N249S)			
Allele	476	390	192
A	411 (86.3)	308 (79.0)	168 (87.5)
G	65 (13.7)	82 (21.0)	24 (12.5)
P		0.0041	0.689
Genotype	238	195	96
AA	176 (74.0)	119 (61.0)	72 (75.0)
AG	59 (24.8)	70 (35.9)	24 (25.0)
GG	3 (1.2)	6 (3.1)	0 (0.0)
P		0.013	0.360

vation, severity of the change in amino acid, location of the residue, and predictions from PolyPhen and SIFT, which analyze conservation of mutated residues across species and the predicted severity of amino acid substitutions (supplemental Table S1). Several nonsense mutations were found in the SNP database that produced truncations of the protein due to frame-shifts and/or introduction of stop codons. All of these were predicted to result in inactive CPA6 protein, based on the requirement for an active site Glu residue known to be involved with catalytic function (13). Missense mutations in CPA6 were visualized within the structural model of the CPA6 protein, as previously described (17) (supplemental Fig. S3). These mutations were organized into several categories based on biochemical properties, the frequency at which they occur in humans,

Carboxypeptidase A6 Mutations




FIGURE 1. Mutations identified in the CPA6 protein. This study examines the biochemical properties of twelve mutations in the CPA6 protein (sticks). These mutations occur throughout the protein, and the residues they alter likely have diverse effects on structure and function.

and the level of conservation of the amino acid that they alter. The F45L, S173C, and N249S mutations, which occur frequently in humans, were predicted to be harmless or minimally damaging to the protein. These mutations represent modest changes in that these amino acids are not highly conserved and the substituted amino acid was not predicted from our analysis to alter the conformation or function of the protein ([supplemental Table S1](#)). The H196R, Q207E, R311E, and A424V mutations were predicted to greatly diminish CPA6 activity. The H196R mutation occurs at a residue that is crucial for the coordination of the zinc ion in the catalytic region of the protein and is necessary for the function of all metallo-carboxypeptidases in the A/B subfamily ([supplemental Fig. S2](#)) (27). The V306I, P342A, and L413F mutations were predicted to be possibly damaging ([supplemental Table S1](#)). These substitutions occur at conserved amino acids but do not cause substitutions that represent severe changes in properties of the residue. Another mutation, P16T, occurs within the signal peptide and was analyzed using SignalP (28); however, this substitution was not predicted to result in a change in signal peptide cleavage ([supplemental Fig. S4](#)). Mutations that were selected for further analysis were visualized within the CPA6 protein structure (Fig. 1). Mutations were subjected to additional analysis to predict whether any would alter mRNA splicing using Human Splicing Finder (29). One SNP, rs17343819 (N249S), occurs within an exon splice enhancer motif in exon 7, a predicted recognition site for 9G8. This SNP results in a modest decrease in the value assigned by Human Splicing Finder for this motif ([supplemental Table 2](#)).

Biochemical Analysis of CPA6 Mutants—Twelve of the naturally occurring mutations were selected for direct testing of the biochemical properties. Two of these, G267R and A270V, were mutations we had previously tested for enzyme activity (11); these were further characterized in the present study. Although not identified in humans, the active site mutation E398Q, which has no detectable carboxypeptidase activity (13), was included as a negative control.

Mutants were expressed in HEK293 cells, and ECM samples were assayed for carboxypeptidase activity (Fig. 2). The F45L mutation was the only mutation that showed no statistically significant difference in activity from WT. The N249S mutation retained 89% of WT activity. Surprisingly, the S173C mutation showed only approximately half of the WT activity. The H196R and A424V mutations had no detectable activity, whereas the Q207E mutant had 11% of WT activity. We estimate that we would be able to detect activity comparable to 5% of WT activity by our assay, indicating that these mutations remove >95% of carboxypeptidase activity. The G267R and E398Q mutations also resulted in a similar near total reduction in carboxypeptidase activity, as previously reported (11). The remaining mutations resulted in an intermediate level of retained carboxypeptidase activity. When we examined expression levels of CPA6 protein in the ECM, we found that many of the mutants that showed a reduction in carboxypeptidase activity were expressed at reduced levels in the ECM. For the A270V and V306I mutants, reduced expression likely accounts for the observed reduction in activity. Although it cannot be ruled out, this is unlikely for the N249S mutant, which shows a measurable reduction in activity but not in expression levels in the ECM. Notably, some mutations such as the A424V had no detectable carboxypeptidase activity but an intermediate level of protein expression in the ECM. Conversely, the S173C and V306I mutations showed a vast reduction in ECM expression and an intermediate reduction in carboxypeptidase activity. No CPA6 protein was detectable in the media for WT or any of the mutant proteins (Fig. 2B), and when heparin was added to displace protein from the ECM into the media, mutant protein expression in the media was detected in approximately the same ratio as it was in the ECM (Fig. 2, B–E).

From the comparison of activity *versus* protein levels, it appeared that the S173C mutant had more activity than expected based on the amount of CPA6 protein, relative to WT CPA6 (Fig. 2, A and C). A potential explanation is that the WT CPA6 in the ECM was not fully active, whereas a greater proportion of the S173C mutant had full activity. One mechanism by which WT CPA6 activity could be reduced without a corresponding decrease in protein would be through oxidation of the methionine residue present in a key position within the substrate binding pocket. In all species examined, CPA6 contains a methionine in the position that confers specificity toward hydrophobic amino acids ([supplemental Figs. S1 and S2](#)); in other related enzymes (carboxypeptidases A1, A2, A3, A4, and A5) the comparable residue is a leucine, isoleucine, or valine and never a methionine. Methionine can become oxidized into the sulfoxide and sulfone, and this change is predicted to eliminate the ability of CPA6 to bind peptides with hydrophobic C termini. Therefore, we tested whether oxidation reduced CPA6 activity. Treatment of WT CPA6 with hydrogen peroxide lowered the amount of enzyme activity by about 60% (Fig. 3). S173C, which has an outward-facing cysteine residue, did not show a significant reduction in activity after exposure to the oxidizing chemical. As a control, the N249S mutation was also examined. This mutant showed a similar reduction in activity as WT CPA6 upon exposure to hydrogen peroxide. To test the possibility that CPA6 could




FIGURE 2. Activity and expression level of CPA6 mutants. WT CPA6 and CPA6 mutants were expressed in HEK293T cells and tested for activity and extracellular expression levels. Mock transfected cells (-) and cells transfected with an active site mutant, E398Q, were included as negative controls. All CPA6 constructs contained an HA tag on the C terminus for protein detection. *A*, the ECM-bound CPA6 was assayed for enzyme activity using the chromogenic carboxypeptidase substrate FA-Phe-Phe. *B* and *C*, the ECM samples were resolved by SDS-PAGE, and Western blots were performed using an antibody against the HA epitope to determine the relative level of CPA6 protein bound to the ECM. A number of mutants showed a reduction in the amount of protein bound to the ECM (quantified in *panel C*). *D* and *E*, media were analyzed on Western blots. In the absence of heparin, CPA6 protein was undetectable in media. With the addition of 400 μ g/ml heparin, which displaces CPA6 from the ECM, CPA6 was detectable in the media in approximately the same ratio as in the ECM (quantified in *panel E*). Error bars indicate S.E. ($n = 3-25$). Rare mutations found in patients with epilepsy are indicated in *bold*. ND, not detected; ns, not statistically significant.




FIGURE 3. Effects of hydrogen peroxide treatment on CPA6 activity. HEK293T cells were transfected with WT or S173C or N249S CPA6 constructs. ECM-bound CPA6 was incubated with or without 2% hydrogen peroxide for 30 min at 22 °C and assayed for activity using the chromogenic carboxypeptidase substrate FA-Phe-Phe. After treatment with hydrogen peroxide, WT activity was reduced by ~40% ($n = 4, p < 0.001$), whereas S173C showed only a modest reduction that was not statistically significant ($n = 5, p > 0.2$). The N249S mutation was reduced by ~30% ($n = 5, p < 0.01$).

have altered enzymatic activity when the methionine was converted into the sulfoxide or the sulfone, the hydrogen peroxide-treated CPA6 was tested with residues representing bulky hydrophobic residues, small hydrophobic residues, and basic residues. However, no change in substrate specificity was observed between treated and untreated CPA6 (supplemental Fig. S5).

The A270V mutation was analyzed for heat stability because of its connection to febrile seizures and to examine possible reasons for its reduced expression in the ECM. A change in heat stability indicates a difference in folding, which could serve as an explanation for the pathological properties of the A270V mutation. This assay has been used to characterize mutants of other carboxypeptidases (30). A modest reduction of carboxypeptidase activity was detected at 48 °C, and the A270V mutant showed a greater reduction than the WT form (Fig. 4A). At temperatures above 48 °C, both WT and A270V enzyme activity were greatly reduced and were not significantly different between WT and A270V. Other mutants with detectable activity were tested after a 48 °C heat treatment; however, none showed a significant difference when compared with WT CPA6 (Fig. 4B), indicating that the A270V may be unique in this property among the mutants tested.

Because proteins that are misfolded are often degraded before secretion, we tested if CPA6 mutants that have no detectable expression in the ECM are more sensitive to trypsin digestion, which could indicate a change in folding pattern and domain structure. All of the mutant proteins tested showed the same bands representing the 50-kDa proCPA6, the ~37-kDa active CPA6, and a ~25-kDa degradation product that likely corresponds to a cleavage at a region rich in basic amino acids (residues 178–187) (Fig. 5, supplemental Fig. S6). Grossly misfolded proteins often show smears due to the accessibility of trypsin to many basic residues, whereas correctly folded proteins will be preferentially cleaved by trypsin at a limited number of basic residues that are accessible to the enzyme. The similar sizes of the trypsin digestion products indicates that the mutant proteins form stable structures that resemble the WT form of CPA6.




FIGURE 4. Heat stability of CPA6 mutants. HEK293T cells were transfected with WT and mutant CPA6. ECM-bound CPA6 was incubated at the temperature indicated for 1 h and then transferred to 37 °C and assayed for CPA6 activity using the chromogenic carboxypeptidase substrate, FA-Phe-Phe. *A*, the A270V CPA6 mutant showed slightly decreased activity when incubated at 48 °C (WT, $n = 12$; A270V, $n = 6$; $p < 0.05$). Other temperatures showed a tendency toward reduced activity, but this was not significant. *B*, other CPA6 mutants with activity were tested by incubating at 48 °C for 1 h and then cooling to 37 °C for enzyme assay. No statistically significant differences were observed between these mutants and WT CPA6. Error bars represent S.E. ($n = 5$). RT, room temperature.




FIGURE 5. Digestion of proCPA6 with trypsin. HEK293T cells were transfected with WT or mutant CPA6. Cells were harvested in PBS, lysed in PBS with 0.1% Nonidet P-40, and sonicated. Samples were then centrifuged, and supernatants were divided into aliquots and incubated for 1 h at 37 °C with varying concentrations of trypsin as indicated. A 1:1 concentration of trypsin represents a stock solution of 0.2 $\mu\text{g}/\mu\text{l}$ trypsin, from which other concentrations were prepared by serial dilution. Both WT CPA6 and mutants, which displayed no detectable CPA6 expression in the ECM, show distinct bands corresponding to 50, 37, and 25 kDa, representing proCPA6, CPA6, and a cleavage product, respectively. The E398Q mutation was included as a control.

To test the possibility of a dominant negative action for CPA6 mutants found in the heterozygous condition, co-expression of WT and mutant CPA6 was performed. For this, WT CPA6 was tagged with the FLAG epitope tag, and mutants were tagged with the HA-His₆ tag. Combinations of the plasmid encoding WT CPA6 with plasmids encoding each of the mutants were transfected into HEK293T and the ECM assayed for enzyme activity (supplemental Fig. S7). The combination of WT and mutant either led to greater activity than WT alone (for those mutants with activity) or no difference with WT alone (for those mutants with no activity). None of the combinations showed activity lower than that of WT alone, indicating that the mutants were not able to act in a dominant negative fashion.

DISCUSSION

Previously, two CPA6 mutations were identified that affected enzymatic activity and protein levels; one mutation was linked to familial FS and TLE, and the other mutation was associated with sporadic TLE (11). A major finding of the present study is that a number of additional mutations within CPA6 similarly affect the protein and are found in patients with epilepsy. Because both of the previously identified mutations were located in the same substrate binding loop of the CPA6 protein, it was unclear if the mutations acted in a related and specific way or if other mutations affecting different parts of the protein could also be associated with epilepsy. The new mutations identified in the present study extend the previous results and suggest that loss-of-function mutations in CPA6 are related to epilepsy and seizures. Our finding that the A270V mutation and hydrogen peroxide treatment, each manipulations that affect the region of the protein involved in substrate binding, had no effect on the substrate specificity of the enzyme suggests that it is unlikely that other mutations that occur far from the substrate binding loop affect specificity of CPA6. However, because all of the other mutants were examined using only one substrate, it remains a possibility that some mutations in CPA6 alter the selectivity of the enzyme. Both gain-of-function (31, 32) and loss-of-function (33–35) mutations of other genes have been found to cause epilepsy. We expect that both types of mutations in CPA6 could lead to epilepsy; however, *in vitro* experiments present difficulties in discriminating between the two, as it is not always clear how these mutations will affect the human brain *in vivo* (36).

Generally, common variations in the CPA6 gene were not found to markedly reduce the activity or expression of the enzyme and were not found to be associated with epilepsy. However, our finding that the SNP rs17343819 (A→G → N249S) shows a small but statistically significant decrease in enzyme activity and is more common in FE patients than in

controls is consistent with the “common disease-common variant” hypothesis. Previous studies that identified susceptibility markers for complex epilepsy predicted considerable underlying polygenic heterogeneity (37). It is possible that the modest reduction in carboxypeptidase activity of the N249S variant contributes to seizures. The cause of the reduction in enzyme activity of this mutant is not known, and although ECM levels of mutant CPA6 protein were comparable to WT CPA6 protein, we cannot rule out a small decrease in expression of the N249S mutant. Alternatively, this mutation may cause changes in protein function or expression that were not detected in the present study. For example, Asn-249 is the last residue coded by exon 7 and the adenine to guanine substitution could affect splicing because it occurs within an exon splice enhancer, offering a potential alternative explanation for the consequences of this mutation. Although the reduction in the score assigned by Human Splicing Finder is modest (supplemental Table S2), this is consistent with a subtle effect of this SNP. Such results have already been found in some epileptic contexts. The well-studied SNP rs3812718 (IV5N+5 G→A), which modifies the proportion of two *SCN1A* alternative transcripts, is a susceptibility common variant for epilepsy. This SNP has been associated with FS and FE with history of FS (38, 39). Moreover, rare non-synonymous mutations in *SCN1A* have been linked to severe myoclonic epilepsy in infancy (40). Collectively, these data demonstrate that common variants and rare mutations of the same gene are implicated in related epileptic syndromes. This is consistent with the findings of the present study, as rare missense mutations and one common marker, rs17343819 (N249S), are detected in FE patients.

Several rare mutations were identified in patients with epilepsy that greatly reduce CPA6 activity and expression. The missense mutation (G267R) was previously found in three TLE patients (11). In the present study we found that one of these three patients also showed another missense mutation, Q207E. We could not confirm if these pathogenic variants were *de novo* because the parents of the individual harboring these mutations were not available. We were also unable to determine whether or not the two mutant alleles were located on the same parental chromosome. Because of this uncertainty, this patient is considered to be compound heterozygous for G267R and Q207E (11). Because each of these mutations alone causes a >95% reduction in CPA6 protein levels in the ECM, it is likely that a compound heterozygote for these mutations would show a complete loss of functional protein. Analogous events have been described in patients with missense mutations in *POLG*, the gene encoding mitochondrial DNA polymerase γ (41). Patients with these mutations display ataxia syndrome, which is a progressive neurological disorder characterized by FE as a primary feature (41). Some compound heterozygotes have been reported among these patients, who harbor both A467T and T748S mutations in *POLG* (42, 43). One patient, heterozygote for T748S only, also showed a similar neurological phenotype with epilepsy (42).

Except for the N249S polymorphism discussed above, all other CPA6 polymorphisms that occur frequently in the population did not show any association with epilepsy. Our finding that the SNP rs10957393 (T→C → F45L) is not significantly

associated with FE and FS groups is consistent with the observation that the F45L mutation showed normal levels of enzyme activity when expressed in cells. The S173C mutation is another relatively common variant in the general population. Although the S173C mutation significantly reduced CPA6 protein levels in the ECM, it had no apparent association with epilepsy. Notably, the S173C mutation has never been reported in the homozygous state, and it is possible that like the A270V mutation, which also causes a partial reduction in ECM expression of CPA6, the S173C mutation may be found to be associated with epilepsy only when both alleles are mutated (11).

Although most of the mutants tested in the present study showed a reduction in enzyme activity relative to the amount of protein present in the ECM, the S173C enzyme appears to be more active than WT CPA6 when enzyme activity is adjusted for protein level. It is possible that CPA6 exists in a variety of states, some more active than others. We hypothesized that CPA6 may be regulated by oxidation and that some percentage of CPA6 in the ECM may be oxidized at a given time. This prediction comes from the observation that CPA6 has a methionine residue at the position that confers substrate specificity; this methionine is conserved throughout evolution for all species where a clear CPA6 orthologue has been identified. Methionine is not present in the comparable position in any other member of the M14 metalloproteinase family. Because methionine is less hydrophobic when oxidized to either the sulfoxide or sulfone, this oxidation was predicted to reduce the ability of CPA6 to cleave peptides with hydrophobic C termini. This prediction was confirmed; CPA6 activity is diminished when treated with hydrogen peroxide, a standard technique to oxidize methionines in proteins. In contrast, the S173C mutant is resistant to hydrogen peroxide and does not lose enzyme activity, possibly because the free Cys residue is oxidized preferentially, protecting the critical methionine from oxidation. If the WT CPA6 in the ECM represents a mixture of protein with reduced and oxidized methionine residues, whereas the S173C is primarily the reduced form, this could account for the observation that the mutant has more activity than expected due to the low level of protein.

Several of the mutations in CPA6 were predicted to affect protein folding based on the steric hindrance imposed by the mutation or the loss of a salt bridge. Consistent with this prediction, we found a number of the mutations had no enzymatic activity and were not secreted from cells. Misfolded proteins are often degraded before being routed to the trans Golgi complex for secretion, potentially providing an explanation for the lack of secretion of some CPA6 mutants. However, all of the mutants presumably have the same general domain structures as they are all cleaved at the same basic regions by trypsin. Furthermore, except for the A270V mutation, none of the mutants showed greater thermo-instability, further supporting similar folding patterns to WT CPA6. Although the A270V mutant showed a significant decrease after being heated to 48 °C, indicating it may be less stable than the WT variant, this temperature is not physiological even during extreme fever. Similar results were found for a naturally occurring mutation of carboxypeptidase E, with greater sensitivity to elevated temper-

atures presumably reflecting decreased stability of the mutant (30).

The mechanism by which CPA6 affects predisposition to seizures is not known. CPA6 is an extracellular protease that cleaves C-terminal amino acids with a preference for large aromatic and aliphatic residues (17). When tested in *in vitro* assays, CPA6 cleaves neurotensin, Met-enkephalin, and Leu-enkephalin into inactive products but converts inactive angiotensin I into angiotensin II, the biologically active form. Many other peptides are cleaved by CPA6; the effect on biological activity is not currently known (13). The working hypothesis is that CPA6 plays a role in the activation of peptides that have neuroprotective effects or in the inactivation of peptides that cause neuronal stimulation and that the reduction of CPA6 activity leads to an imbalance of excitation and inhibition. Alternatively, it is possible that substrates or products of CPA6 play a role in development such that the reduction of CPA6 activity leads to a nervous system that is more prone to seizures. Previously, CPA6 deficiency was found in one patient with Duane Syndrome, an eye movement disorder caused by the failure of cranial nerve VI to innervate the lateral rectus muscle (44). Interestingly, CPA6 is expressed in the region posterior to the developing eye. However, reduction of *cpa6* levels in zebrafish embryos had no effect on eye tracking or cranial nerve pathfinding (45). Furthermore, there are no reports of eye movement disorders in the FS or FE patients found to have point mutations within CPA6. Thus, the link between CPA6 and Duane Syndrome is based on a single patient, in contrast to the link between CPA6 and epilepsy, which has now been extended to several mutations.

REFERENCES

- Hauser, W. A., Annegers, J. F., and Rocca, W. A. (1996) Descriptive epidemiology of epilepsy. Contributions of population-based studies from Rochester, Minnesota. *Mayo Clin. Proc.* **71**, 576–586
- Sander, J. W., Hart, Y. M., Johnson, A. L., and Shorvon, S. D. (1990) National General Practice Study of Epilepsy. Newly diagnosed epileptic seizures in a general population. *Lancet* **336**, 1267–1271
- Commission on Classification and Terminology of the International League Against Epilepsy. (1989) Proposal for revised classification of epilepsies and epileptic syndromes. *Epilepsia* **30**, 389–399
- Falconer, M. A., Serafetinides, E. A., and Corsellis, J. A. (1964) Etiology and Pathogenesis of Temporal Lobe Epilepsy. *Arch Neurol.* **10**, 233–248
- Baulac, S., Gourfinkel-An, I., Nabbout, R., Huberfeld, G., Serratosa, J., Leguern, E., and Baulac, M. (2004) Fever, genes, and epilepsy. *Lancet Neurol.* **3**, 421–430
- Hauser, W. A. (1994) The prevalence and incidence of convulsive disorders in children. *Epilepsia* **35**, S1–S6
- Rich, S. S., Annegers, J. F., Hauser, W. A., and Anderson, V. E. (1987) Complex segregation analysis of febrile convulsions. *Am. J. Hum. Genet.* **41**, 249–257
- Ottman, R. (1989) Genetics of the partial epilepsies. A review. *Epilepsia* **30**, 107–111
- Nakayama, J. (2009) Progress in searching for the febrile seizure susceptibility genes. *Brain Dev.* **31**, 359–365
- Salzmann, A., Malafosse, A. (2012) Genetics of temporal lobe epilepsy. A review. *Epilepsy Res. Treat.* 863702
- Salzmann, A., Guipponi, M., Lyons, P. J., Fricker, L. D., Sapio, M., Lamberty, C., Buresi, C., Ouled Amar Bencheikh, B., Lahjouji, F., Ouazzani, R., Crespel, A., Chaigne, D., and Malafosse, A. (2012) Carboxypeptidase A6 gene (CPA6) mutations in a recessive familial form of febrile seizures and temporal lobe epilepsy and in sporadic temporal lobe epilepsy. *Hum. Mutat.* **33**, 124–135
- Wei, S., Segura, S., Vendrell, J., Aviles, F. X., Lanoue, E., Day, R., Feng, Y., and Fricker, L. D. (2002) Identification and characterization of three members of the human metallo-carboxypeptidase gene family. *J. Biol. Chem.* **277**, 14954–14964
- Lyons, P. J., Callaway, M. B., and Fricker, L. D. (2008) Characterization of carboxypeptidase A6, an extracellular matrix peptidase. *J. Biol. Chem.* **283**, 7054–7063
- Vendrell, J., Querol, E., and Avilés, F. X. (2000) Metallo-carboxypeptidases and their protein inhibitors. Structure, function, and biomedical properties. *Biochim. Biophys. Acta* **1477**, 284–298
- Phillips, M. A., and Rutter, W. J. (1996) Role of the prodomain in folding and secretion of rat pancreatic carboxypeptidase A1. *Biochemistry* **35**, 6771–6776
- Gomis-Rüth, F. X. (2008) Structure and mechanism of metallo-carboxypeptidases. *Crit. Rev. Biochem. Mol. Biol.* **43**, 319–345
- Lyons, P. J., and Fricker, L. D. (2010) Substrate specificity of human carboxypeptidase A6. *J. Biol. Chem.* **285**, 38234–38242
- Fontenele-Neto, J. D., Kalinina, E., Feng, Y., and Fricker, L. D. (2005) Identification and distribution of mouse carboxypeptidase A-6. *Brain Res. Mol. Brain Res.* **137**, 132–142
- Consensus statement. (1980) Febrile seizures. Long-term management of children with fever-associated seizures. *Pediatrics* **66**, 1009–1012
- Montgomery, J., Wittwer, C. T., Palais, R., and Zhou, L. (2007) Simultaneous mutation scanning and genotyping by high resolution DNA melting analysis. *Nat. Protoc.* **2**, 59–66
- Taylor, C. F. (2009) Mutation scanning using high resolution melting. *Biochem. Soc. Trans.* **37**, 433–437
- Wojdacz, T. K., Dobrovic, A., and Hansen, L. L. (2008) Methylation-sensitive high resolution melting. *Nat. Protoc.* **3**, 1903–1908
- Dudbridge, F., and Koeleman, B. P. (2003) Rank truncated product of P-values, with application to genomewide association scans. *Genet. Epidemiol.* **25**, 360–366
- Wallace, H., Shorvon, S., and Tallis, R. (1998) Age-specific incidence and prevalence rates of treated epilepsy in an unselected population of 2,052,922 and age-specific fertility rates of women with epilepsy. *Lancet* **352**, 1970–1973
- Kim, Y. H., Berry, A. H., Spencer, D. S., and Stites, W. E. (2001) Comparing the effect on protein stability of methionine oxidation versus mutagenesis. Steps toward engineering oxidative resistance in proteins. *Protein Eng.* **14**, 343–347
- Stern, C. (1943) The Hardy-Weinberg Law. *Science* **97**, 137–138
- Gomis-Rüth, F. X., Botelho, T. O., and Bode, W. (2012) A standard orientation for metallo-peptidases. *Biochim. Biophys. Acta* **1824**, 157–163
- Petersen, T. N., Brunak, S., von Heijne, G., and Nielsen, H. (2011) SignalP 4.0. Discriminating signal peptides from transmembrane regions. *Nat. Methods* **8**, 785–786
- Desmet, F. O., Hamroun, D., Lalande, M., Collod-Bérout, G., Claustres, M., and Bérout, C. (2009) Human Splicing Finder. An online bioinformatics tool to predict splicing signals. *Nucleic Acids Res.* **37**, e67
- Chen, H., Jawahar, S., Qian, Y., Duong, Q., Chan, G., Parker, A., Meyer, J. M., Moore, K. J., Chayen, S., Gross, D. J., Glaser, B., Permutt, M. A., and Fricker, L. D. (2001) Missense polymorphism in the human carboxypeptidase E gene alters enzymatic activity. *Hum. Mutat.* **18**, 120–131
- Dibbens, L. M., Reid, C. A., Hodgson, B., Thomas, E. A., Phillips, A. M., Gazina, E., Cromer, B. A., Clarke, A. L., Baram, T. Z., Scheffer, I. E., Berkovic, S. F., and Petrou, S. (2010) Augmented currents of an HCN2 variant in patients with febrile seizure syndromes. *Ann. Neurol.* **67**, 542–546
- Volkers, L., Kahlig, K. M., Verbeek, N. E., Das, J. H., van Kempen, M. J., Stroink, H., Augustijn, P., van Nieuwenhuizen, O., Lindhout, D., George, A. L., Jr., Koeleman, B. P., and Rook, M. B. (2011) Nav 1.1 dysfunction in genetic epilepsy with febrile seizures-plus or Dravet syndrome. *Eur. J. Neurosci.* **34**, 1268–1275
- Escayg, A., and Goldin, A. L. (2010) Sodium channel SCN1A and epilepsy. Mutations and mechanisms. *Epilepsia* **51**, 1650–1658
- Kang, J. Q., Shen, W., and Macdonald, R. L. (2009) The GABRG2 mutation, Q351X, associated with generalized epilepsy with febrile seizures plus, has both loss of function and dominant-negative suppression. *J. Neurosci.* **29**, 2845–2856

35. Liao, W. P., Shi, Y. W., Long, Y. S., Zeng, Y., Li, T., Yu, M. J., Su, T., Deng, P., Lei, Z. G., Xu, S. J., Deng, W. Y., Liu, X. R., Sun, W. W., Yi, Y. H., Xu, Z. C., and Duan, S. (2010) Partial epilepsy with antecedent febrile seizures and seizure aggravation by antiepileptic drugs. Associated with loss of function of Na(v) 1.1. *Epilepsia* **51**, 1669–1678
36. Moulard, B., Picard, F., le Hellard, S., Agulhon, C., Weiland, S., Favre, I., Bertrand, S., Malafosse, A., and Bertrand, D. (2001) Ion channel variation causes epilepsies. *Brain Res. Brain Res. Rev.* **36**, 275–284
37. Mulley, J. C., Scheffer, I. E., Harkin, L. A., Berkovic, S. F., and Dibbens, L. M. (2005) Susceptibility genes for complex epilepsy. *Hum. Mol. Genet.* **14** Spec No. 2, R243–R249
38. Schlachter, K., Gruber-Sedlmayr, U., Stogmann, E., Lausecker, M., Hotzy, C., Balzar, J., Schuh, E., Baumgartner, C., Mueller, J. C., Illig, T., Wichmann, H. E., Lichtner, P., Meitinger, T., Strom, T. M., Zimprich, A., and Zimprich, F. (2009) A splice site variant in the sodium channel gene SCN1A confers risk of febrile seizures. *Neurology* **72**, 974–978
39. Le Gal, F., Salzmann, A., Crespel, A., and Malafosse, A. (2011) Replication of association between a SCN1A splice variant and febrile seizures. *Epilepsia* **52**, e135–138
40. Lossin, C. (2009) A catalog of SCN1A variants. *Brain Dev.* **31**, 114–130
41. Winterthun, S., Ferrari, G., He, L., Taylor, R. W., Zeviani, M., Turnbull, D. M., Engelsen, B. A., Moen, G., and Bindoff, L. A. (2005) Autosomal recessive mitochondrial ataxic syndrome due to mitochondrial polymerase γ mutations. *Neurology* **64**, 1204–1208
42. Tzoulis, C., Engelsen, B. A., Telstad, W., Aasly, J., Zeviani, M., Winterthun, S., Ferrari, G., Aarseth, J. H., and Bindoff, L. A. (2006) The spectrum of clinical disease caused by the A467T and W748S POLG mutations. A study of 26 cases. *Brain* **129**, 1685–1692
43. Roshal, D., Glosner, D., and Zangaladze, A. (2011) Parieto-occipital lobe epilepsy caused by a POLG1 compound heterozygous A467T/W748S genotype. *Epilepsy Behav.* **21**, 206–210
44. Pizzuti, A., Calabrese, G., Bozzali, M., Telvi, L., Morizio, E., Guida, V., Gatta, V., Stuppia, L., Ion, A., Palka, G., and Dallapiccola, B. (2002) A peptidase gene in chromosome 8q is disrupted by a balanced translocation in a duane syndrome patient. *Invest. Ophthalmol. Vis. Sci.* **43**, 3609–3612
45. Lyons, P. J., Ma, L. H., Baker, R., and Fricker, L. D. (2010) Carboxypeptidase A6 in zebrafish development and implications for V1th cranial nerve pathfinding. *PLoS One* **5**, e12967