

A8

doi: 10.14232/ syrpharmacognosy.2021.a8


Isolation and synthesis of 6-gingerol and 6-gingerdione derivatives

Sara Hassan Hassan Ahmed

Email: sara.hassan@pharm.u-szeged.hu

Ginger (*Zingiber officinale* Roscoe) has been used as a spice and a traditional remedy since ancient times, especially in Chinese herbal medicine [1,2]. A wide array of bioactivities were reported regarding ginger extract, e.g. antioxidant, anti-inflammatory, antiemetic, anticancer activity [3]. Most of the bioactivities had been correlated to gingerols and shogaols, the most abundant secondary metabolites in ginger [4].

In our work, we aimed to investigate the chemical space around 6-gingerol by introducing different changes in the skeleton and investigating the bioactivity of the resulting derivatives. The current work also focuses on the total synthesis of 6-gingerdione, another important compound in ginger root, and its heterocyclic analogues. So far, six molecules have been synthesized and characterized; the final purification and structure elucidation of further derivatives are currently in progress.



View metadata and citation for this article at www.researchprotocols.org/2021/1/e28282.html

provided by University of Szeged
powered by COBE

Acknowledgements:

This work was supported by the National Research, Development and Innovation Office, Hungary (NKFIH; K-134704), and the Ministry of Human Capacities, Hungary grant 20391-3/2018/FEKUSTRAT.

References

- [1] Leoni, A., et al. *Natural Product Research* 2018; 32(18): 2139–2146.
- [2] Sana, N., et al. *Anti-Cancer Agents in Medicinal Chemistry* 2021; 21(4): 428–432.
- [3] Mao, Q.Q., et al. *Foods* 2019; 8(6).
- [4] de Lima, R.M.T., et al. *Phytother Res* 2018; 32(10): 1885–1907.